# Combinatorial Problems and Exercises by László Lovász

Akadémiai Kiadó Budapest

### COMBINATORIAL PROBLEMS AND EXERCISES

by LÁSZLÓ LOVÁSZ


Combinatorics is one of the fastest-growing fields of mathematics. This is reflected by its application in many fields (like computer science and operations research) but also by the fact that the methods of combinatorics are becoming more complex, and several other mathematical disciplines are applied in it. Thus, for the solution of combinatorial problems, no matter how elementarily they are formulated, ingenuity in itself will no longer suffice: a thorough knowledge of the technical machinery of combinatorics is needed.

The aim of the book is to introduce part of the methods of combinatorics for those who want to apply these methods in the solution of practical and theoretical problems. This is achieved by showing various tricks and techniques through exercises and exercise series. The solutions of these series lead to some of the most important results of combinatorics. In a separate part, a hint is given to the solution of each problem, and a third part contains all solutions in detail. There is also a dictionary containing the definitions of the combinatorial notions occurring in the book.


Two of the main chapters of combinatorics are treated in greater detail: enumeration (combinatorics in the classical sense) and graph theory. But many other important fields are also touched on in connection with these, fields such as permutation groups and hypergraphs.







#### Combinatorial Problems and Exercises



## Combinatorial Problems and Exercises

by

#### L. Lovász

Professor of Mathematics József Attila University, Szeged, Hungary



Akadémiai Kiadó • Budapest 1979

Joint edition published by
Akadémiai Kiadó
Budapest
and
North-Holland
Publishing Company
Amsterdam · New York · Oxford

ISBN 963 05 1469 9

© Akadémiai Kiadó • Budapest 1979

Printed in Hungary



#### Contents

Pr	efa	ce			8
			Problems	Hints	Solutions
8	1.	Basic enumeration (partitions of sets and numbers, recurrence relations and generating functions, combinatorial identities)	13	93	141
·		The sieve (inclusion-exclusion, Selberg sieve, second moment method, Moebius function)	19	97	175
8	3.	Permutations (cycle index polynomial, Hall-Rényi coding, Pólya-Redfield method)	25	99	196
8	4.	Two classical enumeration problems in graph theory (labelled and unlabelled trees, spanning trees of a graph; 1-factors, the Ising problem, restricted permutations)	29	101	216
8	5.	Parity and duality (Eulerian graphs, planar duality, Sperner's Lemma, the linear space of cuts and cycles, planarity criteria)	34	105	248
8	6.	Connectivity (trees, ear-structures, Menger's Theorem, a calculus of cutsets, flow theory)	39	108	267
§	7.	Factors of graphs (König's Theorem, Tutte's Theorem, the structure of factors of graphs, realizability of			
8	8.	Independent sets of points (kernel, games on graphs,	48	113	303
§	9.	α-critical graphs)	55	116	338
8	10.	matic polynomial, planar graphs)	58	118	349
		joint circuits; existence of subdivisions, paths and Hamiltonian circuits; Turán's Theorem)	64	122	376
		Spectra of graphs (relations to the structure and automorphism group, strongly regular graphs)	70	125	405
§	12.	Automorphisms of graphs (Frucht's Theorem, transitive groups, topological problems, endomorphisms)	74	127	421

The Control of	Problems	Hints	Solutions
§ 13. Hypergraphs (circuits, transversal theory, intersection properties and extremal problems, 2-colorability, integral and fractional packing and covering, normal hypergraphs and perfect graphs)	76	129	439
§ 14. Ramsey Theory (Ramsey's Theorem, monochromatic paths and circuits in graphs, geometrical configurations, Van der Waerden's Theorem, monotony and convexity)	84	134	478
§ 15. Reconstruction (line-graphs, the Reconstruction Con-	89	137	506
Dictionary of the combinatorial phrases and concepts used .			528
Notation			541
Index of the abbreviations of textbooks and monographs			543
Subject index			545
Author index	aut chtean		549

#### Preface

Having vegetated on the fringes of mathematical science for centuries, combinatorics has now burgeoned into one of the fastest growing branches of mathematics — undoubtedly so if we consider the number of publications in this field, its applications in other branches of mathematics and in other sciences, and also, the interest of scientists, economists and engineers in combinatorial structures. The mathematical world had been attracted by the successes of algebra and analysis and only in recent years has it become clear, due largely to problems arising from economics, statistics, electrical engineering and other applied sciences, that combinatorics, the study of finite sets and finite structures, has its own problems and principles. These are independent of those in algebra and analysis but match them in difficulty, practical and theoretical interest and beauty.

Yet the opinion of many first-class mathematicians about combinatorics is still in the pejorative. While accepting its interest and difficulty, they deny its depth. It is often forcefully stated that combinatorics is a collection of problems, which may be interesting in themselves but are not linked and do not constitute a theory. It is easy to obtain new results in combinatorics or graph theory because there are few techniques to learn, and this results in a fast-growing

number of publications.

The above accusations are clearly characteristic of any field of science at an early stage of its development — at the stage of collecting data. As long as the main questions have not been formulated and the abstractions to a general level have not been carried through, there is no way to distinguish between interesting and less interesting results — except on an aesthetic basis, which is, of course, too subjective. Those techniques whose absence has been disapproved of above await their discoverers. So underdevelopment is not a case against, but rather for, directing young scientists toward a given field.

In my opinion, combinatorics is now growing out of this early stage. There are techniques to learn: enumeration techniques, matroid theory, the probabilistic method, linear programming, block design constructions, etc. There are branches which consist of theorems forming a hierarchy and which contain central structure theorems forming the backbone of study: connectivity of graphs (network flows) or factors of graphs, just to pick two examples from graph theory. There are notions abstracted from many non-trivial results, which unify large parts of the theory, such as matroids or the concept of good characterization (see below). My feeling is that it is no longer possible to obtain significant results

10 PREFACE

without the knowledge of these facts, concepts and techniques. (Of course, exceptions may occur, since the field is destined to cover such a large part of the world of mathematics that entirely new problems may still arise.)

The reader will forgive I hope the insertion of some general ideas which tend to play the role of systematizing and unifying concepts. The first of these is the notion of a good characterization, formulated by J. Edmonds. A property P of graphs is well-characterized if (making use of the different formulations of the definition and the equivalent condition) we are able to efficiently prove P if it holds and to efficiently disprove it if it does not. For example Kuratowski's classical characterization of planar graphs provides a good characterization of planarity: if a graph is planar, we easily establish this by drawing it in the plane; if it is non-planar, we can show this by exhibiting one of Kuratowski's graphs in it (see problem 5.37). A good characterization reflects a deep underlying logical duality of the property and, as the reader may convice himself by comparing good and "non-good" characterizations occurring throughout this book, often amounts to "the" solution of the problem. Of course, this does not mean that "non-good" characterizations may not be deep and useful theorems.

The existence of good characterizations tends to go hand in hand with the existence of good decision algorithms (by a "good" or "efficient" algorithm we mean one whose running time in the worst case is only a polynomial in the input data; this again does not directly affect its practical value). Several combinatorial properties are known for which there exists a "good" algorithm to decide whether a given structure has the property, but its existence is by no means obvious (e.g. having a 1-factor). An interesting theoretical result, due to S. A. Cook, R. M. Karp and L. A. Levin is the following. Several properties of graphs (for example, the existence of Hamiltonian circuits, independence number, chromatic number, the existence of a kernel etc.) are equivalent in the sense that if any one of them could be solved by a "good" algorithm, all of them could, and in this case the "good" algorithmic solvability of very many other problems in different fields of mathematics would follow (just to mention a very distant one: testing an n-digit number for primarility). It is unlikely that all of these could be solved efficiently, but there is as yet no proof of this.

Another idea which has proved very fruitful is that combinatorial optimization problems can generally be formulated as linear programming problems with integrality constraints. If one could disregard the constraints, the Duality Theorem of linear programming would provide the solution. So the solution of these problems is connected with investigating the effect of the integrality constraints on the behavior of optima. For example, if we can show that they do not change the optimum, we obtain a minimax theorem. Several instances of this idea can be found in § 13 (Hypergraphs).

<sup>&</sup>lt;sup>†</sup> For a detailed description see e.g. V. Chvátal, *Discrete Math.* 4 (1973) 305–337, or A.V. Aho, J.E. Hopcroft, J.D. Ullman, *The Design and Analysis of Computer Algorithms*, Addison-Wesley, 1974, Chap. 10.

<sup>&</sup>lt;sup>‡</sup> See e.g. R.M. Karp, in: Complexity of Computer Computations (R.E. Miller, J.W. Thatcher, ed.) Plenum Press, 1972, 85–104 or A.V. Aho et al., ibid.

PREFACE 11

Last but not least we mention the use of *linear algebra*. This ranges from applications of matrix calculus to the introduction of homology and cohomology groups. A common background of many applications of linear algebra is *Matroid Theory*, which is now a flourishing branch of combinatorics itself.

\*

The main purpose of this book is to provide help in learning existing techniques in combinatorics. The most effective (but admittedly very time-consuming) way of learning such techniques is to solve (appropriately chosen) exercises

and problems.

This book presents all the material in the form of problems and series of problems (apart from some general comments at the beginning of each chapter). We hope that it will be useful to those students who intend to start research in graph theory, combinatorics or their applications, and for those who feel that combinatorial techniques might help them with their work in other branches of mathematics, management science, electrical engineering and so on. For background, only the elements of linear algebra, group theory, probability and calculus are needed.

When selecting the material I have had to restrict the topics covered. I feel that a more detailed analysis of a few basic notions is more useful than touching of all possible fields of research. So in this volume only enumeration problems, graphs and set-systems are discussed. Some fields have had to be completely omitted: random structures (here the reader is advised to read the book *Probabilistic Methods in Combinatorics* by P. Erdős and J. Spencer, Akadémiai Kiadó, Budapest and Academic Press, New York-London, 1974), integer programming, matroids (combinatorial geometries), finite geometries, block designs, lattice geometry etc. I hope eventually to write a sequel to this volume covering some of these latter topics.

The book consists of three major parts: Problems, Hints and Solutions. A reader with less experience may read the hint given to a problem before trying to solve it; those problems with one or two asterisks are thought to be difficult and the reader may read the hint given right away unless he is prepared to sacrifice several days to the problem (some of them are worth it, I venture to say). Even having solved a problem the reader is advised to compare his solution with the one given here: it may be that the idea occurring in our solution will be basic in a later series of problems. Here it should be pointed out that problems come in series and previous problems often serve as steps to the last, deepest result in the series. Also note that the solution often uses notation or properties introduced in the hint.

As references I have preferred to give those where further development of the subject can best be seen. Thus, for those results reproduced in textbooks or monographs, usually the latter are given as reference. No reference means either that the assertion of the exercise is so well known that it would have been impossible or superfluous to trace it back or that the problem is believed to be new. A list of those textbooks and monographs most often cited is given at the end of the volume. A dictionary containing the definitions of combinatorial concepts used in the book and a list of symbols as well as an Author Index

and a Subject Index are included.

12 PREFACE

Acknowledgements. First of all, I wish to express my gratitude to Professors P. Erdős and T. Gallai, through whom I have come to love combinatorics, and from whom in principle I have learnt almost everything included in this book. I have to mention in the same group Professors Vera T. Sós, A. Hajnal and many other graph theorists and combinatorists, who (directly or indirectly) contributed to this book. I am particularly indebted to J.C. Ault, L. Babai, A. Bondy, A. Hajnal, G. Katona, M.D. Plummer and M. Simonovits, who having read the manuscript or parts of it, made many valuable remarks, suggestions and corrections; and to L. Babai and his students, in particular E. Boros and Z. Füredi for their invaluable help in proofreading.

I want to express my thanks to Prof. I. Rábai for the idea of such a problem book and for encouraging its writing; also to Mrs. A. Porubszky for the careful typing of the manuscript; and to Akadémiai Kiadó, in particular Mrs. Á.

Sulvok for the competent and thorough editorial work.

But above all I have to thank my wife Kati; her unbroken encouragement and professional, technical and moral support have been the firm basis of my work.

#### I. Problems

#### § 1. Basic enumeration

There is no rule which says that enumeration problems, even the simplest ones, must have solutions expressible as closed formulas. Some have, of course, and one important thing to be learnt here is how to recognize such problems. Another approach, avoiding the difficulty of trying to produce a closed formula, is to look for "substitute" solutions in other forms such as formulas involving summations, recurrence relations or generating functions. A typical (but not unique or universal) technique for solving an enumeration problem, depending on one or more parameters, is to find a recurrence relation, deduce a formula for the generating function (the recurrence relation is usually equivalent to a differential equation involving this function) and finally, where possible, to obtain the coefficients of the Taylor expansion of the generating function.

However, it should be pointed out that, in many cases, elementary transformations of the problem may lead to another problem already solved. For example it may be possible by such transformations to represent each element to be counted as the result of n consecutive decisions such that there are  $a_i$  possible choices at the i<sup>th</sup> step. The answer would then be  $a_1a_2 \ldots a_n$ . This is particularly useful when each decision is independent of all the previous decisions. Finding such a situation equivalent to the given problem is usually

difficult and a matter of luck combined with experience.

- 1. In a shop there are k kinds of postcards. We want to send postcards to n friends. How many different ways can this be done? What happens if we want to send them different cards? What happens if we want to send two different cards to each of them (but different persons may get the same card)?
- 2. We have k distinct postcards and want to send them all to our n friends (a friend can get any number of postcards, including 0). How many ways can this be done? What happens if we want to send at least one card to each friend?
- 3. How many anagrams can be formed from the word CHARACTERIZATION? (An anagram is a word having the same letters, each occurring the same number of times; this second word does not need to have a meaning.)
- **4.** (a) How many possibilities are there to distribute k forints<sup>†</sup> among n people so that each receives at least one?

<sup>†</sup> forint: Hungarian currency.

(b) Suppose we do not insist that each person receives something. What will be the number of distributions in this case?

1.5

- 5. There are k kinds of postcards, but only in a limited number of each, there being  $a_i$  copies of the  $i^{th}$  one. What is the number of possible ways of sending them to n friends? (We may send more than one copy of the same postcard to the same person).
- **6.** Find a recurrence relation for the second-order Stirling numbers S(n, k) and tabulate them for  $n \leq 6$ .
- 7. Prove that

$$\sum_{k=1}^{n} S(n, k) x(x-1) \dots (x-k+1) = x^{n}.$$

8. Prove that

$$S(n, k) = \frac{1}{k!} \sum_{j=1}^{k} (-1)^{k-j} {k \choose j} j^n.$$

In particular, the right-hand side is 0 for k > n (cf. also 2.4).

**9.** (a) Let  $P_n$  denote the  $n^{\text{th}}$  Bell number, i.e. the number of all partitions of n objects. Prove the formula

$$P_n = \frac{1}{e} \sum_{k=0}^{\infty} \frac{k^n}{k!}.$$

(b)\* 
$$P_n \sim \frac{1}{\sqrt{n}} \lambda(n)^{n+\frac{1}{2}} e^{\lambda(n)-n-1}$$

where  $\lambda(n)$  is defined by  $\lambda(n) \log \lambda(n) = n$ .

- 10. Find a recurrence relation for  $P_n$ .
- 11. Let p(x) be the exponential generating function of the sequence  $P_n$ , i.e.

$$p(x) = \sum_{n=0}^{\infty} \frac{P_n}{n!} x^n.$$

Determine p(x).

12. (a) Prove that

$$P_n = \sum_{\substack{k_1, \dots, k_n \ge 0 \\ k_1 + 2k_2 + \dots + nk_n = n}} \frac{n!}{k_1! (1!)^{k_1} k_2! (2!)^{k_2} \dots k_n! (n!)^{k_n}}.$$

Deduce the formula for p(x) from this expression.

(b) What do we get (asymptotically) if the summation is extended over all systems  $(k_1, \ldots, k_n)$  with  $k_1 + k_2 + \ldots + k_n = n$ ?

13. Give another proof of

$$P_n = \frac{1}{e} \sum_{k=0}^{\infty} \frac{k^n}{k!} \,.$$

14. (a) Let  $Q_n$  denote the number of partitions of an *n*-element set into an even number of classes  $(Q_0 = 1)$ . Determine

$$q(x) = \sum_{n=0}^{\infty} \frac{Q_n}{n!} x^n,$$

and find an analogue of the formula of 1.9a.

(b) Denote by  $R_n$  the number of partitions of an *n*-set into classes of even cardinality  $(R_0 = 1)$ . Determine

$$r(x) = \sum_{n=0}^{\infty} \frac{R_n}{n!} x^n.$$

15. Let  $S_n$  denote the number of mappings  $f: \{1, \ldots, n\} \to \{1, \ldots, n\}$  such that if f takes a value i then it also takes each value j,  $1 \le j \le i$ . Let  $S_0 = 1$ .

(a) Prove that

$$S_n = \sum_{k=0}^{\infty} \frac{k^n}{2^{k+1}}$$

and determine the generating function

$$s(x) = \sum_{n=0}^{\infty} \frac{S_n}{n!} x^n.$$

(b) Prove the asymptotics

$$\frac{S_n(\log 2)^{n+1}}{n!} \to 1 \quad (n \to \infty).$$

\*

16. The number of partitions of the number n into (a sum of) no more than r terms is equal to the number of partitions of n into any number of terms, each at most r.

17. The number of partitions of a number n into exactly m terms is equal to the number of partitions of n-m into no more than m terms. Find a similar identity involving the number of partitions of n into exactly m distinct terms.

18\*. The number of partitions of n into (any number of) distinct terms is equal to the number of partitions of n into odd terms.

19\*. (Pentagonal Numbers Theorem) If  $n \neq \frac{3k^2 \pm k}{2}$  then the number of partitions of n into an odd number of distinct terms is the same as the number of

its partitions into an even number of distinct terms.

- **20.** Let  $\pi_n$  be the number of partitions of the number n. Determine the generating function of the sequence  $\pi_n$ .
- **21.** Prove that the number of partitions of n into distinct terms is equal to the number of partitions of n into odd terms (1.18), by calculating the generating functions of both sides.
- 22. Which identity follows from problem 1.19?
- 23.\* Prove the following identities by combinatorial considerations:

(a) 
$$(1+x)(1+x^3)(1+x^5)\dots = \sum_{k=0}^{\infty} \frac{x^{k^2}}{(1-x^2)(1-x^4)\dots(1-x^{2k})}$$

(b) 
$$(1+x^2)(1+x^4)(1+x^6)\dots = \sum_{k=0}^{\infty} \frac{x^{k^2+k}}{(1-x^2)(1-x^4)\dots(1-x^{2k})}$$

- **24.** We look for those partitions of the number n which have the property that each number between 1 and n is uniquely representable as a partial sum of the partition. When will the trivial partition  $n = 1 + \ldots + 1$  be the only solution?
- 25. The number of non-congruent triangles with circumference 2n and integer sides is equal to the number of partitions of n into exactly three terms. Determine this number.
- **26.** We have n forints. Every day we buy exactly one of the following products: pretzel (1 forint), candy (2 forints), icecream (2 forints). What is the number  $B_n$  of possible ways of spending all the money?
- **27.** What is the number  $A_n$  of ways of going up n stairs, if we may take one or two steps at a time? Determine

$$\sum_{n=0}^{\infty} A_n x^n$$

**28.** (a) We have n forints and each day we make exactly one purchase. There are  $a_i$  sorts of goods which can be bought for i forints  $(i = 1, \ldots, k)$ . Suppose the polynomial  $x^k - a_1 x^{k-1} - \ldots - a_k = 0$  has distinct roots  $\theta_1, \ldots, \theta_k$ . Prove that the number of possible ways of spending our money is

(b) Determine the generating function of  $C_n$ , and thereby a formula for  $C_n$ .

29. Determine the eigenvalues of the matrix

0	1					0
1	0.	1.				
	1.	٠.				
0	)			1	•	
			n			_

- **30.** How many sequences of length n can be composed from a, b, c, d in such a way that a and b are never neighboring elements?
- 31. What is the number of k-tuples chosen from  $1, \ldots, n$  containing no two consecutive integers?
- **32.** What is the number of monotone increasing functions mapping  $\{1, \ldots, n\}$  into itself?
- **33.** (a) How many monotonic mappings f of  $\{1, \ldots, n\}$  into itself satisfy the condition  $f(x) \le x$  for every  $1 \le x \le n$ ?
- (b) What is the number of sequences of n 0's and n 1's such that there are at least as many 0's as 1's among the first k digits for each  $1 \le k \le 2n$ ?
- **34.** Prove that the number of sequences  $(x_1, \ldots, x_r)$   $(1 \le x_i \le n)$  containing less than i entries  $\le i$  for each  $i = 1, \ldots, n$  is  $(n r)n^{r-1}$   $(1 \le r \le n)$ .
- **35.** A sequence of partitions of the *n*-element set S is constructed as follows: We start with S itself and then we obtain the  $(i+1)^{\rm st}$  partition from the  $i^{\rm th}$  by splitting any one of its classes having more than one element into two non-empty subsets. Thus the partition into one-element classes will be reached after n steps. What is the number of ways of carrying out this procedure?
- **36.** The previous problem is modified to the extent that the  $(i+1)^{st}$  partition is obtained from the  $i^{th}$  by splitting all classes having more than one element into two non-empty subsets. What is the number of ways of carrying out the procedure now?
- 37. We want to break a stick of length n into n pieces of unit length. What is the number of ways of doing so if
- (a) at each step, we break one of the pieces with a length greater than 1 into two,
  - (b) at each step, we break all pieces with a length greater than 1 into two.
- **38.** In how many ways can one put brackets into the product  $x_1 \cdot x_2 \cdot \ldots \cdot x_r$  (in the sense that any bracket includes a product of two factors)?
- **39.** What is the number  $D_n$  of triangulations of a convex n-gon? (A triangulation is a set of n-3 diagonals no two of which intersect internally and which, therefore, divide the n-gon into n-2 triangles.)

- **40.** Determine, without using the preceding result, the generating function of  $D_n$  and prove the result of the preceding problem from this.
- 41. In how many ways can one divide a convex n-gon into triangles by n-3 non-intersecting diagonals, in such a way that each triangle has an edge in common with the convex n-gon?

42. Find a closed formula for each of the following expressions:

(a) 
$$\sum_{k=0}^{\lfloor n/2\rfloor} \binom{n}{2k},$$

(b) 
$$\sum_{k=0}^{m} {n-k \choose m-k},$$

(c) 
$$\sum_{k=0}^{m} {u \choose k} {v \choose m-k},$$

(d) 
$$\sum_{k=0}^{m} (-1)^k \binom{u}{k} \binom{u}{m-k},$$

(e) 
$$\sum_{k=m}^{n} {k \choose m} {n \choose k}$$
,

(f) 
$$\sum_{k=0}^{\lfloor n/7\rfloor} \binom{n}{7k},$$

(g) 
$$\sum_{k=0}^{\lfloor n/2\rfloor} {n-k \choose k} z^k,$$

(h) 
$$\sum_{k=0}^{m} (-1)^k \binom{n}{k},$$

(i) 
$$\sum_{k=0}^{m} {u+k \choose k} {v-k \choose m-k}.$$

43. Prove the following identities:

(a) 
$$\sum_{k=0}^{m} {m \choose k} {n+k \choose m} = \sum_{k=0}^{m} {m \choose k} {n \choose k} 2^k$$
,

(b) 
$$\sum_{k=0}^{m} {m \choose k} {n+k \choose m} = (-1)^m \sum_{k=0}^{m} {m \choose k} {n+k \choose k} (-2)^k$$
,

(c) 
$$\sum_{k=0}^{p} {p \choose k} {q \choose k} {n+k \choose p+q} = {n \choose p} {n \choose q}$$
,

(d) 
$$\sum_{k=0}^{p} \binom{p}{k} \binom{q}{k} \binom{n+p+q-k}{p+q} = \binom{n+p}{p} \binom{n+q}{q}.$$

(e) 
$$\frac{d}{dx} \binom{x}{n} = \sum_{k=1}^{n} \frac{(-1)^{k+1}}{k} \binom{x}{n-k} = \sum_{k=1}^{n} \frac{1}{k} \binom{x-k}{n-k}$$
.

44\*. Prove the following Abel identities:

(a) 
$$\sum_{k=0}^{n} {n \choose k} x(x+k)^{k-1} (y+n-k)^{n-k} = (x+y+n)^n$$
,

(b) 
$$\sum_{k=0}^{n} {n \choose k} (x+k)^{k-1} (y+n-k)^{n-k-1} = \left(\frac{1}{x} + \frac{1}{y}\right) (x+y+n)^{n-1},$$

(c) 
$$\sum_{k=1}^{n-1} {n \choose k} k^{k-1} (n-k)^{n-k-1} = 2(n-1) n^{n-2}$$
,

45. Let

$$f_n(x) = x(x-1)...(x-n+1).$$

Prove that

$$f_n(x + y) = \sum_{k=0}^n {n \choose k} f_k(x) f_{n-k}(y)$$
.

Find other examples of such sequences of polynomials.

#### § 2. The sieve

One powerful tool in the theory of enumeration as well as in prime number theory is the inclusion-exclusion principle (sieve of Eratosthenes). This relates the cardinality of the union of certain sets to the cardinalities of intersections of some of them, these latter cardinalities often being easier to handle. However, the formula does have some handicaps: it contains terms alternating in sign, and in general it has too many of them!

A natural setting for the sieve is in the language of probability theory. Of course, this only means a division by the cardinality of the underlying set, but it has the advantage that independence of occurring events can be defined. Situations in which events are almost independent are extremely important in number theory and also arise in certain combinatorial applications. Number theorists have developed ingenious methods to estimate the formula when the events (usually divisibilities by certain primes) are almost independent. We give here the combinatorial background of some of these methods. Their actual use, however, rests upon complicated number-theoretic considerations which are here illustrated only by two problems.

It should be emphasized that the sieve formula has many applications in quite different situations. These are scattered throughout the book, but § 15 (Reconstruction) in particular may be pointed out in this connection.

A beautiful general theory of inclusion-exclusion, usually referred to as the theory of the Moebius function, is due to L. Weisner, P. Hall and G.-C. Rota and the second half of this section is devoted to this.

1. In a high school class of 30 pupils, 12 pupils like mathematics, 14 like physics and 13 chemistry, 5 pupils like both mathematics and physics, 7 both

physics and chemistry, 4 pupils like mathematics and chemistry. There are 3 who like all three subjects. How many pupils do not like any of them?

**2.** (a) (The Sieve Formula) Let  $A_1, \ldots, A_n$  be arbitrary events of a probability space  $(\Omega, P)$ . For each  $I \subseteq \{1, \ldots, n\}$ , let

$$A_I = \prod_{i \in I} A_i; \ A_0 = \Omega;$$

and let

$$\sigma_k = \sum_{|I|=k} P(A_I), \quad \sigma_0 = 1.$$

Then

$$P(A_1 + \ldots + A_n) = \sum_{j=1}^{n} (-1)^{j-1} \sigma_j.$$

(b) (Inclusion-Exclusion Formula) Let  $A_1, \ldots, A_n \subseteq S$  where S is a finite set, and let

$$A_I = \bigcap_{i \in I} A_i; \quad A_0 = S.$$

Then

$$|S - (A_1 \cup \ldots \cup A_n)| = \sum_{I \subseteq \{1,\ldots,n\}} (-1)^{|I|} |A_I|.$$

- **3.** Determine the number  $\varphi(n)$  of integers between 1 and n coprime to n, given the prime factorization  $p_1^{\alpha_1} \dots p_r^{\alpha_r}$  of n.
- 4. Prove the identity

$$\sum_{i=0}^{n} (-1)^{i} {n \choose i} i^{k} = \begin{cases} 0 & \text{if } 0 \le k < n, \\ (-1)^{n} n! & \text{if } k = n. \end{cases}$$

What do we get if k > n?

5. Let  $p(x_1, \ldots, x_n)$  be a polynomial of degree m, and denote by  $\sigma^k p$  the polynomial obtained by substituting 0's for k of the variables in p in every possible combination and summing the arising  $\binom{n}{k}$  polynomials  $(\sigma^0 p = p)$ . Prove that

$$\sigma^0 p - \sigma^1 p + \sigma^2 p - \ldots = \begin{cases} 0 & \text{if } m < n \\ c \cdot x_1 \dots x_n & \text{if } m = n. \end{cases}$$

What is c?

**6.** Let  $A_1, \ldots, A_n$  be any events,  $B_i = f_i(A_1, \ldots, A_n)$   $(i = 1, \ldots, k)$  polynomials in  $A_1, \ldots, A_n$  and  $c_1, \ldots, c_n$  reals. Then

$$\sum_{i=1}^{k} c_i P(B_i) \ge 0$$

holds for every  $A_1, \ldots, A_k$ , provided it holds in those cases when  $P(A_j) = 0$  or 1 for  $j = 1, \ldots, n$ .

7. (a) (K. Jordán's Formula) Let  $A_1, \ldots, A_n$  be events as in 2.2a. The probability that exactly q of them occur is

$$\sum_{j=q}^n (-1)^{j+q} inom{j}{q} \sigma_j$$
 .

(b) Let  $\eta$  denote the number of  $A_i$ 's that occur (this is a random variable). Then

$$E(x^{\eta}) = \sum_{j=0}^{n} (x-1)^{j} \sigma_{j}.$$

- 8. Express by a similar formula the probability of the event  $D_p$  that at least p of  $A_1, \ldots, A_n$  occur.
- 9. The partial sums of

$$P(\bar{A}_1 \ldots \bar{A}_n) - \sigma_0 + \sigma_1 - \sigma_2 + \ldots$$

are alternating in sign.

10. Prove that

$$\sigma_r \leq \frac{n-r+1}{r} \ \sigma_{r-1};$$

sharpen this inequality if  $\sigma_{r+1} = 0$ .

11. For each  $I \subseteq \{1 \dots n\}$ , let a number  $p_I$  be given. When can one find events  $A_1, \dots, A_n$  such that

$$P(A_I) = p_I$$
?

\*

12. Let G be a simple graph on  $V(G) = \{1, \ldots, n\}$ . Denote by  $\mathfrak{I}_0$  the collection of odd independent subsets of V(G) and by  $\mathfrak{I}'_0$  the collection of those even subsets which span at most one edge. Then

$$P(\bar{A}_1 \dots \bar{A}_n) \leq \sum_{I \in \vartheta'_0} P(A_I) - \sum_{I \in \vartheta_0} P(A_I).$$

Find an analogous lower estimate. (Note that for  $E(G) = \emptyset$  we obtain the sieve formula.)

13. (Brun's Sieve) Let  $f(k) \ge 0$  be any integer-valued function defined on  $\{1, \ldots, n\}$  and set

$$\emptyset = \{I \subseteq \{1, \ldots, n\} : |I \cap \{1, \ldots, k\}| \le 2f(k), \ k = 1, \ldots, n\}.$$

Then

$$P(\bar{A}_1 \ldots \bar{A}_n) \leq \sum_{I \in \mathfrak{F}} (-1)^{|I|} P(A_I).$$

Find an analogous lower estimate.

14\*. (Selberg's Sieve) For each  $I \subseteq \{1, \ldots, n\}$ , let  $\lambda_I$  be a real number, and  $\lambda_0 = 1$ .

Then

$$P(\bar{A}_1 \dots \bar{A}_n) \leq \sum_{I,J \subseteq \{1,\dots,n\}} \lambda_I \lambda_J P(A_{I \cup J}).$$

Find an analogous lower estimate.

15\*. (Cont'd) Let  $p_i = P(A_i)$  and

$$p_I = \prod_{i \in I} p_i.$$

Determine the minimum and the minimizing system  $\{\lambda_I\}$  of

$$\sum_{I, J\subseteq\{1,\ldots,n\}} \lambda_I \, \lambda_J \, p_{I\cup J}$$

under the constraints

$$\lambda_{\emptyset} = 1, \ \lambda_{I} = 0 \text{ for } |I| > k,$$

where  $p_1, p_2, \ldots, p_n$  and k are supposed given.

16.\* (Cont'd) Let  $\varepsilon > 0$ ,  $M \ge 1$ ,  $0 < p_i < 1$  and

$$S = \sum p_1^{l_1} \dots p_n^{l_n},$$

where the summation extends over those systems  $l_1, \ldots, l_n \ge 0$  with  $p_1^{l_1} \ldots p_n^{l_n} \ge \frac{1}{M}$ . Assume that

$$|P(A_K) - p_K| < \varepsilon \quad (K \subseteq \{1, \ldots, n\}).$$

Then

$$P(\bar{A}_1 \dots \bar{A}_n) \leq \frac{1}{S} + \frac{\varepsilon M^2}{(1-p_1)^2 \dots (1-p_n)^2}.$$

17\*. Prove that, for large enough x, the number of primes in the sequence  $l, k + l, \ldots, k(x - 1) + l$  (0 < l < k) is not greater than

$$3 \cdot \frac{k}{\varphi(k)} \cdot \frac{x}{\log x}$$
.

18\*. Let G be a simple graph on  $V(G) = \{1, \ldots, n\}$  with all degrees at most d and let an event  $A_i$  be associated with each point i. Suppose that

(i) 
$$P(A_i) \leq \frac{1}{4d}$$
, and

(ii) every  $A_i$  is independent of the set of all  $A_j$ 's for which j is not adjacent to i. Then

$$P(\bar{A}_1 \ldots \bar{A}_n) > 0.$$

19. (Second Moment Method) Prove the inequality

$$P(\bar{A}_1 \ldots \bar{A}_n) \leq \frac{\sigma_1 + 2\sigma_2}{\sigma_1^2} - 1$$
.

**20.** Let  $P(A_i) = p$ , and assume any two events are independent. Estimate

$$P(\bar{A}_1 \ldots \bar{A}_n)$$

from the above using each of 2.9, Selberg's method (2.14–16) and the preceding formula and compare the results.

**21.** Let  $V = \{x_1, \ldots, x_n\}$  be a set partially ordered by a relation  $\leq$ . Call an  $n \times n$ -matrix  $(a_{ij})$  compatible if

$$a_{ij} \neq 0 \Rightarrow x_i \leq x_j$$
.

Show that the sum, the product and (if it exists) the inverse of compatible matrices is compatible.

**22.** Find a function  $\mu$  defined on  $V \times V$  such that

$$\mu(x, y) = 0 \text{ if } x \nleq y,$$

$$\mu(x, x) = 1,$$

$$\sum_{x \leqslant y \leqslant z} \mu(x, y) = 0 \quad (x < z)$$

(the Moebius function of V).

- **23.** Evaluate the function  $\mu(x, y)$  if V is
  - (a) the lattice of all subsets of a set S (ordered by inclusion),
- (b) an arborescence (here  $x \leq y$  means that the unique path going from the root to y passes through x),
  - (c) the set of integers  $1, \ldots, n$  where  $x \leq y$  means  $x \mid y$ .
- **24.** Turn the partially ordered set "upside down" i.e. consider  $(V, \leq *)$  where  $x \leq *y$  iff  $y \leq x$ . Let  $\mu^*$  denote the Moebius function of  $(V, \leq *)$ . Then

(\*) 
$$\mu^*(x, y) = \mu(y, x).$$

- **25.** Write M as a polynomial in Z (see the solution of 2.22).
- **26.** (Moebius Inversion Formula) Let f(x) be any function defined on V, and set

$$g(x) = \sum_{z \le x} f(z) .$$

Then

$$f(x) = \sum_{z \le x} g(z) \, \mu(z, x) \, .$$

Show that the sieve formula is a special case of this statement.

**27.** Let V be a lattice,  $a \leq b \in V$ , and denote by 0 and 1 the minimal and maximal elements of V. Then

(a) 
$$\sum_{x \vee a = b} \mu(0, x) = 0$$
 (if  $a > 0$ );

(b) 
$$\sum_{x \wedge b=a} \mu(x, 1) = 0$$
 (if  $b < 1$ ).

**28.** Let V be a lattice again and  $x \in V$  such that x is not representable as the union of atoms. Then  $\mu(0, x) = 0$ .

**29.** Let V be a lattice and  $\{A, B, C\}$  a partition of V such that if  $x \in A$  and  $y \leq x$  then  $y \in A$ ; on the other hand, if  $x \in C$  and  $x \leq y$  then  $y \in C$ . Then

$$1 + \sum_{x \in A} \sum_{y \in C} \mu(x, y) = \sum_{x, y \in B} \mu(x, y).$$

**30.** Let V be the lattice of all partitions of the set  $S = \{1, \ldots, n\}$ , where  $\{A_1, \ldots, A_p\} \leq \{B_1, \ldots, B_q\}$  means that each  $A_i$  is contained in some  $B_j$ . Denote by 0 the partition into one-element sets and by 1 the partition into a single set. Then

$$\mu(0, 1) = (-1)^{n-1} (n-1)!$$

31. Let  $(V, \leq)$  be a lattice,  $V = \{x_1, \ldots, x_n\}$ , f(x) any function on V and  $g(x) = \sum_{y \leq x} f(x)$ .

Let  $g_{ij} = g(x_i \wedge x_j)$ , then

$$\det(g_{ij}) = f(x_1) \dots f(x_n).$$

Generalize this formula for partially ordered sets.

32. Evaluate the determinants

$$\begin{vmatrix} (1,1)(1,2)\dots(1,n) \\ (2,1)(2,2)\dots(2,n) \\ (n,1)(n,2)\dots(n,n) \end{vmatrix}; \qquad \begin{vmatrix} (2,2)(2,3)\dots(2,n) \\ (3,2)(3,3)\dots(3,n) \\ (n,2)(n,3)\dots(n,n) \end{vmatrix},$$

where (i, j) denotes the greatest common divisor of i and j.

**33\*.** Let *T* be a tree,  $V = V(T) = \{x_1, ..., x_n\}$  and

$$d_{ij} = d(x_i, x_j)$$

the distance of  $x_i$  from  $x_i$  in T. Then

$$\det(d_{ij})_{i,j=1}^n = -(-2)^{n-2} (n-1).$$

**34.** Let V be a lattice and let  $p_k$  denote the number of chains  $a_0 = 0, a_1, \ldots, a_k = 1$ . Then

$$\mu(0,1) = -p_1 + p_2 - p_3 + \dots$$

35. Let V be a lattice and let R be the set of its atoms. Denote by  $q_k$  the number of those k-tuples in R whose union is  $1 (q_0 = 0 \text{ except when } |V| : \mathbb{A}|1|$ . Then

$$\mu(0, 1) = \sum_{i=0}^{n} (-1)^{i} q_{i}.$$

**36\*.** Let C be a set of pairwise incomparable elements of the lattice V such that every maximal chain (linearly ordered subset) contains an element of C. Denote by  $q_k$  the number of those k-tuples from C whose union is 1 and whose intersection is 0. Then

$$\mu(0, 1) = \sum_{i=0}^{n} (-1)^{i} q_{i}.$$

**37.** Let V be a (finite) geometric lattice of rank r, then  $\mu(0, 1) \neq 0$  and has sign  $(-1)^r$ .

#### § 3. Permutations

Permutations play a role in combinatorics mainly as symmetries of combinatorial structures. Various properties of the symmetric group and its subgroups are important. We shall focus our attention on a few questions here which do not involve extensive algebraic machinery (for example, group representations). So we discuss simple enumeration problems, and two important tools in handling such problems: the cycle index polynomial and a coding of permutations due to M. Hall and A. and C. Rényi.

When we enumerate certain structures we sometimes want to consider some of them as not being essentially different (for example, they may be isomorphic). This usually means that a certain permutation group acts on the underlying set of structures, and we do not want to distinguish two structures if some element of this group maps one onto the other. Pólya's method, discussed in the second part of this section, is a beautiful way to handle such situations.

Many other combinatorial properties of permutation groups, only touched upon here, can be found, for example in Wielandt's Finite Permutation Groups,

Academic Press, 1966.

- 1. What is the number of conjugacy classes in the symmetric group  $S_n$ ?
- **2.** Determine the number of the permutations in  $S_n$ 
  - (a) having no fixed points,
  - (b) consisting of one cycle only.
- **3.** We select a permutation of  $\{1, \ldots, n\}$  at random. What is the probability that the cycle containing 1 has length k? (In this and all subsequent problems all permutations are supposed to have an equal probability of selection.)
- **4.** What is the probability that in a permutation of  $\{1, \ldots, n\}$ , chosen at random, 1 and 2 belong to the same cycle?
- 5. We choose a permutation of  $\{1, \ldots, n\}$  at random. What is the expected number of cycles?
- 6. We have n savings boxes, with different keys. Someone locks the boxes and throws the keys into them at random, putting one key in each box. We break open k boxes. What is the probability that we are now able to open all the rest? (In mathematical form: Whe choose a permutation  $\pi$  of  $\{1, \ldots n,\}$  at random. What is the probability that the cycles of  $\pi$  containing  $1, \ldots, k$  cover all points?)

7. Denote by  $p_n(x_1, \ldots, x_n)$  the cycle index of  $S_n$ , the symmetric group on n elements, and put  $p_0 = 1$ . Prove that

$$\sum_{n=0}^{\infty} p_n(x_1, \ldots, x_n) y^n = \exp \left( x_1 y + \frac{x_2}{2} y^2 + \ldots + \frac{x_k}{k} y^k + \ldots \right).$$

8. Determine the cycle index of

(a) the group of rotations of a regular n-gon;

(b) the group of permutations of nk elements which keep a given partition P into k n-element classes invariant.

Generalize these results.

- 9. Two permutation groups (acting effectively) have the same cycle index. Are they necessarily isomorphic?
- 10. Denote by  $q_n(x_1, \ldots, x_n)$  the cycle index of the alternating group  $A_n(q_0 = 2)$ . Determine the generating function

$$\sum_{n=0}^{\infty} q_n(x_1,\ldots,x_n) y^n$$

11. Prove that if all but a finite number of the numbers  $x_1, x_2, \ldots$  are equal to 1, then

$$\lim_{n\to\infty}p_n(x_1,\ldots,x_n)$$

exists, and determine its value.

12. (a) Let  $s_{n,k}$  denote the number of those permutations of  $\{1,\ldots,n\}$  having exactly k cycles. Find a closed form of the polynomial

$$f_n(x) = \sum_{k=1}^n s_{n,k} x^k.$$

What is  $(-1)^{n-k}s_{n,k}$ ?

- (b) Give a new proof of 3.5.
- 13. What is the exponential generating function of the number  $g_n$  of 2-regular simple graphs on n given points?
- 14. In how many ways can one partition the 2n letters  $\{1, 1, 2, 2, \ldots, n, n\}$  into n pairs if

(a) the order of elements in a pair does not count,

- (b) the order of elements in a pair does count? (Determine the generating function.)
- 15. Give a new solution of 3.2, based on the cycle index of  $S_n$ .
- 16. Choose a permutation  $\pi$  at random. Denote by  $\overline{\pi}(i)$  the number of those integers  $1 \leq j \leq i$  with  $\pi(j) \geq \pi(i)$ . Then  $\overline{\pi}(1), \ldots, \overline{\pi}(n)$  are random variables. Prove that they are independent.

- 17. (a) What is the expected number of (on-going) records at a long jump competition, in which each of the n competitors has one jump, these jumps are all different and the starting order is drawn at random?
  - (b) What is the probability that there are exactly k records?
- 18\*. Imagine a long jump competition, in which no two of the n competitors jump the same distance. They jump in a drawn order. One can bet according to the following rule: After a jump, one can say "This is best of all." We arrive on the scene after the k<sup>th</sup> jump.
- (a) What is the probability  $p_k$  that, following the best strategy, we will guess the winner correctly (information about the previous results is available)?
- (b) What strategy should we follow in order to have the largest probability of winning (provided we arrive in time)?
- 19. What is the number of permutations  $\pi$  of  $\{1, \ldots, n\}$  so that there is no triple i < j < k with

$$\pi(j) < \pi(i) < \pi(k)?$$

**20.** Let  $a_{n,k}$  denote the number of those permutations  $\pi$  of  $\{1, \ldots, n\}$  which have k inversions (pairs i < j with  $\pi(i) > \pi(j)$ ). Prove that

$$\sum_{k=1}^{\binom{n}{2}} a_{n,k} x^k = (1+x) (1+x+x^2) \dots (1+x+\dots+x^{n-1}).$$

- 21. Along a speed track there are some gas-stations. The total amount of gasoline available in them is equal to what our car needs for going around the track. Prove that there is a gas-station such that if we start there with an empty tank, we shall be able to go around the track without running out of gasoline.
- **22.** (a) Let  $x_1, \ldots, x_n$  be real numbers. For each permutation  $\pi$  of  $\{1, \ldots, n\}$ , define

$$a(\pi) = \max\{0, x_{\pi(1)}, x_{\pi(1)} + x_{\pi(2)}, \dots, x_{\pi(1)} + \dots + x_{\pi(n)}\}.$$

Also consider the cycles  $C_1, \ldots, C_k$  of  $\pi$  and set

$$b(\pi) = \sum_{i=1}^{k} \max \left(0, \sum_{i \in C_i} x_i\right).$$

Then  $\{a(\varrho): \varrho \in S_n\}$  and  $\{b(\pi): \pi \in S_n\}$  are equal as collections.

(b) m boys and m girls form "rings" for a dance at random (there may be pairs or even singles: two and one-element "rings"). Prove that the probability that in each "ring" the number of boys is equal to the number of girls is  $\frac{1}{m+1}$ .

\*

**23.** (a) How many convex k-gons can be formed from the vertices of a regular n-gon, if two k-gons are considered essentially different when they do not arise from each other by rotation?

- (b) What is the number of k-colorations of the vertices of a regular n-gon, if two k-colorations are not considered as essentially different when they arise by rotation from each other?
- **24.** (Burnside's Lemma) Let  $\Gamma$  be a group of permutations of  $\{1, \ldots, n\}$  with k orbits. Then the average number of fixed points of permutations in  $\Gamma$  is k.
- **25.** Let  $\Gamma$  be a permutation group acting on a set  $\Omega$ . Suppose a "weight" w(x) is associated with each  $x \in \Omega$  such that w(x) is invariant under  $\Gamma$ , i.e. for any given orbit  $\Theta$  of  $\Gamma$ , w(x) takes the same value  $w(\Theta)$  for all  $x \in \Theta$ . Prove that

$$\sum_{\Theta} w(\Theta) = \frac{1}{|\Gamma|} \sum_{\pi \in \Gamma} \sum_{\pi(x) = x} w(x).$$

**26.** (Pólya–Redfield Enumeration Method I) Let  $\Gamma$  be a permutation group with cycle index  $F(x_1, \ldots, x_n)$  on a set D and let R be another set. We say that the mappings  $f, g: D \to R$  are essentially different if no  $\pi \in \Gamma$  can be found such that

$$\pi f = g$$
.

What is the number of essentially different mappings of D into R?

27\*. (Cont'd) What happens if a permutation group  $\Gamma_1$  is given on R and f, g are only considered essentially different if for no  $\pi \in \Gamma$ ,  $\varrho \in \Gamma_1$ 

$$\pi f = g \rho$$
?

(Express this number by the cycle indices F, G of  $\Gamma$  and  $\Gamma_1$ .)

- 28\*. (Cont'd) Count the number of essentially different injections (one-to-one mappings) of D into R.
- **29.** (Pólya–Redfield Method II) Let D be a finite set, R arbitrary; let  $\Gamma$  be a permutation group acting on D with cycle index F. Assume each  $y \in R$  has a positive integral weight w(y). Denote by  $r_n$  the number of elements of R with weight n (suppose this is finite) and set

$$r(x) = \sum_{n=1}^{\infty} r_n x^n.$$

Let  $a_n$  denote the number of essentially different mappings  $f: D \to R$  (in the sense of 4.26) such that

$$\sum_{x \in D} w(f(x)) = n.$$

Prove that

$$\sum_{n=1}^{\infty} a_n x^n = F(r(x), r(x^2), \ldots).$$

- **30.** Solve the problem of distributing n forints to k persons using the above technique.
- 31. Give a new proof of the formula for the exponential generating function of the number of partitions of a set (1.11), using the Pólya-Redfield method.

#### § 4. Two classical enumeration problems in graph theory

I have long felt it a very surprising fact that the number of trees on n labelled points is  $n^{n-2}$ . The reason why this is surprising is that one does not see any direct way to enumerate trees and one would expect a less compact result. The difficulty and beauty of this problem have challenged several authors to find proofs, usually by dealing with a well-chosen generalization of the problem. Some of these are given here; the reader may test his knowledge of much basic enumeration technique.

Another theory, with less general results — but quite ingenious methods — is the enumeration of 1-factors in graphs. The special case when the graphs are bipartite is equivalent to the problem of "restricted permutations", that is, the problem of enumerating those permutations of  $\{1, \ldots, n\}$  which map each element i onto one of the elements of a specified subset  $A_i$ . The methods here also use different enumeration techniques but they usually work for certain

special cases only.

It is also interesting to note that the theory of enumeration of trees is important in solving the Kirchhoff equations in electrical network theory and that enumeration of 1-factors plays an important role in ferromagnetism. (See Seshu-Read, Linear Graphs and Electrical Networks, Addison-Wesley, 1961; E.W. Montroll, in: Applied Combinatorial Mathematics, E. F. Beckenbach, ed. Wiley, 1964.)

1. Let  $v_1, \ldots, v_n$  be given points and  $d_1, \ldots, d_n$  be given numbers such that  $\sum_{i=1}^n d_i = 2n-2, d_i \ge 1$ . Prove that the number of trees on the set  $\{v_1, \ldots, v_n\}$  in which  $v_i$  has degree  $d_i$   $(i=1,\ldots,n)$  is

$$\frac{(n-2)\,!}{(d_1-1)\,!\ldots(d_n-1)\,!}\,.$$

**2.** (Cayley Formula) Prove that the number of all trees on n points is  $n^{n-2}$ .

3. Let

$$p_n(x_1, \ldots, x_n) = \sum x_1^{d_T(v_1)-1} \ldots x_n^{d_T(v_n)-1}$$

where  $d_T(v_i)$  denotes the degree of the point  $v_i$  in the tree T and the summation extends over all trees on  $\{v_1, \ldots, v_n\}$ . Prove, without using 4.1, that

$$p_n(x_1,\ldots,x_n)=(x_1+\ldots+x_n)^{n-2}.$$

Deduce the Cayley formula from this.

- 4. Let  $T_1, \ldots, T_r$  be trees on disjoint sets of points and  $V = V(T_1) \cup \ldots \cup V(T_r)$ . What is the number of trees on V containing  $T_1, \ldots, T_r$ ?
- 5. Let T be a tree on points  $v_1, \ldots, v_n$ . Delate the endpoint having the least index and write down the index of its neighbor. Repeat this procedure with the resulting tree, until a tree with only one point remains. This associates a sequence of n-1 numbers with T, called the *Prüfer Code* of T. Prove that

- (a) the Prüfer code of T uniquely characterizes T,
- (b) given any sequence  $(a_1, \ldots, a_{n-1})$  such that  $1 \le a_i \le n$ ,  $a_{n-1} = n$ , there is a (unique) tree with this Prüfer code.
- (c) Deduce the Cayley formula.
- **6.** Denote by  $T_n$  the number of trees on points  $v_1, \ldots, v_n$ . Prove that

$$T_{n} = \sum_{k=1}^{n-1} k \binom{n-2}{k-1} T_{k} T_{n-k}$$

and prove Cayley's formula from this identity.

7. (a) The exponential generating function

$$t(x) = \sum_{n=1}^{\infty} \frac{T_n}{(n-1)!} x^n$$

satisfies the functional identity

$$t(x) e^{-t(x)} = x.$$

- (b) Prove the Cayley formula from this identity.
- 8. What is the number of all trees on n points with exactly n-l endpoints?
- **9.** (a) Let G be a digraph,  $V(G) = \{v_1, \ldots, v_n\}$  and  $E(G) = \{e_1, \ldots, e_m\}$ . Let A be the point-edge incidence matrix of G, i.e. the matrix  $A = (a_{ij})$  defined by

$$a_{ij} = \begin{cases} 1 & \text{if } v_i \text{ is the head of } e_j, \\ -1 & \text{if } v_i \text{ is the tail of } e_j, \\ 0 & \text{otherwise.} \end{cases}$$

Remove any row from A and let  $A_0$  be the remaining matrix. Prove that the number  $T(\hat{G})$  of spanning trees of G is det  $A_0A_0^T$ .

- (b) What are the entries of  $A_0A_0^T$ ?
- (c) Deduce the Cayley formula from this.
- 10. Let

$$p_G(x_1,\ldots,x_n)=\sum x_1^{d_T(v_1)}\ldots x_n^{d_T(v_n)}$$

where G is a graph on  $\{v_1, \ldots, v_n\}$  and the summation extends over all spanning trees of G (cf. 4.3). Define

$$a_{ij}(x_1\ldots,x_n) = \begin{cases} -x_ix_j & \text{if } v_i,v_j \text{ are adjacent,} \\ 0 & \text{if } i\neq j \text{ and } v_i,v_j \text{ are non adjacent,} \\ x_i \cdot \sum_{v_\nu \text{ adjacent to } v_j} x_\nu & \text{if } i=j; \end{cases}$$
 and

and

$$D = (a_{ij})_{i=1}^{n-1} \stackrel{n-1}{i=1}$$
.

Prove that

$$p_G(x_1,\ldots,x_n)=\det D.$$

- 11. What is the number of trees on  $\{v_1, \ldots, v_n; w_1, \ldots, w_m\}$  in which each edge joins a  $v_i$  to a  $w_i$ ?
- 12. (a) The number T(G) of spanning trees of a graph G is given by

$$T(G) = \Sigma (-1)^{n-r} T(\overline{G}[X_1]) \dots T(\overline{G}[X_r]) |X_1| |\dots |X_r| n^{r-2}$$

where the summation extends over all partitions  $(X_1, \ldots, X_r)$  of V(G) and n = |V(G)|.

Determine T(G) when G is

- (b) the complement of a graph consisting of q independent edges and n-2q isolated points,
- (c) the complement of a graph consisting of q edges adjacent to a point v and n-q-1 isolated points.
- 13. (a) What is the number of "binary plane trees", i.e. those trees on 2n points which are embedded in the plane and each point has degree 1 or 3 with some endpoint specified as root? (Two plane trees are the "same" if there is an isomorphism between them which keeps the cyclic ordering of edges in the star of each point.)
  - (b) What is the number of plane trees on n points rooted at an endpoint?
- 14. What is the number E(n, k) of forests F on points  $v_1, \ldots, v_n$  having k components and such that  $v_1, \ldots, v_k$  belong to distinct components?
- 15. Let G be an acyclic digraph with a specified root a and suppose G has a spanning arborescence rooted at a. What is the number of spanning arborescences of G?
- 16. Let G be a digraph without loops,  $V(G) = \{v_1, \ldots, v_n\}$ . Let  $a_{ij}$  be the number of edges joining  $v_i$  to  $v_j$ , and let  $d_i^-$  be the indegree of  $v_i$ .
- (a) The number of spanning arborescences of G rooted at  $v_n$  is equal to the determinant

$$\Delta(G) = \begin{vmatrix} d_1^- & -a_{12} & \dots - a_{1,n-1} \\ -a_{21} & d_2^- & -a_{2,n-1} \\ \vdots & & \vdots \\ -a_{n-1,1} & -a_{n-1,2} & \dots & d_{n-1}^- \end{vmatrix}.$$

(b) Find a similar formula for the polynomial

$$\sum x_1^{d_T^+(\nu_1)} \dots x_n^{d_T^+(\nu_n)},$$

where T ranges over all spanning arborescences of G rooted at  $v_n$  (cf. 4.10).

17\*. Let n be odd and let  $\pi$  be a permutation of the n-element set V. The number of trees on V admitting  $\pi$  as an automorphism is

0 if  $\pi$  has no fixed points,

$$k_1^{k_1-2} \prod_{i=2}^n \left(\sum_{d\mid i} dk_d\right)^{k_i-1} \left(\sum_{\substack{d\mid i\\d\neq i}} dk_d\right)$$
 if  $\pi$  has a fixed point,

where  $k_i = k_i(\pi)$  is the number of *i*-cycles in  $\pi$ .

18. Let  $W_n$  denote the number of non-isomorphic trees on n points with a specified point called *root*. Prove that

$$2^n < W_n < 4^n \ (n \ge 6).$$

19\*. Set

$$w(x) = \sum_{n=1}^{\infty} W_n x^n.$$

(a) Prove the identity

$$w(x) = x(1-x)^{-W_1}(1-x^2)^{-W_2}\dots(1-x^n)^{-W_n}\dots$$

(b) Deduce the following identities from both (a) and the Pólya–Redfield method:

$$w(x) = x \cdot \exp\left[\frac{w(x)}{1} + \frac{w(x^2)}{2} + \dots\right]$$
$$w(x) = x \sum_{n=0}^{\infty} p_n(w(x), w(x^2), \dots).$$

where  $p_n$  is the cycle index of the symmetric group  $S_n$ .

20\*\*. (a) Show that w(x) has a radius of convergence  $0 < \tau < 1$  and has a unique singularity on the circle  $|x| = \tau$  at  $x = \tau$ . Moreover, w(x) is an analytic function of  $\sqrt{\tau - x}$  near this singularity.

(b) Prove that there is a positive constant c such that

$$W_n \sim c n^{-3/2} \tau^{-n}$$
.

\*

**21.** Let G be a bipartite graph with 2-coloration  $\{U, V\}$ ,  $U = \{u_1, \ldots, u_n\}$ ,  $V = \{v_1, \ldots, v_n\}$  and let  $a_{ij}$  denote the number of  $(u_i, v_j)$ -edges in G. Set

$$A = (a_{ij})_{i=1}^n {n \atop j=1}$$
.

Then the number of 1-factors of G is per A.

22. Determine the number of 1-factors in the "ladder" graph shown in Fig. 1.




Fig. 1

**23.** What is the number of 1-factors in  $K_{n,n}$  minus n disjoint edges? (That is, in how many ways can a company consisting of n married couples dance if everyone dances but nobody dances with his spouse?)

**24.** Let G be a simple graph and  $\vec{G}$  an orientation of G with the property that, in any even circuit C of G, the number of edges directed one way (or the other) around the circuit is odd. Let  $V(G) = \{v_1, \ldots, v_n\}$  and G be defined by

$$B = (b_{ij}), \ b_{ij} = \begin{cases} 1 & \text{if} \quad (v_i, v_j) \in E(\vec{G}), \\ -1 & \text{if} \quad (v_j, v_i) \in E(\vec{G}), \\ 0 & \text{otherwise.} \end{cases}$$

Prove that the number of 1-factors of G is |Pf B|.

- **25.** (Cont'd) For which orientations of a graph G is |Pf B| the number of 1-factors?
- **26.** Let B be any skew symmetric matrix. Then

$$\det B = (\operatorname{Pf} B)^2.$$

- 27. Let G be a 2-connected planar graph which has a 1-factor F. Prove that G has an orientation  $\vec{G}$  such that every circuit C which alternates with respect to F, contains an odd number of edges oriented in some given direction.
- 28. Determine the number of 1-factors of the "ladder" (4.19) from the preceding results.
- **29\***. Let  $a_n$  be the number of ways to cover a  $(2n) \times (2n)$  chessboard by dominoes. Prove the formulas

$$(a) \quad a_{n} = 2^{2n^{3}} \prod_{k=1}^{n} \prod_{l=1}^{n} \left( \cos^{2} \frac{k\pi}{2n+1} + \cos^{2} \frac{l\pi}{2n+1} \right)$$

$$(b) \quad \begin{vmatrix} 2n \\ 0 \end{vmatrix} & \binom{2n-2}{2} & \binom{2n-4}{4} & \dots \\ 0 & \binom{2n}{0} & \binom{2n-2}{2} & \dots \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & \binom{2n}{0} & \binom{2n-2}{2} & \dots \\ \binom{2n-1}{1} & \binom{2n-3}{3} & \dots & \vdots \\ 0 & \dots & \binom{2n-1}{1} & \binom{2n-3}{3} & \dots \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & \binom{2n-1}{1} & \binom{2n-3}{3} & \dots \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & \binom{2n-1}{1} & \binom{2n-3}{3} & \dots \\ \frac{n-1}{2} & \text{columns} \end{vmatrix}$$

(c) Determine

$$\lim_{n\to\infty}\frac{\log a_n}{n^2}.$$

- **30.** Prove that the number of spanning trees of the  $n \times n$  "chessboard" (i.e. the graph whose points are the squares and two of them are adjacent iff they share an edge) is equal to the number of ways of covering a  $(2n-1)\times(2n-1)$  chessboard minus its upper left-hand corner by dominoes.
- 31. What is the number of k-element matchings in the following bipartite graph:

$$V(G) = \{v_1, \dots, v_n; u_1, \dots, u_n\};$$
  
 $E(G) = \{(u_i, v_j) : i < j\}.$ 

32. What is the number of those permutations  $\pi$  of  $\{1, \ldots, n\}$  having the property that

$$|\pi(k) - k| \le 1$$
  $(k = 1, ..., n)$ ?

33\*. Let  $a_n$  denote the number of permutations  $\pi$  of  $\{1,\ldots,n\}$  such that

$$|\pi(k)-k|\leq 2.$$

Determine

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

**34.** Determine the number  $U_{n,p}$  of those permutations  $\pi$  of  $\{1, 2, \ldots, n\}$  which satisfy

$$\pi(k) \le k + p - 1 \quad (k = 1, \ldots, n).$$

35. Find a formula for the number of permutations  $\pi$  such that

$$|\pi(k)-k| \leq n-p \quad (k=1,\ldots,n),$$

where p is small.

**36.** Denote by  $p_n$  the number of those permutations  $\pi$  of  $\{1,\ldots,2n\}$  satisfying

$$|\pi(i)-i|\leq n.$$

Prove that

$$p_n = \sum_{k=0}^n S(n, k)^2 (k!)^2.$$

## § 5. Parity and duality

Parity considerations are applied in graph theory quite often. This loosely connected section contains various examples of parity problems, together with questions of different sorts to which these exercises lead. The first graph-theoretic result, Euler's solution of the Kőnigsberg bridge problem, is typical of the role of parity. Studying Euler trails will lead to another classical problem, finding the way out of a labyrinth.

The most important applications of parity considerations in graph theory are linear spaces of circuits and cuts. This is the starting point of matroid theory on the one hand, algebraic topology on the other. Basic results in algebraic topology such as Sperner's lemma (5.29) depend on parity arguments. We also note that certain existence results follow by showing that the number of objects in question is odd (for example, 5.20 and, more convincingly, 5.30). Linear spaces of circuits and cuts play a role in the characterization of planar graphs and in the study of the relationship between a planar graph and its dual.

Having finished with enumeration problems, we shall more frequently encounter the notion of a good characterization (see the introduction). Problems 5.3, 5.4, 5.6 are simple examples of well-characterizing necessary and sufficient conditions. A more important example is Kuratowski's criterion for planarity. Note that the MacLane and Whitney criteria are not "good characterizations" in this sense. (This shows at the same time that the name "good characterization" may be misleading; it does not have anything to do with the depth of the theorem.)

- 1. (a) Is there a graph with degrees 3, 3, 3, 3, 5, 6, 6, 6, 6, 6, 6?
- (b) Is there a bipartite graph with degrees 3, 3, 3, 3, 5, 6, 6, 6, 6, 6, 6, 6, 6, 6?
  - (c) Is there a simple graph with degrees 1, 1, 3, 3, 3, 3, 5, 6, 8, 9?
- 2. Which numbers occur as orders of k-regular simple graphs?
- 3. A graph is bipartite iff every circuit of it is even. Is the statement (with cycle instead of circuit) true for digraphs? Or strongly connected digraphs?
- 4. We associate a value v(e) with each edge of a digraph G. This may be considered as the work needed to go from the tail of e to the head of e. To go conversely we need -v(e) work. We want to find a "potential", i.e. a function p(x) defined on V(G) such that if e = (x, y) then v(e) = p(y) p(x). Show that this is possible if and only if going around any circuit, the total work needed is 0.
- 5. The points of a strongly connected digraph G can be 2-colored so that each point is joined to a point of opposite color by a directed edge if and only if G has an even cycle.
- **6.** If G is a weakly connected digraph with  $d_G^-(x) = d_G^+(x)$  for each x then G has an Euler trail.
- 7. Let  $G_{k,n}(k \geq 2)$  be defined as follows. The points of  $G_{k,n}$  are vectors of dimension k composed from  $1, \ldots, n$ . The vectors  $(a_1, \ldots, a_k)$  and  $(b_1, \ldots, b_k)$  are joined by a directed edge if  $a_2 = b_1$ ,  $a_3 = b_2$ , ...,  $a_k = b_{k-1}$ . Show that  $G_{k,n}$  is
  - (a) Eulerian;
  - (b) Hamiltonian.
- 8. Show that we can associate 0 or 1 with each point of a  $2^k$ -cycle ( $k \ge 2$ ) in such a way that all arcs of length k of the cycle give different 01-sequences.
- 9. If G has a point of outdegree at least 3, then the number of its Euler trails is even. (Two Euler trails are not considered to be different if they give the same cyclic permutation of the edges.)

- 10. Let G be a digraph in which the outdegree of each point is equal to its indegree. Let T be a spanning inverse arborescence with root  $x_0$ . We start from  $x_0$  and walk along the edges of G according to the following rule:
- (\*) From each point x we go to the next one along an edge (starting from x) not used previously. We use the edges of T only if we have no other choice.

Show that we get stuck at  $x_0$  and that, by this time we have traversed an Euler trail.

11. Let  $V(G) = \{x_0, \ldots, x_{n-1}\}, d_G^+(x_i) = d_G^-(x_i) = d_i$ . Show that the number of Euler trails in G is a multiple of

$$(d_0-1)!\dots(d_{n-1}-1)!$$

- 12. (Labyrinth Problem). Starting at a point  $x_0$  we walk along the edges of a connected graph G according to the following rules:
  - (\*) We never use the same edge twice in the same direction.
- (\*\*) Whenever we arrive at a point  $x \neq x_0$  not previously visited by us, we mark the edge along which we entered x. We use the marked edge to leave x only if we must, i.e. if we have used all the other edges before.

Show that we get stuck at  $x_0$  and that, by then, every edge has been traversed in both directions.

- 13. If G is a graph with even degrees, then G can be oriented in such a way that every point of the resulting digraph  $\vec{G}$  has the same outdegree as indegree.
- 14. (a) A graph G has an Euler trail iff it is connected and has even degrees.
- (b) If a connected graph G has 2k points of odd degrees, then it is the union of k edge-disjoint open trails ( $k \ge 1$ ).
- 15. If a planar map has an Euler trail then the map can be drawn without the lifting pencil from the paper, without going over the same line twice and without crossing our line (at most touching it).
- 16. The number of subgraphs G' of a connected graph G which have even degrees and V(G') = V(G) is equal to  $2^{m-n+1}$  (m = |E(G)|, n = |V(G)|).
- 17\*. (a) The set of points of any graph G can be partitioned into two classes  $V_1$ ,  $V_2$  such that both  $V_1$  and  $V_2$  span subgraphs with even degrees.
- (b) For any graph G, V(G) can be partitioned into two classes  $V_1$ ,  $V_2$  such that  $V_1$  spans a subgraph with even degrees and  $V_2$  spans a subgraph with odd degrees.
- 18\*. (a) Let G be a graph and denote by  $m_r(G)$  the number of r-element matchings of G. Express  $m_1(G), m_2(G), \ldots$  in terms of |V(G)| = n and  $m_1(\overline{G}), m_2(\overline{G}), \ldots$
- (b) If |V(G)| is even and  $\overline{G}$  has an odd number of matchings then G has a 1-factor.
- (c) G has an even number of 1-factors iff there is a non-empty set  $S \subseteq V(G)$  such that every point is adjacent to an even number of points of S (e.g. Eulerian graphs).

19. If G is a simple digraph and h(G) denotes the number of Hamiltonian paths in G, then

$$h(G) \equiv h(\overline{G}) \pmod{2}$$
.

Is this true for undirected graphs?

- 20. Any tournament has an odd number of Hamiltonian paths.
- **21\***. Every edge e of a 3-regular graph G is contained in an even number of Hamiltonian circuits.
- 22. A 3-regular bipartite graph with at least 4 points has an even number of Hamiltonian circuits.

\*

- **23.** Let G be a connected planar map and  $G^*$  its dual. Show that G and  $G^*$  have the same number of spanning trees.
- **24.** (Euler's Formula) Show that if G is a connected planar map then it has m-n+2 faces, where

$$m=|E(G)|,\ n=|V(G)|.$$

- 25. (a) A simple planar graph with n points has at most 3n-6 edges.
- (b) A simple triangle-free planar graph with n points has at most 2n-4 edges.
- 26. If a planar map has even degrees then the faces can be 2-colored in such a way that faces with a common edge on their boundary have different colors.
- 27. If G is a 4-regular planar map, then it can be oriented in such a way that each point has two edges coming in, two edges going out, and these two pairs separate each other in the given embedding.
- 28. (a) Can you draw a planar map in the interior of a pentagon so that the faces are triangles (except, of course, the outermost one), and each point has even degree?
- (b) Suppose we split the interior of a pentagon into triangles by a map such that all points except those on the pentagon have even degree. Which points of the pentagon have odd degree?
- **29.** Let G be a planar map with triangular faces. Suppose the points of G are 3-colored. Show that the number of faces which get all three colors is even.
- **30\***. Let G be a planar map whose faces are triangles except for the exterior face abcd. Let  $V(G) = V_1 \cup V_2$ ,  $a, c \in V_1$ ,  $b, d \in V_2$ . Then either  $V_1$  contains an (a, c)-path or  $V_2$  contains a (b, d)-path.
- **31.** Let V be the space of vectors  $(a_1, \ldots, a_n)^T$  where the  $a_i$ 's are from a field F. For  $\mathbf{a} = (a_1, \ldots, a_n)^T$ ,  $\mathbf{b} = (b_1, \ldots, b_n)^T$ , define their inner product

by  $\mathbf{a}^T\mathbf{b} = a_1b_1 + \ldots + a_nb_n$ . Further, let M be a subspace of V and A a linear transformation of V. Decide whether the following assertions are true<sup>†</sup>:

- (a) If A is non-singular, so is its transpose  $A^T$ .
- (b) More generally, if A is non-singular, so is  $A^{T}A$ .
- (c) Even more generally,  $\operatorname{Ker} A^T A = \operatorname{Ker} A$ .
- (d)  $M \cap M^{\perp} = \{0\}$  (where  $M^{\perp}$  is the subspace orthogonal to M i.e.  $M^{\perp} = \{\mathbf{a} : \mathbf{a}^T \mathbf{b} = 0 \text{ for each } \mathbf{b} \in M\}$ .
  - (e)  $\langle M \cup M^{\perp} \rangle = V(\langle X \rangle$  denotes the subspace generated by X).
  - (f) dim  $M + \dim M^{\perp} = n$ .
  - (g)  $(M^{\perp})^{\perp} = M$ .
- **32.** Let V be the n-dimensional vector space over GF(2), and M a subspace of V. Show that

$$\mathbf{i} = (1, \dots, 1)^T \in \langle M \cup M^{\perp} \rangle.$$

- **33.** Let  $E(G) = \{e_1, \ldots, e_m\}$ . Represent each subset of E(G) by a 01-vector, where the  $j^{\text{th}}$  entry is 1 iff  $e_j$  belongs to the subset. This way we identify the subsets of E(G) with the elements of an m-dimensional vector space  $V_G$  over GF(2).
- (a) Determine the subspace  $U_G$  generated by the stars, and the orthogonal subspace  $W_G$ . Give new proofs of 5.3, 5.16 and 5.17, using these linear spaces.
- (b) Prove that the decomposition in 5.17a is unique iff G has an odd number of spanning trees.
- **34.** Let G be a 2-connected planar map. Show that the boundaries of all finite faces form a basis for the space  $W_G$ .
- **35\***. Let G be a graph,  $C_1, \ldots, C_f$  circuits of G and suppose
  - (1) each edge of G is contained in at most two of  $C_1, \ldots, C_f$ ;
  - (2)  $C_1, \ldots, C_f$  constitute a basis of  $W_G$ . Show that
    - (a) if  $K = \sum_{i \in I} C_i$  is a circuit and


$$\sum_{i \in I} C_i \subseteq \bigcup_{i \in I} C_i$$

then  $J \subseteq I$   $(I, J \subseteq \{1, \ldots, f\})$ ;

(b) either all blocks of G are single circuits or there are two  $C_i$ 's, say  $C_1$  and  $C_2$  such that  $C_1 + C_2$  is a circuit;

 $<sup>\</sup>dagger$  This exercise really belongs to linear algebra. Most textbooks, however, do not consider inner products over finite fields which is the important case for us. (Note that, if F is the field of complex numbers, our inner product is not the same as the one usually introduced.) The reader will find it instructive to clear up the similarities and differences through this exercise. (All statements are true for the real field.)

- (c) G is planar, and  $C_1, \ldots, C_f$  are faces.
- (d) (MacLane's Criterion) A graph G is planar iff  $W_G$  has a basis such that each edge belongs to at most two elements of it.
- **36\***. (Withney's Criterion) A connected graph G is planar iff there exist another graph G\* and a one-to-one mapping  $\varphi$  of E(G) onto  $E(G^*)$  such that whenever T is a spanning tree of G,  $\varphi(E(G)-E(T))$  gives a spanning tree in G\* and vice versa.
- $37^*$ . Let G be a minimal non-planar graph (i.e. each proper subgraph of G is planar) with degrees at least 3. Then
  - (a) G is 3-connected,
  - (b) G contains a circuit with a chord,
  - (c)  $G \simeq K_5$  or  $K_{3,3}$ ,
- (d) (Kuratowski's Criterion) A graph is planar if and only if it contains no subdivision of  $K_5$  or  $K_{3,3}$ .






Fig. 2

38\*. If G is a simple planar graph, then G has an embedding in the plane such that all edges are straight segments.

# § 6. Connectivity

Several central concepts of graph theory belong to this chapter; separating sets of points and edges, connected components, blocks, paths, circuits, trees and forests. These concepts occur in most fields of graph theory and hence

they belong to the basic techniques a graph theorist uses.

The theory of connectivity is one of the most developed branches of graph theory (together with factorization problems, which are discussed in the next chapter). One main reason for this is Menger's theorem, which links connectivity defined in terms of separation to connectivity defined in terms of connecting paths. This gives a "good characterization" of k-connected graphs. Closely related to Menger's theorem is the so-called Max-Flow-Min-Cut theorem, which plays a fundamental role in flow theory. This puts the duality between connection and separation in a very graphic form.

Problems involving connectivity between two specified points are usually settled without difficulty using Menger's theorem. On the other hand, connectivities between more than two points are more difficult to handle and are,

to a large extent, independent of Menger's theorem. Such problems arise in the study of minimal k-connected graphs, multicommodity flows, safe communication networks, etc. Their solutions are difficult but some typical manipulations with cuts occur repeatedly and these may lead to ideas for a general approach.

Some of the strongest results in the field are structure theorems, which prove that certain classes of graphs can be constructed by repeated application of some simple transformation, e.g. 2-connected graphs by repeatedly attaching

"ears" (cf. 6.27, 6.28, 6.33, 6.52, 6.53, 6.64).

1. Show that

$$c(G) + |E(G)| \ge |V(G)|$$

holds for every graph G.

2. (a) Let  $G_1, G_2$  be two graphs with  $V(G_1) = V(G_2)$ . Show that

$$c(G_1) + c(G_2) \le c(G_1 \cup G_2) + c(G_1 \cap G_2).$$

- (b) This holds without assuming  $V(G_1) = V(G_2)$ .
- 3. Let  $d_1 \leq \ldots \leq d_n$  be the degrees of the simple graph G and assume that  $d_k \geq k$  for every  $k \leq n d_n 1$ . Then G is connected.
- **4.** Show that  $G_1 \times G_2$  is connected if and only if  $G_1$ ,  $G_2$  are connected and one of them contains an odd circuit.
- 5. A connected k-regular bipartite graph is 2-connected.
- **6.** (a) Every connected graph G has a point whose removal does not disconnect the graph. What is the situation for strongly connected digraphs?
- (b) Let G be a connected graph which does not contain a "cherry", i.e. two points of degree 1 with a common neighbor. Show that one can remove two adjacent points without disconnecting G.
- 7. (a) Let  $T_1$ ,  $T_2$  be two spanning trees of a connected graph G. Prove that  $T_1$  can be transformed into  $T_2$  through a sequence of "intermediate" trees, each arising from the previous one by removing an edge and adding another.
- (b) Suppose G is 2-connected. Then to obtain  $T_1$  from  $T_2$  it suffices repeatedly to apply the following transformation: we remove the edge adjacent to some endpoint x of the tree and connect x to the rest by some other edge of G.
- 8. (a) Let G be a 2-connected graph on n points and  $n_1 + n_2 = n$ . Then V(G) has a 2-coloration  $\{A_1, A_2\}$  such that  $|A_i| = n_i$  and  $A_1, A_2$  induce connected subgraphs.
- (b) Let G be a 2-connected non-bipartite graph on 2m points. Then V(G) has a partition  $\{A_1, A_2\}$  with  $|A_i| = m$  such that the  $(A_1, A_2)$ -edges form a connected spanning subgraph.
- **9.** A digraph G is strongly connected iff there is at least one edge leaving each set  $X \subset V(G)$ ,  $X \neq \emptyset$ .
- 10. Let G be a digraph,  $x, y \in V(G)$  and suppose the edges of G are colored red, green and black. Then exactly one of the following two assertions holds:

- (i) There is a black and red undirected (x, y)-path P in G such that all black edges of P are oriented from x to y,
- (ii) There is a set  $S \subset V(G)$ ,  $x \in S$ ,  $y \notin S$ , such that no red edge connects S and V(G)-S in any direction and no black edge goes from S to V(G)-S.
- 11. (a) Suppose we can disconnect the strongly connected digraph G by removing  $\leq k$  edges. Prove that we can also disconnect it by inverting  $\leq k$  edges.
- (b) Suppose G is a digraph without isthmuses and we can get a strongly connected digraph from G by contracting at most k edges. Then we can also get a strongly connected graph by inverting at most k edges.
- (c) If we can destroy all cycles of the digraph G having no loops by removing at most k edges, then we can also destroy them by inverting at most k edges.
- 12. A tournament is strongly connected if and only if it contains a Hamiltonian cycle.
- 13. A strongly connected tournament T on  $n \ge 4$  points contains at least two points x such that T x is strongly connected.
- 14. Let G be a directed tree and  $F \subseteq E(G)$ . Prove that there is a point  $x \in V(G)$  such that the edges of F incident with x have their head at x and the other edges incident with x have their tail at x.
- 15. Let G be a tree and let

$$\varphi:V(G)\to V(G)$$

be a mapping such that, whenever  $(x, y) \in E(G)$ , then

$$\varphi(x) = \varphi(y)$$
 or  $(\varphi(x), \varphi(y)) \in E(G)$ .

Prove that  $\varphi$  has a fixed point or a fixed edge.

- 16. If the intersection of a set of subtrees of a tree G is non-empty, then it is a subtree.
- 17. (a) If G is a connected graph, then any two paths of maximum length have a point in common.
  - (b) If G is a tree, then all maximum paths of G have a point in common.
- 18. If  $G_1, \ldots, G_k$  are subtrees of the tree G such that any two intersect, then they have a point in common. (This yields a new proof of part (b) of the preceding problem.)
- 19. Prove that, in a connected graph G, the distance d(x, y) of the points x, y as well as the maximum length D(x, y) of paths connecting x and y are metrics. That is
  - (a)  $d(x, y) \ge 0$ , with equality iff x = y,
  - (b) d(x, y) = d(y, x),
  - (c)  $d(x, y) + d(y, z) \ge d(x, z)$ ,

and similarly for D(x, y).

**20.** If G is a tree and  $p_1, \ldots, p_n, q_1, \ldots, q_{n+1}$  are points of it, then

$$\sum_{1 \le i < j \le n} d(p_i, p_j) + \sum_{1 \le i < j \le n+1} d(q_i, q_j) \le \sum_{i=1}^n \sum_{j=1}^{n+1} d(p_i, q_j).$$

- 21. (a) Let  $\tilde{d}(x) = \max_{y} d(x, y)$ , where x, y are points of a tree. Show that the minimum of  $\tilde{d}(x)$  is attained at one point (the *center* of G) or at two adjacent points (the *bicenter* of G).
- (b) Show further that  $\tilde{d}(x)$  is a convex function in the sense that, if y, z are neighbors of x, then

$$2\tilde{d}(x) \leq \tilde{d}(y) + \tilde{d}(z).$$

**22.** (a) Let  $s(x) = \sum_{y} d(x, y)$ , where x, y are points of a tree G. Show that s(x) is *strictly convex* in the sense that, whenever x is a point and y, z are two neighbors of it, then

$$2s(x) < s(y) + s(z).$$

- (b) Show that s(x) assumes its minimum at one point (the baricenter) or at two adjacent points of G.
- (c) Construct a tree which has a center and a baricenter whose distance apart is greater than 1000.
- **23.** Determine those trees G on n points for which  $\sum_{x,y} d(x,y)$  is maximal and, respectively, minimal.
- **24.** In a tree on n points of diameter at least 2k-3, there are at least n-k paths of length k.
- 25. Given n cities, we want to build a connected telephone network between them. The expense v(e) of a line e between any two cities is known, and we want to minimize the total expense. Thus, we want to find a tree on the given points such that the sum of the expenses of edges is minimal. Show that the following algorithm produces the desired result: At the i<sup>th</sup> step, choose a component  $G_i$  of the graph formed by the edges already selected, and select an edge connecting  $G_i$  to a point not in  $G_i$  with minimum expense. Stop when  $G_i$  contains every point.
- 26. (Cont'd) Show that if the edges have distinct expenses, then the optimal tree is unique.
- 27. Call two edges of a 2-edge-connected graph *G equivalent* if they are equal or their removal disconnects the graph. Show that
  - (a) this is an equivalence-relation,
  - (b) all edges of an equivalence-class lie on a circuit,
- (c) removing the edges of an equivalence-class P, the components of the remaining graph are 2-edge-connected,
  - (d) contracting the components of G P, we obtain a circuit.

28. Given a 2-edge-connected graph G we can "build it up" as follows: G = $=G_1 \cup \ldots \cup G_r$ , where  $G_1$  is a circuit and  $G_{i+1}$  is either a path which has exactly its endpoints in common with  $G_1 \cup \ldots \cup G_i$  or a circuit which has one point in common with  $G_1 \cup \ldots \cup G_i$  (Fig. 3).




Fig. 3

29. A graph G can be oriented so that the resulting digraph G is strongly connected iff G is 2-edge-connected.

**30\***. We say that an edge is well fitted with the circuit C if it lies on C or has no point in common with C.

Let  $e_1, \ldots, e_k$  be independent edges of a 2-edge-connected graph G. Show that there is a circuit C into which all the edges  $e_1, \ldots, e_k$  are well fitted.

31. Show that "edges e, f are equal or lie on a circuit" is an equivalence relation.

**32.** The following properties of a graph G are equivalent:

- (i) G is 2-connected,
- (ii) any two points of G lie on a circuit,
- (iii) any two edges of G lie on a circuit, and G has no isolated points.

**33.** Let G be a 2-connected graph, which is not a circuit. Then there is a path P in G such that the inner points of the path P (if any) are of degree 2, and removing the edges and inner points of P the remaining graph is 2-connected.

**34.** Let p, q be two points of a 2-connected graph G. Show that G can be oriented in such a way that each edge is contained in a (p, q)-path.

**35.** Let G be a 2-connected graph. Then the following properties are equivalent:

- (i) G is critically 2-connected,
- (ii) no circuit of G has a chord.

 $36^*$ . Let G be a critically 2-connected graph. Show that each circuit C of G has a point of degree 2.

37\*. Let G be a critically 2-connected graph. Remove all points of degree 2 from G. Show that the resulting graph G' is a disconnected forest.

**38.** Construct a critically 2-connected graph G such that some  $x \in V(G)$ is at distance at least 1000 from every point of degree 2.

- **39\***. (Menger's Theorem). Let G be a digraph and  $a, b \in V(G)$ . Prove that
- (a) there are k edge-disjoint (a, b)-paths iff G is k-edge-connected between a and b;
  - (b) there are k independent (a, b)-paths iff G is k-connected between a and b;
  - (c) analogous statements hold for undirected graphs.
- **40.** Let A, B be disjoint subsets of V(G) and assume that any set X which has a point from each (A, B)-path has at least k elements. Show that there are k vertex-disjoint (A, B)-paths.
- 41. Formulate and prove a common generalization of the undirected vertex-connectivity version of Menger's theorem 6.39b and 6.40.
- **42\***. Let  $B \subset V(G)$  and  $a \in V(G) B$ ; suppose we are given k independent (a, B)-paths  $P_1, \ldots, P_k$  and k + r independent (a, B)-paths  $Q_1, \ldots, Q_{k+r}$ . Show that there are  $1 \leq j_1 < \ldots < j_r \leq k + r$  and (a, B)-paths  $R_1, \ldots, R_k$ ,  $R_i$  ending in the same points as  $P_i$ , such that  $R_1, \ldots, R_k, Q_j, \ldots, Q_j$  are independent.
- **43.** Let a, b, c be distinct points of G. Assume there are k independent (a, b)-paths  $P_1, \ldots, P_k$  and a (b, c)-path  $P_0$  independent of them. Also, there are k (other) independent (a, b)-paths  $Q_1, \ldots, Q_k$  and an (a, c)-path  $Q_0$  independent of them. Show that there are k+1 independent (c, b)-paths in G. (Try to avoid the use of Menger's theorem.)
- **44.** Without using Menger's theorem, describe a graph G having two non-adjacent points a, b such that
  - (i) there are at most k independent (a, b)-paths in G, and
- (ii) if any new edge is added to G, there will be at least k+1 independent (a, b)-paths in the resulting graph.

Deduce the undirected independent-path version of Menger's theorem from this.

- **45.** Let G be a digraph and  $a, b \in V(G)$ . Decide whether the following statements are true:
- (i) If any two paths which connect one of a and b to the other have a common edge, then there is an edge which is contained in all of these paths.
- (ii) If G is k-edge-connected between a and b, then it is k-edge-connected between b and a.
- (iii) If G is k-edge-connected between a and b and also between b and a, then there are k (a, b)-paths  $P_1, \ldots, P_k$  and k (b, a)-paths  $Q_1, \ldots, Q_k$  which are mutually edge-disjoint.
- **46.** Assume that each point  $x \neq a$ , b of G has the same indegree and outdegree and that  $d_G^+(a) d_G^-(a) = k > 0$ . Prove that there are k edge-disjoint (a, b)-paths in G.
- 47. If the outdegree of each point of G is equal to its indegree, then (i), (ii) and (iii) of 6.45 are true.

- **48.** Let G be a graph and X, Y,  $Z \subseteq V(G)$ . Establish the inequalities:
  - (a)  $\delta_G(X \cup Y) + \delta_G(X \cap Y) \leq \delta_G(X) + \delta_G(Y)$ ;
  - (b)  $\delta_G(X Y) + \delta_G(Y X) \leq \delta_G(X) + \delta_G(Y)$ ;
- (c)  $\delta_G(X-Y-Z) + \delta_G(Y-Z-X) + \delta_G(Z-X-Y) + \delta_G(X\cap Y\cap Z) \le \delta_G(X) + \delta_G(Y) + \delta_G(Z)$ .
- **49\***. Prove that a critically k-edge-connected graph has a point of degree k.
- **50.** Let G be a k-edge-connected graph and let  $F_1, \ldots, F_m$  be k-element cutsets of edges. Prove that  $G = (F_1 \cup \ldots \cup F_m)$  has at most 2m connected components.
- **51\***. Let G be an Eulerian graph,  $x \in V(G)$  and suppose G is k-edge-connected between any two points  $u, v \neq x$ . Then we can find two neighbors y, z of x such that, if we remove two edges (x, y) and (x, z) but join y to z by a new edge, the resulting graph is still k-edge-connected between any two points  $u, v \neq x$ .
- **52\***. Let G be a 2k-edge-connected, 2k-regular graph. Prove that G can be obtained by the following construction:
  - I. We start with two points joined by 2k edges.
- II. Having constructed a graph, we select k edges, subdivide them by one point each, and identify the new points.
- **53\*.** Prove that, provided  $k \geq 2$  the assertion of 6.51 holds for non-Eulerian graphs as well.
- **54.** (a) Let G be a graph and  $\vec{G}$  an orientation of G. If  $\vec{G}$  is k-edge-connected, then G is 2k-edge-connected.
- (b)\* Conversely, if G is 2k-edge-connected, then it has an orientation  $\vec{G}$  which is k-edge-connected (generalization of 6.29).
- **55.** Let  $\alpha$ ,  $\beta$  be integers, G a graph and  $\alpha$ ,  $\alpha'$ , b, b' four points in G. We want to find  $\alpha$   $(a, \alpha')$ -paths  $P_1, \ldots, P_{\alpha}$  and  $\beta$  (b, b')-paths  $Q_1, \ldots, Q_{\beta}$  such that  $P_1, \ldots, P_{\alpha}, Q_1, \ldots, Q_{\beta}$  are edge-disjoint. Show that the following conditions are necessary: (\*) We need to remove at least
  - $\alpha$  edges to separate a from a',
  - $\beta$  edges to separate b from b', and
  - $\alpha + \beta$  edges to separate  $\{a, b\}$  from  $\{a', b'\}$  or  $\{a, b'\}$  from  $\{a', l\}$ .

Show by an example that these conditions are not always sufficient.

**56\***. Let a, a', b, b' be points of an Eulerian graph G, and  $\alpha$ ,  $\beta$  positive even integers. Assume that condition (\*) formulated in the previous problem is satisfied.

<sup>†</sup> If x has only one neighbor, y and z may be identical; but this case is trivial.

(a) Show that G has an orientation  $\vec{G}$  such that

$$\begin{split} d^{+}_{\vec{G}}(a') - d^{-}_{\vec{G}}(a') &= -\alpha, \quad d^{+}_{\vec{G}}(b') - d^{-}_{\vec{G}}(b') = -\beta, \\ d^{+}_{\vec{G}}(a) - d^{-}_{\vec{G}}(a) &= \alpha, \qquad d^{+}_{\vec{G}}(b) - d^{-}_{\vec{G}}(b) = \beta, \\ d^{+}_{\vec{G}}(x) &= d^{-}_{\vec{G}}(x) \quad \text{for} \quad x \in V(G) - \{a, b, a', b'\}. \end{split}$$

- (b) Prove that G contains  $\alpha$  (a, a')-paths and  $\beta$  (b, b')-paths, all edge-disjoint.
- 57. Construct a 5-connected graph G and 4 points a, b, c, d of it such that each (a, b)-path has a point in common with each (c, d)-path.
- 58. Let A, B be k-element sets of points of a k-connected graph G, separating a and b. Show that the set C of those points in  $A \cup B$  which are the endpoints of  $(a, A \cup B)$ -paths is a k-element set separating a and b.
- **59.** Let G be a critically k-connected graph and H a k-connected subgraph of it. Show that H is critically k-connected.
- **60\***. (a) Let G be a k-connected graph, A a k-element separating set of it,  $G_1$  a component of G-A and assume A is chosen so that  $|V(G_1)|$  is minimal. Show that for any k-element separating set B, either  $V(G_1) \subseteq B$  or  $V(G_1) \cap B = G$ . Managery,  $|V(G_1)| \subseteq k$  in the former sees.
- $= \emptyset$ . Moreover,  $|V(G_1)| \le \frac{k}{2}$  in the former case.
  - (b) Every critically k-connected graph G has a point of degree k.
- **61\***. Let G be a simple 3-connected graph and assume that the endpoints of the edge e are both of degree at least 4. Then one of G/e, G-e is 3-connected.
- **62.** Let G be a critically 3-connected graph, and e an edge of it which connects two points of degree at least 4. Then G/e is a critically 3-connected graph.
- **63.** If G is a critically 3-connected graph, then every circuit of G contains at least two points of degree 3.
- **64.** Suppose that a 3-connected graph G is such that for every edge e, G-e and G/e are not 3-connected. Show that  $G \cong K_4$ .
- **65.** Construct a critically 3-connected graph and a point x in it such that every point at distance at most 1000 from x has degree at least 1000.
- **66.** In a k-connected graph G, any k points are on a circuit.
- **67\*.** Let  $e_1$ ,  $e_2$ ,  $e_3$  be independent edges of a 3-connected graph G. Then there is a circuit containing  $e_1$ ,  $e_2$ ,  $e_3$  except when  $e_1$ ,  $e_2$ ,  $e_3$  form a separating system.
- **68.** Let  $a, b, x_1, \ldots, x_k$  be distinct points of a (k+1)-connected graph G. Prove that there is an (a, b)-path which contains  $x_1, \ldots, x_k$ .

- **69.** Show that a circuit C in a simple planar 3-connected map has exactly one bridge iff it is the boundary of a face. Consequently, 3-connected simple planar graphs have an essentially unique embedding in the plane.
- 70\*. Every 3-connected graph G has a circuit C with exactly one bridge.

\*

**71.** We associate a value  $v(e) \ge 0$  with each edge e of a strongly connected digraph G. Let a, b be two points of G and let P and C run over all (a, b)-paths and (a, b)-cuts, respectively. Prove that

$$\max_{P} \min_{e \in P} v(e) = \min_{C} \max_{e \in C} v(e).$$

**72.** Let  $\varphi$  denote a function on V(G) satisfying (\*)  $\varphi(a) = 0$ ,  $\varphi(y) - \varphi(x) \le v(x, y)$  ( $(x, y) \in E(G)$ ). Now define the value u(P) of an (a, b)-path P to be  $\sum_{e \in E(P)} v(e)$ . Then

$$\min_{P} \ u(P) = \max_{\varphi} \ \varphi(b).$$

73. Let f be an (a, b)-flow and C an (a, b)-cut. Show that

$$w(f) = \sum_{e \in C} f(e) - \sum_{e \in C^*} f(e).$$

74. (Max-Flow-Min-Cut Theorem) Show that the maximum value of an (a, b)-flow f satisfying  $f(e) \leq v(e)$  is given by

$$\min_{C} \sum_{e \in C} v(e),$$

where C runs over all (a, b)-cuts.

**75\***. The preceding solution suggests the following algorithm to construct a maximum (a, b)-flow.

Let  $f_k$  be an (a, b)-flow and  $P_k$  an (undirected) (a, b)-path. Let A, B be the sets of edges of  $P_k$  directed toward a and b, respectively. Suppose  $P_k$  is such that f(e) > 0 for  $e \in A$  and f(e) < v(e) for  $e \in B$  (if no such path  $P_k$  exists we know  $f_k$  is optimal). Set

$$\varepsilon = \min_{e \in A, e' \in B} (f_k(e), v(e') - f_k(e')).$$

Increase the value  $f_k(e)$  by  $\varepsilon$  for  $e \in B$  and decrease it by  $\varepsilon$  for  $e \in A$ , to get a new flow  $f_{k+1}$ . Prove the following assertions:

- (a) Even repeating the above procedure infinitely many times, the values of the flows obtained do not necessarily converge to the values of a maximum flow.
- (b) If  $P_k$  is always chosen to be a path with minimum possible length then, repeating the above procedure, the length of  $P_k$  does not decrease.
- (c) If  $P_k$  is always a path with minimum length, we get an optimum flow in at most  $n^3$  steps (n = |V(G)|).

**76.** Suppose there is a value u(x) prescribed for each point x, and a value v(e) prescribed for each edge e. When does a function f(e) exist such that

$$0 \le f(e) \le v(e); \sum_{e=(a, y)} f(e) - \sum_{e=(y, a)} f(e) = u(a)$$

for every a?

77. (a) Assume the capacities v(e) are integers. Show that there is a maximum value (a, b)-flow with integral entries.

(b) Show that the Max-Flow-Min-Cut theorem is a consequence of the

directed-edge-version of Menger's theorem 6.39.

78. Let f be an integral (a, b)-flow of value  $w(f) = w_1 + \ldots + w_k$  ( $w_i = 1, 2, \ldots$ ). Show that f is the sum of k integral (a, b)-flows  $f_1, \ldots, f_k$ , of values  $w_1, \ldots, w_k$ , respectively.

#### § 7. Factors of graphs

Here we consider the general question: given a graph, does there exist a subgraph whose degrees meet certain prescribed conditions? The characteristic example of such a question is the problem of the existence of 1-factors (perfect matchings), the solution of which (the König-Hall theorem for bipartite graphs and Tutte's theorem for the general case) is an outstanding result making this probably the most developed field of graph theory.

In addition to the question of their existence, the problem of describing the structure of all factors is also of importance. We touch only briefly on this

problem (cf. also § 4).

One question which can be handled with the help of factorization results is whether a given sequence of integers can be the degree sequence of a graph (digraph, simple graph). This leads in fact to factorization problems for the complete graph and the answers are, correspondingly, much simpler.

Factorization problems constitute a class of integer programming problems for which there is a satisfactory solution. The basic idea of the correspondence between these two fields will be given in § 13, for the general case of hypergraphs. However, we shall not be able to go into the details in this volume.

- 1. Let G be a graph without isolated points. Then  $\nu(G) + \varrho(G) = |V(G)|$ .
- **2.** (König's Theorem) Let G be a bipartite graph. Then  $\nu(G) = \tau(G)$ , and  $\varrho(G) = \alpha(G)$ .
- **3.** (Hungarian Method) Let G be a bipartite graph with 2-coloration  $\{A, B\}$  and let M be a matching of G. Let  $A_1$  and  $B_1$  be the sets of points in A and B not covered by M. Form a maximal forest  $F \subseteq G$  with the following properties:
- (\*) each point x of F in B has degree 2 and one of the edges adjacent to x belongs to M;
  - (\*\*) each component of F contains a point of  $A_1$ .

Prove that M is a maximum matching iff no point of  $B_1$  is adjacent to any point of F. Deduce König's theorem 7.2 from this. Derive an algorithm to find a maximum matching in a bipartite graph from this result.

- **4.** (a) Let G be a bipartite graph with bipartition  $\{A, B\}$ . Suppose that every  $X \subseteq A$  is adjacent to at least |X| points of B. Prove that G has a matching which matches all the points of A with (certain) points of B. Find different proofs which use and respectively, do not use König's theorem.
  - (b) When does a bipartite graph G have a 1-factor?
- 5. Let G be a bipartite graph with bicoloration  $\{A, B\}$  and

$$\delta = \max_{X \subseteq A} \{ |X| - |\Gamma_G(X)| \}.$$

Prove that

$$v(G) = |A| - \delta.$$

**6.** (a) Suppose G is a bipartite graph with bicoloration  $\{A, B\}$  and let  $k \geq 0$  be a given integer such that

$$|\Gamma_G(X)| \ge |X| + k$$

holds for each  $X \subseteq A$ ,  $X \neq \emptyset$ . Let  $X_1$ ,  $X_2$  be two sets that satisfy (\*) with equality and suppose  $X_1 \cap X_2 \neq \emptyset$ . Show that  $X_1 \cap X_2$  also gives equality in (\*).

- (b) Show that G as defined in part (a) has a spanning subgraph  $G_1$  such that
  - (1)  $d_{G_i}(x) = k + 1$  for all  $x \in A$ ,
  - (2)  $|\Gamma_{G_1}(X)| \ge |X| + k$  for all  $X \subseteq A$ ,  $X \ne \emptyset$ .
- 7. Show that for any bipartite graph G with bicoloration  $\{A, B\}$ , the following three statements are equivalent:
  - (i) G is connected and each edge of G is contained in a 1-factor,
  - (ii) For any  $x \in A$  and  $y \in B$ , G x y has a 1-factor,
  - (iii) |A| = |B| and for each  $\emptyset \neq X \subset A$ ,  $|\Gamma(X)| > |X|$ .

(Such a graph is called an elementary bipartite graph.)

8. Prove that a bipartite graph G is elementary if and only if it can be written in the form

$$G = G_0 \cup P_1 \cup \ldots \cup P_k,$$

where  $G_0$  consists of two points and an edge joining them and  $P_i$  is an odd path which joins two points of  $G_0 \cup P_1 \cup \ldots \cup P_{i-1}$  in different color classes and has no other point in common with  $G_0 \cup P_1 \cup \ldots \cup P_{i-1}$ .

- **9.** Let G be an elementary bipartite graph which will not remain elementary if we remove any edge of it. Show that it has a point of degree 2. Does every edge of it have an endpoint of degree 2?
- 10. A bipartite graph G with maximum degree r = d(G) is the union of r matchings (i.e. has chromatic index r).

<sup>†</sup> In general, the chromatic index of a graph G is bounded by  $\left[\frac{3}{2}d(G)\right]$  [Shannon] and by d(G) + 1, if G is simple [Vizing], [see B].

<sup>4</sup> Lovász

11. Let G be any bipartite graph and suppose  $k \geq 1$ . Then G is the union of k edge-disjoint spanning subgraphs  $G_1, \ldots, G_k$  such that

$$\left\lceil rac{d(x)}{k} 
ight
ceil \leq d_{G_i}(x) \leq \left\lceil rac{d(x)}{k} 
ight
ceil^* \qquad ext{for each } x \in V(G).$$

- 12. Let G be a bipartite graph with minimum degree r. Then G is the union of r disjoint edge-covers.
- 13. Determine the least number r = r(n) with the property that every r-regular bipartite graph G with 2n points has a 1-factor such that each edge of this 1-factor has a parallel in G.
- 14. Let G be an r-regular bipartite graph  $(r \ge 2)$  with 2-coloration  $\{A, B\}$ , where  $A = \{a_1, \ldots, a_n\}$  and  $B = \{b_1, \ldots, b_n\}$ . Suppose a point x to be moving in V(G) and transforming the graph as follows:
- (1) x starts at  $a_1$  and moves along any  $(a_1, b_i)$  to  $b_i$  at which time this edge is removed.
- (2) If x is at  $b_i$ , which it has just entered from  $a_{\mu}$ , then it moves to the *first* point  $a_{\nu}$  such that  $a_{\mu} \neq a_{\nu}$  and  $a_{\nu}$  is adjacent to  $b_i$ , and the edge  $(a_{\nu}, b_i)$  is doubled. (Note that G may have multiple edges and thus,  $a_{\mu}$  may still be adjacent to  $b_i$  though one  $(a_{\mu}, b_i)$ -edge is removed.)
- (3) If x is at  $a_{\nu}$  which it entered from  $b_i$ , then it moves over to the first neighbor  $b_j \neq b_i$  of  $a_{\nu}$  and an  $(a_{\nu}, b_j)$ -edge is removed.

Prove that, in a finite time, the procedure terminates with the situation that  $x = a_1$  and  $a_1$  is adjacent to some  $b_i$  by r parallel edges (and to no other point). Use this procedure to get a 1-factor.

- 15. (a) Suppose the simple bipartite graph G with bipartition  $\{A, B\}$  has a 1-factor and every  $x \in A$  has degree at least k. Show that G has at least k! 1-factors.
- (b) An elementary bipartite graph with n points and m edges has at least m-n+2 1-factors.
- 16. Let G be a bipartite graph with bipartition  $\{A, B\}$  and  $f(x) \geq 0$  an integer valued function on V(G). Show that G has an f-factor if and only if

(i) 
$$\sum_{x \in A} f(x) = \sum_{y \in B} f(y)$$

and

(ii) 
$$\sum_{x \in X} f(x) \le m(X, Y) + \sum_{y \in B-Y} f(y)$$

for all  $X \subseteq A$ , and  $Y \subseteq B$ , where m(X, Y) is the number of edges connecting X to Y.

17. Let G be a bipartite graph with n points and m edges and maximum degree d. Show that G can be embedded as an induced subgraph in a d-regular bipartite graph with  $2n-2\left[\frac{m}{d}\right]$  points, but no less.

- 18. Let G be a simple bipartite graph with bipartition  $\{A, B\}$  such that |A| = |B| = n, and with maximum degree  $d < \frac{n}{2}$ . Show that G can be embedded in a *simple* regular bipartite graph on the same set V(G) of points with degree 2d.
- 19. Let  $A = (a_{ij})$  be a non-negative  $n \times n$  matrix such that all row- and column-sums of A are 1 (such a matrix is called *doubly stochastic*). Prove that per A > 0, i.e. the determinant of A has a non-zero expansion term.
- **20.** (a) Let G be a simple bipartite graph with 2-coloration  $\{A, B\}$ ,  $A = \{a_1, \ldots, a_n\}$  and  $B = \{b_1, \ldots, b_n\}$ . Let  $a_{ij} \neq 0$  iff  $(a_i, b_j) \in E(G)$  and assume the non-zero  $a_{ij}$ 's are algebraically independent transcendentals. Prove that G has a 1-factor iff det  $(a_{ij}) \neq 0$ .
- (b) Prove the König-Hall criterion (7.4a) for the existence of a 1-factor in a bipartite graph from this observation.
- **21.** Let G be a connected graph with n points and  $E(G) = \{e_1, \ldots, e_m\}$ . Consider all vectors  $(y_1, \ldots, y_m)$  such that

$$\sum_{e_i \ni v} y_i = 0$$

for each point v. What is the dimension of the space spanned by them?

- **22.** Let G be a simple graph on 2n points with all degrees at least n. Show that G has a 1-factor.
- **23.** Let  $F_0$  be any matching of G. Then G has a maximum matching which covers all points covered by  $F_0$ .
- 24. (a) Construct a graph with a unique 1-factor having all its degrees at least k.
  - (b) Prove that a graph with a unique 1-factor has a cutting edge.
  - (c) If a simple graph G on 2n points has a unique 1-factor, then

$$|E(G)| \leq n^2.$$

25. If G is a graph without isolated points and with maximum degree d, then

$$u(G) \ge rac{|V(G)|}{d+1}.$$

- **26\***. Let G be a connected graph such that, for each  $x \in V(G)$ , v(G x) = v(G). Show that G is factor-critical.
- 27\*. (a). Show that for any graph G,

$$\delta(G) \stackrel{\mathrm{def}}{=} |V(G)| - 2\nu(G) = \max_{X \subseteq V(G)} \left\{ c_1(G - X) - |X| \right\}.$$

- (b) (Tutte's Theorem) A graph has a 1-factor if and only if  $c_1(G X) \le |X|$  for every  $X \subseteq V(G)$ .
- **28.** (a) Let G be a simple graph such that
  - (i) G has no 1-factor, but

(ii) joining any two non-adjacent points of G by an edge results in a graph with a 1-factor.

Show (without using Tutte's theorem) that G has the following structure: it has a set  $V_1$  of points joined to every other point and the remaining points span disjoint complete graphs.

- (b) Give a new proof of Tutte's theorem 7.27b based on (a).
- 29. (a) (Petersen's Theorem) A 2-connected 3-regular graph has a 1-factor.
  - (b) Construct a simple 3-regular graph having no 1-factor.
- **30.** Let G be a (k-1)-edge-connected, k-regular graph with an even number of points. Remove k-1 edges of G. Show that the remaining graph G' has a 1-factor.
- 31. Let G be a factor-critical graph. Show that it can be expressed in the form

$$G = P_0 \cup P_1 \cup \ldots \cup P_k$$

where  $P_0$  is an odd circuit and  $P_{i+1}$  is either an odd path with both endpoints in  $P_0 \cup P_1 \cup \ldots \cup P_i$  but no inner point in  $P_0 \cup P_1 \cup \ldots \cup P_i$  or an odd circuit having exactly one point in common with  $P_0 \cup P_1 \cup \ldots \cup P_i$   $(i = 0, \ldots, k-1)$ .

- 32\*. (The Gallai–Edmonds Structure Theorem) Let G be a graph. Denote by  $D_G$  the set of those points which are not covered by at least one maximum matching. Let  $A_G$  be the set of neighbors of  $D_G$  and  $C_G = V(G) A_G D_G$ . Prove:
  - (a) Removing a point x of  $A_G$ , the sets  $D_G$  and  $C_G$  do not change.
- (b) Every component of the graph induced by  $D_G$  is factor-critical. The graph induced by  $C_G$  has a 1-factor.
- (c) If M is any maximum matching, it contains a maximum matching of each component of the graph induced by  $D_G$  and a 1-factor of the graph induced by  $C_G$ .

(d) 
$$v(G) = \frac{1}{2} \{ |V(G)| - c(D_G) + |A_G| \}.$$

- (e) Tutte's theorem follows from this.
- (f) Determine the sets A, C, D for the graph shown in Fig. 4.





Fig. 4

- **33.** Let M be a matching of G and C a circuit of length 2k + 1 which contains k edges of M, but meets no other edge in M. Let G' be the graph obtained by contracting C and M' = M E(C). Then M is a maximum matching of G iff M' is a maximum matching of G'.
- **34\***. (Edmonds' Matching Algorithm) (a) Let G be a graph, M be a matching of G and consider a maximal forest F with the following properties:
- (\*) Each component of F contains exactly one point not covered by M, called a root.
- (\*\*) Calling those points of F being at an odd distance from a root inner points and the other points, including the roots) outer points, each inner point has degree 2 and one of the two edges incident with it belongs to M (Fig. 5).



Then, if two outer points in distinct components are adjacent, M is not a maximum matching. Design an algorithm to find maximum matchings based on this observation and the preceding problem.

- (b) Test your algorithm on the graph in Fig. 4.
- **35.** Assume that the k-regular graph G on 2m points has the property that any two odd circuits of it either intersect or are joined by an edge. Show that G has a 1-factor.
- **36.** (a) If G has no 1-factor, then it has a point x such that each edge adjacent to x is contained in some maximum matching.
- (b)\* If G is a 2-connected graph with a 1-factor then it has a point such that each edge adjacent to this point occurs in some 1-factor. (Such a point is often said to be totally covered.)

**37.** The maximum size of a 2-matching of G is

$$|V(G)| = \max_{\substack{X \subseteq V(G) \\ X \text{ independent}}} \{|X| - |\Gamma_G(X)|\}.$$

- 38\*. Every graph G has a maximum matching which is contained in a maximum 2-matching.
- **39.** Let G be a connected 2d-regular graph with an even number of edges. Prove that G has a d-factor.
- 40. A 2d-regular graph is the union of d 2-factors.
- 41. Which connected graphs G have a spanning subgraph F with

$$d_F(x) = \left\lceil \frac{d(x)}{2} \right\rceil \text{ or } \left\lceil \frac{d(x)}{2} \right\rceil^*$$

for each x?

- **42.** (a) Which connected graphs have a spanning subgraph with all degrees odd?
- (b) Every 2-edge-connected graph G with all degrees at least 3 has a spanning subgraph in which each degree is positive and even.
- **43\***. Let G be a graph and let us associate a non-negative integer f(x) with each point of G. Construct a graph G' which has a 1-factor if and only if
  - (a) G has a perfect f-matching,
  - (b) G has an f-factor.
- **44.** We associate a pair (f(x), g(x)) of non-negative integers with each point x of a digraph G. When does G have a spanning subgraph with indegree f(x) and outdegree g(x) at each point x?
- **45.** Given a graph G, when can one orient its edges in such a way that the outdegree of every point x is equal to a prescribed number f(x)?
- **46.** (a) Let G be a connected graph which contains a circuit. We associate a number g(x) with each point x of G. Show that G has a spanning subgraph F whose degree differs from g(x) at every point x.
- (b) Let G be a tree, and let us associate an integer g(x) with each point x. Show that exactly one of the following statements is true:
  - (i) G has an orientation with indegrees g(x);
  - (ii) G has a spanning subgraph with all degrees different from g(x).

**47.** Let  $0 < d_1 \le \ldots \le d_n$  be integers. Show that there exists a tree with degrees  $d_1, \ldots, d_n$  if and only if

$$d_1+\ldots+d_n=2n-2.$$

- **48.** Let  $0 \le d_1 \le \ldots \le d_n$  be integers. Show that they are the degrees of a graph without loops iff
  - (1)  $d_1 + \ldots + d_n$  is even, and
  - $(2) \ d_n \leq d_1 + \ldots + d_{n-1}.$
- **49.** Let  $f_1, \ldots, f_n, g_1, \ldots, g_n \ge 0$  be integers. Prove that there exists a simple digraph on  $\{v_1, \ldots, v_n\}$  without loops in which  $d_H^+(v_i) = f_i, d_H^-(v_i) = g_i$  iff
  - (a)  $f_1 + \ldots + f_n = g_1 + \ldots + g_n$ , and

(b) 
$$\sum_{i \in I} f_i \le \sum_{j \in J} g_j + |I| (n - |J|) - |I - J|$$
  $(I, J \subseteq \{1, \dots, n\}).$ 

Simplify the condition if  $f_1 \leq \ldots \leq f_n$  and  $g_1 \leq \ldots \leq g_n$ .

**50.** Prove that there exists a simple graph with degrees  $d_1 \leq d_2 \leq \ldots \leq d_n$  if and only if there exists one with degrees  $d'_1, \ldots, d'_{n-1}$ , where

$$d_k' = \begin{cases} d_k & \text{for } k = 1, \dots, n - d_n - 1, \\ d_k - 1 & \text{for } k = n - d_n, \dots, n - 1. \end{cases}$$

- **51.** (a) Suppose  $d_i \geq 0$   $(i = 1, \ldots, n)$  and  $d_1 + \ldots + d_n$  is even. Suppose there is a simple digraph H without loops on  $V = \{v_1, \ldots, v_n\}$  such that  $d_H^+(v_i) = d_H^-(v_i) = d_i$   $(i = 1, \ldots, n)$ . Then there exists a simple graph G on V with  $d_G(v_i) = d_i$   $(i = 1, \ldots, n)$ .
- (b) Let  $0 \le d_1 \le \ldots \le d_n$ . Prove that there exists a graph with degrees  $d_1, \ldots, d_n$  if and only if
  - (1)  $d_1 + \ldots + d_n$  is even, and
  - $(2) \sum_{i=n-k+1}^{n} d_i \le k(k-1) + \sum_{i=1}^{n-k} \min (d_i, k)$

for each  $1 \le k \le n$ .

- **52.** Let  $0 \le d_1 \le d_2 \le \ldots \le d_n$  be given integers. Show that they are the degrees of a simple connected graph if and only if
  - (1) they satisfy the conditions of 7.51b and moreover
  - (2)  $d_1 > 0$ , and

$$(3) \sum_{i=1}^{n} d_i \ge 2(n-1).$$

**53\***. Let  $d_1, \ldots, d_n$  be given integers. Prove that they are the degrees of a simple graph with a 1-factor if and only if  $d_1, \ldots, d_n$  as well as  $d_1 - 1, \ldots d_n - 1$  are the degrees of some simple graph.

# § 8. Independent sets of points

The concept of an independent set of points is analogous to, and sounds simpler than the concept of an independent set of edges. However, it is more difficult to handle and much less is known. There is no "good" way to determine the

maximum number of independent points; there are some fundamental reasons for this (see the work of Cook, Karp and Levin quoted in the introduction).

The problem of considering graphs critical with respect to independence number has proved easier to tackle. These graphs have many nice properties and, extending problem 8.25 it is even possible to find a classification of them. The quantity and depth of results concerning  $\alpha$ -critical graphs is considerably larger than those concerning chromatic-critical graphs (see the next section).

A related problem is that of the problem of games on graphs, which is a very general setting for game theory. The reader is advised to translate the games of chess, tic-tac-toe etc. into games on directed graphs as in problem 8.8, and design games to which the surprising results of these problems can be applied (e.g. chess with "page")

applied (e.g. chess with "pass").

1. A graph with all degrees at most d satisfies

$$\alpha(G) \ge \frac{|V(G)|}{d+1}.$$

- 2. The points of a graph G can be covered by not more than  $\alpha(G)$  vertex-disjoint paths.
- 3. The points of a graph G can be covered by not more than  $\alpha(G)$  disjoint circuits, edges and points.
- **4\*.** Let G be a digraph and S a subset of V(G) such that there are disjoint (directed) paths starting from the points of S and covering V(G). Show that S contains a subset  $S_0$  with the same property such that

$$|S_0| \leq \alpha(G)$$
:

- 5. (a) Every symmetric digraph has a kernel.
  - (b) An acyclic digraph G has a unique kernel.
  - (c) If every cycle of G has even length, then G has a kernel.
- **6.** Every tournament T has a point from which every other point can be reached on a (directed) path of length at most 2.
- 7. Every digraph G contains an independent set  $S \subseteq V(G)$  such that every point can be reached from S by a (directed) path of length at most 2.
- 8. Let G be a digraph and let players A, B play the following game. A occupies a point and then, taking turns, they each occupy a point which is accessible from the opponents last move and still unoccupied. A player loses whenever he has no possible move on his turn.
- (a) Suppose G has no cycles. Show that A has a winning strategy and determine the set of points he can start with.
- (b) Now drop the assumption that G is acyclic but assume it has a point  $x_0$  of indegree 0. Show that the opening player A still has a winning strategy.
- (c) Now let G be an arbitrary digraph. Show that the second player has a winning strategy if and only if he has a winning strategy on each strong component of G.

**9.** (Cont'd) Let G be a graph (or, equivalently, a symmetric digraph). Show that the opening player has a winning strategy if and only if G has no 1-factor.

10. Let  $T_1, \ldots, T_k$  be maximum independent sets in a graph G. Show that

$$|T_1 \cup \ldots \cup T_k| + |T_1 \cap \ldots \cap T_k| \geq 2\alpha(G)$$
.

11. Let  $T_1, \ldots, T_k$  be maximum independent sets of G and X any independent set. Let

$$S = X \cap T_1 \cap \ldots \cap T_k.$$

Then

$$|\Gamma(S)| - |S| \le |\Gamma(X)| - |X|.$$

- 12. Let G be an  $\alpha$ -critical graph without isolated points. Then every point x is contained in some maximum independent set, but not in all. If x, y are two points which do not constitute a component of G, then there is a maximum independent set containing exactly one of them. If x, y are adjacent then there is another maximum independent set missing both of them.
- 13. Substituting a complete graph for each point of an  $\alpha$ -critical graph, we get an  $\alpha$ -critical graph.
- 14. Every graph is an induced subgraph of an α-critical graph.
- 15. (a) Find infinitely many connected r-regular  $\alpha$ -critical graphs  $(r \geq 2)$ .
  - (b) Which Platonic bodies are α-critical?
- 16. Which bipartite graphs are α-critical?
- 17\*. Any two adjacent edges of an  $\alpha$ -critical graph are contained in a chordless odd circuit.
- 18. An  $\alpha$ -critical graph has no cutset, spanning a complete graph (in particular it has no cutpoint).
- 19\*. (a) Let  $G_1$ ,  $G_2$  be connected  $\alpha$ -critical graphs other than  $K_2$ . Split a point  $x \in V(G_1)$  into  $x_1$ ,  $x_2$  and remove an edge  $(y_1, y_2)$  of  $G_2$ . Identify  $x_i$  with  $y_i$ . Show that the resulting graph is  $\alpha$ -critical.
- (b) Show further that every connected but not 3-connected  $\alpha$ -critical graph arises in this way.
- **20.** An  $\alpha$ -critical graph G without isolated points has at least  $2\alpha(G)$  points.
- **21.** In an  $\alpha$ -critical graph  $|\Gamma(X)| \geq |X|$  for any independent set X.
- 22. Let X be an independent set of an  $\alpha$ -critical graph G and  $x \in X$ . Then

$$d_G(x) \le |\Gamma(X)| - |X| + 1.$$

- 23. The maximum degree of an  $\alpha$ -critical graph without isolated points is at most  $|V(G)| 2\alpha(G) + 1$ .
- **24\***. Which connected  $\alpha$ -critical graphs G are regular of degree  $|V(G)| 2\alpha(G) + 1$ ?

- **25.** Characterize the connected  $\alpha$ -critical graphs with  $|V(G)| 2\alpha(G) = 0, 1, 2$ .
- **26.** Assume that  $\alpha(G x y) = \alpha(G)$  for every  $x, y \in V(G)$  and  $|V(G)| = 2\alpha(G) + 1$ . Show that G is an odd circuit.
- **27.** Let G be a graph such that  $\alpha(G \{x, y, z\}) = \alpha(G)$  for all  $x, y, z \in V(G)$ . Show that either  $|V(G)| \ge 2\alpha(G) + 3$  or  $G \simeq K_4$ .

#### § 9. Chromatic number

The chromatic number is the most famous graphical invariant; its fame being mainly due to the Four Color Conjecture, which asserts that all planar graphs are 4-colorable. This has been the most challenging problem of combinatorics for over a century and has contributed more to the development of the field than any other single problem. Although today chromatic number attracts attention for several other reasons too, many of which arise from applied mathematical fields such as operations research, the Four Color Conjecture is still the main motivation for its investigation.

Despite its popularity, we know very little about chromatic number. It is difficult to compute it; there is no good characterization of k-chromatic graphs except for the trivial case k=2 (cf. the non-trivial but "non-good" characterizations given in problems 9.6, 9.9, 9.11, 9.15, 9.16); and chromatic critical graphs have much weaker properties than  $\alpha$ -critical graphs. The most difficult

proofs often concern counterexamples.

The collection of problems presented here is an attempt to cover the most important ideas developed so far in this area. None of these have lead to a satisfactory theory of chromatic number as yet (in particular, none of them has enabled us to settle the Four Color Conjecture),† but they may become elements of a more advanced theory. The main ideas dealt with in this section are; graphs with small degrees have small chromatic number; attempts to generalize the idea of "potential" used for bipartite graphs in § 5; the consideration of critical and saturated graphs; perfect graphs; chromatic polynomials, a promising algebraic approach; and Kempe chaining.

- 1. If every point of a graph G has degree at most k, then  $\chi(G) \leq k+1$ .
- 2. Prove that for any graph G, we can find a partition  $V(G) = V_1 \cup V_2 \cup (V_1, V_2 \neq \emptyset)$  such that

$$\chi(G[V_1]) + \chi(G[V_2]) = \chi(G).$$

Further, if G is not complete, we can find a partition  $V(G) = V_1 \cup V_2$  with

$$\chi(G[V_1]) + \chi(G[V_2]) > \chi(G).$$

- **3.** Establish the inequality  $\chi(G_1 \cup G_2) \leq \chi(G_1) \cdot \chi(G_2)$ .
- **4.** Suppose G has a (good) coloration in which each color occurs at least twice. Show that G has such a coloration with  $\chi(G)$  colors.

<sup>†</sup> Since I wrote these sentences Appel and Haken have given a proof of the Four Color Conjecture. They use different methods, among others computers very extensively.

**5.** (a) Suppose |V(G)| = n and V(G) has a partition  $\{V_1, \ldots, V_k\}$  such that, for each  $1 \le i < j \le k$ , there exists an  $x \in V_i$  and a  $y \in V_j$  which are non-adjacent. Then

 $\chi(G) \leq n - k + 1.$ 

(b) Let G be a simple graph on n points. Then

$$\chi(G) + \chi(\overline{G}) \leq n + 1, \ \chi(G) \cdot \chi(\overline{G}) \geq n.$$

**6.** Show that the chromatic number of a graph G is equal to the least number m such that

$$\alpha(G \oplus K_m) = |V(G)|.$$

- $(G \oplus K_m)$  is the Cartesian product of G and the complete m-graph).
- 7. (a)  $\chi(G_1 \times G_2) \leq \min (\chi(G_1), \chi(G_2))$ .
  - (b)  $\chi(G \times G) = \chi(G)$ .
  - (c)  $\chi(G \times K_n) = \min (\chi(G), n)$ .
- (d)\* Show that if G is connected and  $\chi(G) > n$  then  $G \times K_n$  has a unique n-coloration (up to permuting colors).
  - (e) How many n-colorations can a graph have?
- 8. Let S be a set of points. If  $S \subseteq V(G)$ , then any coloration of G induces a partition of S. Embed S into some graph G in such a way that the partitions of S induced by k-colorations of G ( $k \ge 3$ ) are
- (a) the partition of S into 1-element classes only (this makes sense, of course, only if  $|S| \leq k$ );
  - (b) all partitions except the previous one;
  - (c) the partition  $\{S\}$  only;
  - (d) all partitions except the previous one;
  - (e)\* a given set  $\{P_1, \ldots, P_N\}$  of partitions.

\*

- **9.** If a digraph G has no path of length m then  $\chi(G) \leq m$ .
- 10. When can we color the points of a digraph G by any number of colors  $1, 2, \ldots$  in such a way that every edge joins a point of color i to a point of color i+1  $(1 \le i)$ ?
- 11\*. Prove that G is k-colorable if and only if it can be oriented in such a way that for any circuit C of G and any given direction on C, at least  $\frac{|E(C)|}{k}$  edges of C are oriented in this direction.

\*

12. (a) Let G be a connected graph and suppose there is a set C(x) of colors associated with each point x of G. Assume  $|C(x)| \ge d_G(x)$  for every x and that strict inequality holds for at least one  $x_0$ . Then we can find a (good) coloration of G which uses one of the prescribed colors at each point.


- (b) Suppose again that a set C(x) of colors is associated with each point x of a graph G, but now let  $|C(x)| = d_G(x)$  for each point and  $C(a) \neq C(b)$  for some a, b. Further assume G to be 2-connected. Show that G admits a (good) coloration which uses an element of C(x) to color x for each point x. (Note the important special case when G is a circuit; cf. the solution of 9.8d).
- 13\*. (Brooks' Theorem) (a) Show that if every point of G has degree at most  $k \ge 3$  and G is 3-connected but is not a complete (k+1)-graph, then  $\chi(G) \le k$  (i.e. in this case the condition that C(x) is not the same at every point can be dropped).
- (b) Show that the condition of 3-connectivity can be replaced by connectivity.
- 14\*. (a) Let G be an infinite<sup>†</sup> graph all of whose finite subgraphs are k-colorable, but if we join any pair of non-adjacent points by a new edge in G, it will have a finite subgraph which is not k-colorable. Prove that V(G) has a partition such that two points are adjacent iff they belong to distinct classes.
- (b) (Erdős–de Bruijn Theorem) Suppose all finite subgraphs of an infinite graph G are k-colorable. Prove that G itself is k-colorable.
- (c) For each point  $v \in V(G)$ , let  $T_v$  be a copy of the discrete topological space on points  $1, \ldots, k$ . Then the colorations of V(G) with k colors can be considered as points of the topological product space  $\underset{v \in V(G)}{\times} T_v$ . Prove that the legitimate colorations of G form a closed subset. Give another proof of the Erdősde Bruijn theorem.
- 15. Let  $\Re \neq \emptyset$  be a class of simple graphs satisfying the following conditions:
  - (i) If  $G \in \mathbb{H}$  and G has a homomorphism into G', then  $G' \in \mathbb{H}$ .
- (ii) If G is a graph,  $a, b, c \in V(G)$  are such that a, c are adjacent but b is not adjacent to either of them, and if  $G + (a, b) \in \mathcal{H}$  and  $G + (b, c) \in \mathcal{H}$ , then  $G \in \mathcal{H}$ .

Prove that  $\mathcal{K}$  consists of all non-k-colorable graphs for some  $k \geq 0$ .

- 16\*. (Hajós' Construction) Consider the following operations on simple graphs:
  - (α) Addition of edges and/or points to the graph,
- $(\beta)$  Identification of two non-adjacent points (and cancellation of the resulting multiplicities of edges),
- $(\gamma)$  For two graphs  $G_1, G_2$  and  $(x_i, y_i) \in E(G_i)$ , removal of  $(x_i, y_i)$  (i = 1, 2), addition of a new edge  $(y_1, y_2)$ , and identification of  $x_1$  and  $x_2$ .

Prove that these operations produce non-k-colorable graphs from non-k-colorable ones and that every non-k-colorable graph arises by the repetition of these operations, from the initial graph  $K_{k+1}$  (see in Fig. 6 how the 5-wheel is obtained from  $K_4$ 's).

<sup>&</sup>lt;sup>†</sup> This exercise shows that problems concerning infinite graphs with finite chromatic number can usually be reduced to the finite case. This is why we include it here, cf. 14.19.



\*

- 17. (a) Which graphs are critically 3-chromatic?
- (b) Construct critically 4-chromatic graphs on 4n points in which the number of edges is at least  $n^2$ .
- (c) Construct critically 6-chromatic graphs on 2n points in which each point has degree at least n.
- 18. We associate a new point x' with each point x of a  $\chi$ -critical graph G, and join it to all neighbors of x in G. We take another new point y and join it to the points x' ( $x \in V(G)$ ). Show that the resulting graph G' is  $\chi$ -critical with  $\chi(G') = \chi(G) + 1$ .
- 19. (a) Is the graph in Fig. 7 an induced subgraph of some critically 4-chromatic graph?




Fig. 7

- (b) A graph  $G_0$  is a subgraph (or induced subgraph) of a critically (k+1)-chromatic graph G iff  $\chi(G_0/e) \leq k$  for every edge e.
- **20.** If we split a point of a critically (k+1)-chromatic graph G, the resulting graph G' is either k-chromatic or critically (k+1)-chromatic. For which values of k can the latter happen?
- **21.** A critically (k+1)-chromatic graph is at least k-edge-connected.
- 22. Every χ-critical graph is 2-connected. Which ones are not 3-connected?
- 23. Show that if G is a  $\chi$ -critical graph,  $\chi(G) = k + 1$  and S is a separating set of points of cardinality m, then the number of components of G S is not more than the number  $P_{m,k}$  of partitions of m objects into at most k classes. Also show that this is best possible.
- **24.** Let G be a critically (k+1)-chromatic graph. Prove that every pair of adjacent points can be connected by k-1 edge-disjoint chordless even paths.

- 25. (a) Construct a k-chromatic graph without triangles.
  - (b)\* Construct a k-chromatic graph without 3-, 4- and 5-circuits.
- **26.** Let G be a digraph and L(G) its line-graph. Prove that

$$\chi(L(G)) \ge \log_2 \chi(G).$$

- 27. Construct graphs having no odd circuits shorter than 2s + 1 and with chromatic number greater than  $k^{\dagger}$ .
- **28.** Let  $I_1, \ldots, I_n$  be closed intervals on a line. Define a graph G on a set  $\{x_1, \ldots, x_n\}$  by joining  $x_{\nu}$  to  $x_{\mu}$  iff  $I_{\nu} \cap I_{\mu} \neq \emptyset$ .
  - (a) Show that the resulting graph G has  $\chi(G) = \omega(G)$ .
  - (b) Show that the complement of this graph  $\overline{G}$  also satisfies  $\chi(\overline{G}) = \omega(\overline{G})$ .
  - (c) Show that every circuit C of G of length greater than 3 has a chord.
- **29.** (a) Let G be a graph such that any circuit of G of length greater than 3 has a chord. Show that every (inclusionwise) minimal cutset S of G induces a complete graph.
- (b) A graph G has the property that every circuit of length greater than 3 has a chord if and only if it can be represented as follows: Let  $F_1, \ldots, F_n$  be subtrees of a tree T; set  $V(G) = \{x_1, \ldots, x_n\}$  and join  $x_i$  to  $x_j$  if and only if  $F_i \cap F_j \neq \emptyset$ .
- (c) If G has no chordless circuits of length greater than 3, then G has  $\chi(G) = \omega(G)$  and the same holds for its complement.
- **30.** Let G be a bipartite graph. Then  $\chi(\overline{G}) = \omega(\overline{G})$ .
- **31.** If  $G_1$  and  $G_2$  both satisfy  $\chi(G_i) = \omega(G_i)$ , then so does their strong product  $G_1 \cdot G_2$ .
- **32.** (a) Let P be a partially ordered set and define a graph G on P by joining  $x, y \in P$  if and only if  $x \leq y$  or  $y \leq x$ .
  - (a) Show that  $\chi(G) = \omega(G)$ .
  - (b) Show the same for the complement of G.
- 33. Which of the previous exercises yield perfect graphs<sup>‡</sup>?
- 34. A graph G is perfect if and only if every induced subgraph G' of it contains an independent set which meets all maximum cliques of G'.
- **35\*.** Show that, substituting a perfect graph  $G_x$  for each point x of a perfect graph G produces a perfect graph G'.

† In fact, there exist graphs with arbitrarily large girth and chromatic number. [P. Erdős; see ES]

<sup>&</sup>lt;sup>‡</sup> Problems 9.28–32 suggest that the complement of any perfect graph is perfect. This is in fact true, and a proof of it will follow from the theory of hypergraphs. See 13.55–57.

**36.** Show that the chromatic polynomial of a graph G is a polynomial in the number  $\lambda$  of colors, and determine its degree.

37. Show that

$$P_{G}(\lambda) = \sum_{T \subseteq E(G)} (-1)^{|T|} \lambda^{c(T)},$$

where c(T) = c(V(G); T).

38. Show that

$$P_G(\lambda) = P_{G-e}(\lambda) - P_{G/e}(\lambda)$$
.

- 39. Determine the chromatic polynomials of complete graphs, trees, circuits and wheels.
- 40. Express the chromatic polynomial of G given the chromatic polynomials of
  - (a) its components, and
  - (b) its blocks.
- 41. Determine the chromatic polynomial of an interval graph.
- **42.** If G has no loops then

$$P_G(x) = x^n - a_{n-1}x^{n-1} + a_{n-2}x^{n-2} - \dots + (-1)^{n-1}a_1x$$

where  $a_i \geq 0$ . What is  $a_{n-1}$ ? Find an interpretation for  $a_1$ . (Here and later n is the number of points.)

- **43.** If G is connected, then  $a_i > 0$  (i = 1, ..., n 1) and  $a_{n-1} < a_{n-2} < ... < a_{\lfloor n/2 \rfloor + 1}$ .
- **44.** Let  $x_0$  be the largest real root of  $P_G(\lambda)$ . Show that  $x_0 \leq n-1$ .
- **45.** Show that  $P_G(\lambda)$  has no root in the interval (0, 1).
- 46. (a) Determine the meaning of the multiplicities of the roots 0 and 1.
  - (b) Prove that  $|P_G(-1)|$  is the number of acyclic orientations of G.
- 47\*. (a) Let G be a planar map with a quadrilateral face F. Let  $e_1$ ,  $e_2$  be the two diagonals of F, and  $G_i = G + e_i$  (i = 1, 2). Find a relation between  $P_{G_1}(\lambda)$ ,  $P_{G_2/e_1}(\lambda)$ ,  $P_{G_1/e_1}(\lambda)$ ,  $P_{G_2/e_2}(\lambda)$ .
  - (b) Is there any linear relation

$$aP_G(\lambda) + bP_{G_1}(\lambda) + cP_{G_2}(\lambda) = 0$$

with some fixed integers a, b, c not all 0, which holds for every G and  $\lambda$ ? Show that, except for at most five cases, there is no such relation even for a fixed value of  $\lambda$ .

(c) Set 
$$\tau = \frac{\sqrt{5}+1}{2}$$
. Then

$$P_G(\tau+1) = (\tau+1)(P_{G_1}(\tau+1) + P_{G_2}(\tau+1)).$$

48. Show that  $\tau + 1$  is not a root of any chromatic polynomial.

**49\*.** (The Golden Ratio Theorem) (a) If G is a planar triangulation with n points, then

$$P_G(\tau+2) = (\tau+2) \, \tau^{3n-10} \, P_G^2(\tau+1).$$

(b) For any planar graph G,

$$P_G(\tau+2)>0.$$

- 50. A planar graph G without loops has chromatic number at most 5.
- 51. Given a planar 3-regular 2-connected map, we can 4-color the faces (in such a way that adjacent faces get different colors) if and only if its chromatic index is 3.
- 52. Let G be a 3-regular 2-connected planar map. Then the faces of G are 4-colorable if and only if we can associate one of the numbers +1, -1 with each point in such a way that the sum of numbers associated with the points of any face is divisible by 3.
- 53. Assume that a planar map has a Hamiltonian circuit. Show that its faces are 4-colorable.
- 54\*. (a) The connected planar graph G has all degrees at most 5, and at least one point of degree at most 4. Prove that G is 4-colorable.
  - (b) Prove that all 5-regular planar graphs are also 4-colorable.
- 55. We can two-color the points of every simple planar map G in such a way that each face receives both colors.
- 56. A planar triangulation is 3-colorable if and only if every point has even degree.
- 57. Let us draw some lines in the plane, such that no 3 are concurrent. Consider their points of intersection as the points of a graph and the segments between neighboring intersection points as edges. Prove that the resulting planar graph is 3-colorable.

## § 10. Extremal problems for graphs

Turán raised and solved the following problem in 1943: How many edges guarantee that a graph with this number of edges (and n given points) has a complete subgraph with k points? This type of question has been extensively investigated since then. Actually, any problem of the form "what is the extremal value of a given parameter for a given class of graphs?" is an "extremal problem"; in this sense, almost everything in combinatorics could be included in this chapter. However, we shall mainly restrict ourselves to the case when we are looking for the maximum number of edges among graphs on n points

<sup>†</sup> The Four Color theorem asserts  $P_G(4) > 0$  for every planar graph. Since  $\tau + 2 = 3, 6...$  is quite close to 4,  $P_G(\tau + 2) > 0$  is interesting from this point of view.


not containing certain subgraphs. Still, these subgraphs may only be prescribed up to isomorphism, up to homeomorphism (if, for example, circuits are excluded) or in some other way (if for example Hamiltonian cycles are excluded), and this leads to investigations with quite different methods.


One feature common to all of them however is that they can all be translated from questions involving the number of edges into ones involving instead the degrees of the points. There are several well-known tricks for interpreting infor-


mation from one formulation in terms of the other.

Again we are able to select only a few characteristic problems from this large field. The reader will find other such problems in all graph theory books and a large variety of solved and unsolved problems in papers of P. Erdős.<sup>†</sup>

- 1. Suppose that the simple graph G on n points has more than 3(n-1)/2 edges. Prove that it contains a  $\Theta$ -graph, i.e. three independent paths connecting the same pair of points.
- 2. (a) If every point of a simple graph G has degree at least 3, then G contains a circuit with a chord.
- (b) If G is a simple graph with  $n \ge 4$  points and 2n-3 edges, then G contains a circuit with a chord.
- 3. (a) Suppose that the simple graph G has all degrees at least 3. Prove that it contains a subdivision of  $K_4$ .
- (b) Suppose that G is a simple graph with n points and 2n-2 edges. Prove that G contains a subdivision of  $K_4$ .
- **4\***. Let G be a simple graph with all degrees at least 3 and containing no two disjoint circuits. Prove that G is one of the following graphs: (i)  $K_5$ , (ii) a wheel, (iii)  $K_{3,n-3}$  with any set of edges connecting points in the 3-element class added (Fig. 8).







- 5. (a) Every simple 2-connected graph with  $n \ge 5$  points and 2n-2 edges contains a subdivision of  $K_{2,3}$ .
- (b) Every simple 3-connected graph with  $n \ge 6$  points and 3n-5 edges contains a subdivision of  $K_{3,3}$ .

<sup>†</sup> See: P. Erdős, The Art of Counting, Selected Writings, The MIT Press, Cambridge-London, 1973.

- LEMS 10.6
- (c) How many edges guarantee the existence of a subdivision of  $K_{2,3}$  and  $K_{3,3}$  respectively, without the above connectivity assumptions?
- **6.** (a) If a simple graph G can be contracted onto  $K_4$ , it also contains a subdivision of  $K_4$ .
  - (b) This is not true for  $K_5$  instead of  $K_4$ . Find a 4-connected counterexample!
- 7. Let G be a simple graph with m edges and n points. Prove that it contains a connected (non-spanning) subgraph  $G_1$  such that the neighbors of  $G_1$  (i.e. those points of  $V(G) V(G_1)$  adjacent to some point of  $G_1$ ) induce a graph with all degrees at least  $\frac{m}{n} 1$ .
- 8. Let G be a simple graph on n points,  $m \geq 3$  and  $|E(G)| \geq 2^{m-3}n$ . Then G can be contracted onto  $K_m$ .
- 9. Let F be any graph with k edges and no loops or isolates. Prove that if G is a simple graph with

$$|E(G)| \ge 2^k |V(G)|,$$

then G contains a subdivision of F.

\*

- 10. Determine the smallest 3-regular graphs with girths 4 and 5.
- 11. If G is an r-regular graph with girth g, then

$$|V(G)| \ge egin{cases} 1 + r + r(r-1) + \dots + r(r-1)^{rac{g-3}{2}} & ext{if} & g ext{ is odd,} \ 2ig(1 + (r-1) + \dots + (r-1)^{rac{g}{2}-1}ig) & ext{if} & g ext{ is even.} \end{cases}$$

- 12. Let  $r \geq 2$  and  $g \geq 2$  be given. Then there exists an r-regular graph with girth g.
- 13. Let G be an r-regular graph with girth at least g having the least number of points. Prove the following assertions:
  - (a) The diameter of G is at most g.
  - (b) The girth of G is g.

(c) 
$$|V(G)| \le \frac{r}{r-2} (r-1)^g$$
.

- 14. If G is an r-regular graph  $(r \ge 2)$  with girth at least g, and G has 2n points, where  $n \ge 2r^g$  then G can be embedded in an (r + 1)-regular graph with girth at least g on the same set of points.
- 15. (a) Construct a (p+1)-regular graph on  $2(p^2+p+1)$  points with girth 6 (p prime).
- (b)\* Construct a (p+1)-regular graph on  $2(p^3+p^2+p+1)$  points with girth 8 (cf. 10.11).

- **16.** (a) Let G be a graph on n points such that  $3 \le d_G(x) \le d$  for each point x. Then, to represent all circuits of G, we need at least  $\frac{n+2}{d+1}$  points.
- (b) If G has girth g and minimum degree at least 3 then to represent all circuits of G we need at least  $\frac{3}{8}2^{g/2}$  points.
- 17\*. Let G be a graph with all degrees at least 3 and girth  $g \ge 3$ . Prove that G contains  $\frac{3}{8g}$   $2^{g/2}$  disjoint circuits.
- 18. Let G be a graph, let  $\nu$  denote the maximum number of disjoint circuits in G and  $\tau$  the minimum number of points that represent all circuits of G. Prove that
  - (a) if v = 1, then  $\tau \leq 3$ ,
  - (b)  $\tau \ge \nu \ge \frac{\tau}{4 \log \tau}$ ,
  - (c) for infinitely many values of  $\tau$  there are graphs G with

$$v \leq \frac{4\tau}{\log \tau}.$$

\*

- 19. Let G be a 2-connected graph,  $x, y \in V(G)$  and assume that each point different from x, y has degree at least k. Prove that x, y can be connected by a path of length at least k.
- **20\*.** Let G be a simple graph such that, for each  $X \subset V(G)$  with  $|X| \leq k$ .

$$| \Gamma(X) - X | \ge 2 |X| - 1.$$

Prove that G contains a path of length 3k-2.

- **21.** Let G be a simple graph on n points  $x_1, \ldots, x_n$  with degrees  $d_1 \leq d_2 \leq \ldots \leq d_n$ . Prove that G has a Hamiltonian circuit if it satisfies any one of the following:
  - (a) (Dirac's Condition)  $d_1 \ge n/2$ ,
  - (b) (Pósa's Condition)  $d_k \ge k + 1$ , for  $k < \frac{n}{2}$ ,
  - (c) (Bondy's Condition)  $d_l \leq l$ ,  $d_k \leq k$  implies that  $d_l + d_k \geq n \ (k \neq l)$ ,
  - (d) (Chvátal's Condition)  $d_k \le k < \frac{n}{2}$  implies that  $d_{n-k} \ge n-k$ .
- **22.** If G is a simple graph on n points with all degrees at least  $\frac{n+q}{2}$ , then any set F of q edges which form disjoint paths is contained in a Hamiltonian circuit.

- **23\***. Let G be an n-regular simple graph on 2n + 1 points. Prove that G has a Hamiltonian circuit.
- **24.** In the simple graph G on  $n \ge 2$  points, every point has degree greater than n/2. Show that any two points of G can be joined by a Hamiltonian path.
- **25.** Let G be a simple digraph on n points with all indegrees and outdegrees at least n/2. Prove that G contains a Hamiltonian cycle (cf. 10.21a).
- **26.** Let G be a k-connected simple graph such that G contains no set of k+1 independent points  $(k \ge 2)$ . Prove that G has a Hamiltonian circuit.
- 27. Let G be a simple graph on n points and with all degrees at least k. Then
  - (a) G contains a circuit of length at least k+1,
- (b)\* if G is 2-connected, it contains a circuit of length at least 2k or a Hamiltonian circuit.
- **28.** Let G be a simple graph on n points with more than  $\frac{k(n-1)}{2}$  edges. Then G contains a circuit of length at least k+1.
- **29\***. Let G be a 2-connected simple graph and let l be the maximum length of circuits in G. Then G contains no path of length  $\left\lceil \frac{l^2}{4} \right\rceil + 1$ .
- **30.** The simple graph G on n points contains no triangles. Prove that

$$|E(G)| \leq \frac{n^2}{4}.$$

**31.** If G is a simple graph without triangles, then

$$|E(G)| \leq \alpha(G) \cdot \tau(G).$$

Prove the result in the preceding problem from this.

**32.** (a) If G is a simple k-regular graph on n points, then the total number of triangles in G and  $\overline{G}$  is

$$\binom{n}{3} - \frac{n}{2}k(n-k-1).$$

(b) Any simple graph G on n points and its complement  $\overline{G}$  contain together at least

$$\frac{n(n-1)(n-5)}{24}$$

triangles.

33. The number of triangles in a graph with n points and m edges is at least

$$\frac{4m}{3n}\left(m-\frac{n^2}{4}\right).$$

- **34.** (Turán's Theorem) Let G be a simple graph with mk points and more than  $\binom{k}{2}m^2$  edges. Prove that G contains a complete (k+1)-graph. Generalize for the case n=mk+r,  $1\leq r\leq k-1$ .
- **35.** Assume G is a simple graph containing no complete (k + 1)-graph. Prove that there is a simple k-chromatic graph H on V(G) such that

$$d_H(x) \ge d_G(x)$$
  $(x \in V(G)).$ 

Deduce Turán's theorem from this.

- **36.** (a) Every simple graph on n points with at least  $\frac{n}{4}(1+\sqrt{4n-3})$  edges contains a quadrilateral (4-circuit).
- (b) If  $n=p^2+p+1$  (p prime), then there exists a 4-circuit-free simple graph on n points with  $\frac{1}{2} p(p+1)^2 \sim \frac{1}{2} n^{3/2}$  edges.
- **37.** Assume that the simple graph G with n points and m edges contains no  $K_{rr}$ . Prove that

$$m < C \cdot n^{2 - \frac{1}{r}},$$

where C depends only on r.

- **38\***. (Erdős–Stone Theorem) (a) Let  $\varepsilon > 0$  and  $k, t \ge 1$  be given. Prove that if n is large enough, then every graph on n points and with all degrees at least  $(1-1/k+\varepsilon)n$  contains k+1 disjoint t-sets such that any two points in different t-sets are adjacent.
- (b) Let G be a simple graph with n points and  $(1 1/k + \varepsilon) \frac{n^2}{2}$  edges.

Then, if n is large enough, G contains k+1 disjoint t-sets such that any two points in different t-sets are adjacent.

(c) For each graph  $G_0$ , denote by  $M(n, G_0)$  the maximum number of edges in a simple graph on n points containing no subgraph isomorphic to  $G_0$ . Prove that

$$\lim_{n \to \infty} \frac{M(n,G_0)}{n^2} = \frac{1}{2} \left( 1 - \frac{1}{\chi(G_0) - 1} \right).$$

- **39\***. Let  $G_0$  be a (k+1)-chromatic graph such that  $G_0-e$  is k-colorable for some edge e. Prove that the graph  $H_{n,k}$  in the solution of 11.34 does not contain  $G_0$  but if n is large enough, any other graph with this number of edges contains  $G_0$  as a subgraph. So  $H_{n,k}$  is the unique extremal graph.
- **40\*.** Let G be a simple graph on n points and denote by  $N_k$  the number of complete k-graphs in G. Prove that

(a) 
$$\frac{N_{k+1}}{N_k} \ge \frac{1}{k^2 - 1} \left( k^2 \frac{N_k}{N_{k-1}} - n \right)$$
,

$$\text{(b) If } |E(G)| = \left(1 - \frac{1}{\vartheta}\right) \frac{n^2}{2}, \text{ then } N_k \geq \binom{\vartheta}{k} \left\lfloor \frac{n}{\vartheta} \right\rfloor^k \qquad (k \leq \vartheta + 1, \ \vartheta \ \text{real}).$$

**41.** (a) A tournament T on n points contains at most  $\frac{1}{4} \binom{n+1}{3}$  3-cycles.

(b) A strongly connected tournament T on n points contains at least n-2 3-cycles.

**42.** A strongly connected tournament T on n points contains at least  $\binom{n-1}{2}$  cycles.

**43.** A tournament on n points contains at least one and at most  $n!/2^{n/2}$  Hamiltonian paths.

**44.** (a) Every tournament T on n points contains a transitive subtournament on  $[\log_2 n] + 1$  points.

(b) The number of k-element transitive subtournaments  $(1 \le k \le \log_2 n)$  is at least

$$\prod_{j=0}^{k-1} \left( \frac{n+1}{2^j} - 1 \right)$$
.

(c) If T is strongly connected the number of k-element transitive subtournaments  $(k \ge 3)$  is at most

$$\binom{n}{k} - \binom{n-2}{k-2}$$
.

# § 11. Spectra of graphs

It is a classical program to describe a structure in terms of its numerical "invariants". With properly chosen invariants, the problems considered in the theory may be transformed into numerical or algebraic problems concerning the invariants and powerful methods of classical algebra may then be


brought into play.

The invariants considered in the previous chapters (connectivity, chromatic number, chromatic polynomial) were defined combinatorially and their algebraic properties played only a minor role. On the other hand, introducing the spectrum of a graph as the spectrum of its adjacency matrix, we get an algebraically defined invariant system. The spectrum does not, in general, characterize the graph uniquely, though it does reflect many properties of it and more and more such connection are continually being discovered. Thus the introduction of spectra of graphs is not a universal method to solve all problems, but it does prove to be very powerful in some purely graph-theoretic situations (for example, in classifying strongly regular graphs).

The effective use of spectra in graph theoretic investigations depends on our ability to take two major steps. First we must be able to calculate the spectra of large classes of graphs (or, more generally, we must be able to translate graph theoretic information into spectral information) and, second, we must be able to deduce properties of graphs from their spectral properties. It is interesting that there is a large variety of methods for linking graphs with their spectra in both these senses.

Some properties of the spectra of graphs are special consequences of the Frobenius-Perron theory of non-negative matrices. We shall use without prooor reference the following facts (see, for example, F. R. Gantmacher, Applications of the Theory of Matrices, Interscience, 1959). The maximum eigenvalue of any graph G is non-negative, and there is a non-negative eigenvector belonging to it. Moreover if the graph is connected then the maximum eigenvalue has multiplicity 1 and there is a strictly positive eigenvector belonging to it. Any non-negative unity eigenvector belonging to the maximum eigenvalue maximizes the quadratic form  $\mathbf{x}^T A_G \mathbf{x}$  on the unit sphere. The maximum eigenvalue has maximum absolute value among all eigenvalues.

- 1. Determine the spectra and eigenvectors of a complete n-graph  $K_n$ , a star  $S_n$  and a circuit  $C_n$  on n points.
- 2. Suppose we know the spectrum of a regular graph. Determine the spectrum of
  - (a) its complement  $\overline{G}$ .
  - (b) its line-graph L(G).
  - (c) What is the spectrum of the Petersen-graph (Fig. 9)?



**3.** Let T be a forest and  $e = (x, y) \in E(T)$ . Then

$$p_T(\lambda) = p_{T-e}(\lambda) - p_{T-x-y}(\lambda)$$
.

**4.** Let T be a forest on n points and let  $a_k$  denote the number of k-element matchings in T. Then

$$p_T(\lambda) = \lambda^n - a_1 \lambda^{n-2} + a_2 \lambda^{n-4} - \ldots + (-1)^{\left[\frac{n}{2}\right]} a_{\left[\frac{n}{2}\right]} \lambda^{n-2\left[\frac{n}{2}\right]}.$$

5. Let  $P_n$  denote a path with n points. Show that its characteristic polynomial can be written in either of the following forms:

(a) 
$$p_{P_n}(\lambda) = \lambda^n - \binom{n-1}{1} \lambda^{n-2} + \binom{n-2}{2} \lambda^{n-4} - \binom{n-3}{3} \lambda^{n-6} \dots$$

(b) 
$$p_{P_n}(\lambda) = \frac{1}{\sqrt{\lambda^2 - 4}} \left[ \left( \frac{\lambda + \sqrt{\lambda^2 - 4}}{2} \right)^{n+1} - \left( \frac{\lambda - \sqrt{\lambda^2 - 4}}{2} \right)^{n+1} \right].$$

- (c) Determine the eigenvalues of  $P_n$ .
- 6. Find infinitely many non-isomorphic pairs of trees having the same spectrum.
- 7. Suppose the eigenvalues of  $G_1$  and  $G_2$  are given. Determine those of
  - (a) the Cartesian product of  $G_1$  and  $G_2$ ,
  - (b) the (strong) direct product  $G_1 \cdot G_2$  of  $G_1$  and  $G_2$ .
- 8. Let G be a graph whose automorphism group contains a regular, commutative subgroup  $\Gamma$ . Let  $\gamma_{xy}$  be the element of  $\Gamma$  which moves x to y and put  $V(G) = \{v_1, \ldots, v_n\}$ . Let  $\chi$  be a multiplicative character of  $\Gamma$ . Prove that

$$\sum_{(\nu_1,\,\nu_i)\,\in\,E(G)}\chi(\gamma_{\nu_1\,\nu_i})$$

is an eigenvalue of G.

- **9.** Determine the eigenvalues of the *n*-cube  $Q_n$ .
- 10. Let G, H be two graphs on the point set 1,  $\varepsilon$ ,  $\varepsilon^2$ , ...,  $\varepsilon^{p-1}$ , where  $\varepsilon = e^{2\pi i/p}$  and p is prime, both of which are invariant under the rotation by  $2\pi/p$ . Suppose  $G \cong H$ . Then there is an integer t such that taking the t<sup>th</sup> power of each point maps G onto H isomorphically.
- 11. If the bipartite graph G has no 1-factor then 0 is an eigenvalue of it.
- 12. If the 3-regular graph G has a system of disjoint subgraphs, each isomorphic with the graph in Fig. 10, which covers all points, then 0 is an eigenvalue of G.




Fig. 10

- 13. If  $\Lambda$  and  $\Lambda'$  are the greatest eigenvalues of G and G', respectively, where G' is a subgraph of G, then  $\Lambda \geq \Lambda'$ .
- 14. (a) Let  $\Lambda$  be the largest eigenvalue of the graph G and let d, D be the minimum and maximum degrees of G. Then

$$\max (d, \sqrt[]{D}) \leq \Lambda \leq D.$$

For which connected graphs is  $\Lambda = D$ ?

(b) If G has n points and m edges, then

$$\frac{m}{n} \le A \le \sqrt{\frac{2m(n-1)}{n}}$$
.

- 15. Show that the line-graph of any graph G has all its eigenvalues at least -2, and that, if |E(G)| > |V(G)|, then its least eigenvalue is -2.
- 16\*. Suppose the eigenvalues of G are distinct. Show that every non-identity automorphism of G is of order 2.
- 17\*. Suppose that the eigenvalues of the simple graph G are distinct and that its automorphism group is transitive. Show that  $G \cong K_2$ .
- 18. Let  $\lambda_1 \leq \ldots \leq \lambda_n$  be the eigenvalues of the graph G. Show that

$$\lambda_{\tau(G)+1} \geq 0$$
,  $\lambda_{\alpha(G)} \leq 0$ ,  $\lambda_{\omega(G)-1} \leq -1$ ,  $\lambda_{n-\omega(G)+1} \geq -1$ .

- 19. (a) A connected graph G with maximum eigenvalue  $\Lambda$  is bipartite iff  $-\Lambda$  is an eigenvalue.
- (b) A graph is bipartite iff its spectrum is symmetric with respect to the origin.
- **20.** Let  $\Lambda$  be the maximum eigenvalue of G. Then

$$\chi(G) \leq \Lambda + 1.$$

**21\***. (a) Let  $\lambda_1 \leq \ldots \leq \lambda_n = \Lambda$  be the eigenvalues of G, and set  $k = \chi(G)$ . Then

$$\lambda_1 + \ldots + \lambda_{k-1} \leq -\Lambda$$
.

- (b) Show that, if the graph G is k-colorable in such a way that two points are adjacent iff they have different colors, then we have equality in (a).
- **22\***. Suppose G has k negative and l positive eigenvalues. Also suppose that G is the union of m edge-disjoint complete bipartite graphs. Then

$$m \geq \max(k, l)$$
.

- 23. The number of distinct eigenvalues of a connected graph is greater than its diameter.
- **24.** Given a graph G, we can find a polynomial f(x) such that

$$f(A_G) = J$$

if and only if G is regular and connected.

- **25.** Given a projective plane, its *line-graph* L is formed as follows: Let U be the set of points of the plane and V the set of lines, then  $V(L) = U \cup V$  and  $u \in U$  is joined to  $v \in V$  iff  $u \in v$ . Determine the spectrum of L.
- 26\*. The simple graph G has the following properties:
  - (i) each point has degree d,

- (ii) for each pair of points, there are b > 0 other points adjacent to both.
- (a) Determine the spectrum of G.
- (b) Determine all such graphs with b = 1.
- (c) Find infinitely many non-trivial examples of such graphs (a trivial example being a complete graph).

# § 12. Automorphisms of graphs

Although most graphs have no automorphisms other than the identity, many graphs arising from different combinatorial, geometric or algebraic structures possess, or are characterized by, a large automorphism group. This makes the study of relationships between the structure of graphs and their auto-

morphism groups of some importance.

The early trend in these investigations was to deal with independence results; i.e. results which show that even the imposition of rather strong restrictions on the graphs may not restrict their automorphism groups. This approach involves the construction of graphs, for which fairly standard methods are now available. The other approach, in which properties of the graphs are found which do restrict the properties of their automorphism groups, needs more "ad hoc" methods.

If the automorphism group is not only considered as an abstract group but also as a permutation group, then, of course, we find stricter interrelationships with the graph. For example, the property that the automorphism group is transitive implies many nice properties for the graph.

Investigation of the endomorphism semigroup instead of the automorphism group mostly yields similar results with more complicated constructions. However, there are some suprising differences (see the last exercise).

- 1. What is the automorphism group of the Petersen graph (Fig. 9, p. 71)?
- 2. (a) Show that the automorphism group of the dodecahedron graph (Fig. 11) is isomorphic with  $A_5 \times Z_2$ .
- (b) The automorphism group of the cube is isomorphic with  $S_4 \times \mathbb{Z}_2$ . What are the automorphism groups of the other Platonic bodies?




Fig. 11

- **3.** Construct a graph having automorphism group isomorphic with  $Z_k$ , a cyclic group of order k.
- 4. Let  $\Gamma = \{g_1, \ldots, g_n\}$  be a group. Define a digraph G by joining  $g_i$  to  $g_j$  by an edge of color k if  $g_i g_j^{-1} = g_k$ . What is the automorphism group of the resulting colored digraph?
- 5. (Frucht's Theorem) Given a finite group  $\Gamma$ , there exists a simple graph G with automorphism group isomorphic to  $\Gamma$ .
- **6\***. Let  $\Gamma$  be a group of order n  $(n \ge 6)$ . Construct a simple graph G on 2n points with  $A(G) \simeq \Gamma$ .
- 7. (a) Any tournament has an odd number of automorphisms.
- (b)\* Every group  $\Gamma$  of odd order n is the automorphism group of a tournament on 2n points.
- **8\***. Given a finite group  $\Gamma$ , construct a 3-regular graph with automorphism group  $\Gamma$ .
- **9.** If G is a connected graph with one point  $x_1$  of degree 2 and all other points of degree 3 then the automorphism group of G is of order  $2^k$  for some k.
- 10. Let G be a graph such that A(G) acts semiregularly on E(G). Let  $\Gamma$  be any subgroup of A(G). Then G has an orientation  $\vec{G}$  such that  $A(\vec{G}) = \Gamma$ .
- 11\*. Let  $\Gamma_1$ ,  $\Gamma_2$  be two finite groups. Construct a graph G which has an edge e such that  $A(G) \simeq \Gamma_1$  and  $A(G e) \simeq \Gamma_2$ .
- 12. (a) Let G be a connected graph and  $\Gamma$  a subgroup of A(G) acting semi-regularly. Prove that G can be contracted onto a graph G' such that A(G') has a subgroup  $\Gamma' \cong \Gamma$  which acts regularly.
- (b) Let G be a connected graph and  $\Gamma$  a simple subgroup of A(G). Then some connected subgraph of G can be contracted onto a graph G' such that A(G') has a subgroup  $\Gamma' \cong \Gamma$  which acts edge-transitively on G'. If, in addition, the elements of  $\Gamma$  have no fixed point in common, then  $\Gamma'$  has the same property.
- 13. (a) If the automorphism group of a graph G is commutative and transitive it is isomorphic to the direct product of cyclic groups of order 2.
- (b) Construct a graph whose automorphism group is transitive and isomorphic with the direct product of n cyclic groups of order 2, for any given n.
  - (c)\* Construct a simple graph with this property for large enough n.
- 14\*. The r-regular connected graph G has a transitive automorphism group. Show that G is r-edge-connected.
- 15\*. (a) Let G be a graph with transitive automorphism group which is exactly 3-connected. Show that G is 3-regular. Is this true with 4 instead of 3?
- (b) How large must the connectivity of an r-regular connected graph with transitive automorphism group be?
- (c)A connected simple graph with an edge-transitive automorphism group and with all degrees at least r is r-connected.

- 16.\* A connected graph G with an even number of points and transitive automorphism group has a 1-factor and, in fact, each edge of it belongs to a 1-factor.
- 17. The graph G is connected and A(G) contains a commutative transitive subgroup  $\Gamma$ . Prove that G has a Hamiltonian circuit.
- 18. Which planar graphs have edge-transitive automorphism groups?
- 19\*. Assume that the planar graph G has a non-cyclic simple automorphism group. Prove that  $A(G) \simeq A_5$ .
- **20\***. (a) Assume G is a connected graph and A(G) contains a subgroup  $\Gamma \simeq (Z_p)^3$  (p prime) which acts semiregularly. Then G can be contracted onto  $K_p$ .
- (b) Assume G is a connected graph and  $A(G) > \Gamma' > \Gamma \simeq (\mathbb{Z}_p)^3$ , where p is a prime and  $\Gamma'$  is a simple group without a common fixed point. Then G can be contracted onto  $K_m$ , where  $m = \lceil \log p / \log 8 \rceil$ .
- (c) Let  $\mathcal K$  be a class of graphs, not containing all graphs, such that if  $G\in\mathcal K$  and  $e\in E(G)$ , then both G-e and G/e belong to  $\mathcal K$  (e.g. the class of graphs embeddable in a surface F, or the class of graphs having no subgraph contractible onto  $K_m$ ). Then there exists a group  $\Gamma$  such that no graph of  $\mathcal K$  has automorphism group isomorphic to  $\Gamma$ .
- **21.** Let  $\Gamma$  be a permutation group on a set  $\Omega$ . Construct a simple graph G such that  $V(G) \supseteq \Omega$ ,  $\Omega$  is invariant under A(G) and the restriction of A(G) to  $\Omega$  yields  $\Gamma$ .
- 22. Determine the endomorphisms of the Petersen graph (Fig. 9, p. 71).
- 23. Construct a rigid graph.
- **24\***. (a) Let  $\Sigma$  be a finite monoid (semigroup with identity). Prove that there exists a digraph G with colored edges such that the endomorphism semigroup End  $(G) \simeq \Sigma$ .
  - (b) Prove that there exists such a simple graph as well.
- **25.** (a) If End (G) is isomorphic to the multiplicative semigroup  $\{0, 1, -1\}$  then G e cannot be rigid for any edge e.
- (b)\* For any monoid  $\Sigma$  there exists a simple graph G with End G  $\cong \Sigma$  such that G + e is rigid for some new edge e.

# § 13. Hypergraphs

A hypergraph, being a set with a specified collection of subsets, is a very general structure to consider. Graphs (in which each specified subset has just two elements) and several other combinatorial structures (block designs, matroids) are very special cases. Consequently the investigation of hypergraphs throws up a very large variety of problems and the "theory" is very dispersed. However, when various problems are translated to hypergraphs some concepts such as matching number, covering number and chromatic

number seem to occur very frequently. In my opinion these concepts are almost as general as those of logic: for example, given a combinatorial quantity defined as a minimum, it is usually easy to express it as the covering number of a suitable hypergraph. The duality between matching number and covering number, not yet completely worked out, is the background of many more graph-theoretic and combinatorial results than would appear at first glance.

There are also some general methods in hypergraph theory. The averaging method (cf. problems 12, 21, 27, 28, 32, 41, 52, 53) seems to be trivial, but its direct application has settled problems which were unsolved for several years

(e.g. 32, 57).

Finally we remark that certain special questions concerning hypergraphs give rise to theories which are distinguished from the rest by their more developed methods. We touch on transversal theory in problems 5–7. The theory of block designs (i.e. hypergraphs with high degree of symmetry) is not treated here.

1. A connected hypergraph H contains no circuits if and only if

(\*) 
$$\sum_{E \in E(H)} (|E| - 1) = |V(H)| - 1.$$

- **2\***. (a) Let H be a totally balanced hypergraph with  $|E(H)| \ge 2$ . Then H has two edges E, F such that each point of E F has degree 1.
- (b) Let H be a totally balanced hypergraph and suppose that any two edges have at most p points in common. Then

$$\sum_{E \in E(H)} (|E| - p) \le |V(H)| - p.$$

- **3.** Let P be a path and let  $P_1, \ldots, P_m$  be subpaths (intervals) on P. Setting V = V(P),  $E_i = V(P_i)$ , prove that the hypergraph  $(V; \{E_1, \ldots, E_m\})$  is totally balanced.
- **4.** The 3-uniform hypergraph H (without multiple edges) has |V(H)|-1 edges. Prove that it contains a circuit of length at least 3.
- **5.** (a) (Hall's Theorem) A hypergraph H has a system of distinct representatives iff  $|V(H')| \ge |E(H')|$  for each partial hypergraph H' of H.
- (b) Given a hypergraph H, we want to select a 2-element subset f(E) of each edge E in such a way that the sets f(E), considered as the edges of a graph, form a forest. Prove that this is possible iff  $|V(H')| \ge |E(H')| + 1$  for each partial hypergraph H' of H.
- **6.** Let  $T \subseteq V(H)$ . When does H have a system of distinct representatives containing T?
- **7\***. Let  $H_1$ ,  $H_2$  be two hypergraphs with  $V(H_1) = V(H_2)$  and  $|E(H_1)| = |E(H_2)| = m$ . Then  $H_1$ ,  $H_2$  have a common system of distinct representatives iff

$$|V(H') \cap V(H'')| \ge |E(H')| + |E(H'')| - m$$

for any two partial hypergraphs H', H'' of  $H_1$  and  $H_2$ , respectively.

- 8. Let H be a hypergraph with m distinct edges  $(m \ge 2)$ . Prove the following assertions:
  - (a) The number of sets of the form  $E \triangle F(E, F \in E(H))$ , is at least m.
  - (b)\* The same holds for the sets of the form E F,  $(E, F \in E(H))$ .
- (c)\* The same holds for the sets of the form  $E \cup F$  or  $E \cap F$ ,  $(E, F \in E(H), E \neq F)$ .
- **9\***. Let H be a hypergraph such that if  $E \in E(H)$  then every  $F \subseteq E$  belongs to E(H). Prove that there exists a permutation  $\sigma$  of E(H) such that  $E \cap \sigma(E) = \emptyset$  for each  $E \in E(H)$ .
- 10. (a) The hypergraph H on n points has n distinct edges. Prove that it has a point x such that  $H \setminus x$  has n distinct edges.
- (b) The hypergraph H on n points has  $m \le \frac{3}{2}n$  distinct edges. Prove that it has a point x such that  $H \setminus x$  has at least m-1 distinct edges.
  - (c) The hypergraph H on n points has

$$m > \sum_{i=0}^{k-1} \binom{n}{i}$$

distinct edges. Prove that there is a set  $X \subseteq V(H)$  with |X| = k, such that every subset of X occurs among the sets  $X \cap E$ ,  $E \in E(H)$ .

11. Let H, K be two hypergraphs on the same n-element set V and suppose that

$$A' \subseteq A \in E(H) \Rightarrow A' \in E(H),$$

and

$$V \supseteq B' \supseteq B \in E(K) \Rightarrow B' \in E(K).$$

Prove that

$$|E(H) \cap E(K)| \le \frac{1}{2^n} |E(H)| \cdot |E(K)|.$$

- 12. (a) Let H be any r-uniform hypergraph. Prove that V(H) can be split into two classes such that the number of edges contained in one of the two classes is at most  $\frac{|E(H)|}{2^{r-1}}$ .
- (b) Prove that V(H) can be partitioned into r classes such that at least  $\frac{r!}{r^r}|E(H)|$  edges have the property that they contain exactly one element of each class.
- 13\*. An r-uniform hypergraph has m edges, any two of which meet in at most k points. Prove that it has at least

$$\frac{r^2m}{r+(m-1)\,k}$$

points.

14\*. Let H be a hypergraph with n points and m edges. Denote by d(x) the degree of x and suppose d(x) < m, 0 < |E| < n holds for all edges E and points x. Suppose further that, if  $x \notin E$ , then

$$d(x) \leq |E|$$
.

Prove that  $m \leq n$ .

- 15. (a) Let H be a hypergraph,  $V(H) \neq \emptyset$ , in which any two edges have exactly one element in common, and no point occurs in all edges. Then  $|E(H)| \leq |V(H)|$ .
- (b)\* (Fischer's Inequality) Prove that the same holds if any two edges have exactly  $\lambda$  elements in common (y > 0) and that, the characteristic vectors of edges are linearly independent over the reals.
- **16.** Let H, K be two hypergraphs on the same set of points, and suppose |E(H)| = m, |E(K)| = m', where m' = m or m + 1. Then

$$\sum_{A\in E(H)}\sum_{B\in E(K)}|A\triangle B|\geq \sum_{\{A,A'\}\subseteq E(H)}|A\triangle A'|+\sum_{B,B'\}\subseteq E(K)}|B\triangle B'|.$$

17\*. (a) Let  $F_1, \ldots, F_m$  be sets such that  $|F_i \triangle F_j| = 2k$   $(1 \le i < j \le m)$ . Then the degree d of any point satisfies

$$d(m-d) \leq km$$
.

- (b) If there is a point x with degree  $\neq 0, 1, m-1, m$ , then  $m \leq k^2 + k + 2$ .
- 18. (a) Suppose that any two edges of the simple hypergraph H on n points intersect. Prove that  $|E(H)| \leq 2^{n-1}$ .
  - (b) If, in addition, no two edges cover every point, then  $|E(H)| \leq 2^{n-2}$ .
- **19.** Let H be a hypergraph on n points without multiple edges, such that, for any two edges E,  $F \in E(H)$ , either  $E \subset F$  or  $F \subset E$  or  $E \cap F \neq \emptyset$ . What is the maximum number of edges in H?
- **20.** Divide the set of all subsets of an *n*-element set S into symmetric chains. (By a symmetric chain we mean a chain  $E_r \subset E_{r+1} \subset \ldots \subset E_{n-r}$ , where  $|E_i| = i$  and  $0 \le r \le \frac{n}{2}$ .)
- **21.** (a) (Sperner's Theorem) If H is a clutter on n points (i.e. no two edges of H contain each other), then  $|E(H)| \leq \binom{n}{\binom{n}{2}}$ .
  - (b) For which hypergraphs does equality hold?
- **22\*.** In any partially ordered set  $(S, \leq)$  there is a maximum set of pairwise incomparable elements (a maximum anti-chain) which is invariant under the automorphisms of S.

Give a new proof of Sperner's theorem 13.21 using the above observation.

23\*. Let us call a hypergraph H cross-cutting if each subset of V(H) is comparable with an edge of H.

(a) Does every cross-cutting hypergraph contain a cross-cutting clutter?

13.24

- (b) A minimal cross-cutting hypergraph has at most  $\binom{n}{n}$  edges  $\binom{n}{n}$ .
- **24.** If k is the maximum length of chains  $E_1 \subset E_2 \subset \ldots \subset E_k$  of edges in a hypergraph H on n points without multiple edges, then |E(H)| is not greater than the sum of the k largest binomial coefficients in the  $n^{\text{th}}$  row of the Pascal triangle.
- **25.** (a) Suppose that every set of r+1 edges of an r-uniform hypergraph H has a point in common. Then all edges of H have a point in common.
- (b) Suppose any k edges of an r-uniform hypergraph have a point in common. Then

$$\tau(H) \leq \frac{r-1}{k-1} + 1.$$

- **26.** Let H be an r-uniform hypergraph with  $\nu(H) = 1$  and  $\tau(H) > 1$ . Prove that there exists a set  $S \subseteq V(H)$  such that  $|E \cap S| \ge 2$  for every edge E and  $|S| \le 3r 3$ .
- 27\*. Let H be an r-uniform hypergraph such that any two edges of H intersect. Prove that there is a set  $W \subseteq V(H)$  such that

$$|W| \le (2r-1) \begin{pmatrix} 2r-3 \\ r-1 \end{pmatrix}$$

and any two edges in the subhypergraph  $H_W$  induced by W meet. This is not true with

$$|W| \le \binom{2r-3}{r-1}.$$

- **28.** (a) Let  $A_1, \ldots, A_t$  be distinct arcs of length k of a circuit C ( $|V(C)| \ge 2k$ ) and suppose that any two have an edge in common. Then  $t \le k$ .
- (b) (Erdős–Ko–Rado Theorem) Let H be an r-uniform hypergraph on n points without multiple edges such that any two edges of H intersect. Prove that

$$|E(H)| \le \binom{n-1}{r-1}.$$

**29\***. (a) Let H be an r-uniform  $\nu$ -critical hypergraph with n points and m distinct edges. Then (for r,  $\nu$  fixed)

$$m=O(n^{r-2}).$$

(b) Let H be any hypergraph with n points and m distinct edges. Then for  $n \ge n_0(r, v)$  (v = v(H)) we have

$$m \leq {n-1 \choose r-1} + {n-2 \choose r-1} + \cdots + {n-v \choose r-1}$$
.

 $30^*$ . Let H be a hypergraph and denote by d its maximum degree. Then

$$\tau(H) \le (1 + \log d) \ \tau^*(H).$$

**31\***. (a) Let  $0 < k \le r$  be integers,  $u \ge v \ge w \ge r - 1$  be reals and assume that

$$\begin{pmatrix} u \\ r \end{pmatrix} = \begin{pmatrix} v \\ r \end{pmatrix} + \begin{pmatrix} w \\ r-1 \end{pmatrix}.$$

Prove that

$$\binom{u}{k} \leq \binom{v}{k} + \binom{w}{k-1}.$$

- (b) Let H be an r-uniform hypergraph without multiple edges. Write |E(H)| in the form  $\binom{u}{r}$ , where  $u \ge r$  is real. Prove that the number of k-tuples contained in edges of H is at least  $\binom{u}{k}$   $(1 \le k \le r)$ .
  - (c) Deduce the Erdős-Ko-Rado theorem 13.28b from (b).
- 32\*. (a) Let  $|A_1| = \ldots = |A_m| = p$ ,  $|B_1| = \ldots = |B_m| = q$  and  $A_i \cap B_j = 0 \Leftrightarrow i = j$ . Then

$$m \leq \binom{p+q}{p}$$
.

- (b) An r-uniform  $\tau$ -critical hypergraph H has at most  $\binom{\tau(H)+r-1}{r}$  edges.
- **33.** If no two edges of a hypergraph H have exactly one point in common, then H is 2-colorable.
- **34.** If every point of a connected hypergraph H has degree 2 and H is not a (2-uniform) odd circuit, then H is 2-chromatic.
- 35. Determine all hypergraphs in which any two edges have exactly one point in common and which are not 2-colorable.
- **36.** Let H be a 3-uniform hypergraph on  $n \ge 5$  points in which each pair of points occurs in the same (positive) number of edges. Prove that H is not 2-colorable.
- 37\*. A hypergraph H has a totally unimodular incidence matrix if and only if each subhypergraph  $H_W$  has a bicorolation  $\{A_0, A_1\}$  such that

$$\left\lceil \frac{|E|}{2} \right\rceil \leq |E \cap A_1| \leq \left\lceil \frac{|E|}{2} \right\rceil^*$$

for every  $E \in E(H_W)$ . (The incidence matrix of  $H = (\{v_1, \ldots, v_n\}; \{E_1, \ldots, E_m\})$  is the matrix  $A = (a_{ij})$  defined by

$$a_{ij} = \begin{cases} 1 & \text{if } v_i \in E_j \\ 0 & \text{otherwise.} \end{cases}$$

- A matrix is totally unimodular if every square submatrix has determinant 0 or  $\pm 1.$ )
- **38.** (a) If every circuit of the hypergraph H has even length, then its points can be 2-colored so that the numbers of red and blue points in any edge differ by at most one.
  - (b) The incidence matrix of H is totally unimodular.
- **39.** If the incidence matrix A of a hypergraph  $H = (\{v_1, \ldots, v_n\}; \{E_1, \ldots, E_m\})$  is totally unimodular, then H is balanced.
- 40. A balanced hypergraph in which every edge has at least 2 elements is 2-colorable.
- **41.** If an r-uniform hypergraph has at most  $2^{r-1}$  edges, then it is 2-colorable.
- **42.** Prove, using 13.30, that there exists a non-2-colorable r-uniform hypergraph on  $2r^2$  points with  $c \cdot r^2 2^r$  edges.
- **43\***. If every edge of an r-uniform hypergraph H meets at most  $2^{r-3}$  other edges, then H is 2-chromatic.
- 44\*. Let H be an r-uniform hypergraph with n points and m edges such that two edges have at most one point in common.
  - (a) If  $n \leq 2^{r-4}$ , then H is 2-colorable.
- (b) If H is not 2-colorable, then it contains at least  $2^{r-4}/r$  points of degree at least  $2^{r-4}/r$ .
  - (c) If  $m < 4^{r-4}/r^3$ , then H is 2-colorable.

(For existence problems concerning such hypergraphs cf. 14.24).

- **45.** (a) The hypergraph H has the property that, for all  $k \ge 1$ , the union of any k edges has at least k+1 points. Prove that H is 2-chromatic.
- (b) Construct an r-uniform hypergraph H such that the union of any k edges (k = 1, 2, ...) has at least k points but H is not 2-chromatic.
- **46.** Construct, for infinitely many values of r, r-uniform hypergraphs without multiple edges which are not 2-chromatic but any two edges of which intersect and have
  - (a) more than r! edges,
  - (b) at most  $3^{r-1}$  edges,
  - (c) more than  $3^{r-1}$  points.
- 47. If a simple hypergraph H is r-uniform, 3-chromatic and any two edges of H intersect, then

$$|E(H)| < r^r$$
.

**48.** Prove<sup>†</sup> that, for any hypergraph H,

$$v(H) \leq v^*(H) = \tau^*(H) \leq \tau(H)$$
.

- **49.** (a) Let F be a graph. Then any k-cover is the sum of a 2-cover and of a (k-2)-cover.
  - (b) A similar assertion holds for k-matchings if k is even.
- (c) There are optimal fractional covers and also optimal fractional matchings, where the weights are halves of integers. Consequently,  $2\tau^*(G)$  is an integer.
  - (d) Deduce 7.37 from (c).
- **50.** Let G be a bipartite graph. Prove that every k-cover is the sum of k 1-covers and that every k-matching is the sum of k 1-matchings. Consequently,  $\tau(G) = \tau^*(G)$  and  $\nu(G) = \nu^*(G)$ .
- 51. (a) Prove that

$$\begin{split} \tau^*(G \otimes H) &= \tau^*(G) \ \tau^*(H), \\ \tau(G) \ \tau(H) &\geq \tau(G \otimes H) \geq \tau(G) \ \tau^*(H), \\ v(G) \ v(H) &\leq v(G \otimes H) \leq v(G) \ v^*(H) \end{split}$$

hold for any two hypergraphs G, H.

- (b) Given a hypergraph H,  $\tau(G \otimes H) = \tau(G)$   $\tau(H)$  holds for every hypergraph G iff  $\tau(H) = \tau^*(H)$ .
  - (c) Setting  $H^p = H \otimes H \otimes \ldots \otimes H$  (p factors), determine  $\lim_{p \to \infty} \sqrt[p]{\tau(H^p)}$ .
- 52. If a hypergraph H is  $\tau$ -critical, then either H consists of disjoint edges or  $\tau(H) > \tau^*(H)$ .
- **53.** If a hypergraph H is  $\nu$ -critical, then

$$v(H) < v^*(H)$$
.

- **54.** Every balanced hypergraph H has  $\tau(H) = \nu(H)$ .
- 55. If H is a hypergraph, then the following three assertions are equivalent.
  - (i)  $v(H') = v^*(H')$ , for every partial hypergraph H' of H.
  - (ii)  $\tau(H') = \tau^*(H')$ , for every partial hypergraph H' of H.
  - (iii) H is normal, i.e.  $v(H') = \tau(H')$  for every partial hypergraph H' of H.

<sup>†</sup> One may use the duality theorem of linear programming: If  $\max \mathbf{c}^T \mathbf{x}$  exists under the constraints  $A\mathbf{x} \leq \mathbf{b}$ ,  $\mathbf{x} \geq 0$  then also  $\min \mathbf{b}^T \mathbf{y}$  exists under the constraints  $A^T \mathbf{y} \geq \mathbf{c}$ ,  $\mathbf{y} \geq 0$  and  $\min \mathbf{b}^T \mathbf{y} = \max \mathbf{c}^T \mathbf{x}$  (here A is any  $n \times m$  real matrix and  $\mathbf{b}$  and  $\mathbf{c}$  are n- and m-dimensional real vectors.)

- 56\*. (Cont'd) The following assertions are also equivalent to (i)-(iii):
  - (iv) d(H') = q(H'), for every partial hypergraph H' of H.
- (v) H has the Helly-property; i.e. if edges  $E_1, \ldots, E_m$  meet pairwise, then  $E_1 \cap \ldots \cap E_m \neq \emptyset$  and  $\overline{L(H)}$  is perfect.
- 57\*. (Perfect Graph Theorem) A graph G is perfect iff  $\overline{G}$  is perfect.

#### § 14. Ramsey Theory

Every "irregular" structure, if it is large enough, contains a "regular" substructure of some given size. This phenomenon occurs in many situations. A typical example (the title of this chapter) is that, if we take a large complete graph whose edges are 2-colored (as "irregularly" as we like) there always exists a large monochromatic complete subgraph. However, one can color other objects (points in spaces, numbers, subsets, etc.) and also, one can look for other "regular" substructures (monochromatic subspaces, arithmetic progressions, configurations, etc.) Some of these questions turn out to be very difficult. However, a considerable number of them can be given a general treatment using categories (see problems 14.13–18 for the combinatorial background). The last sequence of problems deals with monotone subsequences of sequences and related problems in geometry and combinatorics. For example the famous problem of finding convex subsets of a set is treated here.

Ramsey theory in the narrower sense, when monochromatic subsets are looked for, is a special case of the theory of chromatic number of hypergraphs. Most of the results here can be formulated as assertions that a certain hyper-

graph "regular" and large enough, has high chromatic number.

We shall not deal with infinite Ramsey theory, although this is more closely related to the finite theory than, perhaps, are the finite and infinite counterparts of any other field in combinatorics. We hope that problems 9.14 and 14.19 will shed light on this connection.

- 1. (a) Every graph on  $\binom{k+l}{k}$  points contains either a complete graph with k+1 points or an independent set of l+1 points.
- (b) Let  $a_1, \ldots, a_k \geq 1$  be given integers  $(k \geq 2)$ . Prove that there exists a (least) natural number  $n = R_k(a_1, \ldots, a_k)$  such that, if we k-color the edges of  $K_n$  in any way, there will be an  $1 \leq i \leq k$  and a complete  $a_i$ -graph all of whose edges are colored with the i<sup>th</sup> color.
- 2. (a) Prove that

$$R_k(3) \stackrel{\text{def}}{=} R_k(3, \ldots, 3) \leq \lceil e \cdot k \rceil + 1;$$

and that equality holds for k = 2, 3.

(b)\* Prove that for  $k \geq 2$ ,

$$2^{k/2} \le R_2(k) < 2^{2k}.$$

3. (a) Ramsey's Theorem Let  $K_n^r$  denote the hypergraph consisting of all r-tuples of n points and let  $a_1, \ldots, a_k \ge 1$ . Prove that there exists a (least) number

 $n = R_k^r(a_1, \ldots, a_k)$  such that, if we k-color the edges of  $K_n^r$ , there will be an  $a_i$ -subset of  $V(K_n^r)$  all of whose k-subsets have the  $i^{\text{th}}$  color, for at least one  $1 \leq i \leq k$ .

(b) Prove that

$$R_k^{r+1}(a) < k^{[R_k^r(a)]^r}$$
.

- **4.** 2-color the edges of a complete n-graph ( $n \ge 3$ ). Prove that there will be a Hamiltonian circuit which either is monochromatic or consists of two monochromatic arcs.
- 5\*. We 2-color the edges of a complete  $\left[\frac{3k+1}{2}\right]$ -graph. Prove that there is a monochromatic path of length k.
- **6.** Let us 2-color the edges of a complete *n*-graph. Prove the following assertions:
- (a) If there is a monochromatic (2k+1)-circuit  $(k \ge 3)$ , then there is a monochromatic 2k-circuit.
- (b) If there is a monochromatic 2k-circuit ( $k \ge 3$ ), then there is either a monochromatic (2k-1)-circuit or two monochromatic disjoint complete k-graphs.
  - (c) If  $n \ge 2m 1$  and  $n \ge 6$ , then there will be a monochromatic *m*-circuit.
- 7. Let us 3-color the points of the plane. Prove that there will be two points at distance 1 with the same color.
- 8\*. We 2-color the points of the plane. Suppose there are three points of the same color forming a regular triangle with side 1 (a monochromatic regular triangle with side 1, for short). Then for each a, b > 0 such that |a b| < < 1 < a + b, we have a monochromatic triangle with sides 1, a, b.
- 9. (a) Show that if we 2-color the points of the plane, then we do not necessarily have a monochromatic regular triangle with side 1.
- (b) Prove that, 2-coloring the points of the plane, we always have a mono chromatic triangle with sides  $\sqrt{2}$ ,  $\sqrt{6}$ ,  $\pi$ .
- 10\*. Does there exist a natural number  $n = n_0(k, R)$  such that, if we k-color the points of n-dimensional Euclidean space, one of the colors will contain a configuration congruent to R, where
  - (a) R is a rectangle,
  - (b) R is a non-rectangular parallelogram.
- 11. Let us split the first n natural numbers into k classes. Prove that if  $n \ge k!e$ , then one of the classes contains three integers x, y, z with x + y = z.

12. Let  $k, r \ge 1$  be given. Then there is an  $n_0(k, r)$  such that if  $n \ge n_0(k, r)$  and  $\{1, \ldots, n\}$  is k-colored, then we can find natural numbers  $a, d_1, \ldots, d_r$  such that all sums

$$a + d_{i_1} + \ldots + d_{i_{\nu}} \quad (1 \le i_1 < \ldots < i_{\nu} \le r, \quad 0 \le \nu \le r)$$

have the same color (and, of course,  $a + d_1 + \ldots + d_r \leq n$ ).

- 13. Let us k-color all non-empty subsets of an n-element set. Prove that if n is large enough, there are two disjoint non-empty subsets X, Y such that X, Y,  $X \cup Y$  have the same color.
- 14\*. (a) Suppose that the set of all subsets of an *n*-element set S is k-colored, where  $n \geq N(k, t)$ . Find disjoint sets  $A_1, B_1, \ldots, A_t, B_t$  such that for any fixed sequence  $1 \leq i_1 < \ldots < i_{\nu} \leq t$ , all unions of the form

$$C_{i_1} \cup \ldots \cup C_{i_n}$$
  $(C_i = A_i \text{ or } B_i \text{ or } A_i \cup B_i)$ 

have the same color.

- (b) Prove that for any given k and r there is an n = n(k, r) with the following property: whenever the set of all subsets of an n-element set S is k-colored, then there exist non-empty disjoint subsets  $X_1, \ldots, X_r \subseteq S$  such that all non-empty unions of any of them have the same color.
- 15. Strengthen 14.12 as follows: For any k and r there exists a natural number n such that, if  $\{1, \ldots, n\}$  is k-colored, then there always exist natural numbers  $d_1, \ldots, d_r$  such that  $d_1 + \ldots + d_r \le n$  and all sums  $d_{i_1} + \ldots + d_{i_{\nu}}$   $(1 \le i_1 < \ldots < i_{\nu} \le r, \ \nu \ge 1)$  have the same color.
- 16\*. Consider the set  $m^S$  of all m-collections of points of the set S, where  $m \geq 2$ . (In an m-collection  $\mathcal A$  the same point may occur more than once but at most m times.) Let  $\alpha$  be a k-coloration of  $m^S$ .
- (a) There is an  $n_1(k, m)$  such that if  $|S| \ge n_1(k, m)$ , then there is a set  $X \subset S$  and an m-collection  $\mathcal{B}$  of points of S X such that  $mX + \mathcal{B}$  and  $(m-1)X + \mathcal{B}$  have the same color.
- (b) There is an  $n_r(k, m)$  such that if  $|S| \ge n_r(k, m)$ , then there are disjoint sets  $X_1, \ldots, X_r \subset S$  and an m-collection  $\mathcal{B}$  of points of  $S X_1 \ldots X_r$  such that each collection

$$\mathcal{A} = \sum_{i=1}^{r} a_i X_i + \mathcal{B} \qquad (0 \le a_i \le m)$$

has the same color as

$$\mathcal{R}' = \sum_{i=1}^r \min(a_i, m-1) X_i + \mathcal{B}.$$

17\*. (Cont'd) Prove that there is an N(k, r, m) such that if  $|S| \ge N(k, r, m)$ , then there are disjoint non-empty subsets  $X_1, \ldots, X_r$  of S and an m-collection  $\mathcal{B}$  of points of  $S - X_1 - \ldots - X_r$  such that all m-collections

$$\mathcal{A} = \sum_{i=1}^{r} a_i X_i + \mathcal{B}, \quad (0 \le a_i \le m)$$

have the same color.

- 18. (Van der Waerden's Theorem) There exists a number w = w(k, m) such that if the natural numbers  $0, 1, \ldots, w$  are k-colored, then there is a monochromatic arithmetic progression of length m.
- 19. Let P be any property of k-colored finite sets of natural numbers (e.g. a set has property P if it is monochromatic and forms an arithmetic progression). Suppose that, k-coloring all natural numbers in any way, some finite subset will have property P. Then there exists a natural number N such that, k-coloring the set  $\{1, \ldots, N\}$  arbitrarily, some subset of it will have property P.
- **20\***. Decide whether the following assertion is true: If the set of all natural numbers is k-colored, one of the classes contains x, y, z with
  - (a) x + y = 3z,
  - (b) x + 2y = z.

21\*. Let

$$a_1x_1+\ldots+a_nx_n=0$$

be an equation with integral coefficients. Then the following two properties are equivalent:

- (i) (\*) has a non trivial (0, 1)-solution.
- (ii) If we color the natural numbers with finitely many colors, (\*) will have a monochromatic solution.
- **22\***. Prove that there exists a function f(k) with the property that, if S is any set of integers with  $|S| \ge f(k)$ , then the set of all integers can be k-colored so that S+j meets all the colors for every integer j  $(S+j=\{s+j: s\in S\})$ . The k-coloration can even be required to be periodic.
- **23.** Let q be a prime power and  $k \ge 1$ ,  $r \ge 0$ .
- (a) Let  $n \ge N(k, r, q 1)$ . Then, if we k-color the points of the affine n-dimensional space over GF(q), then some r-dimensional subspace will be monochromatic.
  - (b) Prove an analogous theorem for projective spaces.
- **24.** Give two constructions, based on Ramsey's theorem and 14.23a, of an r-uniform hypergraph H which has chromatic number at least k+1 and in which any two edges have at most one point in common.
- **25.** Let  $(a_1, \ldots, a_{k^2+1})$  be any sequence of integers. Prove that it contains a monotone subsequence of length k+1.
- **26.** (a) Let f be an integral-valued function defined on  $\{1,\ldots,2^{n-1}\}$ ,  $1\leq f(i)\leq i$ , for  $i=1,\ldots,2^{n-1}$ . Prove that there is a sequence

$$1 = a_1 < \ldots < a_n \le 2^{n-1}$$

<sup>&</sup>lt;sup>†</sup> This assertion is a very special case of a general principle called *compactness*. This principle would allow for example the deduction of Ramsey's theorem 14.2 from an infinite version of it. Since we cannot go into set-theory, this exercise is to serve as an illustration.

with

$$f(a_1) \leq \ldots \leq f(a_n)$$
.

(b) This is not true for  $2^{n-1} - 1$  instead of  $2^{n-1}$ .

27\*. (a) Let  $T_n$  denote a transitive tournament on n points and let a number f(x,y) be associated with each edge  $(x,y) \in E(T_n)$ . Prove that if  $n > \binom{p+q-2}{p-1}$ , then there is either a (directed) path  $(x_0, \ldots, x_p)$  with

$$f(x_0, x_1) \le f(x_1, x_2) \le \ldots \le f(x_{p-1}, x_p)$$

or a path  $(y_0, \ldots, y_q)$  with

$$f(y_0, y_1) > f(y_1, y_2) > \ldots > f(y_{q-1}, y_q).$$

(b) Prove that the bound is sharp.

28\*. Let us associate an element  $f(X) \in X$  with each non-empty subset  $X \subseteq S$  of the set S.

(a) If |S| = 2m, then there is a set  $T \subseteq S$ , with |T| = m, and an ordering  $(x_1, \ldots, x_m)$  of S - T such that

$$f(S - \{x_1, \ldots, x_i\}) \in T$$
  $(i = 0, \ldots, m).$ 

(b) If  $|S| = 2^n$ , then there is a chain  $X_0 \subset X_1 \subset \ldots \subset X_n \subseteq S$  with  $f(X_1) = \ldots = f(X_n)$ .

(c) This is no longer true, when  $|S| = 2^n - 1$ .

**29.** Given  $N = k^n + 1$  points in the plane, we can always find an "almost straight" broken line, i.e. a sequence  $a_0, a_1, \ldots, a_k$  such that angle

$$\not < a_{i-1}a_i\,a_{i+1} > \left(1 - \frac{1}{n}\right)\pi.$$

**30.** We call a polygon convex from below (above) if it is convex and any semi-line starting from its vertices and parallel to the negative (positive) half of the y-axis has only its endpoint in common with the polygon (Fig. 12).

(a) Prove that any set S of  $\binom{p+q}{p}+1$  points in the plane, in general position (which includes here the property that no line determined by them




Fig. 12

is parallel to either axis), contains either a (p + 2)-gon convex from below or a (q + 2)-gon convex from above.

- (b) This is not true with  $\binom{p+q}{p}$  points.
- 31. (a) There exists a (least) natural number K(m) such that, if we have K(m) points in the plane in general position there always exists a convex m-gon of whose vertices are from the given set.
  - (b)\* Establish the inequalities†

$$2^{m-2} + 1 \le K(m) \le {2m-4 \choose m-2} + 1$$

(c) Show that K(4) = 5 and K(5) = 9.

#### § 15. Reconstruction

The problem of reconstruction is a version of the classical principle of invariants. If we associate some structure A' with each structure A (for example the line-graph of a graph, or the generating function of a sequence, or the Betti numbers of a manifold), we may be interested in the question as to whether A' uniquely determines A. For example, the fact that a power series determines its coefficients is a result of this type. Also we may be interested in characterizing those structures which arise in the form A'.

We restrict ourselves to reconstruction-type problems of combinatorial character; reconstructing graphs from their line-graphs, "matroids", direct products with themselves etc. The best known problem in this area is the Reconstruction Conjecture, which asserts that a graph can be reconstructed from the collection of its maximal proper induced subgraphs. This is solved

in special cases only.

We mention here for the sake of completeness one class of problems in this connection which we shall not be able to deal with in this volume. These are the problems concerning the construction of symmetric combinatorial structures (block designs, partial geometries, latin squares) from finite fields. It is often the case for certain values of parameters that these are the only structures of the given type. This is so for example in the case of geometries over finite fields. However the techniques needed usually involve an extensive use of the theory of block designs and this is why we cannot include them here.

- 1. (a) Let  $G_1$ ,  $G_2$  be two simple graphs with all degrees at least 4 and suppose that their line-graphs  $L(G_1)$  and  $L(G_2)$  are isomorphic. Prove that  $G_1 \cong G_2$ .
- (b) Let  $G_1$ ,  $G_2$  be two connected simple graphs and suppose  $L(G_1) \cong L(G_2)$ , but  $G_1 \ncong G_2$ . What are  $G_1$  and  $G_2$ ?
- (c) Let  $G_1$ ,  $G_2$  be connected simple graphs and  $\varphi: L(G_1) \to L(G_2)$  an isomorphism between  $L(G_1)$  and  $L(G_2)$ . We say that  $\varphi$  is trivial if there is an isomorphism  $\psi: G_1 \to G_2$  such that  $\psi$  induces  $\varphi$  on  $E(G_1)$ . For which graphs are there non-trivial isomorphisms between  $L(G_1)$  and  $L(G_2)$ ?

<sup>&</sup>lt;sup>†</sup> It is conjectured that the lower bound is the exact value of K(m).

- 2. (a) Let  $\alpha$  be an automorphism of the line-graph of the hypergraph  $K'_n$  consisting of all r-tuples of n points  $(n \geq 3r)$ . Then  $\alpha$  is induced by an automorphism of  $K'_n$  (i.e. a permutation of  $V(K'_n)$ ).
  - (b) The preceding result holds for  $3r \ge n > 2r$  but fails for  $n \le 2r$ .
- **3.** Let  $0 \le i < r$  be given. Form the graph  $L_i(K_n^r)$  whose points are the r-tuples in  $K_n^r$  and in which two r-tuples A, B are adjacent iff  $|A \cap B| = i$ . Show that for sufficiently large n, every automorphism of  $L_i(K_n^r)$  is induced by a permutation of  $V(K_n^r)$  in the natural way.
- **4.** (a) Every simple graph is the line-graph of some hypergraph. Describe these hypergraphs.
- (b) Show that every graph G on n points without isolated points is the line-graph of a hypergraph on  $\left[\frac{n^2}{4}\right]$  points.
- 5. (a) Let G be a simple graph with more than 9 points. Suppose that, for each point x, G-x is the line-graph of some simple graph  $H_x$ . Prove that G is the line-graph of a simple graph.
- (b) One can find a finite number of "sample graphs"  $G_1, \ldots, G_k$  so that a simple graph G' is the line graph of a simple graph if and only if it does not contain any of  $G_1, \ldots, G_k$  as an induced subgraph.
- **6.** (a) Call a triangle abc of a graph G odd if there is a point  $x \neq a$ , b, c which is adjacent either to one or to three of the points a, b, c. Show that if the simple graph G is a line-graph, then
  - (i) it does not contain the 3-star as an induced subgraph,
  - (ii) whenever abc, bcd are odd triangles  $(a \neq d)$ , then a is adjacent to d.
- (b) If G is a simple graph such that (i) and (ii) hold, then G is the line-graph of some simple graph.
  - (c) Determine the "sample graphs" in 15.5b.
- 7\*. Show that the statement of 15.5b is not valid for line-graphs of 3-uniform hypergraphs (without multiple edges).
- 8. (a) Construct two isomorphic 2-connected planar graphs whose duals are not isomorphic.
  - (b) If two 3-connected planar graphs are isomorphic so are their duals.
- 9. Let  $G_1$ ,  $G_2$  be two 3-connected graphs and  $\varphi \colon E(G_1) \to E(G_2)$  a bijection between their edges such that edges forming a circuit in  $G_1$  are mapped onto the edges of a circuit in  $G_2$  and conversely.
  - (a) Prove that  $\varphi$  preserves adjacency of edges.
  - (b) Prove<sup>†</sup> that  $G_1 \cong G_2$ .
- 10. Let T, T' be two trees and assume they have the same set  $\{x_1, \ldots, x_r\}$  of endpoints. Suppose  $d_T(x_i, x_j) = d_{T'}(x_i, x_j) \qquad (1 \le i < j \le r).$

 $w_{T}(w_{l},w_{j})=w_{T}(w_{l},w_{j}) \qquad (1 \leq i < j \leq i).$ 

<sup>†</sup> This problem says that 3-connected graphs can be reconstructed from their matroids.

Show that T, T' are isomorphic.

11. Let  $x_1, \ldots, x_r$  be the endpoints of a tree T and set

$$d_{ij} = d(x_i, x_j).$$

Show that

(a) for any three indices i, j, k,

$$d_{ij} + d_{jk} - d_{ik} \ge 0$$
,  $d_{ij} + d_{jk} - d_{ik} \equiv 0 \pmod{2}$ ,

- (b) for any four indices i, j, k, l, two of the numbers  $d_{ij} + d_{ik}$ ,  $d_{ik} + d_{jl}$ ,  $d_{il} + d_{jk}$  are equal and the third one is not greater than these two.
- (c) Suppose  $(d_{ij})_{i,j=1}^r$  is a symmetric matrix consisting of non-negative integers for which  $d_{ij} = 0$  iff i = j, and which satisfies (a) and (b). Then there exists a tree T with r endpoints  $x_1, \ldots, x_r$  such that

$$d(x_i, x_j) = d_{ij}$$

- 12\*. (a) Let  $n = 2^k$ . Construct two collections  $\{a_1, \ldots, a_n\} \neq \{b_1, \ldots, b_n\}$  of integers (the same integer may occur more than once) such that the two collections  $\{a_i + a_j : 1 \leq i < j \leq n\}$  and  $\{b_i + b_j : 1 \leq i < j \leq n\}$  are equal.
- (b) This is not possible when n is not a power of 2; i.e. in this case  $\{a_1, \ldots, a_n\}$  is reconstructible from the collection  $\{a_i + a_j : 1 \le i < j \le n\}$ .
- 13. Let  $(a_{ij})$ ,  $(b_{ij})$  be complex matrices in each of which no two rows are equal and let

$$f(s_1, \ldots, s_k) = \sum_{i=1}^{N} {a_{i1} \choose s_1} \ldots {a_{ik} \choose s_k},$$

$$\sum_{k=1}^{N} {b_{i1} \choose s_k} \ldots {b_{ik} \choose s_k}$$

$$g(s_1,\ldots,s_k) = \sum_{i=1}^N {b_{i1} \choose s_1} \cdots {b_{ik} \choose s_k}.$$

Suppose further that  $f(s_1, \ldots, s_k) = g(s_1, \ldots, s_k)$  for every choice of non-negative integers  $s_1, \ldots, s_k$ . Prove that  $f \equiv g$ , i.e. the matrices  $(a_{ij})$  and  $(b_{ij})$  arise from each other by permutation of the rows.

- 14. Let  $G_1$ ,  $G_2$  be two simple graphs with a common underlying set V, |V| > 3. Suppose  $G_1 x y \cong G_2 x y$  for all  $x, y \in V$ . Show that  $G_1$  and  $G_2$  are identical.
- 15. (a) Is the following statement true? If  $G_1$  and  $G_2$  are simple graphs with a common underlying set V (which is large enough) and  $G_1 x \simeq G_2 x$  for all  $x \in V$ , then  $G_1$  and  $G_2$  are identical.
- (b) Let  $G_1$ ,  $G_2$  be two graphs on a common underlying set V and suppose  $G_1 x \cong G_2 x$  for all  $x \in V$ . Let H be any given graph with less than |V| points. Show that  $G_1$  and  $G_2$  have the same number of subgraphs [induced subgraphs] isomorphic to H.
  - (c) If  $G_1$  is disconnected, then so is  $G_2$  and they are isomorphic.

<sup>†</sup> Ulam's famous conjecture states that this assertion holds for any pair of graphs.

**16\***. Let  $T_1$ ,  $T_2$  be two trees on the same set V and suppose that for each  $x \in V$ ,

$$T_1 - x \cong T_2 - x$$
.

- (a) Prove that  $T_1$ ,  $T_2$  have the same diameter d.
- (b) Prove further that  $T_1 \simeq T_2$ .
- 17\*. (a) Let  $G_1, G_2$  be two simple graphs with  $|V(G_1)| = |V(G_2)| = n$ ,  $E(G_1) = \{e_1, \ldots, e_m\}$ ,  $E(G_2) = \{f_1, \ldots, f_m\}$ . Assume  $m > \frac{1}{2} \binom{n}{2}$  and  $G_1 e_i \cong G_2 f_i$  for  $i = 1, \ldots, m$ . Then  $G_1 \cong G_2$ .
  - (b) Prove the same result when  $m > \frac{1}{2} \binom{n}{2}$  is replaced by  $m > n \log n$ .

18. Let  $H_1$ ,  $H_2$  be two r-uniform hypergraphs on the same set V, and let  $r \leq k \leq |V| - r$ . Suppose that

$$|E(H_1 - X)| = |E(H_2 - X)|$$

for all  $X \subseteq V$ , with |X| = k. Then  $H_1$  and  $H_2$  are identical.

19. Let  $H_1$ ,  $H_2$  be hypergraphs on the same set V and suppose that, for each  $x \in V$ ,

$$H_1 \setminus x = H_2 \setminus x$$
.

Show that either  $H_1 = H_2$  or  $H_1$  contains all even-element subsets of V and  $H_2$  contains all odd-element subsets of V (or conversely).

- **20\***. (a) Let  $G_1$ ,  $G_2$  be two simple graphs and suppose they have the same number of homomorphisms into any third graph H. Then  $G_1 \simeq G_2$ .
- (b) Suppose  $G_1$ ,  $G_2$  are two simple graphs and every graph H has the same number of homomorphisms into  $G_1$  as into  $G_2$ . Then  $G_1 \simeq G_2$ .
- **21.** If  $G_1 \times G_1 \simeq G_2 \times G_2$ , then  $G_1 \simeq G_2$ .
- **22.** (a) Construct simple graphs  $G_1$ ,  $G_2$ , F without isolates such that  $G_1 \ncong G_2$ , but  $G_1 \times F \cong G_2 \times F$ .
  - (b) Suppose  $G_1 \times F \cong G_2 \times F$ , where  $G_1$ ,  $G_2$ , F are simple graphs. Then

$$G_1 \times F_0 \simeq G_2 \times F_0$$

for all subgraphs  $F_0$  of F.

- (c) Prove that the cancellation law holds for strong direct products, i.e. if  $G_1 \cdot H \simeq G_2 \cdot H$ , then  $G_1 \simeq G_2$ .
- (d) The same cancellation law holds for direct products of finite groups and rings.

# II. Hints

# § 1

- 1. Decide about the persons one by one.
- **2.** Decide now about the postcards. The answer to the second question is n!S(k, n).
- 3. If we form all permutations of the letters, how often does the same word occur?
- **4.** (a) Imagine k 1-forint coins in a row and suppose the people come one by one and pick up forints as long as you allow them.
  - (b) Reduce to the previous case by borrowing one forint from each person.
- 5. One can decide about the different kinds of postcards independently.
- 6. A recurrence relation is

$$S(n, k) = S(n - 1, k - 1) + kS(n - 1, k)$$

- 7. If x is an integer, both sides count the number of mappings of an n-element set into an x-element set.
- 8. Substitute

$$j^n = \sum_{r=0}^n S(n, r) j(j-1) \dots (j-r+1).$$

9. (a) Use

$$P_n = \sum_{k=0}^{\infty} S(n, k).$$

(b) Prove that the distribution of terms in the sum in (a) is asymptotically normal; if we set

$$g_n(y) = \frac{1}{\sqrt{2\pi}} y^{n-y-1/2} e^{y-1} \qquad (\sim y^n/ey!),$$

then

$$rac{g_nig(\lambda(n)\,+\,y\,\lambda(n)/\sqrt{n}ig)}{g_nig(\lambda(n)ig)} o e^{-y^2/2} \qquad (n o\infty).$$

- 10. Consider the partition class containing a given element.
- 11. Use the previous recurrence relation to derive a differential equation for p(x).
- 12. (a) For a partition, consider all permutations in which the classes are arranged in order of non-decreasing lengths.
  - (b) The result is  $\approx (e-1)^n$ .
- 13. Consider the Taylor expansion of  $e^{e^x}$ .
- 14. (a) The second derivation of the formula for p(x) (1.12) can be modified to apply here.
- (b) The method of the first proof of the formula for p(x) (1.11) easily extends to this case.
- 15. (a) Use the recurrence relation

$$S_n = \sum_{k=1}^n \binom{n}{k} S_{n-k}.$$

(b) Use

$$S_n = \oint_{|z|=\varepsilon} \frac{s(z)}{z^n} dz.$$

- 16. Represent the partition  $n = a_1 + \ldots + a_s$ ,  $a_1 \ge \ldots \ge a_s$  by a diagram in which the  $i^{\text{th}}$  column consists of  $a_i$  dots.
- 17. Associate the partitions  $n = a_1 + \ldots + a_m$  and  $n m = (a_1 1) + a_m$  $+\ldots+(a_m-1)$  with each other.
- 18. Let

$$n = a_1 + \ldots + a_m$$

be a partition of n into distinct terms. Write

$$a_i = 2^{\beta_i} b_i$$
, where  $b_i$  is odd.

19. If

$$n = a_1 + \ldots + a_m, \ a_1 > \ldots > a_m \ge 1$$

is a partition of n, then try to form a partition into distinct terms with one more term by subtracting 1 from  $a_1$  and taking a new term 1. This will not work in general as we may have  $a_1 - 1 = a_2$ . In that case, also subtract 1 from  $a_2$  and let the new term be 2, etc.

20. It is

$$S(x) = \frac{1}{(1-x)(1-x^2)(1-x^3)\dots}.$$

21. Represent each of the generating functions as a product similarly as in the preceding solution.

22. The difference between the numbers of partitions of n into an odd and an even number of distinct terms is the coefficient of  $x^n$  in

$$(1-x)(1-x^2)(1-x^3)\dots$$

- 23. Identity (a) expresses the following fact that the number of partitions into distinct odd terms is equal to the number of partitions with symmetric Ferrer diagrams. (Symmetric means axially symmetric with respect to the line starting from the left lower corner and ascending at 45°).
- 24. Decide, starting with 1, which numbers with what multiplicity must occur in such a partition. If 1 occurs  $k_1$  times, the next least number occurring is  $k_1 + 1$ .

25. If

$$n = x + y + z$$

is a partition of n, then x + y, y + z, x + z are the sides of a triangle with circumference 2n.

- **26.** Find a 2-step recurrence relation on  $B_n$ ; guess a 1-step recurrence relation and prove it by induction.
- **27.** Find a recurrence relation for  $A_n$ .
- 28. Deduce a recurrence relation for  $C_n$  and prove that the upper determinant satisfies it as well as the initial values. As initial values, take  $C_{-k+1} = \ldots =$  $=C_{-1}=0, C_0=1.$ 
  - (b) Use the recurrence relation in the same way as in 1.27.
- 29. Prove the recurrence relation

$$p_n(\lambda) = \lambda p_{n-1}(\lambda) - p_{n-2}(\lambda)$$

for the characteristic polynomial

$$p_n(\lambda) = \det (\lambda I - A).$$

- **30.** Denote by  $x_n$  and  $y_n$  the number of such sequences starting with a or b and c or d, respectively, and find recurrence relations on these numbers.
- 31. Use ideas similar to those in the solution of 1.17.
- 32. The graph of such a function can be considered as a polygon, joining the point (1,1) to the point (n,n), moving to the right or one upwards at each step.
- **33.** It is more convenient to deal with the monotonic mappings f of  $\{1, \ldots, n\}$ into  $\{0, \ldots, n-1\}$  with f(x) < x. The mappings not having this property are in a one-to-one correspondence with step-functions connecting (0, 1) to (n+1, n-1). The result is  $\frac{1}{n+1} \binom{2n}{n}$ .

to 
$$(n+1, n-1)$$
. The result is  $\frac{1}{n+1} \binom{2n}{n}$  .

34. Denote by g(n, r) the number in question. Establish and then use the recurrence relation

$$g(n, r) = \sum_{k=0}^{r} {r \choose k} g(n-1, k).$$

**35.** Consider the procedure backwards.

**36.** Use induction as follows: remove an element of S, then the sequence of partitions of the remaining set is almost a sequence arising by the procedure of the problem.

HINTS

- 37. Reduce (b) to the preceding problem.
- 38. Use 1.37 or 1.33.
- **39.** Associate a bracketing of the product  $x_1 cdots x_{n-1}$  with a triangulation.
- 40. Use the recurrence relation

$$D_n = \sum_{k=2}^{n-2} D_k D_{n-k+1}.$$

- 41. The triangles form a chain.
- **42.** (a) Evaluate  $\sum_{k=0}^{\lfloor (n-1)/2 \rfloor} {n \choose 2k+1}$  at the same time.
  - (b) Consider the Pascal triangle.
  - (c) Find a combinatorial interpretation.
  - (d) This is the coefficient of  $x^m$  in  $(1-x)^u(1+x)^u$ .

(e) 
$$\binom{k}{m} \binom{n}{k} = \binom{n}{m} \binom{n-m}{n-k}$$
.

(f) Set  $\varepsilon = e^{\frac{2\pi i}{7}}$  and use

$$\sum_{j=0}^{6} \varepsilon^{kj} = \begin{cases} 7 & \text{if } 7|k, \\ 0 & \text{otherwise} \end{cases}$$

- (g) Find a recurrence relation for these numbers (z fixed, n varies).
- (h) Prove by induction that the result is  $(-1)^m \binom{n-1}{m}$ .
- (i) The result is  $\binom{u+v+1}{m}$ .
- 43. (a) Find a combinatorial interpretation.
  - (b) Substitute -n-1 for n in (a).
  - (c) Substitute

$$\binom{n+k}{p+q} = \sum_{j=0}^k \binom{n}{p+q-j} \binom{k}{j}.$$

- (d) Substitute -n-1 for n in (c).
- (e) Expand  $\binom{x+t}{n} \binom{x}{n}$  by 1.42c and i.
- **44.** To prove (a), differentiate both sides with respect to y and use induction on n. (b), (c) will follow from (a).
- **45.** Use  $f_n(x) = n! \binom{x}{n}$ .

#### \$ 2

- 1. The number of pupils who like either mathematics or physics is not 12 + 14. By how much is 26 too large?
- **2.** Determine the contribution of any atom of the Boolean algebra generated by  $A_1, \ldots, A_n$  on each side.
- 3. Let  $S = \{1, \ldots, n\}$  and let  $A_i$  be the set of integers divisible by  $p_i$ .
- **4.** Count the number of mappings of a set of k given elements onto a set of n given elements.
- **5.**  $p \sigma^1 p + \ldots$  vanishes for  $x_i = 0 \ (i = 1, \ldots, n)$ .
- **6.** Consider the "contribution" of an atom B of the Boolean algebra generated by  $A_1, \ldots, A_n$ .
- 7. (a) Apply the preceding result.
  - (b) Use

$$\sigma_j = E\left(egin{pmatrix} \eta \ j \end{pmatrix}
ight)$$

- 8. Sum the results of the preceding problem for  $p, p + 1, \ldots$
- 9. This goes back to the partial sums of

$$\binom{k}{0} - \binom{k}{1} + \binom{k}{2} - \dots$$

10. To solve the second half, show

$$\sigma_r \leq \frac{1}{r} \, \sigma_{r-1} + (r+1) \, \sigma_{r+1}.$$

11. For each  $J \subseteq \{1, \ldots, n\}$ ,

$$P(A_J \cdot \prod_{i \notin J} \bar{A}_i) \ge 0.$$

12. Denote by  $\mathfrak{I}_1$  and  $\mathfrak{I}_1'$  the set of independent even subsets and the set of odd subsets spanning at most one edge. Prove that

$$|\mathfrak{I}_0'| \ge |\mathfrak{I}_0|, \ |\mathfrak{I}_1'| \ge |\mathfrak{I}_1|$$

simultaneously.

- 13. If  $P(A_1) = \ldots = P(A_n) = 1$ , then the terms  $(-1)^{|I|}$  and  $(-1)^{|J|}$  with  $J = I \cup \{n\}$  cancel out; the remaining terms are positive.
- 14. Use 2.6. A lower estimate is the following:

$$P(\bar{A}_1 \dots \bar{A}_n) \ge 1 - \sum_{\substack{I, J \subseteq \{1, \dots, n\} \\ \max I = \max J}} \lambda_I \lambda_J P(A_{I \cup J}),$$

where  $\lambda_0 = 0$ ,  $\lambda_{\{k\}} = 1$  for  $1 \le k \le n$  and  $\lambda_I$  is an arbitrary real number for  $|I| \ge 2$ .

15. Let  $q_i = 1 - p_i$  and  $q_I = \prod_{i \in I} q_i$  and show that

$$p_{I \cup J} = p_I p_J \sum_{K \subseteq I \cap J} \frac{q_K}{p_K}.$$

16. Set 
$$P(A_K)=p_K+R_K$$
,  $\mathscr{H}=\left\{K:p_K\geq \frac{1}{M}\right\}$ , minimize 
$$\sum_{I,J\in\mathscr{R}}\lambda_I\,\lambda_J\,p_{I\cup J}\qquad (\lambda_{\mathfrak{o}}=1)$$

and estimate

$$\sum_{I,\ I\in\mathcal{R}} \lambda_I \, \lambda_J \, R_{I\cup J}$$

at this place.

17. Let  $M = \sqrt{x}/\log^2 x$  and let  $P_1, \ldots, P_n$  be those primes  $\leq M$  which do not divide k. "Sift out" those numbers from the given sequence which are divisible by one of  $P_1, \ldots, P_n$ , using Selberg's method.

18. Prove that

$$P(A_1|\bar{A}_2\dots\bar{A}_n) \leq \frac{1}{2d}$$

by induction on n.

19. Let  $\xi$  be the number of  $A_i$ 's that occur. Use Chebyshev's (or Markov's) inequality

$$P(\bar{A}_1, \ldots \bar{A}_n) = P(\xi = 0) \le \frac{1}{\sigma_1^2} E((\xi - \sigma_1)^2).$$

- 20. To compare the result of 2.19 with the estimate of Selberg's method, expand the latter as a power series in p.
- **21.** Only the statement concerning the inverse is non-trivial. Towards this, show that  $(a_{ij})$  is invertible iff  $a_{ii} \neq 0$  (i = 1, ..., n).
- **22.** What will the matrix  $M = (m_{ij})$ , where  $m_{ij} = \mu(x_i, x_j)$ , be equal to?
- 23.  $\mu(a, b)$  depends only on the structure of the interval  $\{z : a \le z \le b\}$ .
- 24. Show that  $\mu^*$  defined by (\*) satisfies the identities in the definition of  $\mu$ .
- 25. Use the fact that  $(Z-I)^n=0$ .
- **26.** (For the second part) Choose V to be the set of all subsets of  $\{1, \ldots, n\}$  and

$$f(K) = P(\prod_{i \notin K} A_i \prod_{j \in K} A_j).$$

27. Use

$$\sum_{x \le b} \mu(0, x) = \sum_{a \le b_1 \le b} \sum_{a \lor x = b_1} \mu(0, x).$$

28. Let a be an atom such that  $a \leq x$  and consider

$$\sum_{y \vee a = x} \mu(0, y) = 0.$$

29. Decompose

$$\sum_{x} \sum_{v \geq x} \mu(x, y)^{r}$$

into terms according to whether x, y belong to A, B, C.

30. Use 2.27 and induction on n.

31. Define

$$f_{ij} = \begin{cases} f(x_i) & \text{if} \quad i = j, \\ 0 & \text{otherwise.} \end{cases}$$

What is the relationship between the matrices  $F = (f_{ij})$  and  $G = (g_{ij})$ ?

**32.** Use the set  $\{1, \ldots, n\}$  ordered partially by divisibility and a slight modification of 2.31.

**33.** Find a similar representation  $D = Z^T A Z$  of the matrix  $(d_{ij}) = D$  as for the matrix G in the solution of 2.32.

**34.** Write M as a polynomial in Z-I, in a way similar to the one in 2.25.

**35.** Denote by  $q_k(x)$  the number of those k-tuples in R whose union is x. Show that

$$q_0(x) - q_1(x) + q_2(x) - q_3(x) + \ldots = \mu(0, x).$$

36. Use the identity of 2.29 in an argument similar to the preceding one.

37. Use 2.27 and induction on r.

# § 3

1. Two permutations are conjugate iff their cycles have the same cardinalities.

2. (a) Use inclusion-exclusion;

(b) build up such a permutation, starting from a given point.

3. Count those permutations in which the set of elements in the cycle containing 1 is given.

4. Count similarly as before.

5. Determine the expected number of points in k-cycles.

6. Use the coding of the second solution of 3.3.

7. Prove the recurrence relation

$$np_n(x_1, \ldots, x_n) = \sum_{k=1}^n x_k p_{n-k}(x_1, \ldots, x_{n-k}).$$

- 8. For (b), represent these permutations by a permutation of the set of classes and by k permutations, permuting each of the classes.
- 9. If the corresponding elements of the two groups have the same order, then their regular representations have the same cycle index.
- 10. Observe the identity

$$q_n(x_1,\ldots,x_n)=p_n(x_1,\ldots,x_n)+p_n(x_1,-x_2,\ldots,(-1)^{n-1}x_n).$$

11. Consider

$$\sum_{n=0}^{\infty} (p_n - p_{n-1}) y^n.$$

- 12. (a)  $f_n(x) = n! p_n(x, \ldots, x)$ , where  $p_n(x_1, \ldots, x_n)$  is the cycle index of  $S_n$ .
  - (b) Evaluate  $f'_n(1)$ .
- 13. Such a graph consists of disjoint circuits. If we orient these, we get the graph of the cycle decomposition of a permutation of n elements.
- 14. (a) Join the two numbers in a pair by an edge.
  - (b) How many ways can one orient the preceding graph?
- 15. Express these numbers as certain values of the cycle index.
- 16. The sequence  $\overline{\pi}(1), \ldots, \overline{\pi}(n)$  uniquely characterizes the permutation  $\pi$ .
- 17. If the jumper to jump at the  $i^{\text{th}}$  time executes the  $(\pi(i))^{\text{th}}$  longest jump, then  $\pi$  is a random permutation of  $\{1,\ldots,n\}$ . The  $i^{\text{th}}$  jump is a record iff  $\overline{\pi}(i)=i$ .
- 18. Observe that the optimal strategy to follow is independent of the order of the first k jumps. Determine  $p_{n-1}, p_{n-2}, \ldots$
- **19.** This is equivalent to  $\overline{\pi}(j+1) \geq \overline{\pi}(j)$ .
- **20.** The number of inversions of  $\pi$  is

$$(\overline{\pi}(1)-1)+(\overline{\pi}(2)-1)+\ldots+(\overline{\pi}(n)-1).$$

- 21. Suppose we start with a large reserve in our tank and use it, if running out of gas. Consider the point where the reserve needed is maximal.
- **22.** (a) Call a sequence  $(j_1, \ldots, j_s)$  of different integers  $j_i$   $(1 \le j_i \le n)$  ascending, if
  - (i)  $x_{i_1} + \ldots + x_{i_s} > 0$ , but
  - (ii)  $x_{j_1} + \ldots + x_{j_{\nu}} \le 0$  for  $1 \le \nu < s$ .

(For s=1, the one-element sequence  $(j_1)$  is ascending iff  $x_{j_1} > 0$ .) Call the sequence descending if

- (i')  $x_{i_1} + \ldots + x_{i_s} \leq 0$ , but
- (ii')  $x_{j_{\nu}} + \ldots + x_{j_{s}} > 0$  for  $1 < \nu \le s$ .

(Note that the two definitions are not completely analogous.) Consider partitions of  $\{1, \ldots, n\}$  into ascending and descending sequences, and build up permutations  $\pi$  from them for which  $a(\pi)$  and  $b(\pi)$ , respectively, can be determined easily, cf. also 3.21.

- (b) Put n = 2m,  $x_1 = \ldots = x_m = 1$  and  $x_{m+1} = \ldots = x_{2m} = -1$  in (a).
- **23.** (a) Count the number of k-gons invariant under a given rotation.
  - (b) Use a similar method.
- **24.** Count the number of elements of  $\Gamma$  fixing a given point.
- 25. This is only a slight generalization of 3.23.
- **26.** We want to determine the number of orbits of the permutation group induced by  $\Gamma$  on the set  $\Omega$  of mappings of D into R.
- 27. First find a formula involving F; then use

$$F(u_1, u_2, \ldots) = F\left(\frac{\partial}{\partial z_1}, \frac{\partial}{\partial z_2}, \ldots\right) e^{\sum_{i=1}^{\infty} u_i z_i}\Big|_{z_i=0}.$$

- 28. Use a method similar to the previous one; a one-to-one mapping which is a fixed point of  $(\pi, \varrho)$  must map a cycle of  $\pi$  onto a cycle of  $\varrho$  with the same length.
- **29.** First determine the generating function of the numbers  $q_n(\gamma)$  of mappings f satisfying (\*) and invariant under a given  $\gamma \in \Gamma$ .
- **30.** If D is the set of forints, R the set of people and  $\Gamma$  the symmetric group on D, 3.26 gives the answer.
- **31.** Put |D| = n,  $|R| = N \ge n$ ,  $\Gamma = \{1\}$  and  $\Gamma_1 = S_N$  in 3.27; let  $N \to \infty$ .

#### § 4

- 1. Supposing  $d_n = 1$ , remove the point  $v_n$ .
- 2. Use the preceding result and the binomial theorem.
- 3. Show that

$$p_n(x_1,\ldots,x_{n-1},0)=(x_1+\ldots+x_{n-1})\,p_{n-1}(x_1,\ldots,x_{n-1})$$

and use the identity 2.5.

- **4.** Contract each  $T_i$  onto a single point  $v_i$ .
- 5. Try to reconstruct the sequence  $b_1, \ldots, b_{n-1}$  of removed points; observe that  $b_i$  is the least number not occurring among  $b_1, \ldots, b_{i-1}, a_i, \ldots, a_{n-1}$ .
- 6. Remove one edge of the tree in every possible way.
- 7. (a) Prove that

$$\left(\frac{t(x)}{x}\right)' = t'(x)\frac{t(x)}{x}.$$

(b) Use the fact that

$$T_n = \frac{1}{n} t^{(n)}(0) = \frac{(n-1)!}{2\pi i n} \oint \frac{t'(z)}{z^n} dz,$$

where C is any simple closed curve around the origin.

8. Use 4.1. The result is 
$$\frac{n!}{(n-l)!}S(n-2,l)$$
.

9. (a) Use the Binet-Cauchy formula:

$$\det A_0 A_0^T = \Sigma (\det B)^2,$$

where B ranges over all  $(n-1)\times(n-1)$  submatrices of  $A_0$ .

- (b) Two rows of  $A_0$  have at most one common non-zero entry.
- (c) The number of trees on n points is the number of spanning trees of  $K_n$ , the complete n-graph.
- 10. Write D in the form  $A_0 A_0^T$  and use an argument similar to that in the solution of 4.9.
- 11. This is the number of spanning trees of the complete bipartite graph on  $\{v_1, \ldots, v_n; w_1, \ldots, w_m\}$ .
- 12. (a) Use the inclusion-exclusion principle and 4.4.
  - (b)-(c) Use 4.9 or 4.12a.
- 13. (a) Consider the binary tree with k+1 endpoints as a diagram to calculate a product with k factors.
- (b) Imagine that the edges of such a tree are walls. Starting from the root, walk around the wall, keeping it on your left. Write a 1 when walking along an edge away from the root and a -1 otherwise. Which  $\pm 1$ -sequences arise in this way?
- 14. Use 4.4. Alternatively prove the recurrence relation

$$E(n, k-1) = \left(1 - \frac{1}{k}\right) nE(n, k).$$

- 15. Trivial: the edges entering the points  $\neq a$  can be chosen independently.
- 16. (a) Use induction, by splitting the edges entering  $v_1$  into two classes  $C_1$ ,  $C_2$  and considering the graphs  $G-C_1$ ,  $G-C_2$ .
  - (b) Consider the  $a_{ij}$ 's as variables and substitute  $a_{ij} = x_i$ .
- 17. If there is a fixed point x, orient such a tree T to get an arborescence  $\vec{T}$  rooted at x. Contract all orbits of  $\pi$  and investigate the image of  $\vec{T}$ .
- 18. To prove the upper bound consider rooted plane trees.

19. (a) Write the right-hand side in the form

$$x \prod_{T} (1 + x^{|V(T)|} + x^{2|V(T)|} + \ldots),$$

where T ranges over all isomorphism types of trees.

- (b) Denote by  $W_n^{(d)}$  the number of non-isomorphic rooted trees in which the root has degree d. Express first the generating function of the numbers  $W_n^{(d)}$  for a fixed d in terms of w(x) by the Pólya–Redfield method.
- **20.** (a) w(x) satisfies the equation

$$w(x) e^{-w(x)} = \varphi(x),$$

where

$$arphi(x) = x \exp\left(rac{w(x^2)}{2} + rac{w(x^3)}{3} + \ldots\right)$$


is analytic in a larger circle than w.

(b) Show that

$$w''(x) = B_1(\tau - x)^{-3/2} + B_2(\tau - x)^{-1/2} + h(x),$$

where h(x) is continuous on the (closed) disc  $|x| \leq \tau$ .

- 21. Each expansion term of per A counts the number of 1-factors parallel to a given one.
- 22. Find a recurrence relation.
- 23. Use inclusion-exclusion.
- 24. One has to prove that all terms in pf B have the same sign.
- 25. The even circuits used in the previous solution are all alternating with respect to some 1-factor.
- **26.** Those expansion terms of  $\det B$  which correspond to a permutation with at least one odd cycle will cancel out.
- 27. Prove that there is an orientation such that, if we go around the boundary of any bounded face in the positive sense, we pass an odd number of edges agreeing with its orientation.
- 28. Orient the ladder as in Fig. 13 and apply 4.21 and 4.23.



**29.** (a) Write  $a_n^2$  in the form

 $\det p_n(A)$ ,

where

$$p_n(\lambda) = egin{bmatrix} -\lambda & 1 & 0 \ 1 & -\lambda & & \ & & \ddots & \ & & & 1 \ 0 & & 1 & -\lambda \ \end{bmatrix}, \quad A = egin{bmatrix} 0 & 1 & 0 \ -1 & 0 & & \ & & \ddots & \ & & & \ddots & \ & & & 1 \ 0 & & -1 & 0 \ \end{bmatrix}.$$

Use 1.29 to obtain the roots of  $p_n(\lambda)$ .

- (b) Observe that the formula (a) is the resultant of two polynomials and use its Sylvester determinant form.
  - (c) Using formula (a),  $\frac{\log a_n}{n^2}$  will be a sum approximating

$$\frac{1}{\pi^2} \int_{0}^{\pi} \int_{0}^{\pi} \log (4 \cos^2 x + 4 \cos^2 y) \, dx \, dy$$

- **30.** Consider the  $n \times n$  chessboard as consisting of every second square of the  $(2n-1)\times(2n-1)$  chessboard.
- **31.** Identify  $u_i$  and  $v_i$ .
- 32. Find a recurrence relation.
- **33.** Consider  $a_n$  as the permanent of a matrix and expand it by its first row. Find simultaneous recurrence relations on  $a_n$  and some other analogous numbers.
- 34. This number is

Expand by the first row.

- 35. Use inclusion-exclusion.
- 36. The number is the permanent of the matrix

n	n	
[11 11	0000	
11 11	10 00	
	i	n
11 11	11 10	
01 11	11 11	
		n
00 01	11 11	10
00 00	11 11	

Count the expansion terms having k entries from the upper left block.

- 1. (a) What is the sum of the degrees of a graph?
  - (b) The title of the chapter is misleading, use divisibility by 3.
  - (c) How are the first and last two points joined to each other?
- 2. Use the solution of 5.1a.
- 3. If every circuit is even, then the lengths of any two paths between a pair x, y of points have the same parity.
- **4.** If the total work around any circuit is 0, the work needed to go from x to y is independent of the path used.
- 5. If there is an even cycle C, start defining a 2-coloration of G by first 2-coloring C and extending.
- 6. Take a maximum closed trail in G and show that it contains all the edges.
- 7. (a) Trivial by 6.
  - (b) Note that  $G_{k,n} = L(G_{k-1,n})$ .
- 8. This means the same as " $G_{k,2}$  has a Hamiltonian cycle".
- **9.** Look at the sections of an Euler trail between two consecutive occurrences of a point x of degree at least 3.
- 10. To show that we have used every edge consider an edge of T not used and nearest to  $x_0$ .

- 11. The Euler trails can be considered as starting from  $x_0$  through a specified edge. Show that every Euler trail arises by the algorithm given in 5.10, from some spanning arborescence.
- 12. Call a point "good" if all edges incident with it have been traversed in both directions. Look at the first "bad" point met during the walk.
- 13. Show and then use the fact that a graph with even degrees and without isolated points is the union of edge-disjoint circuits.
- 14. (a) Use 5.13 and 5.6.
  - (b) Take a new point and join it to all points with odd degree.
- 15. Use induction on the number of edges.
- 16. Use induction on the number of edges; observe that if  $G_1$  is a spanning subgraph with even degrees and K is a circuit, then  $(V(G), E(G_1) \triangle E(K))$  is a spanning subgraph with even degrees (briefly; a "good" subgraph) as well.
- 17. (a) Use induction on n. Remove a point of odd degree and take the complement within the set of its neighbors.
  - (b) Add a new point and join it to all points of V(G).
- 18. (a) Take all r-element matchings of the complete graph on V(G) and sieve out those having an edge from G.
  - (b) By (a).
- (c) The number of 1-factors has the same parity as the determinant of the adjacency matrix.
- 19. Use an argument similar to the one above. For undirected graphs, the statement is true if |V(G)| > 3.
- 20. Show that reversing an edge, the parity of the number of Hamiltonian paths does not change.
- **21.** Consider pairs  $(F_1, F_2)$  of 1-factors such that  $F_1 \cap F_2 = \emptyset$ ,  $e \in F_1$ . Show that the number of such pairs is even, and make use of this.
- 22. Reduce the graph as shown in Fig. 14 and distinguish Hamiltonian circuits going through the edges in different directions.




Fig. 14

- **23.** Let F be a spanning tree of G. Show that those edges of  $G^*$  which do not correspond to edges of F form a spanning tree of  $G^*$ .
- 24. Use the previous proof.
- 25. Each face has at least three edges on the boundary.
- 26. Remove the edges on the boundary of a face.
- 27. Use the 2-coloration of the faces.
- 28. (a) Consider a 2-coloration of the faces.
- (b) Take a new point outside the pentagon and join it to the points of odd degree.
- 29. Count the number of edges joining a red point to a blue point.
- **30.** Color a point x red, if  $x \in V_1$  and  $V_1$  contains an (x, a)-path; blue, if  $x \in V_2$  and  $V_2$  contains an (x, b)-path; green otherwise. Show that there is no triangle with all three colors.
- **31.** Only the proof of (f) is somewhat difficult. Take a basis  $\{\mathbf{v}_1, \ldots, \mathbf{v}_k, \mathbf{v}_{k+1}, \ldots, \mathbf{v}_n\}$  of V with  $\langle \mathbf{v}_1, \ldots, \mathbf{v}_k \rangle = M$  and consider the transformation A defined by

$$A: \mathbf{e}_i \to \mathbf{v}_i$$
.

- **32.** Show and then use the fact  $\langle M \cup M^{\perp} \rangle = (M \cap M^{\perp})^{\perp}$ .
- 33. (a) The stars generate the subspace of cuts, the orthogonal subspace consists of those sets of edges which constitute a subgraph with even degrees.
  - (b) Use 4.9.
- 34. Show that each circuit C is the sum of those faces contained in C.
- **35.** (a) Show that  $\sum_{i \in I \cup j} C_i \subseteq K$ .
- (b) Consider a set I such that  $K = \sum_{i \in I} C_i$  is a circuit,  $|I| \ge 2$  and |I| is minimal.
  - (c) Use (b) and induction on f.
  - (d) By (c) and 5.34.
- 36. Use MacLane's criterion.
- **37.** (a) Supposing  $G=G_1\cup G_2,\ V(G_1)\cap V(G_2)=\{x,y\},$  contract  $G_1-x$  and  $G_2-x.$ 
  - (b) Consider a maximum path.
- (c) Remove the chord of the circuit, drow the rest in the plane and choose the circuit so that the number of regions inside be maximal. Prove that the graph has only chords outside the circuit.
  - (d) Clear from (c) and 5.25b.
- **38.** It suffices to consider triangulations. (Prove!) Find an edge *e* which is contained in exactly two triangles. Contract *e* and use induction.

- 1. Addition of an edge decreases the number of components by at most one.
- 2. (a) Let  $T_1, \ldots, T_{c(G_1)}$  be the components of  $G_1, S_1, \ldots, S_{c(G_2)}$  the components of  $G_2$ . Construct a graph  $G^*$  with  $V(G^*) = \{t_1, \ldots, t_{c(G_1)}, s_1, \ldots, s_{c(G_2)}\}$  joining  $t_i$  to  $s_j$  iff  $T_i \cap S_j \neq \emptyset$  and look at the components of this graph.
  - (b) Add the points of  $V(G_1) V(G_2)$  to  $G_1$  as isolated points.
- **3.** Let  $x_i$  have degree  $d_i$ . Consider a component  $G_1$  of G not containing  $x_n$ ; this has points of degree less than  $|V(G_1)|$  only!
- 4. If  $G_1$  contains an odd circuit (say) and is connected, show that  $G_1$  contains a walk of any large enough length connecting any two points, and make use of this fact.
- 5. Cf. 5.1b.
- 6. (a) Consider a longest path.
  - (b) Consider an endpoint of a longest path again.
- 7. (a) One can increase the number of common edges of  $T_1$  and  $T_2$  at each step.
- (b) Use "backwards" induction on the number of edges in a common subtree of  $T_1$  and  $T_2$ .
- 8. (a)-(b) Use 6.7b.
- **9.** To show that there is an (a, b)-path, consider the set X of all points accessible from a.
- 10. Contract the red edges and remove the green ones.
- 11. (a)-(b)-(c). If F is a minimal set whose removal (contraction) yields a digraph with the desired property, then the inversion of F also results in such a graph.
  - (b) Consider the case |F| = 1 first.
- (c) Show first that the points of an acyclic graph G can be ordered in such a way that every edge has larger head than tail.
- 12. To show that a strongly connected tournament contains a Hamiltonian cycle, consider a maximal cycle.
- 13. Consider a longest cycle which is not a Hamiltonian cycle.
- 14. Invert the edges of F.
- 15. Find a proper subtree mapped into itself by  $\varphi$ .
- 16. Consider two points of the intersection and the path connecting them.
- 17. (a) Assuming  $P_1$ ,  $P_2$  are disjoint maximum paths, consider a  $(P_1, P_2)$ -path Q.
- (b) The middle point of a fixed maximum path is contained in every other maximum path.

- **18.** Use induction on k or n. In the first case find an edge separating  $G_k$  from  $G_1 \cap \ldots \cap G_{k-1}$ .
- 19. Only (c) is non-trivial. In the case of d(x, y) consider a minimum (x, y)-path and a minimum (y, z)-path; in the case of D(x, y), consider a maximal (x, z)-path.
- **20.** Take an edge and count how many paths of type  $(p_i, p_j)$ ,  $(p_i, q_j)$ ,  $(q_i, q_j)$  contain it.
- **21.** (a) Show that removing all points of degree 1, d(x) decreases by 1 for all of the remaining points.
  - (b) Consider a longest path with endpoint x.
- **22.** (a) Moving from x to y, for which points z does d(x, z) increase and decrease?
  - (b) Easy from (a).
  - (c) Take a long path and a large "fan" at one of its ends.
- **23.** To determine the minimum of  $\sum_{x,y} d(x,y)$  observe that exactly n-1 terms are equal to 1; can all the others be equal to 2? On the other hand, show that s(x) is maximal when G is a path and x an endpoint of it.
- **24.** Associate, with each of all but k of the points, a path of length k starting from it, such that different points are associated with different paths.
- **25.** Let G be the tree constructed by the algorithm, and H a tree of minimum expense having the largest possible number of edges in common with G. Assuming  $G \neq H$ , consider an edge of G not in H.
- 26. Use an argument similar to the one in 6.25.
- 27. Two edges are equivalent iff they lie on the same circuits!
- **28.** Choose subgraphs  $G_1, \ldots, G_i$  satisfying the condition and show that if  $G_1 \cup \ldots \cup G_i \neq G$  you can choose a  $G_{i+1}$ .
- **29**. The non-trivial part is that if G is 2-edge-connected it has a desired orientation. Use 6.27 or 6.28.
- 30. Find a circuit with which all but one of the edges are well-fitted and contract it, and proceed by induction.
- **31**, To show that if  $e_1$ ,  $e_2$  lie on a circuit  $C_1$  and  $e_2$ ,  $e_3$  lie on a circuit  $C_2$ , then  $e_1$ ,  $e_3$  also lie on a circuit, consider the common points of  $C_1$  and  $C_2$  next to  $e_3$  on  $C_2$  in both directions.
- **32**. In the cycle (i)  $\Rightarrow$  (iii)  $\Rightarrow$  (ii)  $\Rightarrow$  (i) only the first step is non-trivial. Prove (iii) for two adjacent edges first!
- **33.** Find and use a structure for G similar to the one in 6.27. Another possibility is to select 3 paths  $P_1$ ,  $P_2$ ,  $P_3$  connecting some two points such that the length of  $P_1$  is minimal and show that  $P_1$  satisfies the requirements.

- **34.** Arrange the points of G in a sequence such that p is the first point, q is the last point and from any other point there are edges connecting it both to earlier and to later points.
- 35. A chord of a circuit could be removed without violating 2-connectivity.
- **36.** Select a (C, C)-path P with endpoints x, y for which the arc R connecting x and y on C is minimal. Then the neighbor z of x on R has degree 2.
- **37.** If G' were connected, G would contain a circuit whose points of degree 2 would constitute an arc.
- 38. Identify the corresponding endpoints of two isomorphic trees.
- **39.** (a) Use induction on |E(G)|. If there is an  $S \subseteq V(G)$  which defines a k-element (a, b)-cut and  $|S| \ge 2$ ,  $|V(G) S| \ge 2$ , then apply the induction hypothesis on the graphs obtained by contracting S and V(G) S, respectively.
- (b) Split each point  $x \neq a$ , b into two points  $x_1, x_2$ , where  $x_1$  is joined to  $x_2$ , and  $x_2$  is joined to  $y_1$  iff x is joined to y.
  - (c) Replace each (undirected) edge by two oppositely oriented edges.
- **40**. Take two new points a and b; join a to all points of A, b to all points of B.
- **41.** A common generalization is the following. If  $A, B \subseteq V(G)$  are disjoint sets and there is a positive integer "capacity" w(x) associated with each point x, and for every set S meeting all (A, B)-paths we have

$$\sum_{x \in S} w(x) \ge k \,,$$

then there are k(A, B)-paths such that each point x is contained in at most w(x) of them.

**42.** Choose k independent (a, B)-paths  $R_1, \ldots, R_i$  such that  $R_t$  ends at the some point as  $P_t$  and

$$|E(R_1 \cup \ldots \cup R_k) - E(Q_1 \cup \ldots \cup Q_{k+r})|$$

is minimal.

- **43.** Choose  $B = V(P_0)$  and apply 6.42.
- 44. Knowing Menger's theorem it is easy to see that G will consist of two complete graphs  $G_1, G_2$  with k common points,  $a \in V(G_1), b \in V(G_2), a, b \notin V(G_1 \cap G_2)$ . To prove this without Menger's theorem show that every point is connected to a or to b and the points connected to both separate a and b.
- 45. None of them is true.
- 46. Use Menger's theorem.
- 47. For (i): Remove the edges of an (a, b)-path (if any) and apply 6.46.
- 48. By straightforward counting of the edges on both sides.
- **49.** Consider a set X with  $\delta_G(X) = k$  and |X| minimal.
- **50.** Contract those edges not in  $F_1, \ldots, F_m$  and consider the resulting graph.

- 51. Let  $U \subset V(G) \{x\}$  be a maximal set such that  $\delta_G(U) = k$  and a certain neighbor y of x belongs to U. Show that there exists a neighbor z of x not in U.
- **52.** Apply the previous "point-splitting" repeatedly until x disappears.
- 53. Use 6.48c instead of 6.48a.
- **54.** (a) is trivial. For (b), find a point of degree 2k and "split" it as in the solution of 6.52.
- 55. Consider a quadrilateral.
- **56.** (a) Show that there are  $\alpha + \beta$  edge-disjoint paths connecting  $\{a, b\}$  to  $\{a', b'\}$  such that exactly  $\alpha$  of them start at a and exactly  $\alpha$  of them (some other  $\alpha$ , possibly) ends at a'.
  - (b) Use a similar argument to the one as in 6.46 and 6.47.
- 57. Look for a planar example.
- **58.** C obviously separates a and b. To show |C| = k, consider the set D of points accessible from b on a  $(b, A \cup B)$ -path and show that

$$C \cap D \subseteq A \cap B$$
.

As another possibility, use Menger's theorem.

- **59.** If (x, y) is an edge of H and T a (k 1)-element separating set of G (x, y), then  $T \cap V(H)$  separates H (x, y).
- **60.** (a) Assume the statement is false. Let  $G-A=G_1\cup G_2,\ G-B==G_3\cup G_4,\ (G_1\cap G_2=G_3\cap G_4=\emptyset)$  and assume, e.g., that  $G_1\cap G_3\neq\emptyset,\ V(G_1)\cap B\neq\emptyset.$  Let  $a\in G_1\cap G_3,$  and let C be the set of points of  $A\cup B$  accessible from a on an  $(a,A\cup B)$ -path. Show that

$$\begin{split} &1^{\circ} \ |C| > k, \\ &2^{\circ} \ C \subseteq \left(A \cap V(G_3)\right) \cup (A \cap B) \cup \left(B \cap V(G_1)\right), \\ &3^{\circ} \ G_2 \cap G_4 = \varnothing, \\ &4^{\circ} \ |V(G_4)| < |V(G_1)|. \end{split}$$

- (b) Show that  $G_1$  defined in (a) is a one-element set.
- **61.** Set e = (a, b) and assume both G/e and G e are only 2-connected. Show
  - (a) there are two points x, y in G e separating a and b,
  - (b) there is a point u in G such that  $\{a, b, u\}$  separates x and y,
  - (c) if  $\{x, y\}$  separates a from u, then x, y and b are the only neighbors of a.
- **62.** Show that if G is a simple graph and e is an edge of G which connects points with degree at least k and, moreover, G/e is k-connected, then so is G.
- 63. Use induction on the length of the circuit.
- **64.** Let a be a point of degree 3, joined to  $b_1$ ,  $b_2$ ,  $b_3$ . Show that, if  $G \neq K_4$ , there is a point  $u_i$  separating  $b_{i+1}$  and  $b_{i+2}$  in  $G a b_i$  (i = 1, 2, 3;  $b_{j+3} = b_j$ ), and then use this fact.

**65.** Draw a tree T without points of degree 2 in the plane so that its endpoints are on its convex hull; then add the edges of the convex hull C. Show that this graph G is a line-critically 3-connected graph.

6.65

- **66.** Use induction on k. If  $x_1, \ldots, x_{k-1}$  are on C, but  $x_k$  is not, consider k independent  $(x_k, C)$ -paths.
- 67. Assume no circuit contains all the three given edges. Find a circuit which contains two of them and does not touch the third one, and, as in 6.66, investigate the paths joining the third edge to this circuit.
- 68. Use an argument similar to the one in 6.66.
- 69. Show that every bridge of the boundary of a face contains all points of the circuit.
- **70.** Let  $f \in E(G)$  and select a circuit C of G f such that the bridge B of C containing f is maximal.
- 71. Assume  $v(e_1) \ge \ldots \ge v(e_m)$ , where  $E(G) = \{e_1, \ldots, e_m\}$ , and consider the first index k such that  $\{e_1, \ldots, e_k\}$  contains an (a, b)-path.
- 72. Determine a function  $\varphi$  such that, for each point x, there is an (a, x)-path with value  $\varphi(x)$ .
- **73.** Let C be given by  $S \subseteq V(G)$ . Consider

$$\sum_{x \in S} \left( \sum_{e=(x,y)} f(e) - \sum_{e=(y,x)} f(e) \right).$$

**74.** The non-trivial part of the proof is to find an (a, b)-flow f and an (a, b)-cut C such that

$$w(f) = \sum_{e \in C} v(e)$$
.

Consider a flow f of maximum value, satisfying  $f \leq v$ . For each  $e \in E(G)$ , we introduce a new edge e' having the same endpoints but converse orientation, and let

$$v_0(e) = \begin{cases} v(e) - f(e) & \text{if } e \in E(G), \\ f(e_1) & \text{if } e = e_1', e_1 \in E(G). \end{cases}$$

We then consider the digraph  $G_1$  determined by those edges e, e' for which  $v_0(e) > 0$  and  $v_0(e') > 0$ ; respectively. Show that  $G_1$  is not connected between a and b.

- 75. (a) Let  $\omega_n$  be the increase of flow value at the  $n^{\text{th}}$  step; then  $\sum_{n=1}^{\infty} \omega_n$  must be convergent. Try to achieve  $\omega_n = \alpha^n$ ,  $\alpha < 1$ . Also, there must be edges with  $f_k(e) = 0$  or v(e) at each step. How can the value of  $f_k(e)$  change between two successive equal states?
- (b) Show that if we omit those edges which the paths  $P_k$  and  $P_{k+1}$  use in different directions,  $P_k \cup P_{k+1}$  still contains two (a, b)-paths  $Q_1, Q_2$  with  $Q_1 \cap Q_2 \subseteq P_k \cap P_{k+1}$ .

- (c) When an edge is used in opposite directions by  $P_k$  and  $P_{k+l}$ , then  $P_{k+l}$  is longer than  $P_k$ .
- 76. Introduce two new points  $a_0$ ,  $b_0$ ; join  $a_0$  to each point x with  $u(x) \ge 0$  and give the new edge e the capacity v(e) = u(x); join each point x with u(x) < 0 to  $b_0$  and give the new edge e the capacity v(e) = -u(x). The problem is equivalent to finding a suitable  $(a_0, b_0)$ -flow in the resulting digraph  $G_1$ .
- 77. (a) Substitute v(e) parallel edges for every edge e and apply Menger's theorem.
- (b) Demonstrate the existence of an (a, b)-flow  $f(e) \leq v(e)$  and an (a, b)-cut C such that

$$\sum_{e \in C} v(e) = w(f)$$

first for the case of rational capacities v(e).

**78.** One may assume k=2. Consider an integral (a, b)-flow f' such that  $w(f') \ge w_1$ ;  $f'(e) \le f(e)$  and  $\sum f'(e)$  is minimal.

#### \$ 7

- 1. Show that a minimum system of edges which cover all points consists of disjoint stars.
- 2. One can use Menger's theorem (6.39). In order to get a direct proof, consider a minimal subgraph G' of G such that  $\tau(G') = \tau(G)$  and show that G' consists of independent edges.
- 3. From each point y of F in A, there is a path alternating relative to M connecting y to  $A_1$ .
- **4.** To get a direct proof, use induction on |A|; try to find a subset  $\emptyset \neq A_1 \subset A$  such that  $|\Gamma(A_1)| = |A_1|$ .
- 5. Use an argument similar to that in the first solution of 7.4.
- 6. (a) Observe that

$$\Gamma(X_1 \cup X_2) = \Gamma(X_1) \cup \Gamma(X_2), \quad \Gamma(X_1 \cap X_2) \subseteq \Gamma(X_1) \cap \Gamma(X_2).$$

- (b) Consider a minimal subgraph  $G_1$  with  $V(G_1) = V(G)$  and property (2). Observe that (1) means that the one-element sets satisfy (2) with equality.
- 7. The implications (i)  $\Rightarrow$  (iii)  $\Rightarrow$  (ii)  $\Rightarrow$  (i) are easily verified.
- 8. "If" is easy by the preceding result. Conversely, fix a 1-factor F, an edge of F as  $G_0$  and choose the paths  $P_1, \ldots, P_k$  alternating with respect to F (cf. the solution of 6.28).
- **9.**  $P_k$  in the previous problem cannot be a single edge!
- 10. Prove it for r-regular graphs first. Show that G has a 1-factor, and remove the edges of this 1-factor.

- 11. Split each point x into  $\left[\frac{d(x)}{k}\right]$  points of degree k and, if necessary, one point of degree  $d(x) k \left[\frac{d(x)}{k}\right]$ .
- **12**. Use 7.11 with k = r.
- 13. Define a graph  $G_1$  on V(G) by joining two points iff they are joined in G by more than one edge. For which values of r can  $G_1$  fail to satisfy the condition given in 7.4b?
- 14. To show we cannot go into an infinite cycle, consider all edges which x passes infinitely many times and show that every point is adjacent to at most two of them.
- 15. Distinguish between elementary and non-elementary graphs.
- 16. Reduce to a network-flow problem, similar to that in the first solution of 7.2.
- 17. Embedding into a G' of this size is straightforward. To show that smaller G' would not do, count the edges leaving a color-class of G by two different methods.
- 18. It should be shown that, if  $\widetilde{G}$  is the bipartite graph on V(G), where  $x \in A$  is joined to  $y \in B$  iff they are not adjacent in G, then  $\widetilde{G}$  has an (n-2d)-factor.
- 19. Construct a bipartite graph G on  $\{u_1, \ldots, u_n, v_1, \ldots, v_n\}$ , where  $u_i$  is adjacent to  $v_j$  iff  $a_{ij} > 0$ . One has to show that this bipartite graph has a 1-factor.
- **20.** (a) If G has a 1-factor the expansion term corresponding to this cannot cancel out as the entries are algebraically independent.
- (b) If G has no 1-factor the columns of  $(a_{ij})$  are linearly dependent. Consider a minimum set of columns which are linearly dependent.
- **21.** If G is bipartite, the conditions (1) are not linearly independent.
- **22.** Suppose not. Let F be a maximum matching and u, v two points not in F. How many edges can join  $\{u, v\}$  to any edge of F?
- 23. Consider a maximum matching F which has as many edges in common with  $F_0$  as possible.
- **24.** (a) Use induction on k.
  - (b) Consider the decomposition given in 6.27. You may use 6.30 as well.
- (c) Use this sharpening of (b): If G has a unique 1-factor  $F_1$  then it has a cut-edge which belongs to F.
- **25.** Consider a maximum matching F and select an edge from each point not covered by F.
- **26.** Show by induction on the distance between x and y that  $v(G x y) < \langle v(G) \text{ for each } x \neq y \in V(G).$

- **27.** (a)  $\delta(G) \ge \max \ldots$  is easy. To establish  $\le$ , use induction and the previous result.
  - (b) Trivial from (a).
- 28. (a) One has to show that "adjacency" is an equivalence-relation in  $G V_1$ .
  - (b) Adding edges does not disturb the inequalities of the theorem.
- 29. (a) Apply Tutte's theorem.
  - (b) A cut-edge of a 3-regular graph must be contained in every 1-factor.
- **30**. Tutte's condition is satisfied. This follows from counting the edges and removed edges in a way similar to that in the solution of 7.29a.
- 31. Use an argument similar to that in 7.8, selecting paths  $P_0, P_1, \ldots, P_k$  one by one so that they alternate relative to a given maximum matching F.
- **32.** (a) Let  $x \in A_G$  and let  $z \in D_G$  be a neighbor of x. Supposing indirectly that  $y \notin D_G$ , but some maximum matching M' of G x avoids y, consider a maximum matching M of G avoiding z and form  $M \cup M'$ .
  - (b) Remove all points of  $A_G$ , using (a).

For (c) — (e), use (a) and (b).

- 33. From any matching  $M_0$  of G' we get a matching with  $|M_0|+k$  edges in G. For the proof of the converse, consider the Gallai-Edmonds structure described in the preceding problem.
- **34.** If two outer points in the same component are adjacent we have an odd circuit as in 7.33. Otherwise, remove the inner points.
- **35.** Consider the components of the set  $D_G$  in 7.32 and use an argument similar to that in 7.29.
- **36.** (a) Let F be a maximum matching and x a point not covered by F.
- (b) Find an edge e such that e is in some, but not all, 1-factors and G e is 2-connected (cf. 6.36).
- 37. It is easy to see that no 2-matching can be larger than the value given. To find a 2-matching of this size, take two copies G, G' of G, where  $V(G) = \{v_1, \ldots, v_n\}$  and  $V(G') = \{v'_1, \ldots, v'_n\}$  and define a bipartite graph  $G_0$  by joining  $v_i$  to v' iff  $(v_i, v_i) \in E(G)$ .
- 38. First consider factor-critical graphs and then use the Gallai–Edmonds structure theorem 7.32.
- **39.** Consider an Euler trail of G.
- 40. Consider a pseudosymmetric orientation (5.13) and split each point into two, separating the outgoing and incoming edges. Apply 7.10 to this bipartite graph.
  - **41.** Add a new point x and join it to all points of odd degree adding, if necessary, a loop at x.
  - 42. (a) The number of points must be even!
    - (b) G is a contraction of a 2-connected 3-regular graph.

- **43.** (a) Replace each point by f(x) independent points.
- (b) If f(x) = 1 for at least one endpoint of any edge, then an f-matching is an f-factor.
- **44.** Split each point x into two points x', x'' incident with the incoming and outgoing edges, respectively.
- **45.** Subdivide each edge and define f to be 1 for the new points. Look for an f-factor of the resulting graph.
- 46. Use induction.
- 47. Use induction.
- **48.** Show that the sequence  $d_1, \ldots, d_{n-2}, d_{n-1} 1, d_n 1$  satisfies the same conditions.
- 49. Use 7.44.
- **50.** Transform G with degrees  $d_1, \ldots, d_n$  to a graph with the same degrees in which  $v_n$  is adjacent to  $v_{n-d_n}, \ldots, v_{n-1}$ .
- **51.** (a) Choose H with as many pairs of oppositely directed edges as possible. Show that those edges of H whose converse is not in H can form neither an even cycle nor two disjoint odd cycles.
  - (b) Trivial by (a) and 7.49.
- 52. To show the sufficiency of the condition, consider a graph having these degrees and reduce the number of components one by one.
- **53.** Let G be one graph on  $V = \{v_1, \ldots, v_n\}$  with  $d_G(v_i) = d_i$  and G' another with  $d_{G'}(v_i) = d_i 1$ . Choose G' and G as "close" as possible.

### \$ 8

- 1. Consider a maximal independent set.
- 2. Let  $S_1$  be a maximum independent set of G and let  $S_{i+1}$  be a maximum independent set in  $G S_1 \ldots S_i$ . Show that  $S_{i+1}$  has a matching into  $S_i$ .
- 3. Find a circuit or an edge which contains a point x and all neighbors of x.
- **4.** If  $(x, y) \in E(G)$ , where  $x, y \in S$ , consider G x.
- 5. (a) Take any maximal independent set.
  - (b) Use induction, removing a point with indegree 0 and its neighbors.
  - (c) Use 5.3.
- 6. Consider the point of highest outdegree.
- 7. Remove a point and all points which can be reached from it on an edge, and use induction.
- 8. (a) Observe that whether or not the player to move can win depends only on his opponent's last move; call this last move a "winning move" (for the

opponent) if he cannot win. What are the characteristic properties of the set of winning moves?

- (b) The point  $x_0$  is only involved if A starts with it.
- (c) Observe that when they have once left a component they will make no more moves there.
- **9.** If G has no 1-factor, A should select a maximum matching and start with a point not on this.
- 10. Use induction on k.
- 11. Use an argument similar to the one in the previous problem.
- 12. Remove an edge incident with the point x.
- 13. It is enough to show this, if a single point x is replaced by two adjacent points  $x_1$ ,  $x_2$  both joined to the same "old" points as x.
- 14. It is enough to embed the given graph  $G_0$  into a G such that  $\alpha(G-e) > \alpha(G)$  holds for the edges of  $G_0$ .
- 15. (a) For r = 2, odd circuits do. For r > 2, increase the degree by substituting complete graphs for points.
  - (b) The tetrahedron and icosahedron.
- 16. An  $\alpha$ -critical bipartite graph consists of independent edges. Supposing there was an  $x \in V(G)$  joined to two points  $y_1, y_2$ , consider a maximum independent set  $S_i$  of  $G (x, y_i)$  and the subgraph induced by  $S_1 \triangle S_2 \cup \{x\}$  (cf. the solution of 7.2).
- 17. Follow the previous solution.
- 18. If S were a minimal cutset spanning a complete graph and  $x \in S$ , consider two edges joining x to different components of G S.
- 19. (a) Show that  $\alpha(G) = \alpha(G_1) + \alpha(G_2)$ .
- (b) Suppose  $G = G_1 \cup G_2$  with  $V(G_1) \cap V(G_2) = \{x, y\}$ ,  $|V(G_i)| \ge 3$ . Determine, for each  $X \subseteq \{x, y\}$  and i = 1, 2, the maximum size of an independent set in  $G_i$  which meets  $\{x, y\}$  in X, by writing out several inequalities involving these numbers.
- 20. Apply 8.10 with the set of all maximum independent sets.
- 21. Apply 8.11.
- **22.** Let  $T_1, \ldots, T_k$  be the set of all maximum independent sets containing x; apply 8.11 again.
- 23. Use the previous result.
- **24.** These are the complete graphs and odd cycles only; show and use the fact that, if T, S are maximum independent sets with  $T \cap S \neq \emptyset$ , then  $\Gamma(T \cup S) = V(G) T S$ .

- 25. Use 8.19. and the previous problem.
- 26. Consider an α-critical spanning subgraph.
- 27. Again consider an  $\alpha$ -critical spanning subgraph of G.

- 1. Use induction on |V(G)| removing a point.
- 2. To show the second statement, let  $V_1$  be the set of points of a maximum clique.
- **3.** Use pairs of colors to color  $G_1 \cup G_2$ .
- **4.** Consider a coloration  $\alpha$  such that each color occurs at least twice and a coloration  $\beta$  with  $\chi(G)$  colors, so that they have as many color classes in common as possible.
- 5. Use induction on k, and a re-coloration similar to the one in the solution of 9.2.
- **6.** It helps to answer the question: can  $\alpha(G \oplus K_m)$  be larger than |V(G)|?
- 7. (a) If a coloration of  $G_1$  is given, color a point of  $G_1 \times G_2$  according to its first coordinate.
  - (b) Observe that  $G \times G \supseteq G$ .
- (c) If  $\alpha$  is a k-coloration of  $G \times K_n$ , n > k, use the fact that some color occurs twice in the set  $\{(v, x); x \in V(K_n)\}$  for each  $v \in V(G)$ .
- (d) For an *n*-coloration  $\alpha$  of  $G \times K_n$ , show that if  $\{\alpha(v, x) : x \in V(K_n)\}$  are different, then so are  $\{\alpha(u, x) : x \in V(K_n)\}$  for any neighbor u of v.
  - (e) What about  $K_n^m = \underbrace{K_n \times \ldots \times K_n}_{m}$ ?
- 8. For (d), observe that, if we want to 3-color an odd circuit but one of the colors is excluded at each point, then it is possible iff the excluded color is not the same at every point. For (e), consider first the case when all but one partition are allowed.
- 9. First prove this for acyclic graphs; for the general case, see the solution of 6.10.
- 10. If and only if every circuit has the same number of edges oriented in each direction around the circuit.
- 11. If  $\vec{G}$  is an orientation of G with the property mentioned then, for a walk of G, define the expense of it by saying that we gain one if we go along an edge in the right way (from tail to head) but lose k-1 otherwise. Verify and then use the following:
  - (a) the expenses of walks from a to b are bounded from below,
- (b) if a, b are adjacent, then the minimum expenses of  $(x_0, a)$ -walks and  $(x_0, b)$ -walks are different but differ by at most k-1.

- 12. (a) Find an arrangement  $(x_0, \ldots, x_{n-1})$  of the points of G such that  $x_i$  is adjacent to some  $x_i$  (j < i); start by coloring  $x_{n-1}$ .
- (b) Show that the sequence used in the previous solution can be chosen so that  $x_0$  is adjacent to  $x_{n-1}$  and  $C(x_0) \neq C(x_{n-1})$ .
- **13.** (a) Consider  $x_0, x_{n-2}, x_{n-1}$  such that  $(x_{n-2}, x_{n-1}) \notin E(G), (x_0, x_{n-1}) \in E(G), (x_0, x_{n-2}) \in E(G).$
- (b) If  $G = G_1 \cup G_2$  with  $V(G_1) \cap V(G_2) = \{x_1, x_2\}, |V(G_i)| \ge 3$ , consider  $G_1 + (x_1, x_2)$  and  $G_2 + (x_1, x_2)$ .
- 14. (a) One has to show that "to be non-adjacent" is an equivalence relation.
  - (b) Embed G in a maximal graph on the same set with the same property.
- (c) Consider the set of those colorations coloring the endpoints of a given edge *e* differently.
- 15. Let  $K_{k+1}$  be the smallest complete graph in  $\Re$ ; then  $\Re$  will be the class of all non-k-colorable graphs. To show that these are, in fact, in  $\Re$ , consider a counterexample with a fixed number of points but maximum number of edges and prove that "to be non-adjacent" is an equivalence relation on the points.
- 16. Use the preceding result.
- 17. (a) Odd circuits.
  - (b) In every 3-coloration of  $K_{n,n'}$  one of the two classes is monochromatic.
- (c) In every 5-coloration of  $K_{n,n'}$  one of the two classes receives only two colors.
- 18. To show  $\chi(G') > \chi(G)$ , let  $\alpha$  be a coloration of G' with colors  $1, \ldots, \chi(G)$  and  $\alpha(y) = 1$ . "Project"  $\alpha$  to get a coloration of G not using color 1.
- 19. (a) What would a 3-coloration of  $G (x_3, x_4)$  look like?
  - (b) Use 9.8 to show the sufficiency.
- **20.** It can happen for all  $k \geq 3$ .
- **21.** If k-1 edges separate G into two pieces  $G_1$ ,  $G_2$ , consider k-colorations of these and permute the colors to fit the two colorations together.
- 22. If G is critically (k+1)-chromatic and  $G = G_1 \cup G_2$  with  $V(G_1) \cap V(G_2) = \{x, y\}$ , then investigate the k-colorability of the graphs obtained from  $G_1$  and  $G_2$  by connecting or identifying the points x and y.
- **23.** Let  $G_1, \ldots, G_N$  be the components of G S. Consider the partitions of S induced by k-colorations of  $G V(G_i)$   $(i = 1, \ldots, N)$ .
- **24.** In every k-coloration of G-(x, y) the points x, y have the same color. If this color is i, then for any other color j, the set of points with color i and j must contain an (x, y)-path.
- **25.** (a) Use e.g. 9.18.
- (b) If we have  $G_k$  with  $\chi(G_k) = k$  and join each point x of it to a new point x' by an edge, then, n-coloring this piece, the set of new points cannot be monochromatic.

- **26.** A k-coloration of L(G) yields a coloration of the edges of G in which all colors form bipartite graphs.
- 27. Take iterated line-graphs of a transitive tournament.
- 28. (a) Remove an interval whose right endpoint is as far to the left as possible.
- (b) Consider  $I_1$  and a as before, but now remove all intervals containing a from the system.
  - (c) We may assume V(G) = V(C). Consider  $I_1$  and a as before.
- **29.** (a) Consider circuits formed by two arcs joining  $v \in S$  to  $u \in S$  through different components of G S.
- (b) The "if" part follows like 9.28c. For the "only if" part use (a) and induction.
  - (c) Use induction and (a).
- **30.** This is equivalent to  $\alpha(G) = \varrho(G)$ .
- 31. Show that

$$\chi(G_1 \cdot G_2) \leq \chi(G_1) \cdot \chi(G_2),$$
  
$$\omega(G_1 \cdot G_2) \geq \omega(G_1) \cdot \omega(G_2).$$

- 32. Heuristically, you have to partition P into "levels". Take the points on the top as "first level", the next largest points as "second level" etc.
  - (b) Use 8.4.
- 33. 9.31 does not yield a perfect graph, even if  $G_1$  and  $G_2$  are perfect.
- **34.** A color-class of an  $\omega(G)$ -coloration does so.
- 35. It suffices to prove the previous condition for G', in the case when only one  $G_x$  has more than one point.
- **36.** Consider the partitions of V(G) induced by  $\lambda$ -colorations.
- 37. Count the number of  $\lambda$ -colorations using inclusion-exclusion.
- 38. Which  $\lambda$ -colorations of G e are not  $\lambda$ -colorations of G?
- 39. For circuits, use the previous formula.
- **40.** If G has several components, we can color them independently. If it has several blocks, we can color them almost independently.
- 41. Remove the interval whose right endpoint is the farthest to the left.
- 42. Use 9.38.
- **43.** Use induction on |E(G)|, but now starting with a tree.
- 44. Use the formula given in the solution of 9.36.
- **45.** In fact,  $(-1)^{n-1}P_G(\lambda) > 0$  in (0, 1) for connected graphs. Show this by an induction similar to the one in the solution of 9.43.

- **46.** (a) Show that 0 is a simple root iff G is connected, 1 is a simple root iff G is 2-connected or  $\cong K_2$ .
  - (b) Use induction and 9.38.
- **47**. (a) Trivial by 9.38.
- (b) Any linear relation like this ought to be satisfied in the three cases when V(G) = V(F).
- (c) Show that this (in fact, any) linear relation holds generally if it holds for the three cases when V(G) = V(F).

**48.** What about 
$$\frac{3-\sqrt{5}}{2}$$
?

**49.** (a) Show that the relation holds when G is some very simple triangulation and reduce the problem to this case by "switching" the diagonals of quadrilaterals; use the identity

$$P_{G_1}^2(\tau+1) - P_{G_2}^2(\tau+1) = \tau^{-3} \left( P_{G_1|e_1}^2(\tau+1) - P_{G_2|e_2}^2(\tau+1) \right).$$

- (b) Use "backward" induction, removing edges from a triangulation.
- **50.** Show that G has a point of degree at most 5.
- 51. If the faces are colored with colors 1, 2, 3, 4, give color  $\alpha$  to edges adjacent to faces of color 1 and 2 or 3 and 4; color  $\beta$  to edges adjacent to faces of color 1 and 3 or 2 and 4; color  $\gamma$  to the rest.

For the converse, first color the regions of the maps formed by red-green and

red-blue edges in a 3-coloration of the edges.

- **52.** If the edges are 3-colored with 1, 2, 3, associate the number +1 with those points where 1, 2, 3 is the clockwise ordering of the edges incident with the point and associate -1 with the rest. For the converse, consider L(G) and 5.4.
- 53. Use two colors inside and two colors outside the Hamiltonian circuit.
- 54. (a) Use induction, removing a point of degree at most 4, coloring the rest with 4 colors and modifying this coloration to get only three colors in the neighborhood of the removed point.
- (b) Identify two non-adjacent points on the boundary of a face and apply an argument similar to the one in part (a).
- **55.** We may assume G is a triangulation. Then consider a 1-factor in its dual map  $G^*$ .
- **56.** For part "if", orient the edges in such a way that, if 1 unit of work is needed to pass a given edge in the given direction, then 0 (mod 3) units of work are needed to go round each face.
- 57. Suppose no line is parallel to the x-axis. Then color the points one by one, in the order of their ordinates.

- § 10
- 1. Otherwise, each block of G would consist of a single circuit.
- 2. (a) Consider a longest path.
  - (b) Use induction based on (a).
- 3. (a) Instead of (a), prove the slightly stronger statement: If a simple graph G has at most one point with degree at most 2, then G contains a subdivision of  $K_4$ .
  - (b) Reduce to (a) as before.
- **4.** Show and use the fact that G cannot contain a subdivision of  $K_4$  and an edge disjoint from it.
- 5. (a) The graph contains a subdivision of  $K_4$ .
  - (b) Use Kuratowski's theorem 5.39d.
- (c) The only exceptions to (a) when dropping the connectivity assumption are connected graphs whose blocks are complete 4-graphs.
- **6.** The point is that  $K_4$  has degrees at most 3.
- 7. Contract edges forming a connected subgraph as long as

$$\frac{|E(G')|}{|V(G')|} \ge \frac{m}{n}$$


holds for the resulting graph G'.

- 8. Use induction on m, and the preceding problem.
- 9. Use a method similar to the one in 10.8.
- 10. If one builds them up starting with a given point and a given circuit, respectively, when can the branches "grow together"?
- 11. The points which can be reached from a point  $x_0$  on paths of length at most  $\frac{g-1}{2}$  are distinct.
- 12. Supposing G(r, g) and G(r'; g 1) are available, where r' = |V(G(r, g))|, construct G(r + 1, g).
- 13. (a) If two points a, b were at a distance greater than g, remove a, b and match up their neighbors.
- (b) Treat the cases r = 2l, r = 2l + 1 separately. In the first case, there must be a circuit of length g through any given point. In the second case, one of any two adjacent points has this property.
  - (c) Use a counting similar to that in 10.11.
- 14. If G' has girth at least g,  $(u, v) \in E(G')$  and both x, y are at distance at least g 1 from  $\{u, v\}$ , then G' (u, v) + (x, u) + (y, v) has girth at least g.

- 15. (a) Use a finite projective plane.
- (b) Let the points of G be the lines and points on the hypersurface  $x_1^2 + x_2^2 + x_3^2 + x_4^2 + x_5^2 = 0$  in the 4-dimensional projective plane over GF(p) and connect a point to a line iff they are incident.
- 16. (a) Let Z be a minimum set of points representing all circuits and count the number of edges between Z and V(G) Z.
  - (b) Use (a) and an argument similar to that in 10.11.
- 17. Let  $G_1 = G$ ; if  $G_1, \ldots, G_i$  are already defined, then let  $C_i$  be a minimum circuit in  $G_i$ , and let  $G_{i+1}$  be obtained from  $G_i$  by removing the points of  $C_i$  and also all points not in circuits of  $G_i V(C_i)$  and by "smoothing out" points of degree 2. Consider an i for which  $G_i$  has largest girth.
- 18. (a) Use 10.4.
  - (b) By a similar argument to the one in the previous proof.
  - (c) A minimal 3-regular graph with girth  $\log_2 k$  has this property.
- 19. Use induction removing x.
- **20.** Consider a maximum path  $P = (x_0, x_1, \ldots, x_m)$ . We say that the path  $(x_0, \ldots, x_i, x_m, x_{m-1}, \ldots, x_{i+1})$  arises by deformation from P (provided  $(x_i, x_m) \in E(G)$ ). Let X be the set of all endpoints of paths arising by repeated deformation from P.
- 21. (a) One may suppose that adding any edge (x, y) to G a Hamiltonian circuit arises. Consider the neighbors of x and y on this Hamiltonian circuit.
  - (b) (c) prove (d) first.
- (d) The argument in the solution of (a) proves that if  $d_k + d_l \ge n$ , then  $x_k$  and  $x_l$  are adjacent. Consider a missing edge  $(x_k, x_l)$  with k + l maximal.
- 22. Show that if G is saturated and  $x_i$ ,  $x_j$  are non-adjacent points, then

$$d_G(x_i) + d_G(x_i) \le n + q - 1.$$

- 23. Find a circuit of length 2n.
- **24.** Join the two points x, y if necessary and subdivide the edge (x, y).
- **25.** Consider a maximum cycle C. First prove that  $|V(G)| > \frac{n}{2}$ , then consider a maximum path P of G V(C) and try to replace an arc of C by a path going through P.
- **26.** Consider a maximum circuit C, a component  $G_1$  of G V(C) and the neighbors of those points of C adjacent to  $G_1$ .
- 27. (a) Consider an endpoint of a longest path in G.
- (b) Let  $(x_0, \ldots, x_m)$  be a longest path. Consider first the case when there are edges  $(x_0, x_j)$  and  $(x_m, x_i)$  such that i < j.
- **28.** Use induction on n and 7.27b.



- 29. Any path P in G is contained in a subgraph as in Fig. 15.
- **30.** Use the fact that if x, y are adjacent points, then

$$d_G(x) + d_G(y) \leq n$$
.

31. Observe that

$$d_G(x) \leq \alpha(G)$$

for each point x.

- **32.** If a triple  $\{x, y, z\} \subseteq V(G)$  does not span a triangle of G or  $\overline{G}$ , then it contains exactly two points such that the two edges in the triple adjacent to any one of these points do not belong to the same G or  $\overline{G}$ .
- **33.** The edge (x, y) is contained in at least d(x) + d(y) n triangles.
- **34.** Use induction on m, starting with a complete k-graph.
- **35.** Consider the neighborhood of a point with maximum degree and use induction on k.
- **36.** (a) Count the number of "cherries" (3-point trees) in the graph in two different ways.
  - (b) Use a method similar to the one in 10.15.
- 37. Now count the number of  $K_{1,r}$ 's in G.
- **38.** (a) By induction on k, choose k disjoint s-sets  $A_1, \ldots, A_k$  such that any two points in distinct s-sets are adjacent (s is an appropriately large number) and show that many points outside are adjacent to at least t points of every  $A_i$ .
- (b) Remove points with degree  $\left(1 \frac{1}{k} + \frac{\varepsilon}{2}\right) \cdot |V(G)|$  repeatedly; prove that when one gets stuck the number of points is still large.
  - (c) By (b).
- **39.** Prove that if  $G \neq H_{n,k}$  does not contain  $G_0$ , then  $2k \cdot |V(G_0)|$  points can be removed from G so that the remaining graph  $G_1$  satisfies

$$|E(G_1)| - |E(H_{n_1,k})| > |E(G)| - |E(H_{n_1,k})|$$
  $(n_1 = |V(G_1)|).$ 

- **40.** (a) If a given complete k-graph is contained in "few" complete (k+1)-graphs, then we find "many" pairs (A, U) of k-sets of points such that A is a complete graph, U is not, and  $|A \cap U| = k 1$ .
  - (b) First prove that

$$\frac{N_k}{N_{k-1}} \ge \frac{\vartheta - k + 1}{\vartheta} \cdot \frac{n}{k}.$$

**41.** (a) Let  $d_1, \ldots, d_n$  be the outdegrees of points in T. Then T contains

$$\binom{n}{3} - \binom{d_1}{2} - \ldots - \binom{d_n}{2}$$

3-cycles.

- (b) Observe that each point is adjacent to at least one 3-cycle.
- **42.** Prove that it contains at least n-k+1 k-cycles for each  $3 \le k \le n$ .
- **43.** If we specify  $\left\lceil \frac{n}{2} \right\rceil$  independent edges

$$e_1, \ldots, e_{\left[\frac{n}{2}\right]},$$

there is at most one Hamiltonian path whose 1st, 3rd, ... edges are these.

- **44.** (a) The assertion says that, if  $n = 2^{k-1}$ , then there is a transitive subtournament with k points. Prove this by induction on k.
- (b) Using induction on k, observe that the set of points accessible from x contains at least

$$\prod_{i=0}^{k-2} \left[ \frac{d+1}{2^j} - 1 \right]$$

transitive subtournaments with k-1 points, where d is the outdegree of x.

(c) Prove by induction on n that there are at least  $\binom{n-2}{k-2}$  k-tuples that contain a 3-cycle.

- 1. What does an eigenvector mean combinatorially?
- **2.** (a) Use the fact that  $(1, \ldots, 1)$  is an eigenvector and the eigenvectors are pairwise orthogonal.
  - (b) Establish and use the identities

$$A_G = B_G B_G^T - dI, \ A_{L(G)} = B_G^T B_G - 2I.$$

- (c) The Petersen graph is merely  $\overline{L(K_5)}$ .
- **3.** We may assume that  $V(T) = \{x_1, \ldots, x_n\}$ ,  $e = (x_k, x_{k+1})$   $(1 \le k \le n-1)$  and that  $T e = T_1 \cup T_2$ , where  $V(T_1) = \{x_1, \ldots, x_k\}$  and  $V(T_2) = \{x_{k+1}, \ldots, x_n\}$ . Consider the expansion of det  $(\lambda I A)$  by its first k rows.

- 4. Use induction and the previous exercise.
- 5. (a) follows by 11.4 and 1.31. Both (a) and (b) follow by induction. Both (b) and (c) are essentially solved in 1.29.

11.4

- 6. By 11.4, we only have to find two trees with the same number of k-element matchings for every k. Look for them among those trees with no 3 independent edges.
- 7. (a) Prove that

$$p_G(\lambda) = \det p_{G_1}(\lambda I - A_{G_2}).$$

- (b) How does the adjacency matrix  $A_{G_1 \cdot G_2}$  arise?
- 8. The vector  $\mathbf{v} = (\chi(\gamma_{v_1, v_1}), \dots, \chi(\gamma_{v_1, v_n}))^T$  is an eigenvector.
- **9.** The automorphism group of  $Q_n$  contains the commutative regular subgroup  $(Z_2)^n$ ; one can also use 11.7a.
- 10. If  $G \simeq H$ , their eigenvalues are the same.
- 11. Note that 0 is not an eigenvalue iff det  $A \neq 0$ .
- 12. Find an eigenvector which associates a with those points of degree 3 in the covering subgraphs and b with the other points.
- 13. Use the fact that  $A = \max_{|\mathbf{v}|=1} \mathbf{v}^T A_G \mathbf{v}$ .
- 14. (a) Use the previous result; to determine the case of equality use the fact that the adjacency matrix of a connected graph has a (strictly) positive eigenvector.
  - (b) Use the fact that  $\Sigma \lambda_i = 0$  and  $\Sigma \lambda^2 = 2m$ .
- 15. Use the formula  $A_{L(G)} = B^T B 2I$ , where B is the point-edge incidence matrix:

$$B = (b_{ij}), b_{ij} = \begin{cases} 1 & \text{if } v_i \text{ belongs to the } j^{\text{th}} \text{ edge,} \\ 0 & \text{otherwise.} \end{cases}$$

- 16. To any automorphism  $\alpha$  of G there corresponds a permutation matrix P such that  $A_GP = PA_G$ . Also use the fact that  $A = T^{-1}LT$ , where T is an orthogonal matrix and L is a diagonal matrix with the eigenvalues in the diagonal.
- 17. Use the previous result to show that  $\Gamma$  is commutative and regular and apply 11.8 to calculate the eigenvalues.
- 18. Use the fact that if A is a symmetric matrix with eigenvalues  $\lambda_1 \leq \lambda_2 \leq \ldots \leq \lambda_n$  and B is a symmetric submatrix of it with eigenvalues  $\mu_1 \leq \ldots \leq \mu_m$ , then  $\lambda_i \leq \mu_i \leq \lambda_{i+n-m}$ .
- 19. (a) Suppose  $A_G \mathbf{w} = -\Lambda \mathbf{w}$ ,  $\mathbf{w} = (w_1, \dots, w_n)^T$ . Prove that  $\mathbf{w}' = (|w_1|, \dots, |w_n|)^T$  satisfies  $-\mathbf{w}^T A_G \mathbf{w} = \mathbf{w}'^T A_G \mathbf{w}'$ .
  - (b) Consider a component whose least eigenvalue is minimal.
- 20. Use 11.13, 11.14 and induction.

- 21. (a) Show that, with respect to a suitable basis, A has a  $k \times k$  symmetric submatrix D with 0's in the main diagonal and such that  $\Lambda$  is an eigenvalue of D.
  - (b) 0 is an eigenvalue with multiplicity n-k.
- 22. Write  $\mathbf{x}^T A_G \mathbf{x}$  as a sum of m positive and m negative squares.
- 23. If  $f(\lambda)$  is the polynomial whose roots are the distinct eigenvalues of  $A_G$ , then  $f(A_G) = 0$ .
- **24.** Show that both statements are equivalent to "every eigenvector of  $A_G$  is an eigenvector of J".
- 25. Calculate  $A_L^2$ .
- **26.** (a)  $A_G$  satisfies the equation

$$A_G^2 = bJ + (d - b)I$$
.

- (b)  $G = K_3$ .
- (c) Generalize the example of  $K_4 \oplus K_4$ .

- 1. The Petersen graph is the complement of the line-graph of  $K_5$ .
- 2. (a) Consider cubes of the type shown in Fig. 16.
  - (b) For the case of the cube, consider the 4 main diagonals.




Fig. 16

- 3. A k-circuit has some superflows automorphisms; how can we prevent it from changing the direction?
- **4.** Multiplication by any fixed element from the right is an automorphism of G.
- 5. Get rid of the directions and colorations by a method similar to the one in 12.3.
- **6.** Let  $\{h_1,\ldots,h_m\}$  be a minimal set of generators of  $\Gamma$ . Take the elements of  $\Gamma\times\{1,2\}$  as points and connect (g,1) to (g',2) iff  $g'g^{-1}=h_i$  for some i. Add edges within  $\Gamma\times\{1\}$  and  $\Gamma\times\{2\}$  to guarantee that there are no automorphisms other than those induced by the elements of  $\Gamma$  in the natural way.

- 7. (a) A group of even order has an element of order 2.
- (b) Find all tournaments which admit a regular group of automorphisms  $\cong \Gamma$ .

12.7

- 8. Trying the same kind of construction as in 12.5, it is easy to get all points not in  $V(G_0)$  to have degree 3. Split each point of  $G_0$  into points of degree 1 and connect them by a circuit.
- **9.** Let  $x_1, x_2, \ldots, x_n$  be an arrangement of the points of G such that for i > 1  $x_i$  is adjacent to some  $x_j$  with j < i. Consider the subgroups of automorphisms fixing  $x_1, \ldots, x_i$ .
- 10. It suffices to consider one orbit of A(G) on E(G).
- 11. First solve the case  $\Gamma_2 = \{1\}$ . Then consider a certain "product" of examples.
- 12. (a) Consider a maximal tree T such that  $T \cap \gamma(T) = \emptyset$  for every  $\gamma \in \Gamma$ ,  $\gamma \neq 1$ , and contract each  $\gamma(T)$ .
- (b) Let  $e \in E(R)$ ,  $G_1$  the subgraph formed by the edges which are images of e under  $\Gamma$  and try to contract the components of  $G_1$ .
- 13. (a) A commutative and transitive permutation group is regular. Fix an  $x_0 \in V(G)$  and consider the mapping  $\alpha(x_0) \mapsto \alpha^{-1}(x_0)$  ( $\alpha \in A(G)$ ).
  - (b) The *n*-cube almost does it.
- (c) Consider the n-cube  $Q_n$  and a vertex x of it. Add some edges between the neighbors of x and all edges which are forced by those automorphisms of  $Q_n$  preserving the directions of edges. What will the subgraph induced by the neighbors of x be?
- 14. Use an argument similar to the one in the solution of 6.49.
- 15. (a)-(b) Consider a minimal set X which is a component of G-T, for some minimum cutset T. Use 6.60a. The connectivity must be at least  $\frac{2}{3}(r+1)$ .
  - (c) Use 6.60a again.
- 16. Use Tutte's theorem 7.27 and 12.14.
- 17. The Cartesian product of a circuit and a path contains a Hamiltonian circuit.
- 18. The connected ones are the following: circuits, the platonic bodies, the line-graphs of the cube and the dodecahedron, graphs obtained from these by multiplying each edge by the same number, subdivisions of these examples by one point on each edge, the duals of the line-graphs of the cube and the dodecahedron, and the stars.
- 19. Use 12.12b and the previous solution.
- **20.** (a) Show that the Cartesian product of two p-1-paths and a  $K_2$  is contractible onto  $K_p$ .

- (b) One may assume  $\Gamma'$  acts edge-transitively and  $\Gamma$  does not act semiregularly. Use 10.8.
  - (c) Consider a simple group.
- **21.** Construct a simple graph G with  $A(G) \cong \Gamma$  and link it to  $\Omega$  in a suitable manner.
- 22. Every endomorphism of the Petersen graph is an automorphism.
- 23. Use the same idea as in the previous solution: any endomorphism is one-to-one on a minimal odd circuit.
- 24. (a) goes as the solution of 12.4.
- (b) When replacing colored edges by paths with attached pieces use many rigid graphs which have no homomorphisms into each other.
- **25.** (a) Either 0 or -1 remains an endomorphism.
  - (b) Modify the previous construction.

#### § 13

- **1.** Associate a bipartite graph G with H in the following way: Let U = V(H), W = E(H),  $V(G) = U \cup W$ , and join  $u \in U$  to  $w \in W$  iff  $u \in w$ .
- **2.** (a) If E is an edge such that  $H' = (H \{E\}) \setminus E$  is connected (hence  $\emptyset \notin E(H)$ ), then we can find an F with the desired property.
  - (b) Use induction, removing the edge E in the preceding problem.
- 3. Consider any circuit of length > 3 and choose two points which are consecutive on the circuit but there is a further point of the circuit which lies between them in the ordering defined by the path P.
- 4. Otherwise it is totally balanced!
- 5. Apply 7.4 and 7.6.
- **6.** Such a system of distinct representatives exists iff H has a system of distinct representatives and so does the hypergraph

$$V(H_T^*) = E(H),$$
  
 $E(H_T^*) = \{U_x : x \in T\},$ 

where  $U_x$  is the set of edges of H containing x.

- 7. Represent the situation by a flow problem rather than a matching problem.
- **8.** (a) The sets  $E_0 \triangle F$   $(F \in E(H))$  are distinct!
- (b) Let  $x \in V(H)$ , and consider the hypergraphs  $H \setminus x$  and H', where  $V(H') = V(H) \{x\}$ ,

$$E(H') = \{ E \in E(H) : E \cup \{x\} \in E(H), x \notin E \}$$

(c) Consider the hypergraphs  $H_1 = H - x$ ,  $H_2 = H - E(H_1)$ .

<sup>9</sup> Lovász

**9.** Consider the hypergraphs  $H_1 = H - x$ ,  $\overline{H}_2 = H - E(H_1)$ ,  $H_2 = \overline{H}_2 \setminus x$ , for some  $x \in V(H)$ .

13.9

- 10. (a) Prove, by induction on k, that there is a set  $X \subseteq V(H)$  with |X| = k such that  $H_X$  has at least k+1 distinct edges  $(1 \le k \le n-1)$ .
- (b) Form a graph similar to the one in the previous second solution. Prove that it is bipartite and contains no  $\Theta$ -graph.
- (c) Set  $H_1 = H \setminus x$ ,  $V(H_2) = V(H) x$ , and  $E(H_2) = \{E \in E(H) : x \notin E, E \cup \{x\} \in E(H)\}$ . Prove that either  $H_1$  satisfies the condition of the problem with k, or  $H_2$  does so with k-1.
- 11. Consider the edges containing a given  $x \in V$  and use induction.
- 12. (a) Average the number of edges contained in the classes over all partitions of V(H) into two classes.
  - (b) Average now over all r-colorations of V(H).
- 13. Establish and use the inequality

$$\sum_{x \in E} d(x) \le r + (m-1)k \quad (E \in E(H)),$$

where d(x) is the degree of the point x.

14. Suppose indirectly that m > n. Sum the inequalities

$$\frac{d(x)}{m-d(x)} < \frac{|E|}{n-|E|} \quad (x \in E).$$

- 15. (a) H satisfies the conditions of the previous problem.
- (b) If  $\mathbf{a}_1, \ldots, \mathbf{a}_m$  are the characteristic vectors of edges, then  $\mathbf{a}_i \mathbf{a}_j = \lambda$  for  $i \neq j$ .
- 16. How many times is a given element counted on each of the two sides?
- 17. (a) Suppose  $x \in F_1, \ldots, F_d, [x \notin F_{d+1}, \ldots, F_m]$  Take  $F_1 \{x\}, \ldots, F_d \{x\} m d$  times,  $F_{d+1}, \ldots, F_m d$  times and apply 13.16.
- (b) Suppose that  $|F_1| = \ldots = |F_{m-1}| = 2k$ ,  $F_m = \emptyset$ , and (indirectly)  $m = k^2 + k + 3$ . Verify that there are not more than k points of degree at least m k 1, but every  $F_j$   $(1 \le j \le m 1)$  must contain at least k + 1 such points.
- 18. (a) At most one of X and E(H) X can be an edge.
  - (b) Use 13.8b, 13.9, or 13.11.
- 19. Count the pairs  $(E_1, E_2)$  of edges such that  $E_2 \subset E_1$  and there is no  $F \in E(H)$  with  $E_2 \subset F \subset E_1$ .
- **20.** Use induction on n.
- 21. (a) Consider the chains constructed in the previous problem. For another proof, count in two different ways those permutations of V(H) which have an edge of H as beginning section.

- (b) The even case is easy. For n=2k+1, show that each edge must be of size k or k+1 and that, if  $X \subset Y$ , |X|=k, |Y|=k+1, then one of X, Y must belong to E(H).
- **22.** If A and B are maximum antichains, then  $A \lor B = \{x \in A \cup B : x \lessdot y \text{ for all } y \in A \cup B\}$  is a maximum antichain as well  $(x \lessdot y \text{ means } x \leq y \text{ and } x \neq y)$ .
- **23.** To prove (b) consider the subsets  $S_E$  ( $E \in E(H)$ ) which are comparable with the edge E only, and show that, if chosen properly, they form a clutter.

#### 24. Let

$$\binom{n}{p},\ldots,\binom{n}{p+k-1}$$

be the k largest binomial coefficients and let M be the hypergraph on V(H) whose edges are all the  $p, \ldots, (p+k-1)$ -tuples. Compare H with M in each of the symmetric chains of 13.20.

- 25. (a) Fix an edge E. What does it mean if no point of E occurs in all the edges?
- (b) Find j edges whose intersection has at most  $r (j-1)(\tau(H) 1)$  points, for j = 1, 2, ..., k.
- 26. Take two edges with one point in common if possible and a third edge not containing this point.
- 27. Take a minimal W such that any two edges of  $H_W$  meet. For every ordering of W, count the number of those points  $x \in W$  which are both first points and last points of edges of  $H_W$ .
- 28. (a) From any given point, there starts at most one  $A_i$ .
- (b) Count the edges which consist of consecutive points in cyclic permutations of V(H) in two different ways (cf. the second solution of Sperner's theorem 13.21a).
- **29.** (a) Select a set S such that  $|S \cap E| \ge 2$  for each edge E and |S| is bounded.
  - (b) If H is not  $\nu$ -critical, use induction on  $\nu$ .
- **30.** Choose points  $x_1, \ldots, x_t$  such that  $x_{i+1}$  is a point of maximum degree  $\delta_{i+1}$  of  $H x_1 \ldots x_i$ . Show that  $|E(H x_1 \ldots x_i)| \leq \tau^* \delta_{i+1}$ .
- **31.** (a) Write t = w r + 1,

and similarly for  $\binom{v}{r}$ ,  $\binom{w}{r-1}$ ,  $\binom{w}{r-1}$ ,  $\binom{w}{r-1}$ , and compare  $\binom{t+j-1}{j}$  with  $\binom{t+j-1}{j-1}$ .

(b) Consider a point x with minimum degree and consider the hypergraphs H-x, H-E(H-x). Use induction on r and |E(H)| on the basis of (a).

- (c) Suppose any two edges of H meet, where H is r-uniform. How many r-tuples can be contained in the sets V(H) E,  $E \in E(H)$ ?
- **32.** (a) Use an argument similar to the one in 13.21 and 13.27; for a given permutation of the points, there is at most one index i such that every element of  $A_i$  has smaller index than every element of  $B_i$ .
  - (b) This is a special case of (a).
- 33. Color the points one-by-one; when would one be forced to complete a monochromatic edge?
- **34.** Show first that if one point has degree 1 and the others have degree 2, then H is 2-chromatic.
- 35. Suppose H has no 1-element edges and no isolated points. Show that
  - (a) every point has degree at least 2,
  - (b) a point of degree at least 4 belongs to 2-element edges only,
  - (c) any two points belong to an edge.
- **36.** Supposing there was a 2-coloration, count the edges in two different ways; by the monochromatic pairs contained in them and also by the bichromatic pairs contained in them.
- **37.** To show the "if" part find a set  $W \subseteq V(H)$  such that either all of  $|W \cap E_1|, \ldots, |W \cap E_m|$  are even or one of them is odd.
- **38.** (a) Show that replacing an edge E by disjoint subsets  $E_1, \ldots, E_k \subset E$  does not influence the assumption.
- (b) Use the preceding characterization of hypergraphs with totally unimodular incidence matrix.
- 39. The incidence matrix of an odd circuit is not unimodular.
- 40. Show that a minimum counterexample would contain a connected spanning 2-uniform (hyper)graph.
- 41. Show that, coloring the points at random, the probability of getting a monochromatic edge is less than 1 [cf. 13.12a].
- **42.** Let  $|S| = 2r^2$  and let the  $\binom{2r^2}{r}$  r-tuples in S constitute the points of another hypergraph H; for each partition  $P = \{X, Y\}$  of S let those r-tuples contained in X or in Y form an edge  $E_p$  of H. What are  $\tau(H)$  and  $\tau^*(H)$ ?
- 43. Consider a random coloration again but estimate by 2.18.
- 44. (a) Consider a point with maximum degree.
  - (b) Remove the point with largest degree from every edge.
  - (c) Count the edges adjacent to the point of high degree.
- **45.** (a) Select a 2-element subset of each edge such that these subsets form a forest.
  - (b) For r=2, the triangle is such a hypergraph. Use induction on r.

- **46.** For (a), use a modification of the construction in 13.45; for (b), try to "multiply" two smaller constructions; for (c), look for a simple direct example.
- **47.** Select a k-element set X for each  $1 \le k \le r$  such that the number of edges containing X is more than  $\frac{|E(H)|}{r^k}$ .
- **48.** Setting  $V(H)=\{v_1,\ldots,v_n\},\; E(H)=\{E_1,\ldots,E_m\},$   $a_{ij}=\begin{cases} 1 & \text{if} \quad v_i\in E_j,\\ 0 & \text{otherwise,} \end{cases}$

and  $A = (a_{ij})_{i=1,j=1}^n$ , the value  $v^*(H)$  is the maximum of  $\mathbf{1}^T \mathbf{x}$  under the constraints  $A\mathbf{x} \leq 1$ ,  $\mathbf{x} \geq 0$ .


- 49. (a) Decompose directly, but take into consideration the multiplicities of points.
  - (b) Use 7.40.
- (c) There are optimal fractional covers (matchings) with rational weights. Reduce the least common denominator of the weights using (a) and (b).
- (d)  $2v^*(G)$  is the maximum size of 2-matchings.  $2\tau^*(G)$  is the formula given in 7.37 for  $\nu_2(G)$ .
- 50. Use a construction similar to the one in the previous solution.
- **51.** (a) To prove the second inequality observe that if  $S \subseteq V(G \otimes H)$  covers all edges of  $G \otimes H$ , then  $S \cap (V(G) \otimes F)$  has at least  $\tau(G)$  elements for every edge F of H.
- (b) "If" is easy by (a); to prove "only if" assume  $\tau_n(H) < n\tau(H)$  and let G consist of all (N-n)-element subsets of an N-set (N large).
  - (c) Use 13.30 to estimate  $\tau(H^p)$  from above.
- **52.** If a point X is adjacent to two distinct edges  $E_1$ ,  $E_2$ , take the "mean" of  $(\tau 1)$ -element covers of  $H \{E_1\}$  and  $H \{E_2\}$ .
- **53.** Take the "mean" of maximum matchings of  $H v_i$ ,  $v_i \in E_0 \in E(H)$ .
- **54.** One has to show that a  $\tau$ -critical balanced hypergraph consists of disjoint edges. Copy the  $2^{nd}$  solution of 7.2.
- 55. To show (i)  $\Rightarrow$  (iii), note that a minimum counterexample would be  $\tau$ -critical.
- **56.** Show and use the following fact: If a hypergraph H satisfies (iv), then every hypergraph  $H_0$  arising by multiplication of edges also satisfies (iv). (Multiplication includes multiplication by 0, i.e. removal.)
- 57. Suppose G is perfect; associate a hypergraph H with G such that  $L(H) \cong G$ .

## § 14

- 1. (a) A point x is either adjacent to at least  $\binom{k+l-1}{k-1}$  points or non-adjacent to at least  $\binom{k+l-1}{k}$  points.
- (b) The existence of  $R_2(a_1, a_2)$  follows from the preceding result. Use induction on k.
- **2.** (a) [ek!] = k[e(k-1)!] + 1.
- (b) The upper bound is trivial from 14.1a. To prove the lower bound consider the hypergraph H with

 $V(H) = E(K_n)$  and  $E(H) = \{E(A) : A \text{ is a complete } k\text{-subgraph of } K_n\}.$ 

- 3. (a) For r=2 we already know this. Use induction on  $a_1+\ldots+a_k$  and r.
- (b) Select an ordered subset  $(x_1, \ldots, x_m)$  of elements such that for each r-tuple  $1 \leq v_1 < \ldots < v_r < m$  of indices, all (r+1)-sets of the form  $(x_{\nu_1}, \ldots, x_{\nu_r}, x_{\mu})$   $(v_r < \mu \leq m)$  have the same color.
- **4.** Easy by induction on n.
- **5.** Consider a maximum red path P and two disjoint blue paths  $Q_1, Q_2$  such that  $Q_1, Q_2$  consist of (V(P), V(G) V(P))-edges, their endpoints are in V(G) V(P), they do not contain the endpoints of P and their length is maximal. Prove that  $Q_1 \cup Q_2 \cup P$  covers all points.
- **6.** (a) Let  $(x_0, \ldots, x_{2k})$  be a red circuit. Prove that  $(x_i, x_{i+2})$  is blue,  $(x_i, x_{i+4})$  is red and  $(x_i, x_{i+3})$  is blue  $(i = 0, \ldots, 2k; x_{2k+1+j} = x_j)$ .
- (b) Let  $(x_1, \ldots, x_{2k})$  be a red circuit. Prove that  $\{x_1, x_3, \ldots, x_{2k-1}\}$  and  $\{x_2, x_4, \ldots, x_{2k}\}$  induce complete subgraphs.
- (c) Show that there is a monochromatic M-circuit for some  $M \geq m$ , then use (a), (b).
- 7. Consider the configuration in Fig. 17, in which all lines have unit length.






Fig. 18




Fig. 19

- 8. Consider the arrangement of six triangles with sides 1, a, b shown in Fig. 18.
- 9. (a) Take a stripwise coloration.
  - (b) See Fig. 19.
- 10. (a) Consider the Cartesian product of two large simplices.
- (b) One may assume ab = 1, where a, b are the two side vectors of R. Color the point w with  $[w^2]$  modulo 4.
- 11. Let  $\{A_1, \ldots, A_k\}$  be the partition of  $\{1, \ldots, n\}$  in question. Form a complete graph on  $\{1, \ldots, n+1\}$  and color the edge (i, j) with color v iff  $|i-j| \in A_v$ .
- 12. Use induction on r: Find a sequence  $I_1, \ldots, I_r$  of "long" intervals whose first terms constitute an arithmetic progression and which are similarly colored.
- 13. Use 14.2a in a similar way to that in 14.11.
- 14. (a) First select a large set T and  $A_1, B_1 \subseteq S T$  with  $A_1 \cap B_1 = \emptyset$  such that, for every  $X \subseteq T$ , the colors of  $A_1 \cup X$ ,  $B_1 \cup X$ ,  $A_1 \cup B_1 \cup X$  are the same.
- (b) Let t be large in (a) and color the subset  $\{i_1, \ldots, i_v\} \subseteq \{1, \ldots, t\}$  by the color of  $A_{i_1} \cup \ldots \cup A_{i_v}$ .

- 15. Color the subsets of an *n*-element set corresponding to the colors of their cardinalities.
- **16.** (a) Trivial for  $n_1(k, m) = k$ .
- (b) Let  $|T| = k^{m^{r-1}}$ ,  $T \subseteq S$ . Color collections  $\mathcal{A}$  of points of S T with the  $m^{|T|}$ -tuples  $(\alpha(\mathcal{A} + \mathfrak{D}); \mathfrak{D} \in m^T)$ .
- 17. By repeated application of 14.16b, we can find disjoint non-empty sets  $X'_1, \ldots, X'_t \subseteq S$  and an *m*-collection  $\mathcal{B}$  of points of  $S X'_1 \ldots X'_t$  such that the color of every *m*-collection

$$\mathcal{A} = \sum_{i=1}^t c_i X_i' + \mathcal{B}'$$

is the same as the color of the m-collection

$$ar{\mathcal{R}} = \bigcup_{c_i \neq o} X_i' + \mathfrak{B}'.$$

Now apply 14.14b.

- 18. Use a method similar to the one in 14.15.
- 19. Suppose there were a k-coloration of  $\{1, \ldots, N\}$  for each N with no subset having property P. Find such k-colorations which could be combined together.
- **20.** (a) Use the following observation: if x, y, z satisfy (a) and  $x, y, z \equiv 0$  or  $i \pmod{5}$ , then  $x, y, z \equiv 0 \pmod{5}$  (for each fixed  $1 \leq i \leq 4$ ).
  - (b) Use induction on k and Van der Waerden's theorem.
- 21. Translate parts (a) and (b) of the previous solution to the general case.
- 22. Color the points at random and use 2.18 to ensure that every edge meets every color.
- 23. (a) This is a special case of 14.17.
- (b) Given  $r_1,\ldots,r_k\geq 0$ , prove that a number  $\overline{N}(k;r_1,\ldots,r_k;q-1)$  exists such that if  $n\geq \overline{N}$  and all points of the *n*-dimensional projective space over GF(q) are *k*-colored, there always exists an  $r_i$ -dimensional subspace all of whose points have color *i* for some  $1\leq i\leq k$ . Try to reduce this to the affine case by removing the "infinitely distant" hyperplane.
- **24.** For r = q, a prime power, the lines of an affine space over GF(q) of high enough dimension will do it. Otherwise take a prime power  $q \ge r$  and reduce the edges of a q-uniform example.
- **25.** For each  $a_i$ , consider the largest monotonically increasing subsequence starting with  $a_i$ .
- **26.** (a) Let t(x) be the maximum length of a sequence  $x = a_1 < a_2 < \ldots$  with  $f(a_1) \le f(a_2) \le \ldots$  Prove that, if  $t(x) \ge t(1) k$ , then  $f(x) \le 2^k$   $(0 \le k \le t(1) 1)$ .
  - (b) Let f consist of (n-1) monotone decreasing parts.

27. (a) Denote by  $\eta(x, y)$  and  $\varphi(x, y)$  the lengths of the longest monotone increasing and decreasing paths starting with the edge  $(x, y) \in E(T_n)$ . For any two points a, b,

$$\{(\eta(a, x), \varphi(a, x)); (a, x) \in E(T_n)\} \neq \{(\eta(b, x), \varphi(b, x)); (b, x) \in E(T_n)\}.$$

- (b) To prove that the bound is sharp use induction on p and q.
- 28. (a) Let  $x_{i+1}$  be a point different from

$$f(S - \{x_1, \ldots, x_{\nu}\}), \ \nu = 0, \ldots, i \ (0 \le i \le m-1).$$

(b) Prove a common generalization of (a) and (b): There is an ordering  $(x_1, \ldots, x_{2^n})$  of S such that, for each  $1 \leq j \leq n$ ,

$$f({x_1, \ldots, x_{\nu}}) \in {x_1, \ldots, x_{2^j}},$$
  
 $2^j < \nu \le 2^n$ 


holds for at least  $(n-j)2^j$  values of  $\nu$ .

- (c) Look for an f in the following form. Given a function  $\varphi$  with  $1 \leq \varphi(i) \leq 1$ , let f(X) be the  $\varphi(|X|)^{\text{th}}$  element of  $X(X \subseteq S = \{1, \ldots, 2^{n-1} 1\})$ .
- **29.** Direct each segment (p,q) formed by two given points upwards and color it with color i if its angle formed with the positive half of the x-axis is between  $\frac{i}{n}\pi$  and  $\frac{i+1}{n}\pi$ . What can the chromatic number of the graph  $G_i$  formed by the segments of color i be, if it contains no "almost straight" broken line?
- **30.** (a) Associate the value  $\frac{y_1 y_2}{x_1 x_2}$  with the edge  $((x_1, y_1), (x_2, y_2))$  of the complete graph on S; orient this from  $(x_1, y_1)$  toward  $(x_2, y_2)$  if  $x_1 < x_2$  and apply 14.27.
- (b) Carry out the construction in 14.27 with tournaments of the special type considered here.
- 31. (a) Color a quadruple of points red if it forms a convex quadrilateral and blue otherwise; apply Ramsey's theorem.
- (b) The upper bound is immediate from the preceding problem. The lower bound follows by putting together sets with cardinalities  $\binom{m-2}{i}$   $(i=0,\ldots,m-2)$  containing no (i+2)-gons convex from below and no (m-i)-gons convex from above.

- 1. (a) How can one realize the points of G in L(G)?
- (b) There are three edges  $e_1$ ,  $e_2$ ,  $e_3 \in E(G_1)$  starting from some point such that the corresponding edges  $e_1'$ ,  $e_2'$ ,  $e_3' \in E(G_2)$  form a triangle.
  - (c) The solution of (b) contains the answer.

15.2

- HINTS **2.** (a) Show that  $|E \cap F| = |\alpha(E) \cap \alpha(F)|$ , for any two  $E, F \in E(K_n)$ .
- (b) Show that it can be "recognized" in  $L(K_n)$  when two r-tuples have k points in common for k = r - 1; then show this for  $k = 1, 2, \ldots$
- **3.** Again, "recover"  $|E \cap F|$  from  $L_i(K_n^r)$  by counting the r-tuples adjacent to both E and F.
- 4. (a) Consider hypergraphs of stars.
- (b) What we need is a covering of edges by  $\left\lceil \frac{n^2}{4} \right\rceil$  cliques. This can be done by a method similar to the one in Turán's theorem 10.28.
- 5. (a) Use 15.1b to show that the graphs  $H_x$  "fit together".
  - (b) Obvious from the preceding problem.
- **6.** (a) If G = L(H), then those triples of edges of H which form an odd triangle in G belong to a star and conversely.
- (b) Define two edges (a, b) and (c, d) of G to be equivalent if they are identical or they belong to an odd triangle or they span a complete quadrilateral. Show that this is an equivalence relation and that the equivalence classes are complete covering subgraphs as in 15.4a and they will give a desired graph H with L(H) = G in all but a few cases.
  - (c) How can (ii) in (a) be violated?
- 7. Verify that the graph shown in Fig. 20 is not the line-graph of any 3-uniform hypergraph without multiple edges but every proper subgraph of it is such a line-graph.



- 8. (a) This means: find two essentially different embeddings in the plane of the same planar graph.
  - (b) Use 6.69.
- **9.** (a) Consider two edges (x, y), (x, z) and a circuit C through y, z in  $G_1 x$ .
- (b) The solution of 15.1b simplifies since, in this case, no star can correspond to a triangle.
- 10. Use induction on r, showing that T and T' are isomorphic, where  $x_i'$  corresponds to  $x_i$ .

- 11. (a) follows from an identity used in the preceding solution. (b) follows similarly.
  - (c) Construct T by induction. Consider two points  $x_i$ ,  $x_j$  such that

$$d_{ri} + d_{rj} - d_{ij}$$

is minimal.

12. (a) Let 
$$\{a_1, \ldots, a_n\}$$
 consist of  $\binom{k+1}{2l}$  copies of  $2l$ , for  $0 \le l \le \frac{k+1}{2}$  and  $\{b_1, \ldots, b_n\}$  consist of  $\binom{k+1}{2l+1}$  copies of  $2l+1$ ,  $0 \le l \le \frac{k}{2}$ .

(b) Consider the polynomials

$$f(x) = \sum_{i=1}^{n} x^{a_i}, \ g(x) = \sum_{i=1}^{n} x^{b_i}.$$

13. By considering f - g, it suffices to show that

$$\sum_{i=1}^{M} \alpha_i \begin{pmatrix} c_{i1} \\ s_1 \end{pmatrix} \cdot \cdot \cdot \begin{pmatrix} c_{ik} \\ s_k \end{pmatrix}$$

cannot vanish for all integers  $s_1, \ldots, s_k \geq 0$  unless it is identically 0.

- 14. Prove first that  $G_1$ ,  $G_2$  have the same number of edges, then that they have the same degrees.
- 15. (a) Consider two Hamiltonian circuits on V.
  - (b) Count H in all subgraphs G x.
- (c) Prove that each k-point-connected graph occurs among the components of  $G_1$  and  $G_2$  the same number of times, for  $k = |V| 1, \ldots$
- 16. (a) The diameter of  $T_i$  is the maximum of diameters of  $T_i x$ , except when  $T_i$  is a path.
- (b) If  $T_i x$  has diameter d for each endpoint x, then the branches of  $T_i$  relative to its center can be reconstructed using a method similar to the one in 15.9c. If an endpoint  $x_0$  is contained in all paths of length d, then examine whether all (d-1)-paths of  $T_i x_0$  can be covered by one point.
- 17. (a) Express the numbers of isomorphisms of  $G_1$  into  $G_2$  and onto itself by a formula similar to that in 5.18, and observe that the right-hand sides are identical.
  - (b) Use the identity in 2.7b instead of the sieve formula.
- **18.** Show first that if (\*) holds for every |X| = k, then it also holds for every  $|X| \le k$  (provided  $k \le |V| r$ ).
- 19. Suppose  $H_1 \neq H_2$ . Show that  $\emptyset \in E(H_1)$  or  $\emptyset \in E(H_2)$ ; then use induction on k to show that each k-element subset belongs to  $H_1$  or to  $H_2$ , depending on the parity of k.

- **20.** (a) With epimorphisms (homomorphisms onto H) the assertion would be trivial. Express the number of epimorphisms in terms of the number of homomorphisms.
  - (b) "Dualize" the solution of (a).
- **21.** Show that the number of homomorphisms of H into  $G \times G$  is the square of the number of homomorphisms of H into G.
- 22. (a) Try  $F = K_2$ ; observe that  $G_1, G_2$  must have the same degree sequence.
  - (b) Use an argument similar to the one in the preceding solution.
- (c) Let  $G^{\circ}$  denote the graph obtained by adding a loop at each point of G. Then  $(G \cdot H)^{\circ} = G^{\circ} \times H^{\circ}$ .
- (d) Again, if two groups have the property that every group has the same number of homomorphisms into each of them, then they are isomorphic.

# III. Solutions

## § 1. Basic enumeration

- 1. I. The first person may be sent any of the k kinds of postcards. No matter which one he is sent, we may still send the second one any of the k kinds, so there are  $k \cdot k = k^2$  ways to send cards to the first two friends. Again, whatever they are sent, the third friend can still be sent k kinds, etc. So there are  $k^n$  ways to send out the cards.
- II. If they have to be sent different cards, the first person can still be sent any of the k cards. But for any choice of this card, there are only k-1 kinds of cards left for the second person; whatever the first and second friends receive the third one can get one of k-2 postcards, etc. . . . Thus the number of ways to send them postcards is  $k(k-1) \dots (k-n+1)$  (which is, of course, 0 if n > k).
- III. This is the same as the first question but we have  $\binom{k}{2}$  pairs of postcards instead of k postcards. Thus the result is  $\binom{k}{2}^n$ .
- 2. I. We have to decide about the postcards independently. Any postcard can be sent to any of the n friends. Hence, the result is  $n^k$ .
- II. Let  $C_1, \ldots, C_k$  be the cards. The set  $S = \{C_1, \ldots, C_k\}$  must be split into n disjoint non-empty sets  $S_1, \ldots, S_n$ . Thus  $\{S_1, \ldots, S_n\}$  is a partition of S. From any partition of S into n (non-empty) classes we get n! possibilities to send out the postcards. Hence the answer is  $n! \cdot S(k, n)$ .
- 3. There are 16! permutations of the letters of Characterization. However, not all of these give new words; in fact, in any permutation, if we exchange the three A's, the two C's, the two R's, the two I's or the two I's we get the same word. Thus for any permutation, there are  $3! \cdot 2 \cdot 2 \cdot 2 \cdot 2 = 96$  permutations which give the same word, so the result is

$$\frac{16!}{96}$$

In general, if there are  $k_A$  A's,  $k_B$  B's, etc. then the result is

$$\frac{(k_A+k_B+\ldots)!}{k_A!\,k_B!\ldots}.$$

**4.** (a) If we distribute the forints by the procedure described in the hint we have to say "Next please" n-1 times. If we determine at which points (after which coins) we say this we uniquely determine the distribution. There are k-1 possible points to switch and we have to choose n-1 out of these. Hence the result is

$$\binom{k-1}{n-1}$$
.

(b) Borrow one forint from each person. If we distribute the n+k forints we then have in such a way that each person gets at least one, we would then have done the same as if we had distributed the k forints without this requirement. More precisely, distributions of n+k forints among persons so that each one gets at least one are in a one-to-one correspondence with all distributions of n forints among k persons. Hence the answer is

$$\binom{n+k-1}{n-1}$$
.

5. We can distribute the first kind of postcards in

$$\binom{a_1+n-1}{n-1}$$

ways by the preceding solution. Whatever the choice here, there are

$$\binom{a_2+n-1}{n-1}$$

ways to send out the second one, etc. Thus the result is

$$\binom{a_1+n-1}{n-1}\binom{a_2+n-1}{n-1}\cdots\binom{a_k+n-1}{n-1}$$
.

**6.** We interpret S(n, k) as the number of partitions of  $\{1, \ldots, n\}$  into exactly k classes. If we remove the class containing n we get a partition of n-r elements into k-1 classes, where r is the number of elements in this class.

We have  $\binom{n-1}{r-1}$  possible ways of choosing this class. Hence

$$S(n,k) = \sum_{r=1}^{n} {n-1 \choose r-1} S(n-r,k-1) = \sum_{m=0}^{n-1} {n-1 \choose m} S(m,k-1).$$

We can get another recurrence relation if we consider the restrictions of partitions of  $\{1, \ldots, n\}$  onto  $\{1, \ldots, n-1\}$ . This way we obtain each partition of  $1, \ldots, n-1$  into k classes exactly k times, the partitions into k-1 classes exactly once. Hence

$$S(n, k) = S(n - 1, k - 1) + kS(n - 1, k).$$

On the basis of this, it is easy to tabulate the second-order Stirling numbers:

n $k$	0	1	2	3	4	5	6	7
0	1	0	0	0	0	0	0	0
1	0	1	0	0	0	0	0	0
2	0	1	1	0	0	0	0	0
3	0	1	3	1	0	0	0	0
4	0	1	7	6	1	0	0.	0
5	0	1	15	25	10	1	0	0
6	0	1	31	90	65	15	1	0
7	0	1	63	301	350	140	21	1

Table 1
The second-order Stirling numbers

7. Suppose first that x is a positive integer. Let |X| = x, |N| = n. The number of mappings of N into X is  $x^n$  (cf. 1.1). On the other hand, let k denote the cardinality of the range of a mapping of N into X. For k fixed, we can specify S(n, k) ways in which elements of N are mapped onto the same element of X. Once this partition of N is specified, we have to find an image for each class of it, distinct images for distinct classes. This can be done in x(x-1)... (x-k+1) ways. Thus

$$S(n,k) x(x-1) \dots (x-k+1)$$

is the number of mappings of N into X with range of cardinality k. This proves the identity when x is a positive integer. But this means that if we consider x as a variable the polynomials on the two sides have infinitely many values in common. Therefore, they must be identical.

8. 
$$\frac{1}{k!} \sum_{j=0}^{k} (-1)^{k-j} {k \choose j} j^n = \sum_{j=0}^{k} \frac{(-1)^{k-j}}{j!(k-j)!} \sum_{r=0}^{n} S(n,r) j(j-1) \dots (j-r+1) =$$

$$= \sum_{j=0}^{k} \sum_{r=0}^{j} (-1)^{k-j} S(n,r) \frac{1}{(k-j)!(j-r)!} =$$

$$= \sum_{r=0}^{k} \frac{S(n,r)}{(k-r)!} \sum_{j=r}^{k} (-1)^{k-j} {k-r \choose k-j} = \sum_{r=0}^{k} \frac{S(n,r)}{(k-r)!} (1-1)^{k-r} = S(n,k).$$

**9.** (a) Since S(n, k) = 0 for k > n, we have

$$P_n = \sum_{k=0}^{\infty} S(n, k) = \sum_{k=0}^{\infty} \frac{1}{k!} \sum_{j=0}^{k} (-1)^{k-j} {k \choose j} j^n =$$

$$= \sum_{j=0}^{\infty} \frac{j^n}{j!} \sum_{k=j}^{\infty} \frac{(-1)^{k-j}}{(k-j)!} = \sum_{j=0}^{\infty} \frac{j^n}{j!} \cdot \frac{1}{e}.$$

(b) First we remark that

(1) 
$$P_n \sim \frac{1}{\sqrt{2\pi}} \sum_{k=1}^{\infty} k^{n-k-\frac{1}{2}} e^{k-1} \qquad (n \to \infty).$$

In fact, let  $k_0$  be chosen such that for  $k \geq k_0$ ,

$$\left|rac{k\,!}{\sqrt{2\pi\,k}\,(k/e)^k}-1
ight|$$

where  $\varepsilon$  is any given positive number. Since

$$\sum_{k=0}^{k_0-1} \frac{k^n}{k!} = o\left(\frac{k_0^n}{k_0!}\right) = o(P_n),$$

we have

$$P_n \sim \frac{1}{e} \sum_{k=k}^{\infty} \frac{k^n}{k!}$$
.

Similarly,

$$\frac{1}{\sqrt{2\pi}} \sum_{k=1}^{\infty} k^{n-k-1/2} e^{k-1} \sim \frac{1}{\sqrt{2\pi}} \sum_{k=k_0}^{\infty} k^{n-k-1/2} e^{k-1}.$$

By the choice of  $k_0$ , we have

$$1-arepsilon<rac{rac{1}{\sqrt{2\pi}}\sum\limits_{k=k_0}^{\infty}k^{n-k-1/2}\,e^{k-1}}{rac{1}{e}\sum\limits_{k=k_0}^{\infty}k^n/k\,!}<1+arepsilon.$$

This proves (1).

Now put

$$g_n(x) = \begin{cases} \frac{1}{\sqrt{2\pi}} x^{n-x-\frac{1}{2}} e^{x-1} & \text{for } x \ge 0, \\ 0 & \text{for } x \le 0. \end{cases}$$

Then  $g_n(x)$  has a unique maximum at the point  $\lambda(n)$ . Put

$$h_n(y) = \frac{g_n(\lambda(n) (1 + y/\sqrt{n}))}{g_n(\lambda(n))}.$$

Then  $h_n(y)$  has a unique maximum at y = 0, this maximal value is 1 and we have, by the above,

(2) 
$$P_n \sim \left\{ \sum_{k=-\infty}^{\infty} \frac{\sqrt{n}}{\lambda(n)} h_n(y_k) \right\} \cdot \frac{\lambda(n)}{\sqrt{n}} g_n(\lambda(n)),$$

where

$$y_k = k \frac{\sqrt{n}}{\lambda(n)} - \sqrt{n}$$
.

The sum in  $\{\ \}$  is approximating the integral  $\int\limits_{-\infty}^{\infty}h_n(y)\,dy$  and the difference is less than  $(n/\lambda(n))$  max  $h_n(y)=o(1)$ . We shall show that

$$h_n(y) \to e^{-y^2/2} \qquad (n \to \infty)$$

and that the functions  $h_n(y)$  have a common integrable majorant. It will then follow by Lebesgue's theorem that

$$P_n \sim rac{\lambda(n)}{\sqrt{n}} g_nig(\lambda(n)ig) \int\limits_{-\infty}^{\infty} e^{-y^2/2} dy = rac{\lambda(n)}{\sqrt{n}} g_nig(\lambda(n)ig) \sqrt{2\pi} = rac{1}{\sqrt{n}} \lambda(n)^{n+rac{1}{2}} e^{\lambda(n)-n-1} \,.$$

Now (3) follows from the transformation

(4) 
$$\log h_n(y) = -(n - \lambda(n)) \left\{ \frac{y}{\sqrt{n}} - \log\left(1 + \frac{y}{\sqrt{n}}\right) \right\} - \left(\lambda(n) \frac{y}{\sqrt{n}} + \frac{1}{2}\right) \log\left(1 + \frac{y}{\sqrt{n}}\right).$$

The second term is  $\sim \frac{y}{\sqrt{n}} \left( \lambda(n) \frac{y}{\sqrt{n}} + \frac{1}{2} \right) = o(1)$  (as  $n \to \infty$ ), while the first one is equal to

$$(n-\lambda(n))\left(-\frac{y^2}{2n}+\frac{y^3}{3n\sqrt{n}}-\ldots\right)=-\frac{y^2}{2}+o(1)$$

for  $n > y^2$ . This proves (3).

To establish an integrable majorant for  $h_n(y)$ , first let y > 0. Then by (4),

$$\log h_n(y) < -rac{n}{2} \left( rac{y}{\sqrt{n}} - \log \left( 1 + rac{y}{\sqrt{n}} 
ight) 
ight)$$

(for  $n \ge 16$ ). This upper bound is monotone decreasing and so

$$h_n(y) < \exp\left(-2y + 8\log\left(1 + \frac{y}{4}\right)\right) \qquad (n \ge 16).$$

For y < 0 we have similarly

$$\log h_n(y) < -\frac{n}{2} \left( \frac{y}{\sqrt[]{n}} - \log \left( 1 + \frac{y}{\sqrt[]{n}} \right) - \frac{1}{2} \log \left( 1 + \frac{y}{\sqrt[]{n}} \right) \right)$$

(provided  $n > y^2$ ) and then it is easy to see that, for y < -3 the right-hand side is monotone increasing and hence that

$$h_n(y) < e^{-y^2}$$
.

This holds for  $n \le y^2$  as well since then  $h_n(y) = 0$ . Estimating  $h_n(y)$  by 1 in the interval  $-3 \le y \le 0$ , we obtain the desired integrable majorant. [L. Moser-M. Wyman, Trans. Royal Soc. Can. 49 (1955) 49-54.]

10. Let S be the set to be partitioned and  $x \in S$ . If the class containing x has k elements, it can be chosen in  $\binom{n-1}{k-1}$  ways and the remaining n-k elements can be partitioned in  $P_{n-k}$  ways. So the number of partitions in which the class containing x has k elements is

$$\binom{n-1}{k-1}P_{n-k}.$$

This remains true for k = n if we set  $P_0 = 1$ . Thus

$$P_n = \sum_{k=1}^n \binom{n-1}{k-1} P_{n-k} = \sum_{k=0}^{n-1} \binom{n-1}{k} P_k.$$

11. From the recurrence relation in the previous solution, we have

$$p(x) = \sum_{n=0}^{\infty} \frac{P_n}{n!} x^n = 1 + \sum_{n=1}^{\infty} \frac{x^n}{n!} \sum_{k=0}^{n-1} {n-1 \choose k} P_k = 1 + \sum_{k=0}^{\infty} \frac{P_k}{k!} \sum_{n=k+1}^{\infty} \frac{x^n}{n} \frac{1}{(n-k-1)!}$$

and so.

$$p'(x) = \sum_{k=0}^{\infty} \frac{P_k}{k!} \sum_{n=k+1}^{\infty} \frac{x^{n-1}}{(n-k-1)!} = \sum_{k=0}^{\infty} \frac{P_k x^k}{k!} \sum_{r=0}^{\infty} \frac{x^r}{r!} = p(x)e^x$$

or

$$(\log p(x))' = \frac{p'(x)}{p(x)} = e^x, \quad p(x) = e^{e^x + c}$$

for some constant c, which can be determined by setting x = 0:

$$1 = p(0) = e^{e^0 + c}, \qquad c = -1.$$

Thus we have

$$p(x)=e^{e^x-1}.$$

12. (a) Let  $k_i$  denote the number of *i*-element classes in a partition. Then

$$k_1+2k_2+\ldots+nk_n=n.$$

Let us count partitions with  $k_1, \ldots, k_n$  fixed. If we consider any arrangement of the n objects we can get such a partition by taking the first  $k_1$  elements as 1-element classes, the next  $2k_2$  elements a 2-element classes etc. We get any given partition exactly

$$k_1!(1!)^{k_1} \cdot k_2!(2!)^{k_2} \dots k_n!(n!)^{k_n}$$

times; for we can construct the arrangement of the objects by putting the oneelement classes first, then the two-element ones, etc. However there are  $k_i$ ! possible ways to order the *i*-element classes and  $(i!)^{k_i}$  possible ways to order the elements in the *i*-element classes. Thus the number of partitions with  $k_i$  *i*-element classes  $(i = 1, ..., n; k_1 + 2k_2 + ... + nk_n = n)$  is

$$\frac{n!}{k_1!(1!)^{k_1}k_2!(2!)^{k_2}\dots k_n!(n!)^{k_n}}.$$

This proves the assertion. Now

$$p(x) = \sum_{n=0}^{\infty} \frac{P_n}{n!} x^n = \sum_{n=0}^{\infty} \sum_{\substack{k_1, \dots, k_n \ge 0 \\ k_1 + 2k_2 + \dots + nk_n = n}} \frac{1}{k_1! (1!)^{k_1} \dots k_n! (n!)^{k_n}} x^n.$$

We may actually allow infinitely many  $k_i$ 's in each term with the restriction  $k_1 + 2k_2 + \ldots = n$ , since clearly  $k_r = 0$  for r > n. Thus

$$p(x) = \sum_{n=0}^{\infty} \sum_{\substack{k_i \ge 0 \\ k_1 + 2k_2 + \dots = n}} \frac{x^n}{\prod_{i=0}^{\infty} k_i ! (i !)^{k_i}} = \sum_{k_i \ge 0} \frac{x^{\sum_{i=0}^{\infty} i k_i}}{\prod_{i=0}^{\infty} k_i ! (i !)^{k_i}},$$

where the summation extends over all sequences  $k_1, k_2, \ldots$  of non-negative integers containing finitely many non-zero entries. Consequently,

$$p(x) = \prod_{i=1}^{\infty} \left( \sum_{k_i=0}^{\infty} \frac{x^{ik_i}}{k_i! (i!)^{k_i}} \right) = \prod_{i=1}^{\infty} e^{\frac{x^i}{i!}} = e^{e^x - 1}.$$

(b) If we consider

$$\sum_{\substack{k_1,\ldots,k_n\geq 0\\k_1+\ldots+k_n=n}} \frac{n!}{k_1!(1!)^{k_1}k_2!(2!)^{k_2}\ldots(k_n)!(n!)^{k_n}},$$

then we can observe that this is the polynomial expansion of

$$\left(\sum_{i=1}^n \frac{1}{i!}\right)^n \approx (e-1)^n.$$

In fact the difference is

$$(e-1)^n - \left(\sum_{i=1}^n \frac{1}{i!}\right)^n = \left(e-1 - \sum_{i=1}^n \frac{1}{i!}\right) \left(\sum_{k=0}^{n-1} (e-1)^k \left(\sum_{i=1}^n \frac{1}{i!}\right)^{n-k}\right) <$$

$$< \left(\frac{1}{(n+1)!} + \frac{1}{(n+2)!} + \ldots\right) n(e-1)^n \to 0.$$

13. By 1.12,

$$e^{e^x} = \sum_{n=0}^{\infty} \frac{eP_n}{n!} x^n.$$

On the other hand,

$$e^{e^x} = \sum_{k=0}^{\infty} \frac{e^{kx}}{k!} = \sum_{k=0}^{\infty} \frac{1}{k!} \sum_{n=0}^{\infty} \frac{(kx)^n}{n!} = \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{k=0}^{\infty} \frac{k^n}{k!} x^n.$$

Hence

$$\frac{eP_n}{n!} = \frac{1}{n!} \sum_{k=0}^{\infty} \frac{k^n}{k!}$$

which proves the assertion.

We remark that, conversely, the formula in this problem implies 1.11.

#### 14. (a) We have

$$Q_n = \sum_{\substack{k_1,\ldots,k_n \geq 0 \ k_1 + 2k_2 + \ldots + nk_n = n \ k_1 + k_2 + \ldots + k_n \, ext{even}}} rac{n\,!}{k_1\,!\;(1\,!)^{k_1}\ldots k_n\,!\;(n\,!)^{k_n}}$$

and so

$$q(x) = \sum_{n=0}^{\infty} \sum_{\substack{k_1, \dots, k_n \ge 0 \\ k_1 + 2k_2 + \dots + nk_n = n \\ k_1 + \dots + k_n \text{ even}}} \frac{x^n}{k_1! \ (1 \ !)^{k_1} \dots k_n \ ! \ (n \ !)^{k_n}} =$$

$$= \sum_{\substack{k_1, k_2, \dots \ge 0 \\ k_1 + k_2 + \dots \text{ finite and even}}} \prod_{i=1}^{\infty} \frac{x^{ik_i}}{k_i! \ (i \ !)^{k_i}} =$$

$$= \sum_{r=0}^{\infty} \frac{1}{(2r)!} \sum_{\substack{k_1, k_2, \dots \ge 0 \\ k_1 + k_2 + \dots = 2r}} \frac{(2r)!}{k_1! \ k_2! \dots} \prod_{i=1}^{\infty} \left(\frac{x^i}{i!}\right)^{k_i}.$$

The second summation is just the polynomial expansion of

$$\left(\sum_{i=1}^{\infty} \frac{x^i}{i!}\right)^{2r} = (e^x - 1)^{2r}$$

and so

$$q(x) = \sum_{r=0}^{\infty} \frac{1}{(2r)!} (e^x - 1)^{2r} = \operatorname{ch}(e^x - 1).$$

To get an expression for the coefficients we write

$$ch(e^{x}-1) = \frac{e^{e^{x}-1} + e^{1-e^{x}}}{2} = \sum_{n=0}^{\infty} \frac{P_{n} + D_{n}}{2} x^{n},$$

where

$$P_n = \frac{1}{e} \sum_{k=0}^{\infty} \frac{k^n}{k!}$$

is the number of all partitions and

$$D_n = e \sum_{k=0}^{\infty} \frac{(-1)^k k^n}{k!}$$

is the difference between the number of partitions into an even and an odd number of classes [A. Rényi, MTA III. Oszt. Közl. 16 (1966) 77–105].

(b) If the class containing x has 2k elements, then we get the recurrence relation

(1) 
$$R_n = \sum_{k=1}^{\left[\frac{n}{2}\right]} {n-1 \choose 2k-1} R_{n-2k}.$$

We remark that, obviously,  $R_{2m+1} = 0$ .

$$\begin{split} r'(x) &= \sum_{n=1}^{\infty} \frac{R_n}{(n-1)!} x^{n-1} = \sum_{n=1}^{\infty} \frac{x^{n-1}}{(n-1)!} \sum_{k=1}^{\left[\frac{n}{2}\right]} {n-1 \choose 2k-1} R_{n-2k} = \\ &= \sum_{k=1}^{\infty} \frac{1}{(2k-1)!} \sum_{n=2k}^{\infty} \frac{R_{n-2k}}{(n-2k)!} x^{n-1} = \\ &= \sum_{k=1}^{\infty} \frac{x^{2k-1}}{(2k-1)!} \sum_{n=2k}^{\infty} \frac{R_{n-2k}}{(n-2k)!} = r(x) \sum_{k=1}^{\infty} \frac{x^{2k-1}}{(2k-1)!} = r(x) \operatorname{sh} x \,, \end{split}$$

whence

$$r(x) = e^{\operatorname{ch} x - 1}$$
.

(The constant -1 in the exponent follows, as before, by substituting x = 0.) [Rényi, ibid.]

15. (a) If there are exactly  $k \ge 1$  elements mapped onto 1 by f, then these elements can be chosen in  $\binom{n}{k}$  ways and the rest can be mapped into  $2, \ldots, n$  in exactly  $S_{n-k}$  ways. Hence

$$S_n = \sum_{k=1}^n \binom{n}{k} \, S_{n-k}$$

or

$$\frac{2S_n}{n!} = \sum_{k=0}^n \frac{1}{k!} \frac{S_{n-k}}{(n-k)!} \qquad (n \ge 1).$$

Thus

$$egin{aligned} 2s(x) &= \sum_{n=0}^{\infty} rac{2S_n}{n\,!} \, x^n = 1 + \sum_{n=0}^{\infty} \left( \sum_{k=0}^n rac{1}{k\,!} rac{S_{n-k}}{(n-k)\,!} 
ight) x^n = \ &= 1 + \left( \sum_{m=0}^{\infty} rac{S_m}{m\,!} \, x^m 
ight) \left( \sum_{k=0}^{\infty} rac{x^k}{k\,!} 
ight) = 1 + s(x) \, e^{\mathbf{x}}, \end{aligned}$$

whence

$$s(x) = \frac{1}{2 - e^x}.$$

Let us expand this:

$$s(x) = \frac{1}{2} \frac{1}{1 - e^{x}/2} = \frac{1}{2} \sum_{k=0}^{\infty} \left( \frac{e^{x}}{2} \right)^{k} = \sum_{k=0}^{\infty} \frac{1}{2^{k+1}} \sum_{n=0}^{\infty} \frac{k^{n} x^{n}}{n!},$$

whence the formula for  $S_n$  follows by comparing the coefficients of  $x^n$  on both sides.

(b) By the Cauchy formula,

$$rac{S_n}{n\,!}=rac{1}{2\pi\,i}igoplus_{|z|=arepsilon}rac{dz}{(2-e^z)z^{n+1}}$$

for any  $\varepsilon$  small enough. This can be re-written as


$$\frac{1}{2\pi i} \oint\limits_{|z|=2} -\frac{1}{2\pi i} \oint\limits_{|z-\log 2|=\varepsilon},$$

because the only singularities of the integrand are at z=0 and  $z=\log 2+2k\pi i$ , and hence only  $z=\log 2$  lies in the region between the circles |z|=2 and  $|z|=\varepsilon$ . The first term tends to 0 as  $n\to\infty$ . The second term is known to be equal to

$$\left. rac{-1}{z^{n+1} rac{d}{dz} (2-e^z)} \, \right|_{z=\log 2} = rac{1}{2 \left(\log 2 
ight)^{n+1}} \, ( o \infty) \, .$$

This proves the assertion.

16. If  $n = a_1 + \ldots + a_s$  is a partition of  $n, a_1 \ge \ldots \ge a_s$ , then we construct a diagram, the so-called *Ferrer diagram* of the partition as follows: we put  $a_i$  dots in the i<sup>th</sup> column, starting from the bottom (Fig. 21).



Frg. 21

This construction gives a one-to-one correspondence between partitions of the number n and arrays composed of n points, arranged in columns such that the number of points in the successive columns decreases (not strictly). We may also consider these points as lattice points. In that case we will have a collection S of lattice points such that if  $(i, j) \in S$ , then  $i \geq 0, j \geq 0$  and if  $(i, j) \in S$ ,  $0 \leq i' \leq i$ ,  $0 \leq j' \leq j$ , then  $(i', j') \in S$ . Hence it follows that reflecting a Ferrer diagram on the line x = y we get another Ferrer diagram. Now, if a Ferrer diagram has at most r columns, then this reflected Ferrer diagram has at most r points in any column, i.e. it represents a partition into

terms, each at most r. This gives a one-to-one correspondence between partitions into at most r terms and partitions into terms, each at most r.

17. Let  $a_1 + \ldots + a_m = n$  be a partition of n into exactly m terms;  $a_1 \geq 1$  $\geq \ldots \geq a_m$ . Then

$$(1) n - m = (a_1 - 1) + \ldots + (a_m - 1)$$

and here  $a_1 - 1 \ge ... \ge a_m - 1$ . Some terms at the end of (1) may be 0's; discarding these, we obtain a partition of n - m into not more than m terms. Conversely, if we have a partition of n-m into  $r \leq m$  terms, then we can get a partition of n by adding 1 to each term and taking m-r additional 1's. This one-to-one correspondence proves the assertion.

Let  $a_1 + \ldots + a_m = n$  be a partition of n into exactly m distinct terms. Then  $a_1 > a_2 \ldots > a_m \ge 1$ , hence

$$a_1 - (m-1) \ge a_2 - (m-2) \ge \ldots \ge a_{m-1} - 1 \ge a_m \ge 1$$

and so, the decomposition

$$n - {m \choose 2} = (a_1 - (m-1)) + (a_2 - (m-2)) + \ldots + a_m$$

is a partition of  $n - \binom{m}{2}$  into m terms.

Conversely, if we are given a partition of  $n-\binom{m}{2}$  into exactly m terms, then adding  $(m-1), \ldots, 1, 0$  to the terms, we get a partition of n into m distinct

The number of partitions of n into exactly m distinct terms is equal to the number of partitions of  $n - {m \choose 2}$  into exactly m terms.

18. If

(1) 
$$n = a_1 + \ldots + a_m, \quad a_1 > \ldots > a_m \ge 1,$$

then set

$$a_i = 2^{\beta_i} b_i$$
, where  $b_i$  is odd.

Replacing each  $a_i$  by  $2^{\beta_i}$   $b_i$ 's in (1) we get (after rearranging the terms if necessary) a partition of n into odd terms. The number of occurrences of the odd number d is

$$\sum_{b_i=d} 2^{\beta_i} .$$

We show that the above correspondence is a bijection; i.e. any partition of ninto odd terms arises exactly once. In fact, consider a partition of n into odd terms; let there be  $\gamma_1$  1's,  $\gamma_3$  3's . . . If this arises as above, then

$$\gamma_d = \sum_{b_i = d} 2^{eta_i}$$

for any odd number d. Since here the numbers  $\beta_i$  must be distinct (because  $\beta_i = \beta_j$ ,  $b_i = b_j = d$  would imply  $a_i = a_j = 2^{\beta_i}d$ ); it follows that they are uniquely determined by  $\gamma_d$ . Thus a partition of n into odd numbers arises in at most one way. On the other hand,  $\gamma_d$  can be written as a sum of distinct powers of 2:

 $\gamma = \sum 2^{\beta_i(d)},$ 

and then the numbers  $d2^{\beta_i(d)}$  form a partition of n into distinct terms such that the corresponding partition into odd terms is the same as the one we started with.

### 19. Consider a partition

$$(1) n = a_1 + \ldots + a_m, \quad a_1 > \ldots > a_m$$

and its Ferrer diagram. We drow lines through the first and last column, through the lowest row and also a line decreasing at 45° through the dot at the top of the first column (Fig. 22).




Fig. 22

The diagram is contained in the trapezoid formed by these four lines because  $a_1 > \ldots > a_m$ . We denote by p the number of dots on the upper (skew) line. Now we transform the diagram as follows. If  $p < a_m$ , then we take off the last dots of the first p columns and form a new column of p dots at the end (Fig. 22 a  $\rightarrow$  b). The resulting diagram will have distinct rows; this is trivial if  $a_m$  is not decreased, i.e. p < m; and it is also clear in case p = m and  $a_m > p + 1$ . Now  $a_m = p + 1$ , m = p is excluded, because then

$$n = (p+1) + (p+2) + \ldots + (2p) = \frac{3p^2 + p}{2}.$$

If  $a_m \leq p$ , then we remove the last column and add one point to the first

 $a_m$  columns (Fig. 22 b  $\rightarrow$  a). This is possible except when there are no  $a_m$  columns, i.e.  $m-1 < a_m$ . But  $m \ge p \ge a_m$  and hence, this could only arise if  $m=p=a_m$ ; but then

$$n = p + (p + 1) + \ldots + (2p - 1) = \frac{3p^2 - p}{2},$$

which was excluded. Thus we have defined a transformation which associates Ferrer diagrams of partitions into distinct terms with each other. It is easy to check (that is why Fig. 22 could be used to illustrate both cases we considered) that carrying out this transformation twice we get the original diagram, i.e. the transformation pairs off the partitions into distinct terms. Since each pair contains one partition into an odd number of terms and one into an even number of terms, this pairing proves the assertion of the problem. [Euler; cf. also 1.22]

#### 20. If we consider

$$(1+x+x^2+\ldots)(1+x^2+x^4+\ldots)(1+x^3+x^6+\ldots),$$

then the coefficient of  $x^n$  is the number of ways  $x^n$  can be represented as  $x^{k_1}x^{2k_2} \dots x^{nk_n}$ , i.e. the number of partitions of n. Hence

$$S(x) = \sum_{n=0}^{\infty} \pi_n x^n = (1 + x + x^2 + \dots) (1 + x^2 + x^4 + \dots) \dots =$$

$$= \frac{1}{1 - x} \cdot \frac{1}{1 - x^2} \cdot \frac{1}{1 - x^3} \dots = \frac{1}{(1 - x) (1 - x^2) \dots}.$$

(We have skipped the question of convergence of these infinite products; this could easily be settled but they could also be viewed as formal series.)

21. The coefficient of  $x^n$  in the product

$$(1+x)(1+x^2)(1+x^3)\dots$$

is the number of partition of n into distinct terms; the coefficient of  $x^n$  in

$$(1+x+x^2+\ldots)(1+x^3+x^6+\ldots)(1+x^5+x^{10}+\ldots)\ldots$$

is the number of partitions of n into odd terms. Thus we have to show that

$$(1+x) (1+x^2) (1+x^3) \dots =$$
  
=  $(1+x+x^2+\dots) (1+x^3+x^6+\dots) (1+x^5+x^{10}+\dots) \dots$ 

or, equivalently,

(1) 
$$(1+x)(1+x^2)(1+x^3)... = \frac{1}{(1-x)(1-x^3)(1-x^5)...}$$

Now

$$(1-x)(1+x)(1+x^2)(1+x^4)\ldots=1$$

and, substituting  $x^{2k+1}$  for x, we get

$$(1-x^{2k+1})(1+x^{2k+1})(1+x^{2(2k+1)})(1+x^{4(2k+1)})\ldots=1.$$

Multiplying these identities together for  $k = 0, 1, 2, \ldots$ , we get all binomials  $1 - x^{2k+1}$ , also all binomials  $1 + x^n$ ; thus

$$(1-x)(1-x^3)(1-x^5)\dots(1+x)(1+x^2)(1+x^3)\dots=1,$$

which proves (1).

22. If we expand

$$(1-x)(1-x^2)(1-x^3)\ldots$$

we get a  $-x^n$  for any partition of n into an odd number of terms and a  $+x^n$  for any partition of n into an even number of terms. So, by 1.19,  $x^n$  cancels out unless  $n = \frac{3k^2 \pm k}{2}$ .

Let us determine what happens if  $n = \frac{3k^2 + k}{2}$ . The correspondence between

partitions of n constructed in the solution of 1.19, pairs off partitions into an even number of terms with partitions into odd number of terms except for the partition  $n = (k+1) + (k+2) + \ldots + (2k)$ , which will give a coefficient  $(-1)^k$ .

Similarly, if  $n = \frac{3k^2 - k}{2}$ , we get  $(-1)^k x^n$ . Hence

$$(1-x)(1-x^2)(1-x^3)\ldots=1+\sum_{k=1}^{\infty}(-1)^k\left\{x^{\frac{3k^2-k}{2}}+x^{\frac{3k^2+k}{2}}\right\}.$$

[For a tricky analytic proof, which of course also proves 1.19, see e.g. Rademacher, Lectures on Elementary Number Theory, Blaisdell, 1964.]

23. (a) Let us prove the combinatorial assertion formulated in the hint. If we are given a symmetric Ferrer diagram with k elements in the "diagonal", then let  $a_1$  denote the total number of dots in the first row and first column; let  $a_2$  be the number of dots in the second row and column which are not in the first row or column, etc. (Fig. 23). Then  $a_1 > a_2 > \ldots > a_k$  are odd numbers which partition n. Conversely, we can construct from any partition




Fig. 23

into k distinct odd numbers a symmetric Ferrer diagram with k dots in the "diagonal". This proves the assertion of the hint.

Now the number of partitions of n into distinct odd numbers is the coeffi-

cient of  $x^n$  in

$$(1+x)(1+x^3)(1+x^5)\dots$$

The number of symmetric Ferrer diagrams with k elements in the diagonal can be counted as follows. We remove the  $k \times k$  square in the corner and consider the  $i^{\text{th}}$  row above the square plus the  $i^{\text{th}}$  column to the right of the square as a summand. This way we get a partition of n into  $k^2$  and some even numbers  $\leq 2k$ . The number of such partitions is the coefficient of  $x^n$  in

$$x^{k^{2}} (1 + x^{2} + x^{4} + \ldots) (1 + x^{4} + x^{8} + \ldots) \ldots (1 + x^{2k} + x^{4k} + \ldots)$$

$$= \frac{x^{k^{2}}}{(1 - x^{2}) (1 - x^{4}) \ldots (1 - x^{2k})}.$$

Hence

or

$$\sum_{k=0}^{\infty} \frac{x^{k^2}}{(1-x^2)(1-x^4)\dots(1-x^{2k})}$$

in the generating function of the numbers of symmetric Ferrer diagrams with n dots. This proves (a).

(b) Now (b) could be proved similarly but it also follows from another observation. The number of partitions of n into k distinct odd terms is the coefficient of  $x^n y^k$  in

$$(1+xy)(1+x^3y)(1+x^5y)\dots$$

On the other hand, we know that this number is equal to the number of symmetric Ferrer diagrams with n dots and k dots in the "diagonal", i.e. the coefficient of  $x^n$  in (1). This gives the identity

$$(1+xy)(1+x^3y)\ldots = \sum_{k=0}^{\infty} \frac{x^{k^2}y^k}{(1-x^2)(1-x^4)\ldots(1-x^{2k})}$$

Setting y = x we get (b) [Euler].

24. Suppose there are  $k_1$  1's in the partition. The partial sums formed by these 1's represent uniquely each number up to  $k_1$ . So none of  $2, \ldots, k_1$  can occur in the partition. Since  $k_1 + 1$  must be representable as a partial sum, it must occur in the partition with some multiplicity  $k_2 > 0$ . Then all numbers are taken care of till  $k_2(k_1 + 1) + k_1$  (i.e. they all have a unique representation). Going on similarly, we get that the numbers occurring in the partition are  $1, k_1 + 1, (k_1 + 1)(k_2 + 1), \ldots, (k_1 + 1)(k_2 + 1), \ldots (k_m + 1)$ , where  $(k_1 + 1) \ldots (k_i + 1)$  has multiplicity  $k_{i+1}$   $(i = 1, \ldots, m)$ . Since the sum of all numbers must be n, we have

$$n = k_1 \cdot 1 + k_2(k_1 + 1) + \ldots + k_{m+1}(k_1 + 1) \ldots (k_m + 1)$$
  
 $n + 1 = (k_1 + 1) \ldots (k_{m+1} + 1).$ 

If m = 0 we have the trivial partition consisting of 1's; if n + 1 is a prime then this must be the case. On the other hand, if n + 1 is composite then

1.25

$$n+1=(k_1+1)(k_2+1)$$

with  $k_1, k_2 \ge 1$  and then the partition

$$n = \underbrace{1 + \dots + 1}_{k_1} + \underbrace{(k_1 + 1) + \dots + (k_1 + 1)}_{k_2}$$

is a suitable one [see R].

25. If

$$n = x + y + z,$$

then

$$(x + y) + (x + z) + (y + z) = 2n,$$
  
 $(x + y) + (x + z) = y + z + 2x > y + z,$ 

i.e. x + y, y + z, x + z are the sides of a triangle with circumference 2n. Conversely, if a, b, c are sides of a such a triangle (a, b, c) integers) then, setting

$$x = n - a$$
,  $y = n - b$ ,  $z = n - c$ ,

we have

$$x = \frac{a+b-c}{2} > 0, y > 0, z > 0,$$

$$x+y+z=n,$$

and also

$$x + y = c$$
,  $x + z = b$ ,  $y + z = a$ .

Thus, we have a one-to-one correspondence between partitions of n and triangles with circumference 2n (and integer sides).

To determine the number of partitions of n into exactly three terms we observe first that the number of partitions of m into exactly two terms not exceeding a is

$$\begin{cases} 0 & \text{if } a < \frac{m}{2}, \\ a - \left[\frac{m-1}{2}\right] & \text{if } \frac{m}{2} \le a < m, \\ \left[\frac{m}{2}\right] & \text{if } a \ge m. \end{cases}$$

Hence, the number of partitions

$$n = a + b + c$$
,  $a \ge b \ge c$ 

is

$$\sum_{\substack{n-a \\ 2 \leq a < n-a}} \left( a - \left[ \frac{n-a-1}{2} \right] \right) + \sum_{n-a \leq a} \left[ \frac{n-a}{2} \right]$$

or, equivalently,

$$\sum_{\substack{\frac{n}{3} \le a < \frac{n}{2}}} \left( a - \left[ \frac{n-a-1}{2} \right] \right) + \sum_{\substack{\frac{n}{2} \le a \le n}} \left[ \frac{n-a}{2} \right].$$

It is seen that the result depends on the residue of  $n \mod 12$ , e.g. if n = 12k, then the result is  $12k^2$ .

**26.** On the first day, we have three choices: we may buy a pretzel, in which case we have  $B_{n-1}$  further possible ways to spend the remaining n-1 forints; or we may buy candies for 2 forints, and then we can spend the rest in  $B_{n-2}$  ways; similarly we have  $B_{n-2}$  possibilities if we first buy an icecream. Thus we have

$$(1) B_n = B_{n-1} + 2B_{n-2}.$$

We need some values at the beginning and obviously,

$$B_1 = 1, \quad B_2 = 3.$$

Hence by (1)

$$B_3 = 5, B_4 = 11, B_5 = 21, B_6 = 43.$$

We see here and conjecture that

$$(2) B_n = 2B_{n-1} + (-1)^n.$$

In fact, (2) follows easily from (1) by induction:

if

$$B_{n-1} = 2B_{n-2} + (-1)^{n-1}, \ 2B_{n-2} = B_{n-1} + (-1)^n,$$

then

$$B_n = B_{n-1} + 2B_{n-2} = B_{n-1} + (B_{n-1} + (-1)^n) = 2B_{n-1} + (-1)^n.$$

Now we have

$$B_{n} = 2B_{n-1} + (-1)^{n} = 2(2B_{n-2} + (-1)^{n-1}) + (-1)^{n} = 4B_{n-2} + 2(-1)^{n-1} + (-1)^{n} = 8B_{n-3} + 4(-1)^{n-2} + 2(-1)^{n-1} + (-1)^{n} = \dots = 2^{n-1} - 2^{n-2} + 2^{n-3} - \dots + (-1)^{n} = \frac{2^{n} - (-1)^{n}}{2 - (-1)} = \frac{1}{3}(2^{n} + (-1)^{n-1}).$$

Of course, if one guesses this result earlier one can prove it easily by induction from (1), without bringing (2) into consideration.

27. We may begin by taking one or two steps. In the first case, we have  $A_{n-1}$  possibilities to continue; in the second, we have  $A_{n-2}$  possibilities. Thus

$$(1) A_n = A_{n-1} + A_{n-2}.$$

Since we have

$$A_1 = 1, A_2 = 2,$$

the sequence is essentially the sequence of the Fibonacci numbers. We set  $A_0 = 1$  and

$$f(x) = \sum_{n=0}^{\infty} A_n x^n.$$

Then

$$xf(x) = \sum_{n=1}^{\infty} A_{n-1} x^n,$$

$$x^2 f(x) = \sum_{n=2}^{\infty} A_{n-2} x^n,$$

and by (1)

$$f(x) - xf(x) - x^{2}f(x) = A_{0} + (A_{1} - A_{0}) x + \sum_{n=2}^{\infty} (A_{n} - A_{n-1} - A_{n-2}) x^{n} = A_{0} = 1$$

and hence

$$f(x) = \frac{1}{1 - x - x^2}.$$

To get an explicite formula for  $A_n$  we write this as

$$f(x) = \frac{1/\sqrt{5}x}{1 - \frac{1 + \sqrt{5}}{2}x} - \frac{1/\sqrt{5}x}{1 - \frac{1 - \sqrt{5}}{2}x} = \frac{1}{\sqrt{5}x} \sum_{k=0}^{\infty} \left( \left( \frac{1 + \sqrt{5}}{2}x \right)^k - \left( \frac{1 - \sqrt{5}}{2}x \right)^k \right) = \frac{1}{\sqrt{5}} \sum_{n=0}^{\infty} \left\{ \left( \frac{1 + \sqrt{5}}{2} \right)^{n+1} - \left( \frac{1 - \sqrt{5}}{2} \right)^{n+1} \right\} x^n.$$

Hence

$$A_n = \frac{1}{\sqrt{5}} \left\{ \left( \frac{1 + \sqrt{5}}{2} \right)^{n+1} - \left( \frac{1 - \sqrt{5}}{2} \right)^{n+1} \right\}.$$

**28.** (a) If on the first day, we buy a product costing i forints, we have  $C_{n-i}$  possible ways to spend the rest of the money. Hence

(1) 
$$C_n = \sum_{i=1}^k a_i C_{n-i} \quad (n > k).$$

It will be convenient to set  $C_0 = 1, C_{-1} = \ldots = C_{-k+1} = 0$ . These new values will satisfy (1) for  $n \ge 1$ .

Observe that the sequence  $M\vartheta^n_{\nu}$  satisfies (1) for any fixed  $\nu$ , for

$$M\vartheta_{\nu}^{n} = \sum_{i=1}^{k} a_{i} M\vartheta_{\nu}^{n-i}$$

is equivalent to

$$M\vartheta_{\nu}^{n-k}(\vartheta_{\nu}^{k}-a_{1}\vartheta_{\nu}^{k-1}-\ldots-a_{k-1}\vartheta_{\nu}-a_{k})=0,$$

which is clearly satisfied. Hence any sequence

$$(2) x_n = \sum_{\nu=1}^k M_{\nu} \, \vartheta_{\nu}^n$$

satisfies (1). Let us try to look for our sequence  $C_n$  in the form (2). It will be convenient to set

$$x_n = C_{n-k+1}.$$

Then  $x_0 = x_1 = \ldots = x_{k-2} = 0$ ,  $x_{k-1} = 1$ , or

$$M_1 + M_2 + \ldots + M_k = 0$$

$$\vartheta_1 M_1 + \vartheta_2 M_2 + \ldots + \vartheta_k M_k = 0$$

(3)

$$\vartheta_1^{k-2}M_1 + \vartheta_2^{k-2}M_2 + \ldots + \vartheta_k^{k-2}M_k = 0$$
  
$$\vartheta_1^{k-1}M_1 + \vartheta_2^{k-1}M_2 + \ldots + \vartheta_k^{k-1}M_k = 1.$$

Denote the Vandermonde determinant

by D, then, by Cramer's rule, (3) has a solution for  $M_1, \ldots, M_k$ ; in fact

$$M_{m{
u}} = rac{1}{D}egin{bmatrix} 1^{rac{1}{2}} & artheta_1 \ & artheta_2 \ & artheta_2 \ & artheta_1 \ & artheta_2 \ & artheta_2 \ & artheta_1 \ & artheta_2 \ & artheta_2 \ & artheta_1 \ & artheta_2 \ & artheta_$$

(the second form arises by subtracting  $\vartheta_i^{k-1}$  times the  $v^{\text{th}}$  row from the  $i^{\text{th}}$   $(i \neq v)$  and then adding  $\vartheta_v^{j-1}$  times the last column to the  $j^{\text{th}}$   $(j = 1, \ldots, k-1)$ . Hence

$$x_n = \sum_{v=1}^k M_v \, artheta_v^n = rac{1}{D} egin{bmatrix} 1 & artheta_1 \dots & artheta_1^{k-2} & artheta_1^n \ & & & \ & & & \ 1 & artheta_k \dots & artheta_k^{k-2} & artheta_k^n \end{bmatrix}$$

and  $C_n = x_{n+k-1}$  is the value given in the problem.

(b) Set

$$f(x) = \sum_{n=0}^{\infty} C_n x^n.$$

Then

$$f(x) - \sum_{i=1}^{k} a_i x^i f(x) = C_0 + \sum_{n=1}^{\infty} x^n (C_n - a_1 C_{n-1} - \dots - a_k C_{n-k})$$

(where again  $C_{-1} = \ldots = C_{-k+1} = 0$ ,  $C_0 = 1$ ). Here the last sum is 0, so  $f(x) (1 - a_1 x - \ldots - a_k x^k) = 1$ ,

and 
$$f(x) = \frac{1}{1 - a_1 x - \ldots - a_k x^k}$$
.

To get an explicit formula for the coefficients we observe that the roots of the denominator are  $1/\vartheta_1, \ldots, 1/\vartheta_k$  and hence, f(x) can be written as a sum of partial fractions:

$$f(x) = \frac{1}{(1 - \vartheta_1 x) \dots (1 - \vartheta_k x)} = \sum_{\nu=1}^k \frac{A_{\nu}}{1 - \vartheta_{\nu} x}.$$

Multiplying by  $1 - \vartheta_{\nu}x$  and substituting  $x = 1/\vartheta_{\nu}$ , we get

$$A_{\nu} = \prod_{j \neq \nu} \left( 1 - \frac{\vartheta_j}{\vartheta_{\nu}} \right)$$

and so

$$f(x) = \sum_{\nu=1}^k A_{\nu} \frac{1}{1 - \vartheta_{\nu} x} = \sum_{\nu=1}^k A_{\nu} \sum_{n=0}^{\infty} \vartheta_{\nu}^n x^n = \sum_{n=1}^{\infty} \left( \sum_{\nu=1}^k \prod_{j \neq \nu} \left( 1 - \frac{\vartheta_j}{\vartheta_{\nu}} \right) \vartheta_{\nu}^n \right) x^n$$

and hence

$$C_n = \sum_{\nu=1}^k \prod_{j \neq \nu} \left( 1 - \frac{\vartheta_j}{\vartheta_\nu} \right) \vartheta_\nu^n = \sum_{\nu=1}^k \prod_{j \neq \nu} \left( \vartheta_\nu - \vartheta_j \right) \vartheta_\nu^{n-k+1}.$$

It is left to the reader to verify that this formula is equal to the one given in part (a).

**29.** We have  $p(\lambda_n) =$ 

This recurrence relation remains valid for  $n \ge 1$  if we set  $p_0(\lambda) = 1$  and  $p_{-1}(\lambda) = 0$ . As in the solution of 1.27, we denote by  $\theta_1$ ,  $\theta_2$  the roots of

$$x^2 - \lambda x + 1 = 0$$

 $\theta_1$ ,  $\theta_2$  are, of course, functions of  $\lambda$ :

$$\vartheta_1 = \frac{\lambda + \sqrt{\lambda^2 - 4}}{2} \,, \quad \vartheta_2 = \frac{\lambda - \sqrt{\lambda^2 - 4}}{2} \Big) \,.$$

Then

$$p_n(\lambda) = c_1 - \vartheta_1^{n+1} + c_2 \vartheta_2^{n+1}.$$

Taking n = 0 and -1 we get

$$c_1 + c_2 = 0,$$
  
 $c_1 \theta_1 + c_2 \theta_2 = 1.$ 

Hence

$$c_1 = \frac{1}{\sqrt{\lambda^2-4}}\,, \quad c_2 = \frac{-1}{\sqrt{\lambda^2-4}}$$

and

$$p_n(\lambda) = \frac{1}{\sqrt{\lambda^2 - 4}} (\vartheta_1^{n+1} - \vartheta_2^{n+1}).$$

Thus if

$$p_n(\lambda)=0,$$

then

$$\vartheta_1^{n+1} = \vartheta_2^{n+1}$$

or, equivalently,

$$\theta_1 = \varepsilon^2 \theta_2$$

where

$$\varepsilon = e^{\frac{k\pi i}{n+1}}, \quad 0 \le k \le n.$$

Solving  $\vartheta_1 = \varepsilon^2 \vartheta_2$  for  $\lambda$  we obtain

$$\lambda = \pm \left( \varepsilon + \frac{1}{arepsilon} \right) = \pm \ 2 \cos \frac{k\pi}{n+1} \, .$$

Here we may omit  $\pm$  since  $-\cos\frac{k}{n+1} = \cos\frac{n+1-k}{n+1}$ . It is easily seen by substitution that these numbers are roots of  $p_n(\lambda)$  for  $k=1,\ldots,n$ ; therefore, it is not a root for k=0. Thus the eigenvalues of A are

$$2\cos\frac{k\pi}{n+1}, \quad k=1,\ldots,n$$

[cf. also problems 4.28 and 11.5].

- **30.** If we have a sequence of length n-1 in which a, b are not neighbors we can put in front of it
  - (1) c or d or a [b] if it starts with b [a];
  - (2) a, b, c or d if it starts with c or d.

Hence if  $x_n$  and  $y_n$  denote the numbers of sequences of length n starting with a or b and c or d, respectively, we have the recurrence relations

$$x_n = x_{n-1} + 2y_{n-1},$$
  
 $y_n = 2x_{n-1} + 2y_{n-1}.$ 

To solve these recurrence relations, put

$$v_n = \begin{bmatrix} x_n \\ y_n \end{bmatrix}, \quad A = \begin{bmatrix} 1 & 2 \\ 2 & 2 \end{bmatrix}.$$

Then we have

$$v_n = Av_{n-1} = \ldots = A^{n-1} \begin{bmatrix} 2 \\ 2 \end{bmatrix} = A^n \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

We can transform A into a diagonal form by setting

$$L=egin{bmatrix} rac{3+\sqrt{17}}{2} & 0 \ & & \ 0 & rac{3-\sqrt{17}}{2} \end{bmatrix}$$
 ,

$$T = egin{bmatrix} \sqrt{rac{\sqrt{17}-1}{2\,\sqrt{17}}} & -\sqrt{rac{\sqrt{17}+1}{2\,\sqrt{17}}} \ \sqrt{rac{\sqrt{17}+1}{2\,\sqrt{17}}} & \sqrt{rac{\sqrt{17}-1}{2\,\sqrt{17}}} \end{bmatrix};$$

then  $A = TLT^{-1}$  (L has the eigenvalues of A in the diagonal and T is composed of the eigenvectors of A). We want to determine

$$x_n + y_n = [1, 1] A^n \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \frac{1}{2} [0, 1] A^{n+1} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \frac{1}{2} [0, 1] T L^{n+1} T^{-1} \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

Here

$$[0,1] T = \left[ -\sqrt{rac{\sqrt{17}+1}{2\sqrt{17}}}, \sqrt{rac{\sqrt{17}-1}{2\sqrt{17}}} 
ight]$$

whence

$$x_n + y_n = \frac{\sqrt{17} + 1}{4\sqrt{17}} \left(\frac{3 + \sqrt{17}}{2}\right)^{n+1} + \frac{\sqrt{17} - 1}{4\sqrt{17}} \left(\frac{3 - \sqrt{17}}{2}\right)^{n+1}.$$

**31.** Let  $\{a_1, \ldots, a_k\} \subseteq \{1, 2, \ldots, n\}$  be a selection such that  $a_{\nu+1} \ge a_{\nu} + 2$ . Then

$$a_1 < a_2 - 1 < \ldots < a_k - (k - 1),$$

i.e.  $\{a_1, a_2 - 1, \ldots, a_k - (k - 1)\}$  is a k-element subset of  $\{1, \ldots, n - k + 1\}$ . Conversely, for any k-element subset  $\{a'_1, a'_2, \ldots, a'_k\} \subseteq \{1, \ldots, n - k + 1\}$ 

the set  $\{a'_1, a'_2, +1, \ldots, a'_k + (k-1)\}$  is a subset of  $\{1, \ldots, n\}$  containing no two consecutive integers. Hence the answer is

$$\binom{n-k+1}{k}$$
.

32. For any such function f, consider the following polygon: connect (x, f(x)) to (x+1, f(x)) and also connect (x, f(x)), to (x, f(x)-1), to (x, f(x)-2), ..., to (x, f(x-1)) (here f(0)=1). Also join (n+1, f(n)) to (n+1, f(n)+1), ..., to (n+1, n) (Fig. 24). Then we get a polygon connecting (1, 1) to (n+1, n) whose edges join neighboring lattice points, and if we move along it from (1, 1) to (n+1, n), then we always step either to the right or upwards. Call such a polygon a step-polygon. Conversely, a step-polygon connecting (1, 1) to (n+1, n) represents a monotonic function of  $\{1, \ldots, n\}$  into itself. Thus it suffices to enumerate such polygons.




Fig. 24

Let  $e_1, \ldots, e_{2n-1}$  be the edges of a step-polygon. There are n edges which are "horizontal" and n-1 edges which are "vertical". Conversely, if we specify which n edges should be horizontal, then we uniquely determine a step-polygon. Thus the number of step-polygons connecting (1, 1) to (n + 1, n) is

$$\binom{2n-1}{n}$$
.

**33.** (a) We may as well enumerate all monotonic mappings of  $\{1, \ldots, n\}$  into  $\{0, \ldots, n-1\}$  with the property that f(x) < x. Constructing the steppolygon corresponding to such an f, it will connect the points (1,0) and (n+1,n-1) and will not intersect the line x=y.

We know the number of all step-polygons; it is  $\binom{2n-1}{n}$  by the preceding solution. So, if we enumerate those meeting the line x=y we will know the number of those not meeting this line as well.

Let P be such a polygon which meets the line x=y and let (a,a) be their first point of intersection. Reflect the piece between (1,0) and (a,a) in the line x=y. Then we get a step-polygon P' connecting (0,1) and (n+1, n-1).

Conversely, if P' is a step-polygon connecting (0,1) to (n+1, n-1) then, obviously, it must intersect the line x=y and if we again denote by (a,a) their first point of intersection, we get a step-polygon connecting (1,0) to (n+1, n-1) which meets the line x=y. Hence, the number of all step-polygons connecting (0,1) to (n+1, n-1) is equal to the number of step-polygons connecting (1,0) to (n+1, n-1) which intersect x=y (Fig. 25).




Fig. 25

Now the number of step-polygons between (0,1) and (n+1, n-1) can be determined as in the previous solution. Such a polygon has 2n-1 edges, n-2 going upwards. Hence their number is

$$\binom{2n-1}{n-2}$$

and so, the answer to our problem is

$$\binom{2n-1}{n-1} - \binom{2n-1}{n-2} = \frac{1}{n+1} \binom{2n}{n}.$$

- (b) Denoting by f(k) the sum of the first k digits, the problem becomes equivalent to (a).
- **34.** Consider such a sequence and cancel all entries n. The resulting sequence has some length k, say  $0 \le k \le r$  and also satisfies the assumption. Conversely, if a sequence composed of  $1, \ldots, n-1$  of length k has the property that less than i entries of it are  $\le i$   $(i=1,\ldots)$ , then we may insert r-k new entries n (which can be done in  $\binom{r}{k}$  ways), and get a  $(1,\ldots,n)$ -sequence of length r with the same property. Thus,

(1) 
$$g(n, r) = \sum_{k=0}^{r} {r \choose k} g(n-1, k) \quad (1 \le r < n),$$

where g(n, r) denotes the number of  $(1, \ldots, n)$ -sequences of length r with the property that less than i entries are  $\leq i$ ,  $i = 1, 2, \ldots$ 

Now

(2) 
$$g(n, r) = (n - r) n^{r-1}$$

follows easily by induction on n. For n = r, (2) is valid since then, each  $(1, \ldots, n)$ -sequence of length n contains n entries  $\leq n$  and thus, g(n, r) = 0. Let n > r. Then, using (1),

$$g(n,r) = \sum_{k=0}^{r} {r \choose k} g(n-1,k) = \sum_{k=0}^{r} {r \choose k} (n-1-k) (n-1)^{k-1} =$$

$$= \sum_{k=0}^{r} {r \choose k} (n-1)^k - \sum_{k=0}^{r} {r \choose k} k(n-1)^{k-1} = n^r - r \sum_{k=1}^{r} {r-1 \choose k-1} (n-1)^{k-1} =$$

$$= n^r - r \cdot n^{r-1}$$

which proves (2). [H. E. Daniels, Proc. Roy. Soc. A 183, (1945) 405-435.]

**35.** If we write down the partitions in converse order, starting with the identity partition (the partition into one-element sets) and ending with the zero partition  $\{S\}$ , we can observe that each row arises from the previous one by "sticking two classes together". In the  $i^{\text{th}}$  row we will always have n-i+1 classes, thus the number of ways to get the  $(i+1)^{\text{st}}$  row is  $\binom{n-i+1}{2}$ . Hence the number of all such procedures is

$$\binom{n}{2}\binom{n-1}{2}\cdots\binom{2}{2}=\frac{n!(n-1)!}{2^n}.$$

**36.** Let  $P_i$  be the partition arising at the  $i^{\text{th}}$  step,  $P_0 = \{S\}$ ; we call the sequence  $P_0, P_1, \ldots$  a splitting procedure. Denote by  $P'_i$  the partition of  $S - \{x\}$  induced by  $P_i$  ( $x \in S$ ). Then  $P'_{i+1}$  arises from  $P'_i$  by splitting all classes of  $P_i$  with more than one element into two classes, except possibly for a particular value  $i_0$  of i in which case one of the classes of  $P'_{i_0}$ , though with more than one element, may not split: In the corresponding step the class C of  $P_{i_0}$  containing x is split into  $\{x\}$  and  $C - \{x\}$ . Now, let  $P^*_i$  and  $P^{**}_i$  denote the partitions of  $C - \{x\}$  and S - C induced by  $P'_i$  and set

$$Q_i = \begin{cases} P'_i & \text{if } i \leq i_0, \\ P^*_{i+1} \cup P^{**}_i & \text{if } i > i_0. \end{cases}$$

Then  $Q_0 = \{S\}, Q_1, \ldots$  is a splitting procedure of  $S - \{x\}$ .

Conversely, let  $Q_0, Q_1, \ldots$  be such a splitting procedure of  $S - \{x\}$ . We determine how many ways  $Q_0, Q_1, \ldots$  arises from a splitting procedure  $P_0, P_1, \ldots$  of S. We specify the class  $C - \{x\} \in Q_0 \cup Q_1 \cup \ldots$ . Once we have done so we have a unique  $P_0, P_1, \ldots$ . In fact, the index  $i_0$  is the first for which  $C - \{x\} \in Q_{i_0-1}$ . Hence, we get  $P_i$  ( $1 \le i \le i_0$ ) from  $Q_i$  by adding x to the class containing C - x, and we can construct  $P_i(i_0 < i)$  from  $Q_i$  by replacing the partition of  $C - \{x\}$  induced by  $Q_i$  by the partition of  $C - \{x\}$  induced by  $Q_{i-1}$  and adding x as a separate class. It is easy to check that, this way, we get a unique splitting procedure  $P_0, P_1, \ldots$  from any splitting procedure  $Q_0, Q_1, \ldots$  of  $S - \{x\}$  together with a specified member of  $Q_0 \cup Q_1 \cup \ldots$  Let us calculate  $|Q_0 \cup Q_1 \cup \ldots|$ . Luckily, this will depend only on n. Ob-

viously, any one-element subset of  $S-\{x\}$  belongs to  $Q_0 \cup Q_1 \cup \ldots$ , this makes n-1. To each  $C \in Q_0 \cup Q_1 \cup \ldots$ ,  $|C| \geq 2$ , let us observe that there corresponds a splitting and conversely. There are n-2 splittings (since each splitting increases the number of classes by one) and hence, we have

$$|Q_0 \cup Q_1 \cup \ldots| = (n-1) + (n-2) = 2n-3.$$

Thus, the number of splitting procedures of S is 2n-3 times the number of splitting procedures of  $S-\{x\}$ . Hence, by induction, the looked-for number is

$$(2n-3)(2n-5)\dots 1=(2n-3)!!$$

37. (a) We have to break at each of n-1 points; the procedure is determined if we specify the order in which we break at the different places. Hence, the number is (n-1)!

(b) Let us call such a procedure a breaking procedure. Consider any splitting procedure  $P_0, P_1, \ldots$  of an n-element set S (see the previous solution). We may interpret this as follows: whenever we split a class, we specify one half of it and put this in front of the other half. So, the classes of the partition arising at each step will be ordered and the final partition gives an ordering of S; moreover, the splitting procedure corresponds to a breaking procedure of the "stick" consisting of the elements of S in the given order.

Conversely, if a breaking procedure and an ordering of S is given then applying the breaking procedure with the ordered set S we get a splitting procedure. Let  $b_n$  and  $s_n$  denote the number of breaking and splitting procedures, respectively, then the above correspondence between splitting procedures with orderings of the pairs the classes are split into and pairs consisting of an

ordering and a breaking procedure implies

$$2^{n-1}s_n = n \,! \, b_n.$$
  $s_n = (2n-3) \, (2n-5) \, \dots \, 1$ , and therefore  $b_n = rac{2^{n-1} \, (2n-3) \, (2n-5) \, \dots \, 1}{n \,!} = rac{(2n-2) \,!}{n \,! \, (n-1) \,!} = rac{1}{n} inom{2n-2}{n-1}.$ 

**38.** First solution. Consider a bracketed product. The multiplication carried out at the last time means breaking the product into two pieces, where inside each piece there is some bracketing:

$$(x_1 \ldots x_i) (x_{i+1} \ldots x_n).$$

The two multiplications before this last one mean breaking each half of this into two pieces

$$((x_1 \ldots) (\ldots x_i)) ((x_{i+1} \ldots) (\ldots x_n))$$

(some of these brackets may contain a single variable only, in which case it does not occur). This way we see that the number of bracketings of a product with n factors is equal to the number of breaking procedures of a stick of length n, i.e. by the previous result,

$$rac{1}{n}inom{2n-2}{n-1}$$
.

Second solution. Let f(x) denote the number of closing brackets between the first x+1 symbols  $(x=1,2,\ldots,n-1)$ . Then

- (a)  $0 \le f(x) \le n 2$ ,
- (b) f(x) is monotonic and
- (c) f(x) < x as there are at most x 1 pairs of brackets between x + 1 symbols.

Conversely, let f(x) be a function defined on  $\{1, \ldots, n-1\}$  with properties (a), (b), (c), then f(x) uniquely determines a bracketing of the product. It will be more convenient to describe a bracketing as a way to multiply the n elements together (in a non-associative non-commutative structure). Obviously, there must be f(i) - f(i-1) closing brackets between  $x_i$  and  $x_{i+1}$ , and n-2-f(n-1) closing brackets at the end.

Now consider the first closing bracket (from the left), multiply the two elements in front of it and remove this bracket. Again, find the first remaining closing bracket, multiply the two elements in front of it (one of these two may be the result of the previous multiplication), etc. This must be the same order of multiplication as the one prescribed by the bracketing which determines f(x). It is also easy to check that we will be able to carry this out, i.e. we never get a closing bracket with just one element in front of it, by (c). So we have a bijection between functions f with properties (a), (b), (c) and bracketings of products. The functions with properties (a), (b), (c) have been enumerated in 1.33, so the result is

$$\frac{1}{n} \binom{2n-2}{n-1}$$
.

**39.** Let  $v_1, \ldots, v_n$  be the vertices of our *n*-gon. If  $v_i$  is joined to  $v_j$  by a diagonal in a given triangulation, put a pair of brackets into the product:

$$x_1 \ldots (x_i \ldots x_{j-1}) x_j \ldots x_{n-1}$$

Then it is easy to see that this bracketing is a "correct" one and conversely, from each correct bracketing we get a triangulation. Thus, the answer is

$$D_n = \frac{1}{n-1} \begin{pmatrix} 2n-4 \\ n-2 \end{pmatrix},$$

by 1.38.

**40.** Let  $v_1, \ldots, v_n$  be the vertices of our n-gon (in this cyclic order). In any triangulation, the edge  $(v_1, v_n)$  is contained in a unique triangle  $(v_1, v_n, v_i)$ . If k is fixed, we have  $D_k$  possible ways to triangulate the k-gon  $(v_1, \ldots, v_k)$  and  $D_{n-k+1}$  possible ways to triangulate the n-gon  $(v_{k+1}, \ldots, v_n)$ . If we set  $D_2 = 1$  this will also be true for k = 2 and k = n - 1. Thus

$$D_n = \sum_{k=2}^{n-1} D_k D_{n-k+1}$$
  $(n \ge 3),$ 

$$D_2 = 1.$$

Now let

$$D(x) = \sum_{n=2}^{\infty} D_n x^n.$$

Then (1) implies that

$$D(x)^{2} = \sum_{k=2}^{\infty} D_{k} x^{k} \sum_{l=2}^{\infty} D_{l} x^{l} = \sum_{k=2}^{\infty} \sum_{l=2}^{\infty} D_{k} D_{l} x^{k+l} = \sum_{n=4}^{\infty} x^{n} \sum_{k=2}^{n-2} D_{k} D_{n-k} =$$

$$= \sum_{n=4}^{\infty} D_{n-1} x^{n} = x \left( \sum_{n=3}^{\infty} D_{n} x^{n} \right) = x \left( D(x) - x^{2} \right).$$

Hence

$$D(x)^2 - xD(x) + x^3 = 0,$$
  
 $D(x) = \frac{x \pm \sqrt{x^2 - 4x^3}}{2}.$ 

Since D'(0) = 0, we must take the negative sign for x = 0; since D(x) is continuous, we must take the negative root for  $x < \frac{1}{4}$ . Thus

$$D(x) = \frac{1 - \sqrt{1 - 4x}}{2} \cdot x.$$

Expanding this by Newton's formula we get

$$D(x) = \frac{x}{2} \left( 1 - \sum_{k=0}^{\infty} {1/2 \choose k} (-1)^k \, 4^k \, x^k \right) = \sum_{k=1}^{\infty} (-1)^{k-1} {1/2 \choose k} \, 2^{2k-1} \, x^{k+1}.$$

Here

and hence

$$D(x) = \sum_{k=1}^{\infty} \frac{1}{k} \binom{2k-2}{k-1} \cdot x^{k+1} = \sum_{n=0}^{\infty} \frac{1}{n-1} \binom{2n-4}{n-2} x^n.$$

Hence the result follows.

41. We may assume n > 4. Let us consider a vertex  $v_1$ ; there are n possible ways to choose this. Connect the two neighbors of  $v_1$  by a diagonal  $d_1$ . Now the triangle incident with this diagonal from the other side must have an edge in common with the boundary of the n-gon; so its third edge must be one of the two diagonals connecting an endpoint of  $d_1$  to the neighbor of the other endpoint. So we have two possible ways of choosing this second diagonal  $d_2$ . Similarly, if  $d_1, \ldots, d_i$  (i < n - 3) have been chosen, then there are two possibilities for  $d_{i+1}$ . So we have found

$$n \cdot 2^{n-4}$$

ways to choose the point  $x_1$  and a sequence  $d_1, d_2, \ldots, d_{n-3}$  of diagonals dividing the convex n-gon into triangles having an edge in common with the

n-gon. However, every triangulation (with the required property) has two triangles which contain two (adjacent) edges of the n-gon; thus, each is counted twice. So the result is

 $n2^{n-5}$ .

**42.** (a) We have

$$\sum_{k=0}^{\left[\frac{n}{2}\right]} \binom{n}{2k} + \sum_{k=0}^{\left[\frac{n-1}{2}\right]} \binom{n}{2k+1} = \sum_{k=0}^{n} \binom{n}{k} = 2^{n},$$

$$\sum_{k=0}^{\left[\frac{n}{k}\right]} \binom{n}{2k} - \sum_{k=0}^{\left[\frac{n-1}{2}\right]} \binom{n}{2k+1} = \sum_{k=0}^{n} (-1)^{k} \binom{n}{k} = (1-1)^{n} = 0,$$

thus

$$\sum_{k=0}^{\left[\frac{n}{2}\right]} \binom{n}{2k} = \frac{2^n+0}{2} = 2^{n-1} \quad (n \ge 1).$$

(b) We can replace the last term  $\binom{n-m}{0}$  by  $\binom{n-m+1}{0}$ ; then the sum of the last two terms is  $\binom{n-m+1}{0}+\binom{n-m+1}{1}=\binom{m-m+2}{1}$ ; now the sum of the last two terms is  $\binom{n-m+2}{1}+\binom{n-m+2}{2}=\binom{n-m+3}{2}$ , etc. Finally we get  $\binom{n+1}{m}$  (Table 2).

1 2 1

1 3 3 1

1 4 6 4 1

1 5 10 10 5 1

1 6 15 20 15 6 1

1 7 21 35 35 21 7 1

1 8 28 56 70 56 28 8 1

Table 2

(c) First solution. Let |U|=u, |V|=v,  $U\cap V=\emptyset$ . Then,  $\binom{u}{k}$  is the number of k-tuples from U,  $\binom{v}{m-k}$  is the number of (m-k)-tuples from V;  $\sum_{k=0}^{m} \binom{u}{k} \binom{v}{m-k}$  is the number of m-tuples from  $U \cup V$ , i.e.

$$\sum_{k=0}^{m} \binom{u}{k} \binom{v}{m-k} = \binom{u+v}{m}.$$

Second solution. The coefficient of  $x^m$  on each side of the identity

$$(1+x)^{u}(1+x)^{v} = (1+x)^{u+v}$$

is the left- and right-hand side of (1), respectively.

- (d) The coefficient of  $x^m$  in  $(1-x)^u(1+x)^u = (1-x^2)^u$  is 0, if m is odd and  $\begin{pmatrix} u \\ m/2 \end{pmatrix}$  if m is even.
  - (e) We have

Thus

$$\sum_{k=m}^{n} \binom{k}{m} \binom{n}{k} = \sum_{k=m}^{n} \binom{n}{m} \binom{n-m}{n-k} = \binom{n}{m} \sum_{k=m}^{n} \binom{n-m}{n-k} = \binom{n}{m} 2^{n-m}.$$

(f

$$\sum_{k=0}^{\left[rac{n}{7}
ight]} inom{n}{7k} = \sum_{k=0}^{n} \left(rac{1}{7}\sum_{j=0}^{6}arepsilon^{kj}
ight) inom{n}{k} = rac{1}{7}\sum_{j=0}^{6}\sum_{k=0}^{n}inom{n}{k}arepsilon^{kj} = rac{1}{7}\sum_{j=0}^{6}(1+arepsilon^{j})^{n} = rac{2^{n}}{7}igg[1+2\sum_{j=1}^{3}\cosrac{\pi j n}{7}\cos^{n}rac{\pi j}{7}igg].$$

(g) Let

$$a_n = \sum_{k=0}^{\left[\frac{n}{2}\right]} \binom{n-k}{k} z^k.$$

Then

$$a_n=1+\sum_{k=1}^{\left[rac{n}{2}
ight]}\left(inom{n-1-k}{k}+inom{n-k-1}{k-1}
ight)z^k=\ =\sum_{k=0}^{\left[rac{n}{2}
ight]}inom{n-1-k}{k}z^k+\sum_{k=1}^{\left[rac{n}{2}
ight]}inom{n-k-1}{k-1}z^k.$$

The bound  $\left[\frac{n}{2}\right]$  can be replaced by  $\left[\frac{n-1}{2}\right]$  in the first term since  $k=\frac{n}{2}$  (if this is an integer) gives a 0 term. Hence the first term is

$$\sum_{k=0}^{\left[\frac{n-1}{2}\right]} \binom{n-1-k}{k} z^k = a_{n-1}.$$

Similarly, the second term is

$$\sum_{k=1}^{\left[\frac{n}{2}\right]} \binom{n-1-k}{k-1} z^k = z \sum_{k=0}^{\left[\frac{n-2}{2}\right]} \binom{n-2-k}{k} z^k = z a_{n-2}.$$

So we have the recurrence relation

(1) 
$$a_n = a_{n-1} + za_{n-2}, \quad a_0 = a_1 = 1.$$

As in 1.28, (1) remains valid if we put  $a_{-1} = 0$ . Then by 1.28,

$$a_n = egin{array}{c|c} egin{array}{c|c} 1 & lpha^{n+1} \ \hline 1 & eta^n \end{array} = egin{array}{c|c} eta^{n+1} - lpha^{n+1} \ \hline eta - lpha \end{array},$$

where  $\alpha$ ,  $\beta$  are the two roots of the equation

$$x^2 - x - z = 0,$$

i.e.

$$lpha=rac{1-\sqrt{1-4z}}{2}\,,\quad eta=rac{1+\sqrt{1-4z}}{2}$$

and thus

$$a_n=rac{1}{\sqrt{1-4z}}igg(igg(rac{1+\sqrt{1-4z}}{2}igg)^{n+1}-igg(rac{1-\sqrt{1-4z}}{2}igg)^{n+1}igg).$$

This holds if  $z \neq \frac{1}{4}$ , i.e.  $\alpha \neq \beta$ . In case  $z = \frac{1}{4}$  we can rewrite (1) as

$$a_n - \frac{1}{2}a_{n-1} = \frac{1}{2}\left(a_{n-1} - \frac{1}{2}a_{n-2}\right).$$

Hence

$$a_n - \frac{1}{2}a_{n-1} = \frac{c}{2^n}$$

where from  $a_1 - \frac{1}{2}a_0 = 1 - \frac{1}{2} = \frac{1}{2}$  it follows that c = 1, i.e.

$$a_n - \frac{1}{2}a_{n-1} = \frac{1}{2^n}$$
.

Thus

$$a_n = \frac{1}{2^n} + \frac{1}{2}a_{n-1} = \frac{1}{2^n} + \frac{1}{2}\left(\frac{1}{2^{n-1}} + \frac{1}{2}a_{n-2}\right) =$$

$$= \frac{1}{2^n} + \frac{1}{2^n} + \frac{1}{4}a_{n-2} = \dots = \frac{n}{2^n} + \frac{1}{2^n}a_0 = \frac{n+1}{2^n}.$$

(h) The result is  $(-1)^m \binom{n-1}{m}$ . This is clear for m=0. We prove it by induction:

$$\begin{split} \sum_{k=0}^{m} (-1)^k \binom{n}{k} &= \sum_{k=0}^{m-1} (-1)^k \binom{n}{k} + (-1)^m \binom{n}{m} = \\ &= (-1)^{m-1} \binom{n-1}{m-1} + (-1)^m \binom{n}{m} = (-1)^m \binom{n-1}{m}. \end{split}$$

- (i) The number of those (u+v+1-m)-tuples of  $\{1,\ldots,u+v+1\}$  whose  $(u+1)^{\rm st}$  element is u+k+1 is  $\binom{u+k}{k}\binom{v-k}{m-k}$ . Summing for  $k=0,\ldots,m$  we get the result.
- **43.** (a) The left-hand side can be interpreted as follows: we select k points of an m-element set M; then we select m points out of an n-element set N and the k points chosen before. The result is, therefore, a pair (X, Y), where  $X \subseteq M$ ,  $Y \subseteq N \cup X$  and |Y| = m. The number of possible ways of choosing this is

$$\sum_{k=0}^{m} {m \choose k} {n+k \choose m}.$$

The same enumeration problem can be solved by selecting  $Y \cap M$  and  $Y \cap N$  first; if  $|Y \cap N| = j$ , then this can be done in  $\binom{n}{j} \binom{m}{m-j}$  ways; then we have to select on X with  $Y \cap M \subseteq X \subseteq M$ , which can be done in  $2^j$  ways; so the total number of choices is

$$\sum_{j=0}^m \binom{n}{j} \binom{m}{j} 2^j$$
.

(b) Since identity (a) means the equality of two polynomials in n for infinitely many values of n, it follows that the two polynomials are identical. Thus we may substitute -n-1 for n and get, on the left hand side,

$$\sum_{k=0}^{m} \binom{m}{k} \binom{-n-1+k}{m} = \sum_{k=0}^{m} \binom{m}{k} (-1)^m \binom{n+m-k}{m} = (-1)^m \sum_{k=0}^{m} \binom{m}{k} \binom{n+k}{m}.$$

The right-hand side becomes

$$\sum_{k=0}^{m} \binom{n}{k} \binom{-n-1}{k} 2^k = \sum_{k=0}^{m} \binom{m}{k} (-1)^k \binom{n+k}{k} 2^k = \sum_{k=0}^{m} \binom{m}{k} \binom{n+k}{k} (-2)^k.$$

This proves the assertion.

(c)

$$\sum_{k=0}^{\infty} {p \choose k} {q \choose k} {n+k \choose p+q} = \sum_{k=0}^{\infty} {p \choose k} {q \choose k} \sum_{j=0}^{k} {k \choose j} {n \choose p+q-j} =$$

$$= \sum_{j=0}^{\infty} {n \choose p+q-j} \sum_{k=0}^{\infty} {p \choose k} {q \choose k} {k \choose j}.$$

Here

$$\begin{split} \sum_{k=0}^{\infty} \binom{p}{k} \binom{q}{k} \binom{k}{j} &= \sum_{k=0}^{\infty} \binom{p}{k} \frac{q!}{(q-k)! j! (k-j)!} = \\ &= \sum_{k=0}^{\infty} \binom{p}{k} \binom{q}{j} \binom{q-j}{q-k} = \binom{q}{j} \binom{p+q-j}{q} \end{split}$$

by 1.42c. Thus the sum on the left-hand side is

$$\sum_{j=0}^{\infty} {n \choose p+q-j} {q \choose j} {p+q-j \choose q} = \sum_{j=0}^{\infty} \frac{n!}{(n-p-q+j)! (q-j)! (p-j)! j!} = \sum_{j=0}^{\infty} {n \choose p} {p \choose j} {n-p \choose q-j} = {n \choose p} \sum_{j=0}^{\infty} {p \choose j} {n-p \choose q-j} = {n \choose p} {n \choose q}$$

by 1.42c again.

- (d) This follows in exactly the same way as (b).
- (e) We have

$$\binom{x+t}{n} - \binom{x}{n} = \sum_{k=1}^{n} \binom{t}{k} \binom{x}{n-k}$$

by 1.42c. Dividing by t and letting  $t \to 0$  the first expression follows. The second one is obtained analogously by 1.42i and

$${x+t \choose n} - {x \choose n} = \sum_{k=1}^n {t+k-1 \choose k} {x-k \choose n-k}.$$

**44.** We prove (a) by induction on n. For n = 0 it is trivial. Let n > 0. We have

$$\frac{\partial}{\partial y}(x+y+n)^n = n(x+y+n)^{n-1} = n(x+(y+1)+(n-1))^{n-1},$$

$$\frac{\partial}{\partial y}\sum_{k=0}^n \binom{n}{k} x(x+k)^{k-1}(y+n-k)^{n-k} =$$

$$= \sum_{k=0}^n \binom{n}{k} x(x+k)^{k-1}(n-k) (y+n-k)^{n-k-1} =$$

$$= n \sum_{k=0}^{n-1} \binom{n-1}{k} x(x+k)^{k-1} ((y+1)+(n-1)-k)^{(n-1)-k}.$$

The two right-hand sides here are equal by the induction hypothesis. To prove the identity it suffices to show that (a) holds for any one value of y. Choose y = -x - n. Then the right-hand side vanishes while the left-hand side is

$$\begin{split} \sum_{k=0}^{n} \binom{n}{k} x(x+k)^{k-1} (-x-k)^{n-k} &= x \sum_{k=0}^{n} \binom{n}{k} (-1)^{n-k} (x+k)^{n-1} = \\ &= x \sum_{k=0}^{n} \binom{n}{k} (-1)^{n-k} \sum_{j=0}^{n-1} \binom{n-1}{j} k^{j} x^{n-j} = \\ &= x \sum_{j=0}^{n-1} \binom{n-1}{j} x^{n-j} \sum_{k=0}^{n} \binom{n}{k} k^{j} (-1)^{n-k} = \\ &= x \sum_{j=0}^{n-1} \binom{n-1}{j} x^{n-j} S(j,n) = 0 & \text{as } j < n \text{ (by 1.8).} \end{split}$$

(b) The left-hand side of (a) can be rewritten as

$$\sum_{k=0}^{n} \binom{n}{k} x(x+k)^{k-1} (y+n-k)^{n-k-1} (y+n-k) =$$

$$= \sum_{k=0}^{n} \binom{n}{k} x(x+k)^{k-1} y(y+n-k)^{n-k-1} +$$

$$+ \sum_{k=0}^{n} \binom{n}{k} x(x+k)^{k-1} (y+n-k)^{n-k-1} (n-k).$$

Here the second term is

$$\sum_{k=0}^{n-1} n \binom{n-1}{k} x(x+k)^{k-1} ((y+1) + (n-1) - k)^{(n-1)-k} =$$

$$= n(x+(y+1) + (n-1))^{n-1} = n(x+y+n)^{n-1}$$

by (a). Hence

$$\sum_{k=0}^{n} {n \choose k} x(x+k)^{k-1} y(y+n-k)^{n-k-1} = (x+y+n)^n - n(x+y+n)^{n-1} = (x+y) (x+y+n)^{n-1}.$$

Dividing by xy identity (b) follows.

(c) Subtract  $\frac{1}{x}(y+n)^{n-1} + \frac{1}{y}(x+n)^{n-1}$  from both sides of (b) then we get

$$\sum_{k=1}^{n-1} \binom{n}{k} (x+k)^{k-1} (y+n-k)^{n-k-1} = \frac{1}{x} \left\{ (x+y+n)^{n-1} - (y+n)^{n-1} \right\} + \frac{1}{y} \left\{ (x+y+n)^{n-1} - (x+n)^{n-1} \right\}.$$

Letting  $x, y \to 0$  we obtain (c).

45.

$$\sum_{k=0}^{n} {n \choose k} f_k(x) f_{n-k}(y) = \sum_{k=0}^{n} {n \choose k} k! {x \choose k} (n-k)! {y \choose n-k} =$$

$$= \sum_{k=1}^{n} n! {x \choose k} {y \choose n-k} = n! {x+y \choose n}$$

by 1.42c (strictly speaking, the identity

$$\sum_{k=0}^{n} \binom{x}{k} \binom{y}{n-k} = \binom{x+y}{n}$$

was proved for the case when x, y are natural numbers; but if the polynomials on the two sides coincide for every such choice of the variables they are identical).

Other instances of such polynomials:  $f_n(x) = x^n$  (then the formula in the problem is the binomial theorem),  $f_n(x) = x(x+1) \dots (x+n-1)$  (this follows in the same way as  $f_n(x) = x(x-1) \dots (x-n+1)$ ),  $f_n(x) = x(x+n)^{n-1}$  (this follows from identity (b) in the preceding problem). [See also G. C. Rota-R. Mullin, in *Graph Theory and its Applications*, Academic Press, New York 1970, 167-213.]

## § 2. The sieve

1. Let us subtract from 30 the number of pupils who like mathematics, physics, chemistry, respectively:

$$30 - 12 - 14 - 13$$
.

This way, however, a student who likes both mathematics and physics is subtracted twice; so we have to add them back, and also for the two other pairs of subjects:

$$30 - 12 - 14 - 13 + 5 + 7 + 4$$
.

There is still trouble with those who like all three subjects. They were subtracted 3 times, but put back 3 times, so we have to subtract them once more to get the result

$$30 - 12 - 14 - 13 + 5 + 7 + 4 - 3 = 4$$
.

2. (a) Let

$$B = A_1 A_2 \dots A_k \bar{A}_{k+1} \dots \bar{A}_n$$

be any atom of the Boolean algebra generated by  $A_1, \ldots, A_n$  (with an appropriate choice of indices, every atom has such a form.) Every event in the formula is the union of certain (disjoint) atoms; let us express each  $P(A_I)$  and  $P(A_1 + \ldots + A_n)$  as the sum of the probabilities of the corresponding atoms. We show that the probability of any given atom cancels out.

The coefficient of P(B) on the left-hand side is

$$\begin{array}{ll}
1 & \text{if} & k \neq 0, \\
0 & \text{if} & k = 0.
\end{array}$$

B occurs in  $A_I$  iff  $I \subseteq \{1, \ldots, k\}$ , so its coefficient on the right-hand side is

$$\sum_{j=1}^{k} {k \choose j} (-1)^{j-1} = 1 - \sum_{j=0}^{k} {k \choose j} (-1)^{j} = 1 - (1-1)^{k} = \begin{cases} 1 & \text{if } k \neq 0, \\ 0 & \text{if } k = 0. \end{cases}$$

Thus P(B) has the same coefficient on both sides, which proves (a).

To get (b), choose an element x of S by a uniform distribution. Then  $A_i$  can be identified with the event that  $x \in A_i$ , and we have

$$P(A_i) = \frac{|A_i|}{|S|}.$$

So we have, by the above

$$P(A_1 + \ldots + A_n) = \sum_{j=1}^n (-1)^{j-1} \sum_{|I|=j} \frac{|A_I|}{|S|} = \sum_{\emptyset \neq I \subseteq \{1, \ldots, n\}} (-1)^{|I|-1} \frac{|A_I|}{|S|},$$
 or, equivalently,

$$P(\bar{A}_1 \dots \bar{A}_n) = 1 - P(A_1 + \dots + A_n) = 1 - \sum_{\varnothing \neq I \subseteq \{1,\dots,n\}} (-1)^{|I|-1} \frac{|A|}{|S|}.$$

The assertion (b) follows on multiplying by |S|.

**3.** Let  $A_i$  be the set of integers between 1 and n divisible by  $p_i$ . Then, for  $I \subseteq \{1, \ldots, r\}$ ,

, 
$$A_I = \bigcap_{i \in I} A_i = \{k : 1 \leq k \leq n, \prod_{i \in I} p_i \mid k\}$$

and so,

$$|A_I| = rac{n}{\prod\limits_{i \in I} p_i}.$$

Hence,

$$\varphi(n) = \sum_{I \subseteq \{1, \dots, r\}} (-1)^{|I|} \frac{n}{\prod_{i \in I} p_i}.$$

Now  $\prod_{i \in I} p_i$  ranges over divisors of n; over those divisors of n, in fact, which are not divisible by the square of any prime. To write (1) in a neater form, define

 $\mu(k) = \begin{cases} 0 & \text{if } k \text{ is divisible by the square of some prime,} \\ (-1)^i & \text{if } k \text{ has } i \text{ distinct prime factors.} \end{cases}$ 

Then

$$\varphi(n) = \sum_{k|n} \frac{\mu(k)}{k} n.$$

Also, one can observe that (1) is the expansion of the product

$$\varphi(n) = n \prod_{i=1}^r \left(1 - \frac{1}{p_i}\right).$$

4. Let |X| = k,  $Y = \{y_1, \ldots, y_n\}$ . Then the number of those mappings of X into Y which are onto Y is, obviously,

$$0 \quad \text{if} \quad k < n,$$

$$n! \quad \text{if} \quad k = n.$$

On the other hand, let  $A_i$  denote the set of mappings of X into  $Y-y_i$  and let S be the set of all mappings of X into Y. Then we are interested in  $S-(A_1 \cup \ldots \cup A_n)$ . By the inclusion-exclusion formula.

$$|S - (A_1 \cup \ldots \cup A_n)| = \sum_{I \subseteq \{1,\ldots,n\}} (-1)^I |A_I|.$$

Here  $A_I$  is the set of mappings of X into  $Y = \{y_i : i \in I\}$ , hence

$$|A_I| = (n - |I|)^k$$

So

$$|S - (A_1 \cup \ldots \cup A_n)| = \sum_{j=0}^n (-1)^j \binom{n}{j} (n-j)^k = \sum_{j=0}^n (-1)^{n-j} \binom{n}{j} j^k,$$

which proves the assertion.

If k > n, the result of the same inclusion-exclusion procedure is the number of mappings of X into Y, which is n ! S(k, n). Thus the result is a new proof of 1.8.

5. Put  $x_i = 0$  into  $\sigma^0 p - \sigma^1 p + \dots$  Then, if we set

$$\tilde{p}(x_1, \ldots, x_{i-1}, x_{i+1}, \ldots, x_n) =$$

$$= p(x_1, \ldots, x_{i-1}, 0, x_{i+1}, \ldots, x_n),$$

we have  $\sigma^k p|_{x_i=0} = \sigma^{k-1} \tilde{p} + \sigma^k \tilde{p}$  (a k-tuple of variables of p either contains or does not contain  $x_i$ ),

$$\sigma^{0} p|_{\mathbf{x}_{i=0}} = p|_{\mathbf{x}_{i=0}} = \tilde{p} = \sigma^{0} \tilde{p}, \text{ so } (\sigma^{0} p - \sigma^{1} p + \sigma^{2} p - \dots)|_{\mathbf{x}_{i=0}} = \sigma^{0} \tilde{p} - (\sigma^{0} \tilde{p} + \sigma^{1} \tilde{p}) + (\sigma^{1} \tilde{p} + \sigma^{2} \tilde{p}) - \dots = 0.$$

Therefore,  $x_1 ldots x_n$  divides  $\sigma^0 p - \sigma_1 p + \ldots$  Since  $\sigma^0 p - \sigma^1 p + \ldots$  has degree at most m, it follows that

$$\sigma^0 p - \sigma^1 p + \ldots = \begin{cases} 0 & \text{if} \quad m < n, \\ cx_1 \ldots x_n & \text{if} \quad m = n. \end{cases}$$

Also note that  $\sigma^k p$   $(k \ge 1)$  certainly does not contain any term of the form  $x_1 \dots x_n$ ; hence c is the coefficient of  $x_1 \dots x_n$  in  $\sigma^0 p = p$ .

We remark that for  $p(x_1, \ldots, x_n) = (x_1 + \ldots + x_n)^k$  and  $x_1 = \ldots = x_n = 1$  we obtain the preceding result (cf. problem 4.3).

**6.** Let us express P(B) as the sum of probabilities of atoms; we show that each atom will have a non-negative coefficient. Let

$$B = A_1 \dots A_k \bar{A}_{k+1} \dots \bar{A}_n$$

be any atom of the Boolean algebra generated by  $A_1, \ldots, A_n$ . Define  $A_1' = \ldots = A_k' = 1$ ,  $A_{k+1}' = \ldots = A_n' = 0$ , and let  $B_i' = f_i(A_1', \ldots, A_n')$ . Then  $B \subseteq B_i$  iff  $B_i' \neq 0$ . Therefore, the coefficient of P(B) in  $\sum_{i=1}^k c_i P(B_i)$  is

$$\sum_{B_i \supseteq B} c_i = \sum_{i=1}^k c_i P(B_i') \ge 0,$$

by the assumption [Ré].

7. (a) By the preceding result, we may assume that

$$P(A_1) = \ldots = P(A_k) = 1, \quad P(A_{k+1}) = \ldots = P(A_n) = 0.$$

Then the left-hand side is 1 if k = q and is 0 if  $k \neq q$ ; the right-hand side is

$$\sum_{j=q}^{k} (-1)^{j+q} \binom{k}{j} \binom{j}{q}.$$

If q > k, this sum is, obviously, 0. If q = k, the sum is 1. Finally, if q < k, then

$$\binom{k}{j} \binom{j}{q} = \frac{k!}{q! (j-q)! (k-j)!} = \binom{k}{q} \binom{k-q}{j-q}.$$

So we have

$$\sum_{j=q}^{k} (-1)^{j+q} {k \choose q} {k-q \choose j-q} = {k \choose q} \sum_{j=q}^{k} (-1)^{j-q} {k-q \choose j-q} =$$

$$= {k \choose q} (1-1)^{k-q} = 0 \quad \text{[K. Jordán; Ré]}.$$

(b)  $E\left(\binom{\eta}{j}\right)$  is the expected number of j-tuples of the events  $A_1, \ldots, A_n$  such that all j occur. Since the probability of simultaneous occurrence of a given j-tuple  $\{A_{i_1}, \ldots, A_{i_j}\}$  is  $P(A_{i_1}, \ldots, A_{i_j})$ , we have

$$E\left(inom{\eta}{j}\right) = \sum_{1 \leq i_1 < \cdots < i_l \leq n} P(A_i, \ldots A_{i_l}) = \sigma_j.$$

Thus

$$E(x^{\eta}) = E\left(\sum_{j=0}^{\eta} {\eta \choose j} (x-1)^j\right) = \sum_{j=0}^{n} \sigma_j (x-1)^j.$$

8. By the preceding result, the probability to be determined is

$$\sum_{q=p}^{n} \sum_{j=q}^{n} (-1)^{j+q} {j \choose q} \sigma_{j} = \sum_{j=p}^{n} \sigma_{j} \sum_{q=p}^{j} (-1)^{j+q} {j \choose q} = \sum_{j=p}^{n} \sigma_{j} \sum_{r=0}^{j-p} (-1)^{r} {j \choose r} =$$

$$= \sum_{j=p}^{n} \sigma_{j} (-1)^{j-p} {j-1 \choose j-p} = \sum_{j=p}^{n} (-1)^{j+p} {j-1 \choose p-1} \sigma_{j}$$

by 1.42h [see Ré].

**9.** Let j > 0, be odd. Then we want to show that

$$P(\bar{A}_1 \ldots \bar{A}_n) - \sigma_0 + \sigma_1 - \ldots + \sigma_j \geq 0.$$

By 2.6, we may assume that

$$P(A_1) = \ldots = P(A_k) = 1, \ P(A_{k+1}) = \ldots = P(A_n) = 0.$$

The case k=0 is trivial, so let  $k\geq 1$ . Then we have to show that

$$0-1+\binom{k}{1}-\binom{k}{2}+\cdots+\binom{k}{j}\geq 0,$$

if j is odd. This is clear as by 1.42h,

$$1-\binom{k}{1}+\binom{k}{2}-\ldots-\binom{k}{j}=(-1)^{j}\binom{k-1}{j}.$$

The case of even j follows similarly.

10. I. By 2.6 we may assume again that

$$P(A_1) = \ldots = P(A_k) = 1, \ P(A_{k+1}) = \ldots = P(A_n) = 0.$$

Then

$$\sigma_r = \begin{pmatrix} k \\ r \end{pmatrix}, \qquad \sigma_{r-1} = \begin{pmatrix} k \\ r-1 \end{pmatrix},$$

so what we have to prove is

$$\binom{k}{r} \leq \frac{n-r+1}{r} \binom{k}{r-1}$$

or, dividing by  $\binom{k}{r}$ ,

$$1 \leq \frac{n-r+1}{r} \cdot \frac{r}{k-r+1} = \frac{n-r+1}{k-r+1}$$

which is clear [M. Frèchet; Ré].

II. We prove the inequality given in the hint by the usual method. It suffices to show that

$$\binom{k}{r} \leq \frac{1}{r} \binom{k}{r-1} + (r-1) \binom{k}{r+1}.$$

This is clear, if k < r, so suppose  $k \ge r$ . Dividing by  $\binom{k}{r}$ , we obtain

$$1 \le \frac{1}{r} \cdot \frac{r}{k-r+1} + (r+1) \frac{k-r}{r+1} = 1 + \frac{(k-r)^2}{k-r+1}$$

which is obviously true.

So, if 
$$\sigma_{r+1} = 0$$
, we have  $\sigma_r \leq \frac{1}{r} \sigma_{r-1}$ .

11. I. Suppose we have such events  $A_1, \ldots, A_n$  and let  $J \subseteq \{1, \ldots, n\}$ . By the sieve formula 2.2a

$$P(\prod_{i \notin J} \bar{A}_i \cdot A_J) = \sum_{I \subseteq \{1, \dots, n\} - J} (-1)^{|I|} P(A_I A_J) = \sum_{J \subseteq K \subseteq \{1, \dots, n\}} (-1)^{|K-J|} p_K.$$

2.12

Since a probability is non-negative, we have

(1) 
$$\sum_{J \subseteq K \subseteq \{1, \dots, n\}} (-1)^{|K-J|} p_K \ge 0,$$

for every  $J \subseteq \{1, \ldots, n\}$ . Also, obviously,

$$(2) p_o = 1.$$

II. We show that (1) and (2) are sufficient. Take  $2^n$  atoms  $v_J$  ( $J \subseteq \{1, ..., n\}$ ) and define

$$P(v_J) = \sum_{J \subseteq K \subseteq \{1, \dots, n\}} (-1)^{|K-J|} \, p_K,$$
  $A_o = \{v_J : J \subseteq \{1, \dots, n\}\},$   $A_i = \{v_J : i \in J\} \qquad (i = 1, \dots, n).$ 

Then

$$A_I = \prod_{i \in I} A_i = \{v_J : I \subseteq J \subseteq \{1, \ldots, n\}\}.$$

We claim

$$P(X) = \sum_{v_J \in X} P(v_J)$$

defines a probability measure on  $A_0$  and  $P(A_I) = p_I$ . Let us show the second statement first:

$$\begin{split} P(A_I) &= \sum_{I \subseteq J \subseteq \{1, \dots, n\}} P(v_J) = \sum_{I \subseteq J \subseteq \{1, \dots, n\}} \sum_{J \subseteq K \subseteq \{1, \dots, n\}} (-1)^{|K-J|} p_K = \\ &= \sum_{I \subseteq K \subseteq \{1, \dots, n\}} p_K \sum_{I \subseteq J \subseteq K} (-1)^{|K-J|}. \end{split}$$

Here

$$\sum_{I\subseteq J\subseteq K} (-1)^{|K-J|} = \begin{cases} 1 & \text{if } I=K, \\ 0 & \text{otherwise,} \end{cases}$$

so we have

$$P(A_I) = p_I$$

To show P is a probability measure, we merely note that (1) implies  $P \ge 0$  while (2) implies  $P(A_0) = 1$  [Ré].

12. It suffices to prove the inequality if there is a set  $K \subseteq \{1, \ldots, n\}$  such that

$$P(A_j) = \begin{cases} 1 & \text{if } j \in K, \\ 0 & \text{otherwise,} \end{cases}$$

by 2.6. If  $K = \emptyset$  both sides are 1. So let  $K \neq \emptyset$ . Then we may assume without loss of generality that  $K = \{1, \ldots, n\}$ ; otherwise we could consider the subgraph induced by K. Then the inequality has the form

We prove this together with

$$|\mathfrak{I}_1'| \ge |\mathfrak{I}_1|$$

by induction on n. If G is a complete graph, then

$$|ec{artheta}_0|=n,\;\; |ec{artheta}_0'|=inom{n}{2}+1,\;\; |ec{artheta}_1|=0,\;\; |ec{artheta}_1'|=n$$
 ,

thus (1), (2) are satisfied. Suppose G is not complete and let, say,  $n \in V(G)$  be adjacent to  $1, \ldots, h$  but not adjacent to  $h+1, \ldots, n-1$ . Form the subgraphs  $G^*$ ,  $G^{**}$  of G induced by  $\{1, \ldots, n-1\}$  and  $\{h+1, \ldots, n-1\}$ , respectively. Denote by  $\mathfrak{F}_0^*$ ,  $\mathfrak{F}_0^{**}$  etc. the set  $\mathfrak{F}_0$ , etc. for  $G^*$  and  $G^{**}$ . Then an independent set of G is either an independent set of  $G^*$  or consists of G and an independent set of  $G^{**}$ . Hence

$$|\mathfrak{I}_{i}| = |\mathfrak{I}_{i}^{*}| + |\mathfrak{I}_{1-i}^{**}|.$$

Also, each subset of  $G^*$  which spans at most one edge is such a subset of G; and if X is such a subset of  $G^{**}$ , then  $X \cup \{n\}$  is such a subset of G. But now G may have other subsets inducing at most one edge, e.g.  $\{1, n\}$ . Hence, we can only claim

$$|\mathfrak{I}_{i}'| \geq |\mathfrak{I}_{i}'^{*}| + |\mathfrak{I}_{1-i}'^{**}|.$$

Now by the induction hypothesis,

$$|\mathfrak{I}_{i}^{\prime*}| \geq |\mathfrak{I}_{i}^{*}|, \quad |\mathfrak{I}_{1-i}^{\prime**}| \geq |\mathfrak{I}_{1-i}^{**}|,$$

thus (3) and (4) imply (1) and (2).

The lower estimate follows from (2) exactly as the upper one follows from (1) and reads

$$P(\bar{A}_1 \dots \bar{A}_n) \ge \sum_{I \in \vartheta_1} P(A_I) - \sum_{I \in \vartheta_1'} P(A_I).$$

[A. Rényi, J. Math. Pures Appl. 37 (1958) 393-398].

13. Again we may assume

$$P(A_i) = \begin{cases} 1 & \text{if} \quad i \in K, \\ 0 & \text{otherwise} \end{cases}$$

for some  $K \subseteq \{1, \ldots, n\}$ . If  $K = \emptyset$ , both sides are 1, so suppose  $K \neq \emptyset$ . Then, we have to show

(1) 
$$\sum_{\substack{I\in \mathfrak{F}\\I\subseteq K}} (-1)^{|I|} \geq 0.$$

Let x be the largest element of K. In (1), those pairs of terms  $(-1)^{|I|}$ ,  $(-1)^{|J|}$ , where  $J = I \cup \{x\}$   $(x \notin I)$  obviously cancel each other.

If  $J \in \mathbb{S}$  and  $x \in J$ , then, obviously,  $J - \{x\} \in \mathbb{S}$ . Thus, in (1), only those terms  $(-1)^{|I|}$  are left, where  $x \notin I$  and  $I \cup \{x\} \notin \mathbb{S}$ . We show all such terms are positive. In fact,  $I \cup \{x\} \notin \mathbb{S}$  means

$$|(I \cup \{x\}) \cap \{1,\ldots,k\}| > 2f(k)$$

for some k; since  $I \in \mathfrak{I}$  we must have

$$|I \cap \{1,\ldots,k\}| \leq 2f(k),$$

whence  $k \geq x$  and

$$|I \cap \{1,\ldots,k\}| = 2f(k).$$

But  $I \subseteq K \subseteq \{1, \ldots, x\} \subseteq \{1, \ldots, k\}$ , and thus

$$|I \cap \{1, \ldots, k\}| = |I| = 2f(k),$$

i.e.  $(-1)^{|I|} = (-1)^{2f(k)} = 1$  as stated.

The lower estimate is clearly the following: Let

$$\mathfrak{I}' = \{I \subseteq \{1, \ldots, n\} : |I \cap \{1, \ldots, k\}| \le 2f(k) + 1 \quad (k = 1, \ldots, n)\}$$

then

$$P(\bar{A}_1 \ldots \bar{A}_n) \geq \sum_{I \in \mathcal{S}'} (-1)^{|I|} P(A_I).$$

[For an account of sieve methods, from the number theoretic point of view, see H. Halberstam-K.F. Roth, Sequences, Clarendon Press, Oxford, 1966.]

14. By 2.6, we may assume

(1) 
$$P(A_1) = \ldots = P(A_k) = 1, \quad P(A_{k+1}) = \ldots = P(A_n) = 0.$$

If k = 0, then  $P(A_{I \cup J}) = 0$  except when  $I = J = \emptyset$  and then we have to show that

$$1 \leq \lambda_a^2$$

which is clear. If k > 0, then we have to verify that

$$0 \leq \sum_{I, J \subseteq \{1, \dots, k\}} \lambda_I \lambda_J = (\sum_{I \subseteq \{1, \dots, k\}} \lambda_I)^2,$$

which is again clear.

To prove the lower estimate given in the hint, we may again assume that

$$P(A_i) = \begin{cases} 1 & \text{if } i \in K, \\ 0 & \text{otherwise} \end{cases}$$

(since the ordering of the indices is involved in the formula we may not assume this time that K is a section  $\{1, \ldots, k\}$ ). The relation is trivial if  $K = \emptyset$   $(1 \ge 1)$ , so suppose  $K \ne \emptyset$ . Then we have to show

$$0 \ge 1 - \sum_{\substack{I, J \subseteq K \\ \max I = \max J}} \lambda_I \lambda_J.$$

Let

$$K = \{k_1, \ldots, k_{\nu}\}, \quad k_1 < \ldots < k_{\nu},$$

then this inequality can be written as

$$1 \leq \sum_{j=1}^{r} \sum_{\substack{I \subseteq K \\ \max{I} = k_j}} \sum_{\substack{J \subseteq K \\ \max{I} = k_j}} \lambda_I \lambda_J = \sum_{j=1}^{r} \left( \sum_{\substack{I \subseteq K \\ \max{I} = k_j}} \lambda_I \right)^2 = \lambda_{\{k_1\}}^2 + \sum_{j=2}^{r} \left( \sum_{\substack{I \subseteq K \\ \max{I} = k_j}} \lambda_I \right)^2,$$

which is clearly true.

15. We have

$$\sum_{\substack{I,J\subseteq\{1,\cdots,n\}\\|I|\leq k}}\lambda_I\lambda_J\,p_{I\cup J}=\sum_{\substack{|I|\leq k\\|J|\leq k}}\lambda_I\,\lambda_J\,p_{I\cup J}=\sum_{\substack{|I|\leq k\\|J|\leq k}}\lambda_I\,\lambda_J\,p_I\,p_J\sum_{K\subseteq I\cap J}\frac{q_K}{p_K}$$

because

$$p_{I \cup J} = p_I p_J / p_I \cap \mathbf{j}$$

and

$$\sum_{K\subseteq I\cup J} \frac{q_K}{p_K} = \prod_{i\in I\cap J} \left(1 + \frac{q_i}{p_i}\right) = \frac{1}{p_{I\cap J}}.$$

Thus

$$\begin{split} \sum_{\substack{|I| \leq k \\ |J| \leq k}} \lambda_I \lambda_J p_{I \cup J} &= \sum_{|K| \leq k} \frac{q_K}{p_K} \sum_{\substack{K \subseteq I \\ |I| \leq k}} \sum_{\substack{K \subseteq J \\ |J| \leq k}} \lambda_I \lambda_J p_I p_J = \\ &= \sum_{K} \frac{q_K}{p_K} \left( \sum_{\substack{K \subseteq I \\ |I| \leq k}} p_I \lambda_I \right)^2 = \sum_{K} \frac{q_K}{p_K} \mu_K^2 \,, \end{split}$$

where

$$\mu_K = \sum_{\substack{K \subseteq I \\ |I| \le k}} p_I \lambda_I$$

Here we can express  $\lambda_I$  as

(2) 
$$\lambda_{I} = \frac{1}{p_{I}} \sum_{\substack{I \subseteq K \\ |K| \le k}} (-1)^{|K-I|} \mu_{K},$$

because the  $\mu_K$ 's obviously determine the  $\lambda_I$ 's and the  $\lambda_I$ 's defined by (2) satisfy (1):

$$\sum_{\substack{K\subseteq I\\|I|\leq k}}p_I\lambda_I=\sum_{\substack{K\subseteq I\\|I|\leq k}}\sum_{\substack{I\subseteq J\\|J|\leq k}}(-1)^{|J-I|}\,\mu_J=\sum_{\substack{K\subseteq J\\|J|\leq k}}\mu_J\sum_{K\subseteq I\subseteq J}(-1)^{|J-I|}=\mu_K.$$

Thus we have to minimize

$$\sum_{|K| \le k} \frac{q_K}{p_K} \mu_K^2$$

subject to

(4) 
$$\sum_{|K| \le k} (-1)^{|K|} \mu_K = 1.$$

This can be done either by the Lagrange method or simply by the transformation:

(5) 
$$\sum_{|K| \le k} \frac{q_K}{p_K} \mu_K^2 = \sum_{|K| \le k} \frac{p_K}{q_K} \left( \frac{q_K}{p_K} \mu_K - \frac{(-1)^{|K|}}{Q} \right)^2 + \frac{1}{Q},$$

where

$$Q = \sum_{|K| \leq k} rac{p_K}{q_K},$$

which holds under the assumption (4). Since

(6) 
$$\mu_K = \frac{1}{Q} \cdot \frac{p_K}{q_K} \cdot (-1)^{|K|}$$

is a choice of the variables which minimizes the right-hand side of (5) and also satisfies the constraint (4), we get that the minimum of (3) is 1/Q and (6) gives the extremum. Hence

$$\lambda_I = \frac{1}{p_I} \sum_{\substack{I \subseteq K \\ |K| \le k}} (-1)^{|K-I|} \frac{1}{Q} \, \frac{p_K}{q_K} (-1)^{|K|} = \frac{(-1)^{|I|}}{Q q_I} \sum_{\substack{L \subseteq \{1, \dots, n\} - I \\ |L| \le k - |I|}} \frac{p_L}{q_L}.$$

16. By 2.14,

$$P(\bar{A}_1 \dots \bar{A}_n) \leq \sum_{I,J \subseteq \{1,\dots,n\}} \lambda_I \lambda_J P(A_{I \cup J})$$

for any choice of  $\lambda_I$  ( $I \subseteq \{1, \ldots, n\}$  with the only restriction that  $\lambda_0 = 1$ . We will only use  $\lambda_I$ 's that satisfy the requirement  $\lambda_I = 0$  for  $I \notin \mathcal{X}$ . Setting

$$P(A_I) = p_I + R_I,$$

we have

$$(1) P(\bar{A}_1 \dots \bar{A}_n) \leq \sum_{I,J \in \mathcal{X}} \lambda_I \lambda_J p_{I \cup J} + \sum_{I,J \in \mathcal{X}} \lambda_I \lambda_J R_{I \cup J}.$$

As in the preceding solution, the choice

(2) 
$$\lambda_I = \frac{(-1)^{|I|}}{Qq_I} \sum_{\substack{L \subseteq \{1, \dots, n\}-I \\ L \cup I \in \mathcal{X}}} \frac{p_L}{q_L}$$

minimizes the first term and then the value of it is 1/Q, where

$$Q = \sum_{K \in \mathcal{R}} \frac{p_K}{q_K}.$$

We now estimate the second term in (1) when  $\lambda_I$  is given by (2). We have

$$|\lambda_I| \leq \frac{1}{Qq_I} \sum_{\substack{L \subseteq \{1,\dots,n\}-I \ I \perp I \in \mathscr{X}}} \frac{p_L}{q_L} \leq \frac{1}{q_I},$$

because  $L \cup I \in \mathcal{H}$  implies  $L \in \mathcal{H}$ . Thus

$$\begin{split} \left| \sum_{I,J \in \Re} \lambda_I \, \lambda_J R_{I \cup J} \right| &\leq \varepsilon \sum_{I,J \in \Re} \frac{1}{q_I} \, \frac{1}{q_J} = \varepsilon \left( \sum_{I \in \Re} \frac{1}{q_I} \right)^2 \leq \varepsilon \left( \sum_{I \in \Re} \frac{M p_I}{q_I} \right)^2 \leq \\ &\leq \varepsilon \, M^2 \left( \sum_{I \subseteq \{1,\dots,n\}} \frac{p_I}{q_I} \right)^2 = \varepsilon M^2 \left( \prod_{i=1}^n \left( 1 + \frac{p_i}{q_i} \right) \right)^2 = \frac{\varepsilon M^2}{q_1^2 \dots q_n^2} \,. \end{split}$$

The first term can be estimated as follows:

$$Q = \sum_{K \in \mathcal{X}} \frac{p_K}{q_K} = \sum_{K \in \mathcal{X}} \prod_{k \in K} \frac{p_k}{1 - p_k} = \sum_{K \in \mathcal{X}} \prod_{k \in K} (p_k + p_k^2 + \dots).$$

If we multiply out we get exactly those terms  $p_1^{l_1} \dots p_n^{l_n}$  for which  $\prod_{l_i > 0} p_i \ge \frac{1}{M}$ ; so we certainly get all terms of this form which are themselves at least  $\frac{1}{M}$ . Thus

and the assertion follows.

17. Let  $P_1, \ldots, P_n, M$  be defined as in the hint. Let us choose one of the numbers  $l, k+l, \ldots, (x-1)k+l$  at random and let  $A_i$  be the event that it is divisible by  $P_i$ . Set

$$P_I = \prod_{i \in I} P_i \quad (I \subseteq \{1, \ldots, n\}).$$

 $x \cdot P(A_I)$  is the number of multiples of  $P_I$  among  $l, k+l, \ldots, (x-1)$  k+l. Since  $(P_I, k) = 1$ ,  $P_I$  has one multiple among  $\mu P_I \cdot k + l$ ,  $(\mu P_I + 1)$  k + l, ...,  $(\mu P_I + P_I - 1)$  k + l for every  $0 \le \mu \le \frac{x}{P_I} - 1$ . Hence

$$\left[\frac{x}{P_I}\right] \! \leq \! x P(A_I) \leq \! \left[\frac{x}{P_I}\right] + 1$$

and thus

$$\left| P(A_I) - \frac{1}{P_I} \right| \leq \frac{1}{x} \,.$$

Thus if we set  $p_i = \frac{1}{P_i}$ ,  $\varepsilon = \frac{1}{x}$ , the conditions of 2.16 are satisfied and hence,

$$P(\bar{A}_1 \dots \bar{A}_n) \leq \frac{1}{S} + \frac{\varepsilon M^2}{(1-p_1)^2 \dots (1-p_n^2)}$$
.

Now here

$$S = \sum \frac{1}{P_1^{l_1} \dots P_n^{l_n}},$$

where the summation extends over all choices of  $l_1, \ldots, l_n \geq 0$  with  $P_1^{l_1} \ldots P_n^{l_n} \leq M$ . This sum is exactly

$$S = \sum_{\substack{m \le M \\ (m,k)=1}} \frac{1}{m}.$$

To get a lower bound on this, observe that

$$\left(\sum_{\substack{m \leq M \\ (m,k)=1}} \frac{1}{m}\right) \prod_{P \mid k} \left(1 + \frac{1}{P} + \frac{1}{P^2} + \ldots\right) \geq \sum_{m \leq M} \frac{1}{m} > \log M,$$

whence

$$S \ge \log M \, \frac{1}{\displaystyle \prod_{P \mid k} \left(1 + \frac{1}{P} + \ldots\right)} = \log M \, \prod_{P \mid k} \left(1 - \frac{1}{P}\right) = \log M \, \frac{\varphi(k)}{k} \, .$$

On the other hand, it is known that

$$(1-p_1)^2\dots(1-p_n)^2 \ge \prod_{P\le M} \left(1-\frac{1}{P}\right)^2 \ge \frac{c}{\log^2 M}$$

and thus

$$P(\bar{A}_1 \dots \bar{A}_n) \leq \frac{k}{\varphi(k) \cdot \log M} + \frac{M^2 \log^2 M}{cx} \leq \frac{k}{\varphi(k)} \frac{2.5}{\log x}$$

if x is large enough. Thus the number of primes in the sequence  $l, k + l, \ldots, (x-1) k + l$  is

$$\leq n + x P(\bar{A}_1 \dots \bar{A}_n) \leq \sqrt{x} + 2.5 \frac{k}{\varphi(k)} \frac{x}{\log x} \leq 3 \frac{k}{\varphi(k)} \frac{x}{\log x}$$

if x is large enough.

18. We prove that

$$(1) P(A_1 \mid \bar{A}_2 \dots \bar{A}_n) \leq \frac{1}{2d}$$

by induction on n. This, in fact, will imply that

$$P(\bar{A}_1 \dots \bar{A}_n) = P(\bar{A}_2 \dots \bar{A}_n) P(\bar{A}_1 \mid \bar{A}_2 \dots \bar{A}_n) \ge P(\bar{A}_2 \dots \bar{A}_n) \left(1 - \frac{1}{2d}\right) > 0$$

because by the induction hypothesis,  $P(\bar{A}_2 \dots \bar{A}_n) > 0$  (this assures that  $P(A_1 \mid \bar{A}_2 \dots \bar{A}_n)$  is meaningful).

Let, say,  $2, \ldots, h$  be the neighbors of 1. We have

$$P(A_1 \mid \bar{A}_2 \dots \bar{A}_n) = \frac{P(A_1 \bar{A}_2 \dots \bar{A}_h \mid \bar{A}_{h+1} \dots \bar{A}_n)}{P(\bar{A}_2 \dots \bar{A}_h \mid \bar{A}_{h+1} \dots \bar{A}_n)}.$$

Here the numerator is

$$P(A_1 \bar{A}_2 \dots \bar{A}_h \mid \bar{A}_{h+1} \dots \bar{A}_n) \leq P(A_1 \mid \bar{A}_{h+1} \dots \bar{A}_n) = P(A_1) \leq \frac{1}{4d}$$

(as  $A_1$  is independent of  $A_{h+1}, \ldots, A_n$ ) while for the denominator we have

$$P(\bar{A}_{2}....\bar{A}_{h} | \bar{A}_{h+1}...\bar{A}_{n}) = 1 - P(\bar{A}_{2} + ... + \bar{A}_{h} | \bar{A}_{n+1}...\bar{A}_{n}) \geq$$

$$\geq 1 - \sum_{i=2}^h P(A_i \mid \bar{A}_{h+1} \dots \bar{A}_n).$$

Now applying our induction hypothesis with the graph induced by  $\{i, h + 1, \ldots, n\}$  we get

$$P(A_i \mid A_{h+1} \dots A_n) \leq \frac{1}{2d}$$

and so

$$P(\bar{A}_2 \dots \bar{A}_h \mid \bar{A}_{h+1} \dots \bar{A}_n) \ge 1 - \frac{h-1}{2d} \ge \frac{1}{2}$$

as h-1 is the degree of 1 and so it is at most d. Thus

$$P(A_1 \mid ar{A}_2 \dots ar{A}_n) \leq rac{rac{1}{4d}}{rac{1}{2}} = rac{1}{2d}.$$

[P. Erdős-L. Lovász, in: Infinite and Finite Sets, Coll. Math. Soc. J. Bolyai 10 (1974) Bolyai-North Holland, 609-627.]

19. We have

$$P(\bar{A}_1 \dots \bar{A}_n) = P(\zeta = 0) \leq P((\xi - \sigma_1)^2 \geq \sigma_1^2) \leq$$

$$\leq \frac{1}{\sigma_1^2} E((\zeta - \sigma_1)^2) = \frac{1}{\sigma_1^2} (E(\zeta^2) - \sigma_1^2).$$

Now let

$$\zeta_i = egin{cases} 1 & ext{if } A_i ext{ occurs,} \ 0 & ext{otherwise,} \end{cases}$$

then

$$\sum_{i=1}^n \zeta_i = \sum_{i=1}^n \zeta_i$$

and thus,

$$E(\zeta^2) = E\left(\sum_{i=1}^n \zeta_i^2 + 2\sum_{1 \leq i < j \leq n} \zeta_i \zeta_j
ight) = \sigma_1 + 2\sigma_2,$$

whence

$$P(\bar{A}_1 \dots \bar{A}_n) \leq \frac{1}{\sigma_1^2} (\sigma_1 + 2\sigma_2 - \sigma_1^2).$$

The first step is actually Chebyshev's inequality;

$$P(\zeta=0) \leq P\left(\left(|\zeta-E(\zeta)| \geq \frac{E(\zeta)}{D(\zeta)}D(\zeta)\right) \leq \frac{D^2(\zeta)}{E^2(\zeta)} = \frac{E\left((\zeta-\sigma_1)^2\right)}{\sigma_1^2}\right).$$

20. I. By 2.9,

$$P(\bar{A}_1 \dots \bar{A}_n) \leq 1 - np + \binom{n}{2} p^2 = \alpha.$$

II. By Selberg's method,

$$P(\bar{A}_1 \ldots \bar{A}_n) \leq \sum_{|I|,|I| \leq 1} \lambda_I \lambda_J p^{|I \cup J|},$$

where  $\lambda_0 = 1$ ,  $\lambda_{\{i\}}$  arbitrary. By 2.15, the  $\lambda_I$ 's can be chosen so that the right-hand side becomes

$$rac{1}{\sum\limits_{|I| \leq 1} rac{p^{|I|}}{(1-p)^{|I|}}} = rac{1}{1+rac{np}{1-p}} = eta \, .$$

III. The preceding formula gives

$$P(\bar{A}_1 \dots \bar{A}_n) \leq \frac{1-p}{np} = \gamma$$

by direct substitution.

Obviously,  $\gamma \geq \beta$ , thus II is better than III. However, I and II are incomparable. For let p be fixed and  $n \to \infty$ , then  $\beta \to 0$  but  $\alpha \to \infty$ , thus II even III is better if n is large. On the other hand, fix n and let  $p \to 0$ . Then the power series of  $\beta$  is

$$1-np+n(n-1)p^2+\ldots,$$

which is larger than  $\alpha$  if p is very small. 30-600 broaded the

**21.** It is trivial that sum of compatible matrices is compatible. Let  $A=(a_{ij})$ ,  $B=(b_{ij})$  be compatible matrices,  $AB=C=(c_{ij})$  and suppose  $c_{ij}\neq 0$ . Since

$$c_{ij} = \sum_k a_{ik} b_{kj}$$
 ,

there exists a k such that  $a_{ik} \neq 0$  and  $b_{kj} \neq 0$  Hence  $x_i \leq x_k$  and  $x_k \leq x_j$ , and so, by transitivity,  $x_i \leq x_j$ . Thus C is compatible. Let A be an invertible compatible matrix. We may assume without loss

Let A be an invertible compatible matrix. We may assume without loss of generality that  $x_i \leq x_j \Rightarrow i \leq j$  (this can be achieved by indexing appropriately). Then, it is seen that all the non-zero entries of A are above the

diagonal, so det  $A = \prod_{i=1}^{n} a_{ii}$ . So  $a_{ii} \neq 0$ . Now let  $B = (b_{ij})$  be the inverse of

A, and suppose indirectly that it is not compatible, i.e. there are  $x_i \leq x_j$  with  $b_{ij} \neq 0$ . Choose i here to be maximal. We have

$$\sum_{k=1}^n a_{ik} b_{kj} = 0,$$

but

$$a_{ii}b_{ij}\neq 0$$

so there must be a  $k \neq i$  with

$$a_{ik}b_{kj}\neq 0.$$

Since  $(a_{ij})$  is compatible, this implies that  $x_i \leq x_k$  and so, i < k. By the choice of i,  $b_{kj} \neq 0$  implies  $x_k \leq x_j$  and so,  $x_i \leq x_j$ , a contradiction.

[Most problems in the rest of this section are based on G. C. Rota's paper in: Zeitschr. f. Wahrscheinlichkeitstheorie, 2 (1964) 340-368.]

22. Define

$$z_{ij} = \begin{cases} 1 & \text{if } x_i \leq x_j, \\ 0 & \text{otherwise,} \end{cases}$$

 $Z=(z_{ij})$ . Then the two requirements on  $\mu$  can be re-written as

$$ZM = I$$

where  $M = (m_{ij})$  with  $m_{ij} = \mu(x_i, x_j)$  and I is the identity matrix. Since  $z_{ii} \neq 0$ , Z is invertible (see the preceding solution) and so, by the preceding

result,  $M = Z^{-1}$  is a compatible matrix. This defines the desired function  $\mu(x, y)$ .

23. Note that if we know  $\mu(a, y)$  for every y such that  $a \leq y < b$ , then the equation

$$\sum_{a \le y \le b} \mu(a, y) = 0$$

uniquely determines  $\mu(a, b)$ . Therefore, we can calculate  $\mu(a, x)$  for every  $a \le x \le b$  inductively using only elements between a and b. Hence, if  $a \le b \in V$ ,  $a' \le b' \in V'$  and the partially ordered sets  $\{z \in V : a \le z \le b\}$ ,  $\{z' \in V' : a' \le z' \le b'\}$  are isomorphic as partially ordered sets, then

$$\mu(a, b) = \mu(a', b').$$

(a) We claim that  $\mu(X, Y) = (-1)^{|Y-X|}$   $(X \subseteq Y \subseteq S)$ . In fact,

$$\sum_{A \subseteq Y \subseteq B} (-1)^{|Y-A|} = \sum_{k=0}^{|B-A|} {|B-A| \choose k} (-1)^k = \begin{cases} 1 & \text{if } B = A, \\ 0 & \text{if } B \supset A. \end{cases}$$

(b) Here, any interval  $a \le z \le b$  is a chain. So it suffices to calculate  $\mu(a_1, a_n)$  for a chain

$$a_1 < a_2 < \ldots < a_n$$

We have by definition,

$$\mu(a_1,a_1)=1$$
. An investigate of this is  $\lambda$  thus

Also,

$$\mu(a_1, a_1) + \mu(a_1, a_2) = 0,$$

whence

$$\mu(a_1,a_2)=-1.$$

Now, for  $i \geq 3$ ,

$$\mu(a_1, a_1) + \mu(a_1, a_2) + \ldots + \mu(a_1, a_i) = 0,$$

whence, obviously,

$$\mu(a_1, a_3) = \ldots = \mu(a_1, a_n) = 0.$$

Thus

$$\mu(x, y) = \begin{cases} 1 & \text{if } x = y, \\ -1 & \text{if } (x, y) \text{ is an edge,} \end{cases}$$

(c) We claim that

(1) 
$$\mu(x, y) = \mu\left(\frac{y}{x}\right),$$

where  $\mu(k)$  is the (number theoretical) Moebius function: if  $k = p_1^{z_1} \dots p_r^{z_r}$   $(p_1, \dots, p_r$  distinct primes), then

$$\mu(k) = \begin{cases} (-1)^r & \text{if} \quad \alpha_1 = \ldots = \alpha_r = 1, \\ 0 & \text{otherwise.} \end{cases}$$

2.24

Obviously,  $\mu(x, y)$  defined by (1) satisfies  $\mu(a, a) = 1$ . Also, if  $a \neq b$ ,  $a \mid b$ , then

$$\sum_{a|y|b} \mu(a, y) = \sum_{a|y|b} \mu\left(\frac{y}{a}\right) = \sum_{d\mid \frac{b}{a}} \mu(d).$$

Set

$$\frac{b}{a}=k=p_1^{\alpha_1}\dots p_r^{\alpha_r},$$

then

$$\sum_{d|k} \mu(d) = \sum_{1 \leq i_1 < \cdots < i_{\nu} \leq r} \mu(p_{i_1} \cdots p_{i_{\nu}}) = \sum_{\nu=0}^{r} {r \choose \nu} (-1)^{\nu} = 0.$$

24. We have to show

$$\mu^*(a,a) = 1,$$

$$\sum_{a \leq {}^*\mathbf{x} \leq {}^*b} \mu^*(a,x) = 0 \qquad (a \neq b),$$

i.e.

$$\mu(a, a) = 1,$$

$$\sum_{b \le x \le a} \mu(x, b) = 0.$$

The first assertion is trivial. To show the second one, note that, in terms of M and Z, it can be expressed as

$$Z \cdot M = I$$

which is obvious, because  $M = Z^{-1}$ .

**25.** Let  $Z - I = (u_{ij})$ , then

$$u_{ij} = \begin{cases} 1 & \text{if } x_i < x_j, \\ 0 & \text{otherwise.} \end{cases}$$

Then the  $(i, j)^{th}$  entry of  $(Z - I)^n$  is

$$\sum_{k_1,\ldots,k_{n-1}} u_{ik_1} u_{k_1k_2} \ldots u_{k_{n-1}j} = 0,$$

because a non-zero term would correspond to a chain  $x_i < x_{k_1} < x_{k_2} < \dots < x_{k_{n-1}} < x_j$ , which clearly does not exist. Hence

$$(I - Z)^n = 0$$

or, equivalently,

$$\sum_{i=1}^{n} \binom{n}{i} (-1)^{i-1} Z^i = I.$$

Multiplying by M, we get

$$\sum_{i=1}^{n} \binom{n}{i} (-1)^{i-1} Z^{i-1} = M.$$

Note that instead of n, we could consider the maximum length of chains in V.

**26.** 
$$\sum_{z \le x} g(z) \ \mu(z, x) = \sum_{z \le x} \sum_{y \le z} f(y) \ \mu(z, x) = \sum_{y \le x} f(y) \sum_{y \le z \le x} \mu(z, x) = f(x).$$

(Note: setting  $f = (f(x_1), \ldots, f(x_n))$ ,  $g = (g(x_1), \ldots, g(x_n))$ , the definition of g means

$$g = f \cdot Z$$

and the assertion is

$$f = gM = gZ^{-1}.$$

To get the sieve formula let V consist of all subsets of  $\{1, \ldots, n\}$ , and let  $\leq$  mean inclusion. By 2.23,  $\mu(K, L) = (-1)^{|L-K|}$   $(K \subseteq L)$ . Set

$$f(K) = P\left(\prod_{i \in K} A_i \prod_{j \in K} \bar{A}_j\right),\,$$

then

$$g(L) = \sum_{K \subseteq L} f(K) = P\left(\prod_{i \notin L} A_i\right) = P(A_{\overline{L}})$$

 $(\overline{K} = \{1, \ldots, n\} - K)$ , and so,

$$P(\prod_{i=1}^{n} \bar{A}_{i}) = f(\{1, \dots, n\}) = \sum_{K \subseteq \{1, \dots, n\}} (-1)^{n-|K|} P(A_{\overline{K}}) = \sum_{K \subseteq \{1, \dots, n\}} (-1)^{|K|} P(A_{K}).$$

27. (a) We proceed by induction on the number of elements between a and b. If a = b, then the sum looks like

$$\sum_{x\leq b}\mu(0,x)\,,$$

which is 0 by the definition of  $\mu$ . Let b > a. Then

$$\sum_{x \leq b} \mu(0, x) = \sum_{a \leq b_1 \leq b} \left( \sum_{a \vee x = b_1} \mu(0, x) \right).$$

The left-hand side is 0 by the definition of  $\mu$ . All terms on the right-hand side with  $b_1 < b$  are 0's by the induction hypothesis. Thus, the last one is 0 as well.

- (b) follows immediately from (a) and 2.24 by inverting the order.
- 28. We use induction on the number of elements below x. If x = 0 the assertion is true (0 is the union of some atoms). Let x > 0. If  $a \le x$  is any atom, then, by the preceding result,

$$\sum_{y,y,q=x}\mu(0,y)=0.$$

Here, obviously, no y is a union of atoms. So each term with y < x is 0 by the induction hypothesis. It follows that so is  $\mu(0, x)$ .

29. We have

$$\sum_{y \ge a} \mu(a, y) = \begin{cases} 1 & \text{if } a = 1, \\ 0 & \text{otherwise,} \end{cases}$$

SO

$$\sum_{x,y} \mu(x,y) = 1.$$

On the other hand,

(1) 
$$\sum_{x,y} = \sum_{x \in A, y \in C} + \sum_{x,y \in A \cup B} + \sum_{x,y \in B \cup C} - \sum_{x,y \in B}$$

Here

$$\sum_{x,y\in A\cup B}\mu(x,y)=\sum_{y\in A\cup B}\sum_{x\leq y}\mu(x,y)=1$$
 , and the second back

and similarly,

$$\sum_{x,y\in B\cup C}\mu(x,y)=\sum_{x\in B\cup C}\sum_{y\geq x}\mu(x,y)=1.$$

Hence (1) has the form

$$1 = \sum_{x \in A} \sum_{y \in C} \mu(x, y) + 1 + 1 - \sum_{x, y \in B} \mu(x, y),$$

which proves the assertion.

**30.** Let  $b = \{\{1\}, \{2, \ldots, n\}\}$  and apply 2.27:

$$\sum_{x \land b=0} \mu(x, 1) = 0$$

or, equivalently,

(1) 
$$\mu(0,1) = -\sum_{\substack{x \land b = 0 \\ x \neq 0}} \mu(x,1).$$

Determine the partitions x with  $x \wedge b = 0$ ,  $x \neq 0$ . Let  $x = \{A_1, \ldots, A_p\}$  be such a partition. Let  $1 \in A_1$  say. Then for i > 1,  $A_i = A_i \cap \{2, \ldots, n\}$  is a class of  $x \wedge b = 0$ , hence  $|A_i| = 1$ . As  $x \neq 0$ ,  $|A_1| \geq 2$ . But  $A_1 \cap \{2, \ldots, n\}$  is also a class of  $x \wedge b = 0$ , so  $|A_1| = 2$ . So x is of the form

$$x = \{\{1, i\}, \{2\}, \dots, \{i-1\}, \{i+1\}, \dots, \{n\}\}.$$

The interval  $\{z : x \leq z \leq 1\}$  consists of all partitions which do not separate 1 and i. This interval is, obviously, isomorphic to the lattice of all partitions of n-1 elements (1 and i "stick together"). Hence

$$\mu(x, 1) = (-1)^{n-2} (n-2)!$$

by the induction hypothesis. Since there are n-1 partitions x to consider, (1) implies

$$\mu(0, 1) = -(n-1) (-1)^{n-2} (n-2)! = (-1)^{n-1} (n-1)!.$$

[M. P. Schützenberger].

31. If F is defined as in the hint, we have

$$Z^TFZ=G$$
.

In fact,

$$\sum_{k=1}^{n} \sum_{l=1}^{n} z_{ki} f_{kl} z_{lj} = \sum_{k=1}^{n} z_{ki} f_{kk} z_{kj} = \sum_{\substack{k \\ x_k \leq x_i \\ x_k \leq x_j}} f(x_k) = \sum_{\substack{k \\ x_k \leq x_i \land x_j}} f(x_k) = g(x_i \land x_j) = g_{ij}.$$

Hence

$$\det G = \det (Z^T F Z) = (\det Z^T) (\det F) (\det Z) = \det F = f(x_1) \dots (f(x_k)).$$

If V is only a partially ordered set, then we define F as before and consider

$$G = Z^T F Z = (g_{ij}).$$

We have, as before,

$$g_{ij} = \sum_{k=1}^{n} \sum_{l=1}^{n} z_{ki} f_{kl} z_{lj} = \sum_{\substack{x_k \leq x_i \\ x_k \leq x_j}} f(x_k).$$

We cannot rewrite this in terms of  $x_i \wedge x_j$  (as this is not defined) so we have to state the formula as follows: let f(x) be any function on V and set

$$g_{ij} = \sum_{\substack{x \le x_i \\ x \le x_j}} f(x).$$

Then

$$\det (g_{ij}) = f(x_1) \dots f(x_n).$$

Note, that if any two elements  $x_i$ ,  $x_j$  of V have a greatest lower bound  $x_i \wedge x_j$ , then

$$g_{ij} = \sum_{\mathbf{x} \leq \mathbf{x}_i \wedge \mathbf{x}_j} f(\mathbf{x}) = g(\mathbf{x}_i \wedge \mathbf{x}_j),$$

where

$$g(y) = \sum_{x \le y} f(x)$$

[H. S. Wilf, Bull. Amer. Math. Soc. 74 (1968) 960-964].

**32.** I. Let  $V = \{1, \ldots, n\}$ , let the partial order be defined divisibility and g(x) = x. We try to apply the preceding formula  $(V \text{ is not a lattice but it is closed under the operation <math>x \land y = (x, y)$  and so, the same formula holds by the remark at the end of the previous solution). To get a suitable f(x) we use 2.26:

$$f(x) = \sum_{y \le x} \mu(y, x) \ g(y) = \sum_{y \mid x} y \mu\left(\frac{x}{y}\right) = \varphi(x)$$

by the solution of 2.3. So the value of the determinant is  $\varphi(1) \cdot \varphi(2) \dots \varphi(n)$ 

II. This determinant is just  $\varphi(1) \dots \varphi(n) \times \{\text{the upper left entry in } G^{-1}\}$  where, as in the preceding solution

$$G = oldsymbol(i,j)_{i,j=1}^n = Z^T F Z \,,$$
  $F = (f_{ij}) = egin{pmatrix} arphi(1) & 0 \ & \cdot \ & 0 & arphi(n) \end{pmatrix},$   $Z = (z_{ij}), \, z_{ij} = egin{bmatrix} 1 & ext{if} & i \mid j \,, \ 0 & ext{otherwise}. \end{pmatrix}$ 

Thus

$$G^{-1} = (Z^T F Z)^{-1} = M F^{-1} M^T,$$

where  $M = (m_{ij})$ ,

$$m_{ij} = egin{cases} \mu\left(rac{j}{i}
ight) & ext{if} \quad i \mid j, \ 0 & ext{otherwise} \end{cases}$$

Thus the upper left corner entry of  $G^{-1}$  is

$$\sum_{k=1}^{n} \sum_{l=1}^{n} \mu(1, k) (F^{-1})_{kl} \mu(1, l) = \sum_{k=1}^{n} \frac{\mu^{2}(k)}{\varphi(k)}.$$

Hence

$$\begin{vmatrix} (2,2)\ldots(2,n) \\ \vdots \\ (n,2)\ldots(n,n) \end{vmatrix} = \varphi(1)\ldots\varphi(n) \sum_{k=1}^{n} \frac{\mu^{2}(k)}{\varphi(k)}.$$

33. Let

$$A=egin{bmatrix} 0&1&\dots&1\ 1&-2&&&&0\ &\ddots&&&&&\ &\ddots&&&&&\ &1&0&1&\dots&-2 \end{pmatrix}.$$

Consider  $x_1$  as a root, then T can be considered as an arborescence and the defines an ordering of V = V(T):  $x_i \leq x_j$ , if the (unique)  $(x_i, x_j)$ -path contains  $x_i$ . Then let Z be as in 2.22; we claim that

$$Z^TAZ = D.$$

In fact, the (i, j)<sup>th</sup> entry of  $Z^TAZ$  is

$$d'_{ij} = \sum_{k=1}^{n} \sum_{l=1}^{n} z_{ki} a_{kl} z_{lj} = \sum_{\mathbf{x}_k \leq \mathbf{x}_k} \sum_{\mathbf{x}_l \leq \mathbf{x}_j} a_{kl}.$$

Now  $a_{kl}$  is non-zero only if k = l or k = 1 or l = 1 so we have

$$egin{aligned} d'_{ij} &= \sum\limits_{x_k \leq x_i, \, x_j} (-2) + \sum\limits_{x_k \leq x_i} 1 + \sum\limits_{x_l \leq x_j} 1 = \ &= -2[d(x_i \, igwedge \, x_j, \, x_1) + 1] + [d(x_i, \, x_1) + 1] + [d(x_j, \, x_1) + 1], \end{aligned}$$

where  $x_i \wedge x_j$  is the last common point of the  $(x_1, x_i)$  and  $(x_1, x_j)$ -paths. Hence

$$d'_{ij} = d(x_i, x_1) + d(x_j, x_1) - 2d(x_i \wedge x_j, x_1) = d(x_i, x_j)$$

as stated. Thus

$$\det D = \det A = -(n-1)(-2)^{n-2}$$

[R. L. Graham-H. O. Pollak, Bell Sys. Tech. J. 50 (1971) 2495-2519].

34. Set  $U = Z - I = (u_{ij})$ . Then  $U^n = 0$ . Further we have

$$(I+U)(I-U+U^2-U^3+\ldots+(-1)^{n-1}U^{n-1})=I+(-1)^{n-1}U^n=I.$$

So

$$I-U+U^2-U^3+\ldots=(I+U)^{-1}=Z^{-1}=M.$$

Hence

$$\mu(0,1) = \sum_{\nu=1}^{n} (-1)^{\nu} p'_{\nu},$$

where  $p'_{\nu}$  is the  $(0,1)^{\text{th}}$  entry in  $U^{\nu}$ ; but this entry is

$$p'_{\nu} = \sum_{i_1, \dots, i_{\nu}} u_{0i_1} u_{i_1 i_2} \dots u_{i_{\nu-1} 1} = \sum_{0 < x_{i_1} < \dots < x_{i_{\nu-1}} < 1} 1 = p_{v}.$$

35. Set

$$q(x) = q_0(x) - q_1(x) + q_2(x) - q_3(x) + \dots$$

Then

$$\sum_{x \le y} q(x) = \sum_{k=0}^{n} (-1)^k \sum_{x \le y} q_k(x).$$

Now, let  $a_1, \ldots, a_m$  be all the atoms  $\leq y$ . Then  $\sum_{x \leq y} q_k(x)$  counts all k-tuples from  $\{a_1, \ldots, a_m\}$ . Therefore,

, 
$$x \le y$$
  $\sum_{x \le y} q_k(x) = {m \choose k}$ 

and so

$$\sum_{x \le y} q(x) = \sum_{k=0}^{m} (-1)^k \binom{m}{k} = \begin{cases} 1 & \text{if } m = 0, & \text{i.e. } y = 0, \\ 0 & \text{if } m > 0, & \text{i.e. } y > 0. \end{cases}$$

By the Moebius inversion formula 2.26, we have

$$q(x) = \sum_{y \le x} \mu(y, x) \left( \sum_{z \le y} q(z) \right) = \mu(0, x).$$

**36.** There are three types of elements in L:

- (a) elements of C,
- (b) elements such that there is a  $y \in C$ , with x < y,
- (c) elements such that there is a  $y \in C$ , with x > y.
- (a), (b) and (c) partition the elements; in fact, there cannot be any x with two of these properties since this would give two comparable elements of C. Also, if an x did not have any of the properties (a), (b), (c), then no maximal chain containing x would meet C.

Let us write x < C and x > C if (b) and (c) holds, respectively. Consider the number  $q_k(a, b)$  of k-tuples in C whose union is b and meet is a, and let a < C < b. Then we prove that

$$q(a,b) = \sum_{k=2}^{n} (-1)^k q_k(a,b) = \mu(a,b)$$

by induction on the number of elements between a and b. Let  $a_1, \ldots, a_m$  be those elements of C between a and b. We may assume a=0 and b=1, because  $\{a_1,\ldots a_m\}$  forms a set in the sublattice  $\{z:a\leq z\leq b\}$  with the same properties as C. We have

$$\sum_{x < C < y} q_k(x, y) = \binom{m}{k}$$

and so

(1) 
$$\sum_{x < C < y} q(x, y) = {m \choose 2} - {m \choose 3} + \ldots = m - 1.$$

By 2.29,  $\mu$  satisfies the same identity:

(2) 
$$\sum_{x < C < y} \mu(x, y) = \sum_{x, y \in C} \mu(x, y) - 1 = \sum_{x \in C} \mu(x, x) - 1 = m - 1.$$

Now in (1) and (2)  $q(x, y) = \mu(x, y)$  if  $x \neq 0$  or  $y \neq 1$  by the induction hypothesis. So we must have

$$q(0, 1) = \mu(0, 1).$$

37. Let a be any atom, then

$$\sum_{\mathbf{a}\mathbf{v}\mathbf{x}=1}\mu(0,\mathbf{x})=0$$

by 2.27. Since a covers 0, either x = 1 or  $a \lor x = 1$  covers x, so x is a coatom.

If we use induction on r we may assume  $\mu(0, x)$  has sign  $(-1)^{r-1}$  (the rank of the interval  $0 \le z \le x$  is r-1, since any maximal (0, x)-chain extended by 1, forms a maximal (0, 1)-chain). So

$$\mu(0, 1) = -\sum_{\substack{x \neq 1 \\ x \neq a = 1}} \mu(0, x)$$

implies the assertion.

## § 3. Permutations

1. If  $\pi$ ,  $\varrho$  are permutations of  $\{1,\ldots,n\}$ , then

$$\pi' = \varrho^{-1}\pi\varrho$$

is the permutation arising as follows: we draw the "graph" of  $\pi$ , i.e. we connect i to  $\pi(i)$  for  $i=1,\ldots,n$ ; then consider the image of this graph by  $\varrho$  as the graph of a permutation  $\pi'$ . Hence it follows that two permutations are conjugate iff their graphs are isomorphic. These graphs consist of disjoint cycles (corresponding to the cycle decomposition of the permutation). Therefore,

the graphs of two permutations are isomorphic if and only if the cardinalities

of their cycles are the same.

Hence, a conjugacy class can be described by the cardinalities of cycles. These cardinalities form a partition of the number n and conversely, each partition describes a conjugacy class. Hence, the number of conjugacy classes of  $S_n$  is  $\pi_n$ , the number of partitions of the number n (see problem 1.20).

2. (a) By inclusion-exclusion, the number in question is

$$\sum_{X\subseteq\{1,\ldots,n\}}(-1)^{|X|}S_X,$$

where  $S_X$  denotes the number of those permutations which fix all points in X. This number is, obviously (n - |X|)! So we have

$$\sum_{X\subseteq\{1,\ldots,n\}} (-1)^{|X|} (n-|X|) ! = \sum_{k=0}^n (-1)^k \binom{n}{k} (n-k) ! = n ! \sum_{k=0}^n \frac{(-1)^k}{k !} \approx \frac{n !}{e},$$

where the difference at the pprox sign is less than  $rac{1}{n+1}$  .

- (b) If we want to construct a permutation  $\pi$  consisting of a single cycle, we can map 1 onto any of  $2, \ldots n$ . This gives n-1 possibilities. We can map  $\pi(1)$  onto any of n-2 elements etc, if  $\pi(1), \pi^2(1), \ldots, \pi^k(1)$  are specified, we can choose any of the remaining n-k-1 elements for  $\pi(\pi^k(1))$ . Thus the number of such permutations is  $(n-1)(n-2)\ldots 1=(n-1)!$ .
- 3. Let us count the permutations in which 1 is contained in a cycle of length k. There are  $\binom{n-1}{k-1}$  possible ways to choose the elements of this cycle; there are (k-1)! ways to order them in a cycle and (n-k)! ways to permute the rest. Thus we get

$$\binom{n-1}{k-1}(k-1)!(n-k)! = (n-1)!$$

and the probability is

$$\frac{(n-1)!}{n!} = \frac{1}{n},$$

independently of k, a remarkable fact.

Second solution. Given a permutation  $\pi$ , let us write it as a product of cycles (we also write out one-element cycles i.e. fixed points) such that each cycle should end with its least element and these last elements of cycles should be in increasing order. Thus the first cycle ends with 1. The resulting string of numbers can be viewed as another permutation  $\hat{\pi}$ . It is easy to see that  $\pi$  can be recovered from  $\hat{\pi}$  uniquely: the first cycle of  $\pi$  ends with 1; the second cycle ends with the least number not occurring in the first cycle etc.

The length of the cycle of  $\pi$  containing 1 is determined by the position of 1 in  $\hat{\pi}$ . This clearly can be any one of the n positions with the same probability 1/n.

4. We count those permutations in which 1 and 2 belong to distinct cycles. If the cycle containing 1 has length k we have  $\binom{n-2}{k-1}$  possibilities to select its elements and (k-1)! possibilities to order them into a cycle; we have (n-k)! possibilities to order the rest. Hence the number of such permutations is

$$\sum_{k=1}^{n-1} \binom{n-2}{k-1} (k-1)! (n-k)! = (n-2)! \sum_{k=1}^{n-1} (n-k) = (n-2)! \frac{n(n-1)}{2} = \frac{n!}{2},$$

whence the probability in question is 1/2.

Second solution. Consider the coding of permutations introduced in the second solution of 3.3. 1 and 2 belong to the same cycle of  $\pi$  iff 2 occurs in  $\hat{\pi}$  before 1. Obviously this happens with probability 1/2.

5. Let  $1 \leq k \leq n$ , and

$$\zeta_i = \begin{cases} 1 & \text{if } i \text{ is contained in a } k\text{-cycle,} \\ 0 & \text{otherwise.} \end{cases}$$

Then  $\zeta_1 + \ldots + \zeta_n$  is the number of points which are contained in k-cycles and  $\frac{1}{k}(\zeta_1 + \ldots + \zeta_n)$  is the number of k-cycles. Now

$$E\left(\frac{1}{k}\left(\zeta_1+\ldots+\zeta_n\right)\right)=\frac{1}{k}\sum_{i=1}^n E(\zeta_i).$$

Here, by 3.3,

$$E(\zeta_i) = \frac{1}{n}$$

and thus, the expected number of k-cycles is

$$\frac{1}{k}\sum_{i=1}^n E(\zeta_i) = \frac{1}{k}.$$

The expected number of cycles is, therefore,

$$rac{1}{1}+rac{1}{2}+\ldots+rac{1}{n}\!\sim\!\log n.$$

6. Let  $\pi$  denote the permutation defined by

 $\pi(j)$  = the index of the box containing key j.

Suppose we have broken open boxes  $1, \ldots, k$ . Clearly we can open the rest iff each cycle of  $\pi$  contains one of  $1, \ldots, k$ .

Consider the permutation  $\hat{\pi}$  introduced in the second solution of 3.3. Since the cycles of  $\pi$  are ordered in the definition of  $\hat{\pi}$  such that their least elements are in increasing order, it suffices to require that the least element of the last cycle should be at most k; or, by the definition of  $\hat{\pi}$ , that  $\hat{\pi}(n) \leq k$ . The probability that this occurs is clearly k/n.

[J. Bognár – J. Mogyoródi – A. Prékopa – A. Rényi – D. Szász, Problem Book

on Probability (in Hungarian), Tankönyvkiadó, Budapest 1970, p. 56.]

## 7. We have

$$n! p_n(x_1, \ldots, x_n) = \sum_{\pi \in S_n} x_1^{k_1(\pi)} \ldots x_n^{k_n(\pi)},$$

where  $k_i(\pi)$  denotes the number of i-cycles of  $\pi$ . Let us consider a point v of the underlying set and let  $C_{\pi}$  be the cycle of  $\pi$  which contains v; let it have length  $l_{\pi}$ . Also, let  $\varrho_{\pi}$  be the permutation of the elements not in  $C_{\pi}$ , induced by  $\pi$ . Thus

$$n! p_n(x_1, \ldots, x_n) = \sum_{\pi \in S_n} x_{l_{\pi}} x^{k_1(\varrho_{\pi})} \ldots x_{n-l_{\pi}}^{k_{n-l_{\pi}}(\varrho_{n})}.$$

The cycle  $C_{\pi}$  with length  $l_{\pi}$  can be chosen  $(n-1)(n-2)\dots(n-l_{\pi}+1)$  ways (see 3.3); with each of these cycles, we can pair any permutation  $\varrho_{\pi}$ of the remaining  $n - l_{\pi}$  points. Thus

$$n! p_n(x_1, \ldots, x_n) = \sum_{l=1}^n (n-1) \ldots (n-l+1) x_l(n-l)! p_{n-l}(x_1, \ldots, x_{n-l}) =$$

$$= \sum_{l=1}^n (n-1)! x_l p_{n-l}(x_1, \ldots, x_{n-l})$$

or, equivalently,

$$n \cdot p_n(x_1, \dots, x_n) = \sum_{l=1}^n x_l \, p_{n-l}(x_1, \dots, x_{n-l}).$$
 Set

$$f(y) = \sum_{n=0}^{\infty} p_n(x_1, \dots, x_n) y^n,$$

then

$$f'(y) = \sum_{n=0}^{\infty} n \cdot p_n(x_1, \dots, x_n) y^{n-1} = \sum_{n=0}^{\infty} \sum_{l=1}^{n} x_l \, p_{n-l}(x_1, \dots, x_{n-l}) y^{n-1} =$$

$$= \sum_{l=1}^{\infty} x_l \, y^{l-1} \sum_{n=l}^{\infty} p_{n-l}(x_1, \dots, x_{n-l}) y^{n-l} = \left( \sum_{l=1}^{\infty} x_l \, y^{l-1} \right) f(y).$$

$$(\ln f(y))' = x_1 + x_2 y + \ldots + x_k y^{k-1} + \ldots,$$
  
 $\ln f(y) = C + x_1 y + x_2 \frac{y^2}{2} + \ldots + x_k \frac{y^k}{k} + \ldots,$ 

3.8 SOLUTIONS

and substitution of y = 0 yields C = 0. Thus

$$f(y) = \exp\left(x_1y + x_2\frac{y^2}{2} + \ldots + x_k\frac{y^k}{k} + \ldots\right)$$

as stated. [G. Pólya, Acta Math. 68 (1937) 145-254; this reference is relevant for all problems on cycle index and the "Pólya-Redfield method"; also see N. G. de Bruijn in: Applied Combinatorial Mathematics, (E. E. Beckenbach, ed.) Wiley, 1964.]

8. (a) The rotation by m decomposes into (n, m) cycles of length n/(n, m). Hence the cycle index is

$$\sum_{m=1}^{n} x_{n/(n,m)}^{(n,m)} = \sum_{d|n} x_d^{n/d} \sum_{\substack{m \le n \\ (m,n) = \frac{n}{d}}} 1 = \sum_{d|n} \varphi(d) \ x_d^{n/d}.$$

More generally, let G be any group and consider its regular representation as a permutation group, i.e. define, for  $a \in G$  and  $x \in G$ ,

$$a(x) = xa$$

Then a is a permutation of the set G which decomposes into |G|/k cycles of length k, where k is the order of a; because

$$xa^m = x \Leftrightarrow a^m = 1.$$

Hence the cycle index is

$$\sum_{d \mid |G|} x_d^{|G|/d} \varphi(G, d),$$

where  $\varphi(G, d)$  is the number of elements of order d in the group G.

(b) Let  $P = \{C_1, C_2, \ldots, C_k\}$ ,  $C_i = \{c_{i1}, \ldots, c_{in}\}$ . A permutation which keeps P invariant can be described as follows. We take a permutation  $\pi$  of  $\{1,\ldots,k\}$  and k permutations  $\varrho_1,\ldots,\varrho_k$  of  $\{1,\ldots,n\}$ . We denote by  $\langle \pi; \varrho_1, \ldots, \varrho_k \rangle$  the permutation defined by

$$\langle \pi; \varrho_1, \ldots, \varrho_k \rangle (c_{ij}) = c_{\pi(i), \varrho_i(j)}$$
  
 $(1 \leq i \leq k, \quad 1 \leq j \leq n).$ 

Let us determine the cycles of  $\langle \pi; \varrho_1, \ldots, \varrho_k \rangle$ . If, say,  $(1, \ldots, l)$  is a cycle of  $\pi$ , then the iterated images of  $c_{1j}$  by  $\langle \pi; \varrho_1, \ldots, \varrho_k \rangle$  are in  $C_1, \ldots, C_l, C_1, \ldots$ . Thus, the first image which is equal to  $c_{1j}$  is the  $(lm)^{\text{th}}$  image for some m. To determine this m we observe that the  $l^{\text{th}}$  image of  $c_{1j}$  is  $c_{1j'}$  with  $j' = \varrho_l (\ldots \varrho_1(j) \ldots)$ . Hence, m is the length of the cycle of  $\varrho_1 \ldots \varrho_l$  contains ing j. Thus,  $C_1 \cup \ldots \cup C_l$  contains  $k_i(\varrho_1 \ldots \varrho_l)$  cycles with length li. Similar assertions hold for the other cycles of  $\pi$ . The term corresponding to this permutation is

$$\prod x_l^{k_1(\varrho_{i_1}\dots\varrho_{i_l})} x_{2l}^{k_2(\varrho_{i_1}\dots\varrho_{i_l})} \dots x_{nl}^{k_n(\varrho_{i_1}\dots\varrho_{i_l})},$$

where the product extends over all cycles  $(i_1, \ldots, i_l)$  of  $\pi$ . Summing this over all choices of  $\varrho_1, \ldots, \varrho_l$  we get (since for different cycles  $\varrho_{i_1}, \ldots, \varrho_{i_l}$  can be chosen independently),

$$\Pi\left(\sum_{\varrho} x_l^{k_1(\varrho_{i_1}\dots\varrho_{i_l})}\dots x_{nl}^{k_n(\varrho_{i_1}\dots\varrho_{i_l})}\right),$$

where again, the product extends over all cycles  $(i_1 \dots i_l)$  of  $\pi$ . Since  $\varrho_{i_1} \dots \varrho_{i_l}$  represents each permutation  $(n!)^{l-1}$  times,

$$\sum_{\varrho_{i_1},\ldots,\varrho_{i_l}} x_l^{k_1(\varrho_{i_1}\ldots\varrho_{i_l})} \ldots x_{nl}^{k_n(\varrho_{i_1}\ldots\varrho_{i_l})} = (n!)^{l-1} \sum_{\pi \in S_n} x_l^{k_1(\pi)} \ldots x_{nl}^{k_n(\pi)} = (n!)^l \ p_n(x_l,\ldots,x_{nl}),$$

where  $p_n(x_1, \ldots, x_n)$  is the cycle index of  $S_n$ . Thus

$$\prod_{(l_1,\ldots,l_l)} \left( \sum_{e_{l_1}\ldots,e_{l_l}} x_l^{k_1(e_{l_1}\ldots e_{l_l})} \ldots x_n^{k_n(e_{l_1}\ldots e_{l_l})} \right) = \prod_{(l_1,\ldots,l_l)} (n !)^l p_n(x_l,\ldots,x_{nl}) = \\
= \prod_{l=1}^k (n !)^{l k_l(\pi)} p_n(x_l,\ldots,x_{nl})^{k_l(\pi)} = (n !)^k \prod_{l=1}^k p_n(x_l,\ldots,x_{nl})^{k_l(\pi)}.$$

If we sum over  $\pi$ , and divide by  $k!(n!)^k$ , the number of permutations in the group, we get

$$p_k(p_n(x_1,\ldots,x_n), p_n(x_2,\ldots,x_{2n}),\ldots,p_n(x_k,\ldots,x_{kn})).$$

In exactly the same way we prove, that if  $\Gamma$ ,  $\Gamma_1$  are permutation groups with cycle indices  $F(x_1, \ldots, x_k)$ ,  $G(x_1, \ldots, x_n)$  respectively, then the cycle index of their wreath product (the group of permutations  $\langle \pi; \varrho_1, \ldots, \varrho_k \rangle$ , where  $\pi \in \Gamma$ ,  $\varrho_i \in \Gamma_1$ ) is

$$F(G(x_1,\ldots,x_n),G(x_2,\ldots,x_{2n}),\ldots,G(x_k,\ldots,x_{kn})).$$

[G. Pólya, ibid.]

**9.** By (a) of the previous problem, if we can find two groups which have the same number of elements of order k for  $k = 1, 2, \ldots$ , then the regular permutation group representations of these groups yield a counterexample.

We will find two non-isomorphic groups of order  $p^3$  (p > 2, prime) in which each element  $\neq 1$  has order p. One of these is, plausibly,  $G_1 = Z_p \times Z_p \times Z_p$ . Let  $G_2$  consist of all matrices of form

$$\begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix},$$

where  $a, b, c \in GF(p)$ . Since

(1) 
$$\begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & a+x & b+y+az \\ 0 & 1 & c+z \\ 0 & 0 & 1 \end{pmatrix},$$

these matrices form a group with  $p^3$  elements. Moreover,

$$\begin{bmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & ka & kb + \binom{k}{2} & ac \\ 0 & 1 & kc \\ 0 & 0 & 1 \end{bmatrix},$$

which follows easily by induction on k. Thus

$$\begin{bmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix}^p = \begin{bmatrix} 1 & pa & p\left(b + \frac{p-1}{2}ac\right) \\ 0 & 1 & pc \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

i.e. each element other than 1 has order p in this group  $G_2$ . Finally,  $G_1 \neq G_2$  as  $G_2$  is non-commutative: the matrices in (1) commute iff az = cx. [G. Pólya, ibid.]

## 10. We have

$$p_n(x_1, -x_2, \dots, (-1)^n x_n) = \frac{1}{n!} \sum_{\pi \in S_n} (-1)^{k_2(\pi) + k_4(\pi) + \dots} x_1^{k_1(\pi)} \dots x_n^{k_n(\pi)} =$$

$$= \frac{1}{n!} \left( \sum_{\pi \in A_n} x_1^{k_1(\pi)} \dots x_n^{k_n(\pi)} - \sum_{\pi \in S_n - A_n} x_1^{k_1(\pi)} \dots x_n^{k_n(\pi)} \right).$$

Hence

$$p_n(x_1, x_2, \dots x_n) + p_n(x_1, -x_2, \dots, (-1)^n x_n) =$$

$$= \frac{2}{n!} \sum_{\pi \in A_n} x_1^{k_1(\pi)} \dots x_n^{k_n(\pi)} = q_n(x_1, \dots, x_n).$$

Thus

$$\sum_{n=0}^{\infty} q_n(x_1,\ldots,x_n) y^n = e^{x_1 y + x_2 \frac{y^2}{2} + \ldots + x_k \cdot \frac{y^k}{k} + \ldots} + e^{x_1 y - x_2 \cdot \frac{y^2}{2} + \ldots + (-1)^k x_k \cdot \frac{y^k}{k} + \ldots}.$$

11. Let  $p_{-1} = 0$ , then

$$\sum_{n=0}^{\infty} (p_n - p_{n-1}) y^n = (1 - y) \sum_{n=0}^{\infty} p_n y^n = (1 - y) e^{\sum_{k=0}^{\infty} \frac{x_k}{k} y^k} =$$

$$= e^{\sum_{k=0}^{\infty} \frac{x_k}{k} y^k + \ln(1-y)} = e^{\sum_{k=0}^{\infty} \frac{x_k-1}{k} y^k}.$$

By our assumption on the  $x_i$ 's, there is a polynomial in the exponent and hence, this function is an entire function. Therefore, its Taylor series converges at y = 1:

$$\lim_{N \to \infty} \sum_{n=0}^{N} (p_n - p_{n-1}) = \lim_{N \to \infty} p_N = e^{\sum_{k=0}^{\infty} \frac{x_k - 1}{k}}.$$

12. (a) Obviously,

$$n! p_n(x, \ldots, x) = \sum_{\pi \in S_n} x^{k_1(\pi) + \ldots + k_n(\pi)} = \sum_{k=0}^n s_{n,k} x^k.$$

Hence

$$\sum_{n=0}^{\infty} \frac{f_n(x)}{n!} y^n = e^{xy + x \frac{y^2}{2} + \dots + x \frac{y^k}{k} + \dots} = e^{x(-\ln(1-y))} = (1-y)^{-x} =$$

$$= \sum_{n=0}^{\infty} {\binom{-x}{n}} (-1)^k y^n = \sum_{n=0}^{\infty} {\binom{x+n-1}{n}} y^n.$$

Thus

$$f_n(x) = n! {x+n-1 \choose n} = x(x+1) \dots (x+n-1)$$

and

$$s_{n,k} = (-1)^{n-k} s(n, k),$$

where s(n, k) is the Stirling number of first kind.

Second solution. Let x be a positive integer. If a permutation  $\pi$  of a set S has exactly k cycles, then  $x^k$  is the number of x-colorations of S invariant under  $\pi$ . So  $f_n(x)$  sums these numbers for all permutations  $\pi$  of  $S = \{1, \ldots, n\}$ . A given x-coloration f of S is counted  $k_1! \ldots k_x!$  times, where  $k_i$  is the number of elements with color i. The number of occurrences of a given sequence  $k_1, \ldots, k_x$  is  $n!/k_1! \ldots k_s!$  (see 1.3) and so by 1.4b,

$$f_n(x) = \sum_{\substack{k_1,\ldots,k_x \ge 0 \\ k_1+\ldots+k_x=n}} \frac{n!}{k_1!\ldots k_x!} k_1!\ldots k_x! = n! {x+n-1 \choose n}.$$

If this equality between polynomials holds for positive integers x, then it holds for every value.

(b) Since

$$f_n'(1) = \sum_{k=0}^n k s_{n,k}$$

is the sum of numbers of cycles in permutations, the expected number of cycles in permutations is  $\frac{1}{n!}f'_n(1)$ . Evaluating this,

$$\frac{1}{n!}f'_n(x) = \left(\frac{x(x+1)\dots(x+n-1)}{n!}\right)' =$$

$$= \frac{x(x+1)\dots(x+n-1)}{n!} \left(\frac{1}{x} + \dots + \frac{1}{x+n-1}\right)$$

and thus

$$\frac{1}{n!}f'_n(1)=\frac{1}{1}+\ldots+\frac{1}{n}\sim\log n.$$

13. Orient each circuit of such a graph. Let  $\pi$  denote the permutation we get this way. Then  $k_1(\pi) = k_2(\pi) = 0$ . Moreover, we get from the same graph  $2^{k_3(\pi)+\cdots+k_n(\pi)}$  permutations. Thus, if we form the sum

3.13

$$\sum \frac{1}{2^{k_3(\pi)+\ldots+k_n(\pi)}}$$

over the permutations associated with a given graph we get 1; and if we sum over all permutations such that  $k_1(\pi) = k_2(\pi) = 0$  we get the number of 2-regular simple graphs. Thus

$$g_n = \sum_{\substack{n \ k_3(\pi) = k_3(\pi) = 0}} \frac{1}{2^{k_3(\pi) + \dots + k_n(\pi)}} = n! \, p_n \left[ 0, 0, \frac{1}{2}, \dots, \frac{1}{2} \right].$$

Hence

$$\sum_{n=0}^{\infty} \frac{g_n}{n!} y^n = \sum_{n=0}^{\infty} p_n \left( 0, 0, \frac{1}{2}, \dots, \frac{1}{2} \right) y^n = e^{\frac{1}{2} \left( \frac{y^3}{3} + \frac{y^4}{4} + \dots \right)} =$$

$$= e^{\frac{1}{2} \left( -\ln\left( (1-y) \right) - \frac{1}{2} y - \frac{1}{4} y^2} = \frac{1}{\sqrt{1-y} e^{\frac{1}{2} y + \frac{1}{4} y^2}}.$$

14. (a) If we join i and j  $(1 \le i, j \le n)$  by an edge whenever (i, j) is a pair in our partition we get a 2-regular graph; this time double edges and/or loops may occur. Similarly as before, we get that our number is

$$n! p_n \left(1, 1, \frac{1}{2}, \dots, \frac{1}{2}\right)$$

and hence, the exponential generating function is

$$\sum_{n=0}^{\infty} p_n \left( 1, 1, \frac{1}{2}, \dots, \frac{1}{2} \right) y^n = e^{-\frac{1}{2} \ln (1-y) + \frac{y}{2} + \frac{y^2}{4}} = \frac{e^{\frac{y}{2} + \frac{y^2}{4}}}{\sqrt{1-y}}.$$

(b) Now join i to j by a directed edge whenever (i, j) belongs to the partition. By changing the directions in such a way as to get cycles, we associate a permutation with the given partition. As before, we get

$$2^{k_3(\pi)}+\ldots+k_n(\pi)$$

permutations from a given partition; but we obtain

$$3^{k_2(\pi)} 2^{3k_3(\pi)} + 4k_4(\pi) + \ldots + nk_n(\pi)$$

digraphs (i.e. partitions) from a given permutation. Thus the answer is

$$\sum_{\pi \in S_n} \frac{3^{k_2(\pi)} \, 2^{3k_3(\pi) + \ldots + nk_n(\pi)}}{2^{k_3(\pi) + \ldots + k_n(\pi)}} = n \, ! \, p_n(1, 3, 2^2, 2^3, \ldots)$$

and the exponential generating function is

$$p_n(1, 3, 2^2, 2^3, \ldots) y^n = e^{y + \frac{3}{2} \cdot y^2 + \frac{2^2 y^2}{3} + \ldots + \frac{2^{k-1} y^k}{k}} = e^{\frac{y^2}{2} + \frac{1}{2} (-\log(1-2y))} = = \frac{e^{\frac{y^2}{2}}}{\sqrt{1-2y}}$$
 [G. Baróti].

15. (a) Obviously, the number of permutations having no fixed point is

$$n! p_n(0, 1, \ldots 1).$$

The exponential generating function of these numbers is

$$\sum_{n=0}^{\infty} p_n(0, 1, \dots 1) y^n = e^{\frac{y^2}{2} + \frac{y^3}{3} + \dots + \frac{y^k}{k} + \dots} = e^{-y - \ln(1 - y)} =$$

$$= \frac{e^{-y}}{1 - y} = (1 + y + y^2 + \dots) \left( 1 - \frac{y}{1!} + \frac{y^2}{2!} - \dots \right),$$

whence

$$p_n(0, 1, \ldots, 1) = \sum_{k=0}^n \frac{(-1)^k}{k!} \sim \frac{1}{e}.$$

(b) The number of permutations consisting of exactly one cycle is, by 3.10, the coefficient of x in the polynomial

$$x(x+1)\ldots(x+n-1),$$

i.e. it is equal to (n-1)!.

16. Let  $k_1, \ldots, k_n$  be integers,  $1 \le k_i \le i$ . We claim that there exists a unique permutation  $\pi$  with  $\overline{\pi}(k) = k_i$ . Since  $\overline{\pi}(n)$  denotes the number of integers

 $1 \le j \le n$  with  $\pi(j) \ge \pi(n)$ , if follows that  $\pi(n) = n - k_n + 1$ .

Suppose we have determined what the values of  $\pi(n)$ ,  $\pi(n-1)$ ,...,  $\pi(n-k+1)$  are. Then  $\pi(n-k)$  must be the  $(k_{n-k})^{\text{th}}$  number in the sequence obtained from  $(n, n-1, \ldots, 1)$  by deleting  $\pi(n), \ldots, \pi(n-k+1)$ . It is immediately seen that in this way  $\pi(n), \ldots, \pi(1)$  are uniquely determined by  $k_n, \ldots, k_1$  and they form a permutation of  $\{1, \ldots, n\}$ . Moreover, this permutation  $\pi$  will satisfy  $\pi(i) = k_i$   $(i = 1, \ldots, n)$  by the construction.

Let us determine

$$n! P(\overline{\pi}(i_1) = k_{i_1}, \ldots, \overline{\pi}(i_r) = k_{i_r}),$$

where  $1 \le k_i \le i$  are given integers. The values  $\bar{\pi}(i)$ ,  $i \ne i_1, \ldots i_r$  can be chosen arbitrarily; hence

$$n$$
 !  $Pig(ar{\pi}(i_1)=k_{i_1},\ldots,ar{\pi}(i_r)=k_{i_r}ig)=\prod_{\substack{i
eq i_1,\ldots,i_r\ 1\leq i\leq n}}i$ 

and thus

$$P(\bar{\pi}(i_1) = k_{i_1}, \dots, \bar{\pi}(i_r) = k_{i_r}) = \frac{1}{i_1 \dots i_r}.$$

Thus

$$P(\bar{\pi}(i_1)=k_{i_1},\ldots,\bar{\pi}(i_r)=k_{i_r})=\prod_{\nu=1}^r P(\bar{\pi}(i_{\nu})=k_{i_{\nu}}),$$

which shows that  $\bar{\pi}(1), \ldots, \bar{\pi}(n)$  are independent. [C. Rényi and A. Rényi, in: Combinatorial Theory Appl. Coll. Math. Soc. J. Bolyai 4, Bolyai-North-Holland (1970) 945–971.]

17. (a) By the remark in the hint, we have to determine the expected number of integers i with  $\bar{\pi}(i) = i$ . Let

$$\zeta_i = \left\{ egin{array}{ll} 1 & ext{if} & \overline{\pi}(i) = i, \ 0 & ext{otherwise.} \end{array} 
ight.$$

Then we are interested in

$$E(\zeta_1 + \ldots + \zeta_n) = \sum_{i=1}^n E(\zeta_i) = \sum_{i=1}^n P(\overline{\pi}(i) = i) = \sum_{i=1}^n \frac{1}{i} \sim \log n.$$

(b) The probability that  $i_1, \ldots, i_k$  (and only these) are records is, by the preceding solution,

$$\prod_{v=1}^{k} \frac{1}{i_{v}} \cdot \prod_{i \neq i_{1}, \dots, i_{k}} \left( 1 - \frac{1}{i} \right) = \frac{1}{n!} \prod_{i \neq i_{1}, \dots, i_{k}} (i-1)$$

and thus, the probability that there are exactly k records is

$$\frac{1}{n!} \sum_{1 \leq j_1 \leq \dots \leq j_{n-k} \leq n} (j_1 - 1) \dots (j_{n-k} - 1) = \frac{|s(n, k)|}{n!},$$

where s(n, k) is the Stirling number of first kind.

Second solution: Observe that in  $\hat{\pi}$  (see the second solution of 3.3) the last elements of cycles of  $\pi$  are exactly the "negative records", i.e. those elements which are not followed by any smaller element. Thus the number of "negative records" in  $\hat{\pi}$  is equal to the number of cycles in  $\pi$ . Therefore the number of permutations with k "negative records" is the same as the number of those with k cycles, which is |s(n,k)| by 3.12a. The number of permutations with k records is clearly the same. [C. Rényi and A. Rényi, ibid.]

18. (a) If we consider the same random permutation of  $\{1, \ldots, n\}$  as in 3.16,

then the winner is the last i with  $\bar{\pi}(i) = i$ .

A strategy is a function which associates with each sequence  $(\overline{\pi}(1), \ldots, \overline{\pi}(k))$   $(1 \le k \le n)$  one of "yes" or "no". The value of a strategy is the probability that the first "yes" is after the winner. Since there are finitely many strategies, there is a best one. Furthermore, if we are late, i.e. come after the  $k^{\text{th}}$  jump, then we have to follow a strategy which says "no" for the first k jumps. Obviously, there are optimal strategies among these, too.

Let us denote by  $A_k$  the event that a (fixed) optimal strategy (with k "no"s first) wins, and put  $P(A_k) = p_k$ . Observe that  $A_k$  is independent of the order of the first k jumpers. This is heuristically clear and can be shown as

follows: Let

$$P(A_k | \overline{\pi}(1) = l_1, \dots, \overline{\pi}(k) = l_k) = q_1, \dots, l_k$$

and let  $q_{m_1,\ldots,m_k}$  be maximal among the numbers  $q_{l_1,\ldots,l_k}$   $(1 \leq l_i \leq i)$ . Then define a strategy which says "yes" for a sequence  $(\bar{\pi}(1),\ldots,\bar{\pi}(\nu))$  iff our original strategy says "yes" for  $(m_1,\ldots,m_k,\bar{\pi}(k+1),\ldots,\bar{\pi}(\nu))$ . The probability that this new strategy wins is  $q_{m_1,\ldots,m_k}$ , whence

$$q_{m_1...,m_k} \leq \frac{1}{k!} \sum_{l_1,...,l_k} q_{l_1...,l_k} = P(A_k) = p_k.$$

We must have equality here, whence  $q_{l_1...,l_k} = p_{k}$ . Now let k < n. Then

(1) 
$$P(A_k) = P(\bar{\pi}(k+1) = k+1)P(A_k | \bar{\pi}(k+1) = k+1) + P(\bar{\pi}(k+1) < k+1) \cdot P(A_k | \bar{\pi}(k+1) < k+1).$$

Here the second term corresponds to the case when the  $(k+1)^{st}$  jump is not a record. In this case we obviously will not bet on this jump; hence we say "no" and follow the optimal strategy after k+1 jumps, i.e.

$$P(A_k | \bar{\pi}(k+1) < k+1) = P(A_{k+1} | \bar{\pi}(k+1) < k+1) = p_{k+1}.$$

By 3.16,

$$P(\bar{\pi}(k+1) < k+1) = \frac{k}{k+1}.$$

If the  $(k+1)^{st}$  jump is a record we have two choices. Either we say "yes", then we win with probability

$$P(\overline{\pi}(k+2) < k+2,\ldots,\overline{\pi}(n) < n) = \frac{k+1}{k+2}\ldots\frac{n-1}{n} = \frac{k+1}{n}$$

or we say "no", in which case we follow the strategy after k+1 "no"s and have a chance of

$$P(A_{k+1} | \overline{\pi}(k+1) = k+1) = P(A_{k+1}) = p_{k+1}.$$

Thus

$$P_kig(A_k\,|\,\overline{\pi}(k+1)=k+1ig)=\max\left(rac{k+1}{n}\,,p_{k+1}
ight).$$

Since

$$P(\bar{\pi}(k+1) = k+1) = \frac{1}{k+1}$$
,

it follows from (1) that

$$(2) \quad p_k = \max\left(\frac{p_{k+1}}{k+1}\,,\,\frac{1}{n}\right) + \frac{k}{k+1}\,p_{k+1} = \max\left(p_{k+1},\frac{1}{n} + \frac{k}{k+1}\,p_{k+1}\right).$$

This makes it possible to determine  $p_{n-1}, p_{n-2}, \ldots$  recursively. If we arrive after the  $(n-1)^{st}$  jump we have to bet on the last jumper and win with probability

$$p_{n-1} = P(\overline{\pi}(n) = n) = \frac{1}{n}.$$

Now define  $\tilde{p}_k$  for  $k = n - 1, \ldots, 1$  recursively as follows:

$$ilde{p}_{n-1} = rac{1}{n} \,, \, ilde{p}_k = rac{1}{n} + rac{k}{k+1} \, ilde{p}_{k+1} .$$

Since

$$\frac{\tilde{p}_k}{k} = \frac{1}{nk} + \frac{\tilde{p}_{k+1}}{k+1},$$

it follows that

(3) 
$$\frac{\tilde{p}_k}{k} = \frac{1}{nk} + \frac{1}{n(k+1)} + \dots + \frac{1}{n(n-2)} + \frac{1}{n(n-1)},$$

$$\tilde{p}_k = \frac{k}{n} \left( \frac{1}{k} + \dots + \frac{1}{n-1} \right) \sim \frac{k}{n} \log \frac{n}{k}.$$

Now we have  $p_{n-1} = \tilde{p}_{n-1}$ ; let k be the largest index with  $p_k \neq \tilde{p}_k$ . Then

$$p_k \neq \frac{1}{n} + \frac{k}{k+1} p_{k+1} = \frac{1}{n} + \frac{k}{k+1} \tilde{p}_{k+1}$$

and so,

$$p_k = p_{k+1}.$$

(2) then implies that

$$p_k = p_{k-1} = \ldots = p_1.$$

This k is the largest value for which

$$\widetilde{p}_{k+1}>rac{1}{n}+rac{k}{k+1}\,\widetilde{p}_{k+1}=\widetilde{p}_k,$$

i.e. by (3),

$$\frac{1}{k+1} + \dots + \frac{1}{n-1} > 1.$$

This value of k satisfies

$$\log \frac{n}{k+1} \approx 1, \ k+1 \approx \frac{n}{e}.$$

So 
$$p_m = \frac{m}{n} \left( \frac{1}{m} + \dots + \frac{1}{n} \right) \approx \frac{m}{n} \log \frac{n}{m}$$
 for  $m > k$ , and  $p_1 = \dots = p_{k+1} = \frac{k+1}{n} \left( \frac{1}{k+1} + \dots + \frac{1}{n} \right) \approx \frac{1}{e}$ .

(b) The solution above already contains the optimum strategy. If k is such that  $\tilde{p}_k \leq \tilde{p}_{k+1}$ , then we get equality in (2), by the argument before it, if we say "no"; but if  $p_k > p_{k+1}$  and we have a record we have to say "yes". Thus we have to say "no" for the first k cases, where k is the largest number such that (4) holds. After this, we say "yes" to the first record. [J. Bognár et al., loc. cit. at 3.6.]

19. If there is no triple i < j < k with  $\pi(j) < \pi(i) < \pi(k)$ , then each i < j such that  $\pi(i) > \pi(j)$  also satisfies  $\pi(i) > \pi(j+1)$ ; thus,  $\overline{\pi}(j) \le \overline{\pi}(j+1)$ , i.e. the mapping  $\overline{\pi}$  is monotone. Conversely, if  $\overline{\pi}$  is monotone, there exists no i < j < k with  $\pi(j) < \pi(i) < \pi(k)$ . Suppose indirectly that there is such a triple. Consider  $\pi(j)$ ,  $\pi(j+1)$ , . . . ,  $\pi(k)$ . In this sequence there must be two consecutive terms  $\pi(l)$ ,  $\pi(l+1)$  such that  $\pi(l) < \pi(i) < \pi(l+1)$ . Then any v < l+1 with  $\pi(v) > \pi(l+1)$  also satisfies v < l and  $\pi(v) > \pi(l)$ ; moreover, i satisfies i < l and  $\pi(i) > \pi(l)$  but  $\pi(i) < \pi(l+1)$ . Thus  $\overline{\pi}(l) > \overline{\pi}(l+1)$ , a contradiction.

Thus, the number we seek is equal to the number of monotone mappings  $\varphi$  of  $\{1, \ldots, n\}$  into itself such that  $1 \leq \varphi(i) \leq i$ . By 1.33, this number is

$$rac{1}{n+1}inom{2n}{n}$$
 .

**20.** If  $\pi$  is any permutation, then the number of pairs i < j with  $\pi(i) > \pi(j)$  is, for a fixed j, equal to  $\overline{\pi}(j) - 1$ . Hence the total number of inversions is

$$(\overline{\pi}(1)-1)+(\overline{\pi}(2)-1)+\ldots+(\overline{\pi}(n)-1).$$

Thus, the number of permutation with exactly k inversions is the number of solutions of the equation

$$x_1 + \ldots + x_n = k$$

in integers satisfying  $0 \le x_i \le i - 1$  ( $x_i$  will be  $\overline{\pi}(i) - 1$ ). This number is, similarly as in 1.20, the coefficient of  $x^k$  in the product

$$1(1+x)\dots(1+x+x^2+\dots+x^{n-1}).$$

21. Suppose we have a very large tank with sufficient gasoline in it to go around the track. Let us start at an arbitrary point and go around. Stop at each station and buy all the gasoline in it. Then returning to the starting point we end up with the same amount of gasoline we started with, by the hypothesis.

Let A be the point, where the amount of gasoline in our tank was minimal. Then clearly A is at a gas station. Now if we start at A we can clearly go around

the whole track.

22. Define ascending and descending sequences as in the hint. Let  $\pi$  be an arbitrary permutation and  $C_1, \ldots, C_k$  its cycles. Assume  $\sum_{j \in C_i} x_j > 0$  for  $i = 1, \ldots, p$ , but  $\leq 0$  for  $i = p + 1, \ldots, k$ .

We claim each of the cycles  $C_1, \ldots, C_p$  has a unique decomposition into

We claim each of the cycles  $C_1, \ldots, C_p$  has a unique decomposition into arcs which are ascending sequences. Let, e.g.  $C_1 = (1, \ldots, r)$ . Let us imagine  $C_1$  as a speed track (cf. 3.21), in which each positive  $x_i$  corresponds to a gas station containing  $x_i$  liters of gasoline, while each negative  $x_i$  corresponds to a piece of the track which takes  $-x_i$  liters of gasoline to travel (there may be two gas stations immediately after each other or two consecutive pieces of the route with no gas station between them).

Going around the track we gain some gasoline, since  $x_1 + \ldots + x_r > 0$ . Let us keep on going around. Take those gas-stations after which the amount of gasoline in the tank is a record. There is at least one such station, since the amount of gasoline increases at each turn, and trivially if we return to

a record point the amount is a record again. The pieces of the track between two consecutive record points yield the desired decomposition of C into arcs which are ascending sequences. It is also easy to verify that there is no other such decomposition.

Similarly we can decompose each  $C_i$ ,  $p+1 \le i \le k$  into descending arcs. So each permutation  $\pi$  yields a decomposition of  $\{1, \ldots, n\}$  into disjoint

ascending and descending sequences.

Now suppose that we are given a partition of  $\{1,\ldots,n\}$  into u ascending and v descending sequences. Let, e.g.  $1,\ldots,m$  be those numbers occurring in ascending sequences. Compose cycles each element of which is an ascending sequence in this partition and other cycles whose elements are the descending x sequences. The number of ways to do so is clearly u!v!. In each case we obtain a permutation  $\pi$  such that  $b(\pi) = x_1 + \ldots + x_m$ . Now let us arrange the u ascending sequences of the given partition into a

Now let us arrange the u ascending sequences of the given partition into a linear order, followed by the v descending sequences. The number of ways to do so is u!v!. This results in a permutation  $\rho$  of  $\{1, \ldots, n\}$  such that

$$a(\varrho) = x_1 + \ldots + x_m$$
.

It is easy to see that, conversely, every permutation  $\varrho$  is obtained from a unique partition into ascending and descending sequences.

So every partition of  $\{1, ..., n\}$  into u ascending and v descending sequences corresponds to u!v! permutations  $\pi$  and the same number of permutations  $\varrho$ , and  $b(\pi) = a(\varrho)$  for each such  $\pi$  and  $\varrho$ . This proves the assertion.

It is interesting to remark that this result has important applications in the theory of orthogonal series. [F. Spitzer, *Trans. Amer. Math. Soc.* 82 (1956) 323-339.]

(b) Following the hint, take  $x_1 = \ldots = x_m = 1$ ,  $x_{m+1} = \ldots = x_{2m} = -1$  in (a). Let  $\pi$  be a permutation of  $\{1, \ldots, 2m\}$ . This clearly corresponds to the boys and girls forming "rings" to dance. The condition that the number of boys is equal to the number of girls in each "ring" is clearly equivalent to  $b(\pi) = 0$ . The number of  $\pi$ 's with this property is, by (a), the same as the number of permutations  $\varrho$  of  $\{1, \ldots, 2m\}$  such that  $a(\varrho) = 0$ . Each such permutation yields a sequence of 1's and (-1)'s such that all sums of the first r of them  $(r = 1, \ldots, 2m)$  are non-positive. Each such  $\pm 1$ -sequence corresponds to  $(m!)^2$  permutations  $\varrho$  with  $a(\varrho) = 0$ . The number of such  $\pm 1$ -sequences is, by 1.33b,

$$rac{1}{m+1}inom{2m}{m}$$
 .

So the number of permutations  $\varrho$  with  $a(\varrho) = 0$  is

$$\frac{1}{m+1} \binom{2m}{m} (m!)^2 = \frac{(2m)!}{m+1}.$$

Dividing this by the total number (2m)! of permutations we obtain the probability as desired.

**23.** (a) The set of all k-gons can be split into classes, where each class contains k-tuples arising by rotation from each other. We want to determine the number of these classes.

In a class, there are the *n* rotated copies of a *k*-gon *K*. These are not all different however; if c(K) denotes the number of rotations which map *K* onto itself, then the class contains  $\frac{n}{c(K)}$  elements. The sum

$$\sum_{K} c(K)$$

over all convex k-gons counts the class containing K  $c(K) \cdot \frac{n}{c(K)} = n$  times. Thus, we want to determine the number

$$\frac{1}{n} \sum_{K} c(K).$$

Let us consider how many times a given rotation is counted in  $\Sigma c(K)$ . If it is the rotation by m, then we get all convex k-gons invariant under this rotation as follows: we split the n-gon into  $\frac{n}{(m,n)}$  arcs of length (m,n) and specify  $\frac{k(m,n)}{n}$  points on one of these arcs. The other arcs must contain rotated copies of this tuple and therefore we uniquely determined a convex k-gon Thus, the number of convex k-gons invariant under the rotation by m is

$$\begin{pmatrix} \frac{(m, n)}{k(m, n)} \\ \frac{k(m, n)}{n} \end{pmatrix} \text{ if } n|k(m, n)$$
0 otherwise

and thus,

$$\frac{1}{n} \sum_{K} c(K) = \frac{1}{n} \sum_{\substack{m \leq n \\ n \mid k(m,n)}} \left( \frac{(m,n)}{k(m,n)} \right),$$

Introducing here  $d = \frac{n}{(m, n)}$ , we find that d|(n, k) and there are exactly  $\varphi(d)$  terms with the same d, and thus

$$\frac{1}{n} \sum_{K} c(K) = \frac{1}{n} \sum_{d \mid (n,k)} {n/d \choose k/d} \varphi(d).$$

(b) Proceeding as before, we consider the number  $c(\alpha)$  of rotations keeping a given k-coloration  $\alpha$  invariant. Then

$$\frac{1}{n}\sum_{\alpha}c(\alpha)$$

is the number of essentially different k-colorations. If we consider the rotation by m, there will be  $k^{(n,m)}$  k-colorations invariant under it and thus,

$$\frac{1}{n}\sum_{\alpha}c(\alpha)=\frac{1}{n}\sum_{m=1}^{n}k^{(n,m)}=\frac{1}{n}\sum_{d|n}\varphi(d)\,k^{\frac{n}{d}}.$$

**24.** Let  $V_1, \ldots, V_k$  be the orbits of  $\Gamma$ ,  $|V_i| = n_i$ . Let  $x \in V_i$  and denote by  $\Gamma_x$  the subgroup fixing x. Then the right cosets of  $\Gamma_x$  in  $\Gamma$  consist of those permutations in  $\Gamma$  which map x onto a given element of  $V_i$ . Hence  $\Gamma_x$  has index  $n_i$  and

3.24

$$|\Gamma_x| = \frac{|\Gamma|}{n_i}$$
.

Now let  $k_1(\pi)$  be the number of fixed points of the permutation  $\pi$ . Then the average number of fixed points is

$$\frac{1}{|\Gamma|} \sum_{\pi \in \Gamma} k_1(\pi) = \frac{1}{|\Gamma|} \left( \sum_{\pi \in \{1,\dots,n\}} |\Gamma_{\pi}| \right) = \frac{1}{|\Gamma|} \sum_{i=1}^k n_i \frac{|\Gamma|}{n_i} = k.$$

25. Consider

$$\sum_{\pi \in \Gamma} \sum_{\pi(x)=x} w(x) = \sum_{x \in \Omega} w(x) |\Gamma_x|,$$

where  $\Gamma_x$  is the stabilizer of x. Let  $V_x$  be the orbit containing x, then

$$|\Gamma_x| = \frac{|\Gamma|}{|V_x|}$$

and thus

$$\sum_{\pi \in \Gamma} \sum_{\pi(x)=x} w(x) = \sum_{x \in \Omega} w(x) \frac{|\Gamma|}{|V_x|} = \sum_{\theta} \sum_{x \in \theta} w(x) \frac{|\Gamma|}{|\theta|} = |\Gamma| \sum_{\theta} w(\theta).$$

26. If we define

$$\overline{\pi}(f)(x) = f(\pi(x))$$

for  $f: D \to R$  and  $\pi \in \Gamma$ , then  $\overline{\pi}$  can be considered as a permutation of the set  $\Omega$  of mappings of D into R. By Burnside's lemma 3.24, the number of orbits of  $\Gamma$  on  $\Omega$ , i.e. the number of essentially different mappings of D into R, is equal to the average number of fixed points of elements of  $\Gamma$ , acting on  $\Omega$ .

Let  $\pi \in \Gamma$ . Then  $f \in \Omega$  is a fixed point of  $\overline{\pi}$  if

$$f(\pi(x)) = f(x) \qquad (x \in D),$$

i.e. if f is constant on the orbits of  $\Gamma$  in D. Let  $k(\pi)$  be the number of orbits of  $\pi$  in D, then the number of such mappings f is, obviously,  $|R|^{k(\pi)}$  and hence, the average number of fixed points of elements of  $\Gamma$  in  $\Omega$  is

$$\frac{1}{|\Gamma|} \sum_{\pi \in \Gamma} |R|^{k(\pi)} = F(|R|, \dots, |R|).$$

27. Consider the permutation group  $\Gamma_2 = \Gamma \times \Gamma_1$ , acting on  $\Omega$ , defined by

$$((\pi, \varrho)f)(x) = \varrho(f(\pi^{-1}(x))) \quad (\pi \in \Gamma, \varrho \in \Gamma, f \in \Omega, x \in D).$$

Then we have to determine the number of orbits of  $\Gamma_2$ . By Burnside's lemma 3.24, this is equal to

(1) 
$$\frac{1}{|\Gamma| \cdot |\Gamma_1|} \cdot \sum_{\pi \in \Gamma} \sum_{\varrho \in \Gamma_1} k_1(\pi, \varrho),$$

where  $k_1(\pi, \varrho)$  is the number of fixed points of the permutation  $(\pi, \varrho) \in \Gamma_2$ , i.e. the number of mappings  $f \in \Omega$  such that

$$\varrho(f(\pi^{-1}(x))) = f(x)$$

holds for each  $x \in D$ . To determine this number, let  $k_i(\pi) = k_i$ ,  $k_i(\varrho) = l_i$  (i = 1, 2, ...). Let  $C_1, ..., C_k$  be the cycles of  $\pi$  and  $c_i \in C_i$ . Now note that the values  $f(c_1), ..., f(c_k)$  uniquely determine f; for by (2),

(3) 
$$f(\pi^t(c_i)) = \varrho^{-t}(f(c_i))$$

and every element of D can be written as  $\pi^t(c_i)$  with some t and i. Also, (3) must hold for  $t = |C_i|$ :

$$f(c_i) = \varrho^{|C_i|} f(c_i) ,$$

i.e. the length of the cycle of  $\rho$  containing  $f(c_i)$  must divide  $|C_i|$ .

Conversely, if  $f(c_i)$ ,  $i = 1, \ldots, k$  are given such that the length of the cycle of  $\varrho$  containing  $f(c_i)$  divides  $|C_i|$ , then (3) defines a fixed point of  $(\pi, \varrho)$ . To select the image of a point  $c_i$  with  $|C_i| = m$ , we therefore have

$$\sum_{d|m} dl_d$$

possibilities. This can be done independently, hence

$$k_1(\pi, \varrho) = \prod_{m=1}^{|D|} \left( \sum_{d|m} dl_d \right)^{k_m}.$$

Thus (1) can be written as

$$\frac{1}{|\Gamma_1|} \frac{1}{|\Gamma|} \sum_{\varrho \in \Gamma_1} \sum_{\pi \in \Gamma} \prod_{m=1}^{|D|} \left( \sum_{d \mid m} dk_d(\varrho) \right)^{k_m(\pi)} = \frac{1}{|\Gamma_1|} \sum_{\varrho \in \Gamma_1} F(u_1(\varrho), u_2(\varrho), \ldots),$$

where

$$u_m(\varrho) = \sum_{d \mid m} dk_d(\varrho).$$

To express this in terms of F and G, observe that

$$F(u_1(\varrho), u_2(\varrho), \ldots) = F\left[\frac{\partial}{\partial z_1}, \frac{\partial}{\partial z_2}, \ldots\right] e^{u_1(\varrho) z_1 + u_2(\varrho) z_2 \cdots}\Big|_{z_i=0}$$

and here

$$\sum_{m=1}^{\infty} u_m(\varrho) z_m = \sum_{m=1}^{\infty} \sum_{d \mid m} dk_d(\varrho) z_m = \sum_{d=1}^{\infty} dk_d(\varrho) \sum_{s=1}^{\infty} z_{ds}.$$

Thus

$$\begin{split} &\frac{1}{|\varGamma_{1}|} \sum_{\varrho \in \varGamma_{1}} F\left(\frac{\partial}{\partial z_{1}}, \frac{\partial}{\partial z_{2}}, \ldots\right) e^{u_{1}(\varrho)z_{1} + u_{2}(\varrho)z_{2} + \cdots} = \\ &= F\left(\frac{\partial}{\partial z_{1}}, \frac{\partial}{\partial z_{2}}, \ldots\right) \frac{1}{|\varGamma_{1}|} \sum_{\varrho \in \varGamma_{1}} \prod_{d=1}^{\infty} \left(e^{d(z_{d} + z_{2d} + \ldots)}\right)^{k_{d}(\varrho)} = \\ &= F\left(\frac{\partial}{\partial z_{1}}, \frac{\partial}{\partial z_{2}}, \ldots\right) G(e^{z_{1} + z_{2} + \cdots}, e^{2z_{2} + 2z_{4} + \cdots}, \ldots) \end{split}$$

and so, the number of essentially different mappings is

$$F\left(\frac{\partial}{\partial z_1}, \frac{\partial}{\partial z_2}, \ldots\right) G(e^{z_1+z_2+\cdots}, e^{2z_2+2z_4+\cdots}, \ldots) \bigg|_{z_i=0}$$

[N.G. de Bruijn, loc. cit. at 3.7.]

**28.** We use the same notation as before. The group  $\Gamma_2$  keeps invariant the set  $\Omega'$  of all one-to-one mappings of D into R. We want to determine the number of orbits of  $\Gamma_2$  in  $\Omega'$ . As before, this is equal to

$$\frac{1}{|\Gamma|\cdot|\Gamma_1|}\sum_{\pi\in\Gamma}\sum_{\varrho\in\Gamma_1}k_1'(\pi,\,\varrho),$$

where  $k'_1(\pi, \varrho)$  is the number of those one-to-one mappings of D into R which

are kept fixed by  $(\pi, \varrho)$ .

Let  $C_1, \ldots, C_k$  be the cycles of  $\pi$ . Then, again, it suffices to determine the image of a representative system  $\{c_1, \ldots, c_k\}$ ,  $c_i \in C_i$ . Moreover, since f is one-to-one,  $f(c_i)$  must belong to a cycle of  $\varrho$  of length  $|C_i|$ . Let  $C_1, \ldots, C_{k_i}$  be the cycles of  $\pi$  of length i (say), then we have  $i \cdot l_i$  choices for the image of  $c_1 \in C_1$ ;  $c_2 \in C_2$  cannot be mapped onto the same cycle of and thus, we have  $i(l_i-1)$  choices for the image of  $c_2$ . Going on similarly, we will have

$$(il_i)(i(l_i-1))\dots(i(l_i-k_i+1))=i^{k_i}l_i(l_i-1)\dots(l_i-k_i+1)$$

possible ways to map  $c_1, \ldots, c_{k_i}$ . Since the images of cycles of  $\pi$  of distinct length will be automatically disjoint, we can choose them independently and therefore

$$k_1'(\pi,\varrho) = \prod_{i=1}^{|D|} i^{k_i} l_i(l_i-1) \dots (l_i-k_i+1)$$

and hence, the number of essentially different one-to-one mappings of D into R is

$$\frac{1}{|\Gamma|}\frac{1}{|\Gamma_1|}\sum_{\pi\in\Gamma}\sum_{\varrho\in\Gamma_1}\prod_{i=1}^{|D|}i^{k_i(\pi)}k_i(\varrho)\big(k_i(\varrho)-1\big)\dots\big(k_i(\varrho)-k_i(\pi)+1\big).$$

Now observe that

$$egin{aligned} &rac{1}{|ec{\Gamma}|} \sum_{\pi \in ec{\Gamma}} \prod_{i=1}^{|D|} i^{k_i(\pi)} k_i(arrho) \left(k_i(arrho) - 1
ight) \ldots \left(k_i(arrho) - k_i(\pi) + 1
ight) = \ &= F\left(rac{\partial}{\partial z_1}\,, rac{\partial}{\partial z_2}\,, \ldots
ight) (1 + z_1)^{k_1(arrho)} \, (1 + 2 z_2)^{k_2(arrho)} \ldots \, igg|_{z_i = 0} \end{aligned}$$

and thus, the result is

$$egin{aligned} rac{1}{|arGamma_1|} \sum_{arrho \in arGamma_1} F\left(rac{\partial}{\partial z_1}\,,\,rac{\partial}{\partial z_2}\,,\,\ldots
ight) (1+z_1)^{k_1(arrho)}\,(1+2z_2)^{k_2(arrho)}\,\ldots\,igg|_{z_i=0} = \ &= F\left(rac{\partial}{\partial z_1}\,,\,rac{\partial}{\partial z_2}\,,\,\ldots
ight) G(1+z_1,1+2z_2,\,\ldots)\,igg|_{z_i=0}. \end{aligned}$$

**29.** Following the hint, consider a  $\gamma \in \Gamma$ ; let  $C_1, \ldots, C_k$  be its cycles. We prove that

(1) 
$$\sum_{n=1}^{\infty} q_n(\gamma) x^n = \prod_{i=1}^k r(x^{|C_i|}).$$

In fact, the coefficient of  $x^n$  on the right-hand side is

$$\sum_{\substack{v_1,\ldots,v_k>0\\\Sigma v_i\,|C_i|=n}} r_{v_1}\ldots r_{v_k},$$

and clearly this is the same as the number of mappings satisfying (\*) which are constant on each of the sets  $C_1, \ldots, C_k$ .

Now (1) can be rewritten as

$$\sum_{n=1}^{\infty} q_n(\gamma) x^n = \prod_{j=1}^{|D|} r(x^j)^{k_j(\gamma)}.$$

By Burnside's lemma 3.24,

$$a_n = \frac{1}{|\Gamma|} \sum_{\gamma \in \Gamma} q_n(\gamma)$$

and hence

$$\sum_{n=1}^{\infty} a_n x^n = \frac{1}{|\Gamma|} \sum_{\gamma \in \Gamma} q_n(\gamma) x^n = \frac{1}{|\Gamma|} \sum_{\gamma \in \Gamma} \prod_{j=1}^{|D|} r(x^j)^{k_j(\gamma)} = F(r(x), r(x^2), \ldots).$$

[G. Pólya, loc. cit. at 3.7.]

**30.** Let D be the set of n forints, R the set of k people who get them. A distribution of the forints is a mapping of D into R. Since all forints are alike, two distributions f, g are essentially the same if there is a permutation  $\pi$  of D such that

$$f(\pi(x)) = g(x) \quad (x \in D).$$

By 3.26, the number of such distributions is

$$p_n(k,\ldots,k),$$

where  $p_n(x_1, \ldots, x_n)$  is the cycle index of  $S_n$ . By the solution of 3.12,

$$p_n(k,\ldots,k) = \binom{n+k-1}{n}.$$

**31.** Let |D| = n,  $|R| = N \ge n$ . Two mappings  $f, g: D \to R$  induce the same partition of R iff

$$g(x) = \varrho(f(x))$$
  $(x \in D)$ 

for some permutation  $\varrho$  of R. Thus, the number of partitions of D is, by 3.27,

$$P_n = \left(\frac{\partial}{\partial z_1}\right)^n p_N(e^{z_1+z_2+\cdots}, e^{2z_2+2z_4+\cdots}, \ldots) \bigg|_{z_i=0} = \left(\frac{\partial}{\partial z}\right)^n p_N(e^z, 1, \ldots 1) \bigg|_{z=0},$$

i.e.  $\frac{P_n}{n!}$  is the coefficient of  $z^n$  in the Taylor series of  $p_N(e^z, 1, ... 1)$ , for any  $N \ge n$ . Since by 3.11,

SOLUTIONS

$$\lim_{N\to\infty} p_N\left(e^z, 1, \dots 1\right) = e^{e^z-1},$$

 $\frac{P_n}{n!}$  is the coefficient of  $z^n$  in the power series of  $e^{e^z-1}$  as well; i.e.

$$\sum_{n=0}^{\infty} \frac{P_n}{n!} z^n = e^{e^z - 1}.$$

## § 4. Two classical enumeration problems in graph theory

1. We use induction on n. For n=1, 2 the statement is trivial. Since  $\sum_{i=1}^{n} d_i =$ 

=2n-2<2n, there is a  $d_i=1$ . We may assume  $d_n=1$ . Remove  $v_n$ . In any tree under consideration,  $v_n$  is adjacent to some  $v_j, \ 1 \le j \le n-1$  and the removal of  $v_n$  results in another tree on  $\{v_1,\ldots,v_{n-1}\}$  with degrees  $d_1,\ldots,d_{j-1},d_j-1,d_{j+1},\ldots,d_{n-1}$ . Conversely, if we are given a tree on  $\{v_1,\ldots,v_{n-1}\}$  with degrees  $d_1,\ldots,d_{j-1},d_j-1,d_{j+1},\ldots,d_{n-1}$  then joining  $v_j$  to  $v_n$  we get a tree on  $\{v_1,\ldots,v_n\}$  with degrees  $d_1,\ldots,d_n$ . The number of trees on  $\{v_1,\ldots,v_{n-1}\}$  with degrees  $d_1,\ldots,d_{j-1},d_j-1,d_{j+1},\ldots,d_{n-1}$  is

$$\frac{(n-3)!}{(d_1-1)!\dots(d_{j-1}-1)!\frac{(d_j-2)!(d_{j+1}-1)!\dots(d_{n-1}-1)!}}=$$

$$=\frac{(d_j-1)(n-3)!}{(d_1-1)!\dots(d_n-1)!}.$$

This is also valid if  $d_j = 1$  since, then it is 0. Thus, the number of trees on  $\{v_1, \ldots, v_n\}$  with degrees  $d_1, \ldots, d_n$  is

$$\sum_{j=1}^{n-1} \frac{(d_j-1)(n-3)!}{(d_1-1)!\dots(d_n-1)!} = \left(\sum_{j=1}^{n-1} (d_j-1)\right) \frac{(n-3)!}{(d_1-1)!\dots(d_n-1)!} = \frac{(n-2)(n-3)!}{(d_1-1)!\dots(d_n-1)!},$$

which proves the assertion [see B].

**2.** The number of trees on n points is

$$\sum_{\substack{d_1, \dots, d_n \ge 1 \\ d_1 + \dots + d_n = 2n - 2}} \frac{(n-2)!}{(d_1 - 1)! \dots (d_n - 1)!} = \sum_{\substack{k_1, \dots, k_n \ge 0 \\ k_1 + \dots + k_n = n - 2}} \frac{(n-2)!}{k_1! \dots k_n!} = \underbrace{(1 + \dots + 1)^{n-2}}_{n} = \underbrace{(1 + \dots + 1)^{n-2}}_{n} = n^{n-2}$$
[A. Cayley; see B].

3.  $p_1(x_1,\ldots,x_{n-1},0)=\sum x_1^{d_T(v_1)-1}\ldots x_{n-1}^{d_T(v_{n-1})-1}$ , where the summation extends over all trees having an endpoint at  $v_n$ . If we set  $S=T-v_n$ , then S is a tree on  $\{v_1,\ldots,v_{n-1}\}$  and conversely, for any tree S on  $\{v_1,\ldots,v_{n-1}\}$  we can get n-1 trees with an endpoint at  $v_n$  by joining  $v_n$  to a point of S. For these trees T, we have

$$x_1^{d_T(v_1)-1} \dots x_{n-1}^{d_T(v_{n-1})-1} = x_i x_1^{d_S(v_1)-1} \dots x_{n-1}^{d_S(v_{n-1})-1},$$

where j is the index for which  $v_i$  is adjacent to  $v_n$ . Hence

$$\sum_{\substack{T \\ T-v_n=S}} x_1^{d_T(v_1)-1} \dots x_n^{d_T(v_n)-1} = (x_1 + \dots + x_{n-1}) \cdot x_1^{d_S(v_1)-1} \dots x_{n-1}^{d_S(v_{n-1})-1}$$

and, summing over all trees S on  $\{v_1, \ldots, v_{n-1}\}$ , we get

(1) 
$$p_n(x_1,\ldots,x_{n-1},0)=(x_1+\ldots+x_{n-1})\,p_{n-1}(x_1,\ldots,x_{n-1}).$$

Now we prove by induction on n that

(2) 
$$p_n(x_1,\ldots,x_n) = (x_1 + \ldots + x_n)^{n-2}.$$

By (1), we may assume

$$p_n(x_1,\ldots,x_{n-1},0)=(x_1+\ldots+x_{n-1})^{n-2}.$$

Set

$$p_n^*(x_1,\ldots,x_n)=(x_1+\ldots+x_n)^{n-2}.$$

Then

$$p_n(x_1,\ldots,x_{n-1},0)=p_n^*(x_1,\ldots,x_{n-1},0)$$

and similarly, if we substitute 0's for any non-empty set of variables,  $p_n$  and  $p_n^*$  become identical. Hence, using the notation of 2.5,

$$\sigma_k p_n(x_1,\ldots,x_n) = \sigma_k p_n^*(x_1,\ldots,x_n)$$

for k = 1, ..., n. Now by 2.5,

$$p_n - \sigma_1 p_n + \sigma_2 p_n + \ldots + (-1)^n \sigma_n p_n = 0,$$
  
$$p_n^* - \sigma_1 p_n^* + \sigma_2 p_n^* - \ldots + (-1)^n \sigma_n p_n^* = 0$$

because  $p_n$ ,  $p_n^*$  are of degree n-2 < n. This implies that  $p_n = p_n^*$ . The Cayley formula now follows on substituting  $x_1 = \ldots = x_n = 1$  [A. Rényi, in: Combinatorial Str. Appl. Gordon and Breach, 1970, 355–360].

**4.** If we contract each  $T_i$  onto a single point  $v_i$  (i = 1, ..., r), then any tree on V containing  $T_1, ..., T_r$  is mapped onto a tree on  $\{v_1, ..., v_r\}$ . Let us count how many distinct trees on V are mapped onto a fixed tree T' on  $\{v_1, ..., v_r\} = V$ .

If, for each edge  $(v_i, v_j) \in E(T')$ , we choose a  $(T_i, T_j)$ -edge we will get such a tree and, obviously, any tree on V which is mapped onto T' can be obtained

in this way. The number of selections of such trees is

$$\prod_{(v_i, v_j) \in E(T')} |V(T_i)| \cdot |V(T_j)| = \prod_{i=1}^r |V(T_i)|^{d_T(v_i)}$$

and hence, the number of all trees on V containing  $T_1, \ldots, T_r$  is

$$\sum_{T'} \prod_{i=1}^r |V(T_i)|^{d_{T'}(v_i)},$$

where the summation extends over all trees on V'. By the preceding formula, this is equal to

$$|V(T_1)| \dots |V(T_r)| (|V(T_1)| + \dots + |V(T_r)|)^{r-2} = |V(T_1)| \dots |V(T_r)| |V|^{r-2}$$
 [J.W. Moon; see B].

5. (a) Let  $b_1, \ldots, b_{n-1}$  be the indices of removed points. Let us see how to

determine  $b_i$ , if we know the Prüfer code.

 $b_i$  is obviously different from  $b_1, \ldots, b_{i-1}, a_i$ . Also,  $b_i \neq a_j$  for j > i; for  $b_i$  is removed and it cannot be the neighbor of an endpoint at a later step. Conversely, if  $k \notin \{b_1, \ldots, b_{i-1}, a_i, \ldots, a_{n-1}\}$ , then  $v_k$  is an endpoint of  $T - \{v_{b_1}, \ldots, v_{b_{i-1}}\}$ ; otherwise it would be a neighbor of a point removed at a later step. Thus,

(1) 
$$b_i = \min \{k : k \notin \{b_1, \ldots, b_{i-1}, a_i, \ldots, a_{n-1}\}\}.$$

Thus, the Prüfer code uniquely determines the numbers  $b_i$ . Since  $(v_{a_i}, v_{b_i})$  are the edges of T, the Prüfer code uniquely determines T.

(b) Let  $(a_1, \ldots, a_{n-1})$  be any sequence of integers with  $1 \le a_i \le n$ ,  $a_{n-1} = n$ . Define  $b_i$  recursively by (1) and join  $v_{a_i}$  to  $v_{b_i}$  for  $i = 1, \ldots, n-1$ . We claim that the resulting graph T is a tree with Prüfer code  $(a_1, \ldots, a_{n-1})$ . Both assertions will follow, if we show that  $v_{b_i}$  is an endpoint of the graph  $T_i = T - \{v_{b_1}, \ldots v_{b_{i-1}}\}$  and no point with smaller index is endpoint. We have

$$v_{a_i} \in V(T_i)$$

because  $a_i \neq b_1, \ldots, b_{i-1}$  by (1). Thus,  $v_{b_i}$  has a neighbor in  $T_i$ .  $v_{b_i}$  cannot be adjacent to any other point of  $T_i$ ; for suppose  $(v_{a_i}, v_{b_i})$  is another edge of  $T_i$  adjacent to  $b_i$ , then j > i as  $v_{b_i} \in V(T_i)$  and either  $b_i = b_j$  or  $b_i = a_j$ , which both contradict (1). Hence  $v_{b_i}$  is an endpoint of  $T_i$ . This proves that T and all  $T_i$ 's are trees.

Now suppose that  $T_i$  has an endpoint  $v_k$  with  $k < b_i$ . Since k did not come into consideration when defining  $b_i$  by (1), it follows that either  $k = b_v$ , v < i or  $k = a_j$ ,  $j \ge i$ . Now the first possibility does not occur because  $v_{b_v} \in V(T_i)$ , so  $k = a_j$ ,  $j \ge i$ . Since  $a_{n-1} = n \ge b_i > k$ , we have  $j \le n - 2$ . By the argument above,  $v_{b_j}$  is an endpoint of  $T_j$  and its neighbor is  $v_{a_j}$ . But  $v_{a_j} = v_k$  is an endpoint of  $T_i$ , and therefore it must be an endpoint of  $T_j$  too. Hence  $V(T_i) = \{v_{a_i}, v_{b_i}\}$  which is impossible as  $T_i$  has  $n - j + 1 \ge 3$  points.

The Cayley formula follows immediately: The number of sequences  $(a_1, \ldots, a_{n-1})$  with  $1 \le a_i \le n$ ,  $a_{n-1} = n$  is, obviously,  $n^{n-2}$  [A. Prüfer, Archiv

f. Math. u. Phys. 27 (1918) 142-144].

**6.** If T is a tree on  $V = \{v_1, \ldots, v_n\}$  and  $e \in E(T)$ , then T - e consists of two disjoint trees, which together cover V. This way we get (n-1)  $T_n$  such pairs. Let T', T'' be two disjoint trees which together cover V; let |V(T')| = k, |V(T'')| = n - k,  $v_1 \in V(T')$ . Then we have k(n-k) possible ways to add

an edge joining T' to T'' to obtain a tree on V. Thus the pair (T', T'') arises from k(n-k) trees T. The number of such pairs T', T'' is, obviously,

$$\binom{n-1}{k-1}T_kT_{n-k}$$

and hence,

(1) 
$$\sum_{k=1}^{n-1} {n-1 \choose k-1} T_k T_{n-k} k(n-k) = (n-1) T_n.$$

Since

$$\binom{n-1}{k-1} = \frac{n-1}{n-k} \binom{n-2}{k-1},$$

the result follows. Observe that substituting  $T_n = n^{n-2}$  we obtain one of Abel's identities.

To get Cayley's formula note that by changing the index to n-k, we get from (1),

(2) 
$$\sum_{k=1}^{n-1} {n-1 \choose k} T_k T_{n-k} k(n-k) = (n-1) T_n$$

and hence, adding (2) to (1),

(3) 
$$\sum_{k=1}^{n-1} \binom{n}{k} T_k T_{n-k} k(n-k) = 2(n-1) T_n.$$

Using induction on n, we know

$$T_k = k^{k-2}, T_{n-k} = (n-k)^{n-k-2} \quad (1 \le k \le n-1)$$

and so, the left-hand side of (3) is

$$\sum_{k=1}^{n-1} \binom{n}{k} k^{k-1} (n-k)^{n-k-1} = 2(n-1)n^{n-2}$$

by Abel's identity 1.44. This proves that

$$T_n = n^{n-2}$$

- [O. Dziobek, Sitzungsber. Berl. Math. G. 17 (1917) 64-67].
- 7. (a) The recurrence relation in the last problem can be written in the form

$$\frac{T_n}{(n-2)!} = \sum_{k=1}^{n-1} \frac{k \, T_k}{(k-1)!} \, \frac{T_{n-k}}{(n-k-1)!}$$

and thus

(1) 
$$\sum_{n=2}^{\infty} \frac{T_n}{(n-2)!} x^{n-2} = \sum_{n=2}^{\infty} \left( \sum_{k=1}^{n-1} \frac{k \, T_k}{(k-1)!} \, \frac{T_{n-k}}{(n-k-1)!} \right) x^{n-2}.$$

The right-hand side is the product of the power series

$$\sum_{k=1}^{\infty} \frac{k T_k}{(k-1)!} x^{k-1} \text{ and } \sum_{n=1}^{\infty} \frac{T_n}{(n-1)!} x^{n-1}$$

$$\lim_{k \to \infty} \left( \frac{t(x)}{x} \right)' = t'(x) \frac{t(x)}{x}$$

and thus, (1) yields  $\left(\frac{t(x)}{x}\right)' = t'(x)\frac{t(x)}{x}$ 

or, equivalently,

$$\left(\ln \frac{t(x)}{x}\right)' = \frac{\left(\frac{t(x)}{x}\right)'}{\frac{t(x)}{x}} = t'(x),$$

$$\ln \frac{t(x)}{x} = t(x) + C,$$

$$\frac{t(x)}{x} = c \cdot e^{t(x)}.$$

Substituting x = 0 we get c = 1, thus

$$x = t(x)e^{-t(x)}.$$

(b) By the Cauchy integral formula we have for  $n \ge 1$ ,

$$T_n = \frac{1}{n} t^{(n)}(0) = \frac{(n-1)!}{2\pi i n} \oint_C \frac{t'(z)}{z^n} dz = \frac{(n-1)!}{2\pi i n} \oint_C \frac{t'(z)}{\left(t(z)e^{-t(z)}\right)^n} dz,$$

where C is a small circuit around the origin. Since  $t'(0) = 1 \neq 0$ , the mapping t is a homeomorphism in a small neighborhood of 0. Hence t(C) is a simple closed curve around 0 and hence,

(1) 
$$T_n = \frac{(n-1)!}{2\pi i n} \oint_{t(C)} \frac{dw}{(we^{-w})^n} = \frac{(n-1)!}{2\pi i n} \oint_{t(C)} \frac{e^{wn}}{w^n} dw.$$

Here

$$\frac{1}{2\pi i} \oint_{t(C)} \frac{e^{wn}}{w^n} dw$$

is the coefficient of  $x^{n-1}$  in the power series

$$e^{xn} = \sum_{k=0}^{\infty} \frac{(xn)^k}{k!}.$$

Hence

$$T_n = \frac{(n-1)!}{n} \cdot \frac{n^{n-1}}{(n-1)!} = n^{n-2}$$

[G. Pólya, Acta Math. 68 (1937) 145-254].

8. First, let the endpoints  $v_1, \ldots, v_{n-l}$  be specified. Then, by 4.1, the result is

(1) 
$$\sum_{\substack{k_{n-l+1}+\ldots+k_n=n-2\\k_i>1}} \frac{(n-2)!}{k_{n-l+1}!\ldots k_n!}.$$

because  $d_1 = \ldots = d_{n-l} = 1$  in this case. Now observe that

$$\frac{(n-2)!}{k_{n-l+1}!\dots k_n!}$$

is the number of ways to partition n-2 objects into l labelled classes (i.e. there is a first, second, . . . class specified), where the  $i^{\text{th}}$  class contains  $k_{n-i+1}$  elements. Thus, (1) counts the total number of partitions of n-2 objects into l labelled non-empty classes. Thus, (1) is equal to l! S(n-2, l).

Since we have not specified the endpoints, they can be chosen in  $\binom{n}{l}$  ways.

Thus, the number of trees on n points with exactly n-l endpoints is

$$\binom{n}{l} l! S(n-2, l) = \frac{n!}{(n-l)!} S(n-2, l)$$

[A. Rényi, Mat. Kut. Int. Közl. 4 (1959) 73-85].

9. (a) We may assume without loss of generality that the  $n^{th}$  row of A has been removed. Let B be an  $(n-1)\times(n-1)$  submatrix of  $A_0$ . We claim that

$$\det B = \begin{cases} \pm 1 & \text{if the subgraph } G' \text{ formed by edges corresponding to columns in } B \text{ is a spanning tree,} \\ 0 & \text{otherwise.} \end{cases}$$

We use induction on n. Suppose first there is a point  $v_i$   $(i \neq n)$  which has degree 1 in G'. Then the  $i^{th}$  row of B contains exactly one  $\pm 1$  (all other entries in this row belonging to B are 0). Expand det B by this row. The resulting  $(n-2)\times(n-2)$  determinant det B' will correspond to  $G'-v_i$  in the same way as det B does to G'. Since G' is a spanning tree of G iff  $G'-v_i$  is a spanning tree of  $G - v_i$  and  $|\det B| = |\det B'|$  the assertion follows.

Secondly suppose no point of G has degree 1, except possibly  $v_n$ . Then G'is not a spanning tree. Since |E(G')| = n - 1 < n, there must be a point having degree 0 in G'. If this point differs from  $v_n$ , then B has a 0 row and det B = 0. If this point is  $v_n$ , then each column of B contains a 1 and a -1,

thus the sum of rows of B is 0, hence det B = 0.

Now use the Binet-Cauchy formula:

$$\det A_0 A_0^T = \Sigma (\det B)^2,$$

where B ranges over all  $(n-1)\times(n-1)$  submatrices of  $A_0$ . Since by the above,

$$(\det B)^2 = \begin{cases} 1 & \text{if } B \text{ corresponds to a} \\ & \text{spanning tree,} \\ 0 & \text{otherwise,} \end{cases}$$

the assertion follows.

(b) The  $(i, j)^{\text{th}}$  entry of  $A_0 A_0^T$   $(i \neq j)$  is the inner product of the  $i^{\text{th}}$  and  $j^{\text{th}}$  rows of  $A_0$ . In any column of  $A_0$  but at most one, one of the  $i^{\text{th}}$  and  $j^{\text{th}}$  entry

is 0. In the column corresponding to the (i, j)-edge (if any) we have a +1

and a -1. Thus the (i, j)<sup>th</sup> entry of  $A_0 A_0^T$  is -1. The (i, i)<sup>th</sup> entry of  $A_0 A_0^T$  is, obviously, the number of non-zero elements in the i<sup>th</sup> row of  $A_0$ , i.e. the degree of  $v_i$ . Hence the (i, j)<sup>th</sup> entry is

$$\begin{cases} \text{the degree of } v_i & \text{if } i=j, \\ -1 & \text{if } v_i \text{ and } v_j \text{ are adjacent,} \\ 0 & \text{otherwise.} \end{cases}$$

(c) The number of trees on n points is the same as the number of spanning trees of  $K_n$ , i.e.

$$\begin{vmatrix} n-1 & -1 & \dots & -1 \\ -1 & n-1 & \dots & -1 \\ \vdots & \vdots & \ddots & \vdots \\ -1 & -1 & \dots & n-1 \end{vmatrix} = \begin{vmatrix} 1 & 1 & \dots & 1 \\ -1 & n-1 & \dots & -1 \\ \vdots & \vdots & \ddots & \vdots \\ -1 & -1 & \dots & n-1 \end{vmatrix}$$

(by adding all rows to the first one)

$$=egin{bmatrix} 1 & 1 & \dots & 1 \ 0 & n & \dots & 0 \ & & \ddots & & \ddots \ & & \ddots & \ddots & \ddots \ 0 & 0 & \dots & n \ \end{pmatrix}$$

(by adding the first row to the others)

$$= n^{n-2}$$

- [G. Kirchhoff; see Biggs].
- 10. Let us orient G in some way to get a digraph  $\vec{G}$  and define  $E(G) = \{e_1, \dots, e_m\}$ ,

$$a_{ij} = egin{cases} -\sqrt{x_i\,x_k} & ext{if} & e_j = (v_i,\,v_k), \ \sqrt{x_i\,x_k} & ext{if} & e_j = (v_k,\,v_i), \ 0 & ext{if} & e_j ext{ is not adjacent to } v_i \ (1 \leq i \leq n, \ 1 \leq j \leq m), \end{cases}$$

and set

$$A_0 = (a_{ij})_{i=1}^{n-1} {\atop j=1}^m.$$

Then it is easy to check that

$$D = A_0 A_0^T.$$

Let B be an  $(n-1)\times(n-1)$  submatrix of  $A_0$ . We claim that

$$\det B = \begin{cases} \pm \sqrt[n]{x_1}^{d_T(v_1)} \cdots \sqrt[n]{x_n}^{d_T(v_n)} & \text{if the columns of } B \text{ correspond to a spanning tree,} \\ 0 & \text{otherwise.} \end{cases}$$

This follows by induction on n in the same way as in the solution of 4.9. By the Binet-Cauchy formula,

$$\det A_0 A_0^T = \sum (\det B)^2 = \sum_T x_1^{d_T(v_1)} \dots x_n^{d_T(v_n)},$$

where T ranges over all spanning trees of G. This proves the formula.

11. We have to determine the number of spanning trees of the complete bipartite graph  $K_{n,m}$ . It follows from 4.9 (or from 4.10 by setting  $x_1 = \ldots = x_n = 1$ ), that

add all rows to the first one, then add the first row to each of the last m-1 rows, and then we get

$$T(K_{n,m}) = egin{bmatrix} 1 & 1 & \dots & 1 & & 0 & \dots & 0 \ 0 & m & \dots & 0 & & -1 & \dots & -1 \ \vdots & \ddots & \vdots & & \vdots & & \vdots \ 0 & 0 & \dots & m & & -1 & \dots & -1 \ 0 & 0 & \dots & 0 & & n & \dots & 0 \ \vdots & \vdots & & \vdots & & \vdots & & \vdots \ 0 & 0 & \dots & 0 & & 0 & \dots & n \ \end{bmatrix} = m^{n-1}n^{m-1}.$$

12. (a) Let  $E(\overline{G}) = \{e_1, \ldots, e_m\}$  and let  $A_k$  denote the set of those trees on V(G) containing  $e_k$ ; we denote by S the set of all trees on V(G'). Then, by the inclusion-exclusion principle,

(1) 
$$T(G) = \left| S - \bigcup_{k=1}^{m} A_k \right| = \sum_{K \subseteq \{1, \dots, m\}} (-1)^{|K|} |A_K|,$$

where

$$A_{\mathfrak{o}} = S, \quad A_K = \bigcap_{k \in K} A_k.$$

We determine  $|A_K|$ . Set

$$E_K = \{e_k : k \in K\}, G_K = (V(G), E_K)$$

and let  $X_1^{(K)}, \ldots, X_{r_K}^{(K)}$  be the sets of points of components of  $G_K$ . If  $G_K$  is not a forest, then obviously,  $A_K = \emptyset$ . Suppose  $G_K$  is a forest.  $A_K$  consists of all trees on V(G) containing  $G_K$ . By 4.4 the number of such trees is

$$|X_1^{(K)}| \ldots |X_{r_K}^{(K)}| n^{r_K-2}$$
.

Also,  $|K| = n - r_K$  and hence, (1) yields

$$T(G) = \sum_{\substack{K \subseteq \{1, \dots, m\} \\ G_K \text{ is a forest}}} (-1)^{n-r_K} |X_1^{(K)}| \dots |X_{r_K}^{(K)}| \, n^{r_K-2}.$$

For a fixed partition  $\{X_1, \ldots, X_r\}$  the number of terms K with  $\{X_1^{(K)}, \ldots, X_{r_K}^{(K)}\} = \{X_1, \ldots, X_r\}$  is, obviously,

$$T(\bar{G}[X_1]) \dots T(\bar{G}[X_r]);$$

thus, if we sum over all partitions  $X_1, \ldots, X_r$  of V(G), we get

$$\sum (-1)^{n-r} T(\bar{G}[X_1]) \dots T(\bar{G}[X_r]) |X_1| \dots |X_r| n^{r-2}$$

as stated [H.N.V. Temperley; see B].

## (b) First solution. By 4.9,

	2q - 1							n-2q			
	n-2	-1	-1		-1	-1	-1		1		
	-1	n-2	0		-1	-1	-1		-1		
	-1	0	n-2		-1	-1	-1		-1		
						· 17 · 10					
T(G) =	-1	-1			n-2	0	-1		-1		
Age 4. le la	-1	-1	-1		0	n-2	-1		-1		
	-1	-1	-1		-1	-1	n-1		-1		
						:	100		. :		
	-1	-1	-1		-1	-1	-1		n-1		

Add all rows to the first one, then add the first row to all others; we get

	d L	ne decision	2q-1	41			n-2q		
	0	1	1	1	 1	1		1	
T(G) =	-1	n-1	1	0	 0	0		0	
	-1	1	n-1	0	 0	0		0	
	:			•			3 - "	No.	
in the	-1	0	0	0	 0	n		0	
		14 18 12							
of the fee	-1	0	0	0	 0	0		n	

Now add  $\frac{1}{n}$  times all columns to the first one; we get

Second solution. In the formula of 4.12, it suffices to consider those partitions  $(X_1, \ldots, X_r)$  in which every  $X_i$  is one-element or consists of two points

adjacent in 
$$\overline{G}$$
. There are  $\binom{q}{j}$  ways to select  $j$  such classes, thus 
$$T(G) = \sum_{j=0}^q \binom{q}{j} (-1)^{n-(n-j)} 2^j n^{n-j-2} = n^{n-q-2} (n-2)^q \text{ [L. Weinberg, see B].}$$

(c) We shall only give in detail the solution based on (a). Here it suffices to consider partitions  $(X_1, \ldots, X_r)$ , where  $X_1$  contains v and some j points adjacent to v and all other  $X_i$ 's are one-element. The number of such partitions  $\binom{q}{j}$ . Thus we get

$$\begin{split} T(G) &= \sum_{j=0}^{q} \binom{q}{j} (-1)^{n-(n-j)} (j+1) \ n^{n-j-2} = n^{n-2} \big( (1-x)^q x \big)' \bigg|_{\mathbf{x} = \frac{1}{n}} = \\ &= n^{n-2} \left( 1 - \frac{1}{n} \right)^q - n^{n-2} q \left( 1 - \frac{1}{n} \right)^{q-1} \frac{1}{n} = \\ &= (n-q-1) \ (n-1)^{q-1} \ n^{n-q-2} \quad [\text{P.V. O'Neil; see B}]. \end{split}$$

13. (a) Draw the tree so that a is the uppermost point, and departing from a we descend while the other endpoints are on the "ground". It is clear that each binary plane tree can be drawn essentially uniquely in this way. The endpoints different from a have a natural ordering  $v_1, \ldots, v_k$ ; going along the "ground" line in this direction, we have a to our left. Note that, from

$$|E(T)| = \frac{1}{2}(k+1+3(2n-k-1)) = 2n-1,$$

we have k = n.

The tree can be considered as a diagram to calculate the product  $x_1, \ldots x_n$ . The variables are associated with the endpoints on the ground, and every inner point represents the multiplication of the two products corresponding to the two points immediately below it. Thus the number of binary plane trees with 2n points is the same as the number of bracketings of a product of n factors, which is  $\frac{1}{n} \binom{2n-2}{n-1}$  by 1.38.

(b) Consider the  $\pm 1$ -sequence defined in the hint associated with a rooted plane tree. It follows trivially by induction that the sum of the first k entries is equal to our distance from the root after having walked along k edges. This sum is therefore positive, except for k=2n-2, when it is 0. Conversely, given a sequence of n-1 (+1)'s and n-1 (-1)'s such that the partial sums are positive, there is a unique rooted plane tree with this code. In fact, we can build up the walls and the walk around them step-by-step as follows; we start from the root. Assume the walls corresponding to the first k entries have been built up, and we are where we have to be after k steps in the walk. If the next entry is +1, we attach a new wall and build it for a while, always having it to our left. If the next entry is -1, we do not build but just walk along the wall toward the root (keeping it to our left). It is easy to check that this is a correct "decoding". So the number of plane rooted trees on n points is equal to the number of sequences of n-1 (+1)'s and n-1 (-1)'s such that all partial sums are positive. This number is

$$\frac{1}{n-1} \binom{2n-4}{n-2}$$

by 1.33b. [G. Pólya, *Acta Math.* 68 (1937) 145–254. Note the relation between the results of parts (a) and (b); cf. D. A. Klarner, *J. Comb. Theory* 9 (1970) 401–411.]

14. First solution. Draw an arbitrary tree T with points  $v_1, \ldots, v_k$ . Then any forest F in which  $v_1, \ldots, v_k$  are in distinct components and which has k components gives, together with T, a tree containing  $T_0$ ; and conversely. Thus the result is the same as the number of trees on n points containing a given k-point subtree; this number is, by 4.4,

$$(1) E(n,k) = kn^{n-k-1}.$$

Second solution. Let F be one of the forests on  $\{v_1, \ldots, v_n\}$  having k-1 components and such that  $v_1, \ldots, v_{k-1}$  are in distinct components  $T_1, \ldots, T_{k-1}$ . Let  $v_k \in V(T_i)$  and remove the edge adjacent to  $v_k$  which is on the  $(v_k, v_i)$ -path. Then we get a forest F' which has k components and  $v_1, \ldots, v_k$  are in distinct components.

Let us see how many times a given forest F' occurs. Let F' have components  $T'_1, \ldots, T'_k, v_i \in V(T')$ . Then we have to join  $v_k$  to a point of  $V(T'_1 \cup \ldots \cup U'_{k-1})$ , which gives  $|V(T'_1 \cup \ldots \cup T'_{k-1})| = n - |V(T'_k)|$  possibilities.

Now, unfortunately,  $n - |V(T'_k)|$  depends on the special choice of F'. But we can sum this over all permutations of  $v_1, \ldots, v_k$ . Then a given F' will be counted

$$(k-1)! \sum_{i=1}^{k} (n-|V(T_i')|) = (k-1)! (nk-n)$$

times; on the other hand, we get from each forest F with k-1 components  $(v_1, \ldots, v_{k-1})$  in distinct components k! forests F'. Hence

$$k! E(n, k-1) = (k-1)! n(k-1) E(n, k)$$

or, equivalently,

(1) 
$$E(n, k-1) = \frac{k-1}{k} nE(n, k).$$

Hence

$$E(n, k) = \frac{k}{k+1} nE(n, k+1) = \frac{k}{k+1} \frac{k+1}{k+2} n^2 E(n, k+2) =$$

$$= \dots = \frac{k}{k+1} \frac{k+1}{k+2} \dots \frac{n-1}{n} n^{n-k} E(n, n).$$

Since E(n, n) = 1 by definition, this gives

$$E(n, k) = kn^{n-k-1}.$$

Since

$$T(n) = E(n, 1) = n^{n-2},$$

we have obtained a new proof of the Cayley formula too [A. Cayley; see B].

15. Select one edge entering each point  $x \neq a$ . Then we get an arborescence T rooted at a; for we have n-1 edges (n=|V(G)|) and for each x, we can find an edge  $(x_1, x) \in E(T)$ ; then an edge  $(x_2, x_1) \in E(T)$ , etc. The same point cannot occur twice in the sequence  $x, x_1, x_2, \ldots$ , because G is acyclic. Hence there is a unique (a, x)-path in T for every x, and, consequently, T is an arborescence rooted at a.

Conversely, every arborescence rooted at a occurs in this way. Hence the number of such arborescences is

$$\prod_{x \neq a} d^-(x),$$

where  $d^{-}(x)$  is the indegree of x.

16. We use induction on |E(G)|. If |E(G)| < n-1, then  $\Delta(G)$  contains a column which is identically 0. So we may suppose  $|E(G)| \ge n-1$ . Also we may assume no edge enters the root  $v_n$ , because such an edge plays no role.

Suppose first that there is a point, say  $v_1$ , which is the head of at least two edges. Split the edges entering  $v_1$  into two non-empty classes  $C_1$ ,  $C_2$ . Since any arborescence contains exactly one edge which enters  $v_1$ , the number of all (spanning) arborescences of G is equal to the sum of the numbers of arborescences of  $G - C_1$  and  $G - C_2$ . On the other hand, all but one row of  $\Delta(G - C_1)$  are the same as those of  $\Delta(G)$ ; the first rows of  $\Delta(G - C_1)$  and  $\Delta(G - C_2)$  add up to the first row of  $\Delta(G)$ . Hence

$$\Delta(G - C_1) + \Delta(G - C_2) = \Delta(G).$$

Since by the induction hypothesis,  $\Delta(G - C_i)$  is the number of spanning arborescences rooted at  $v_n$  in  $G - C_i$ , the assertion follows.

In the case remaining, exactly one edge enters each point  $v_i$ ,  $1 \le i \le n-1$ . If G is an arborescence, then we have to show

$$\Delta(G)=1.$$

To show this we may renumber the points in such a way that, if  $v_i$  is joined to  $v_j$ , then i > j. Then the matrix  $\Delta(G)$  has 0's above the diagonal and 1's in the diagonal, whence  $\Delta(G) = 1$  follows.

On the other hand, assume G is not an arborescence. Since each point  $\neq v_1$  has indegree 1 and thus, |E(G)| = n - 1, G must be disconnected; let, say,  $v_1, \ldots, v_k$  form a component of G which does not contain  $v_n$ . Then the sum of the first k rows of  $\Delta(G)$  is 0, hence  $\Delta(G) = 0$ . This completes the proof.

(b) Replace each edge e of G by  $x_e$  parallel edges. Then the number of arborescences in the resulting graph G', rooted at  $v_n$  is

$$\sum x_{e_1}\ldots x_{e_{n-1}},$$

where the summation extends over all (n-1)-tuples  $\{e_1, \ldots, e_{n-1}\}$  of edges which form a spanning arborescence. Let  $x_{e_i} = y_v$ , if  $e_i = (v_v, v_\mu)$ . Then

$$\sum x_{e_1} \dots x_{e_{n-1}} = \sum_T \prod_{i=1}^n y_i^{d_T(v_i)};$$

on the other hand, this is equal to

$$\Delta(G) = \begin{vmatrix} \sum_{j\neq 1} a_{j1} y_j & -a_{12} y_1 & \dots & -a_{1,n-1} y_1 \\ -a_{21} y_2 & \sum_{j\neq 2} a_{j2} y_j & \dots & -a_{2,n-1} y_2 \\ \vdots & \vdots & & & & \\ -a_{n-1,1} y_{n-1} & -a_{n-1,2} y_{n-1} & \dots & \sum_{j\neq n-1} a_{j,n-1} y_j \end{vmatrix}$$

Since this equality holds for any choice of  $y_i$  such that  $y_i > 0$  and  $y_i$  is an integer, it holds identically [see B].

17. It is easy to prove that if  $\pi$  has no fixed points, then no tree is invariant under it. We use here the notion of the center (see 6.21) but the reader will find no difficulty in replacing this by other considerations if he wants to avoid reference to later chapters. If our tree has a center this must be a fixed point of  $\pi$ . If it has a bicenter,  $\pi$  must exchange the two points of this. But then it must exchange the two branches of the tree relative to the edge connecting the bicenters, which is only possible if the tree has an even number of points.

II. Now suppose that x is a fixed point of  $\pi$ . Let  $\vec{T}$  be the arborescence arising by orienting T so that x becomes its root. We need some further information on  $\vec{T}$ .

Let  $V_1, \ldots, V_m$  be the underlying sets of the cycles of  $\pi$ ,  $|V_1| \ge |V_2| \ge \ldots \ge |V_m|$ ,  $V_m = \{x\}$ .

- (i)  $\pi$  is an automorphism of  $\tilde{T}$ , obviously.
- (ii) If  $(x, y) \in E(\vec{T})$ ,  $x \in V_i$ ,  $y \in V_j$ , then  $|V_i| \mid |V_j|$ . In fact, if  $|V_i| \nmid |V_j|$ , then there is an image  $(\pi^k(x), \pi^k(y))$  of (x, y) with  $\pi^k(y) = y$  but  $\pi^k(x) \neq x$ . This means two edges of  $\vec{T}$  enter y, which is impossible, because  $\vec{T}$  is an arborescence.
- (iii) If  $(x, y) \in E(\vec{T})$ ,  $x \in V_i$ ,  $y \in V_j$ , then  $i \neq j$ . For there is an edge (z, u) with  $u \in V_j$ ,  $z \notin V_j$ ; a suitable image  $(\pi^k(z), \pi^k(u))$  is such that  $\pi^k(u) = y$ ; thus  $\pi^k(z) = x \notin V_j$ .
- (iv) If we contract each  $V_i$  onto a point  $v_i$ , and cancel the arising multiplicities of edges of  $\vec{T}$ , then  $\vec{T}$  is mapped onto a certain arborescence  $\vec{T}_1$  on  $\{v_1,\ldots,v_m\}$ . In fact, the same argument as in (i) or (ii) shows that all edges of  $\vec{T}$  entering  $V_i$  must come from the same  $V_j$ . Hence exactly one edge of  $\vec{T}_1$  enters each  $v_i$ ,  $i \neq m$ . Since, obviously, each point  $v_i$  is accessible from  $v_m$  on a path of  $\vec{T}_1$ , it follows that  $\vec{T}_1$  is an arborescence rooted at  $v_m$ .
- (v)  $\vec{T}_1$  is a subgraph of the digraph  $G_1$  in which  $(v_i, v_j) \in E(G_1)$  iff  $|V_i| \mid |V_j|$ ; by (i).

Now let us determine, how many trees T yield a given arborescence  $\vec{T}_1$ . Each such T arises by taking one  $(V_i, V_j)$ -edge and its images by  $\pi$  for every  $(v_i, v_j) \in E(\vec{T}_1)$ ; conversely, if we select one  $(V_i, V_j)$ -edge and its images for every  $(v_i, v_j) \in E(\vec{T}_1)$  we get a tree which is invariant under  $\pi$ . The set of all  $(V_i, V_j)$ -

edges consists of  $|V_i|$  orbits (remember  $|V_i| ||V_i|$ ). Thus the number of trees T belonging to a given arborescence  $\vec{T}_1$  is

$$\sum_{(v_i, v_j) \in E(\vec{T}_1)} |V_i| = \prod_{l=1}^m |V_i|^{d_{T_1}^{-}(v_i)}$$

and hence, the number of all trees on V invariant under  $\pi$  is

$$\sum_{\vec{T},} |V_1|^{d_{\vec{T}_1}(v_1)} \dots |V_m|^{d_{\vec{T}_1}(v_m)}.$$

By the preceding problem, this is equal to the determinant

$$\Delta = \begin{bmatrix} \sum_{j \neq 1} a_{j1} \, |V_j| & -a_{12} \, |V_1| & \dots & -a_{1,m-1} \, |V_1| \\ -a_{21} \, |V_2| & \sum_{j \neq 2} a_{j2} \, |V_j| & \dots & -a_{2,m-1} \, |V_2| \\ \vdots & \vdots & \ddots & \vdots \\ -a_{m-1,1} \, |V_{m-1}| & -a_{m-1,2} \, |V_{m-1}| & \dots & \sum_{j \neq m-1} a_{j,m-1} \, |V_j| \end{bmatrix}$$

where

$$a_{ij} = \begin{cases} 1 & \text{if} \quad |V_i| \mid |V_j| \quad (1 \leq i, j \leq m), \\ 0 & \text{otherwise.} \end{cases}$$

This determinant can be evaluated as follows. It has the form

0	0	0	$\left. \left. \left. \right  k_n \right  \right. \right.$
1///	0	0	$\left\{k_{n-1}\right\}$
	1///	0	:
			$\left. \begin{cases} k_1-1, \end{cases} \right.$

since  $|V_i| |V_j|$ ,  $i \ge j$  implies  $|V_i| = |V_j|$ . Hence  $\Delta$  is equal to the product of the shaded subdeterminants. Consider the block corresponding to  $k_i$ . It has

$$\alpha_i = \sum_{d \mid i} dk_d - i$$

in the diagonal, so its value is

for i > 1 and  $k_i \ge 1$ , and

$$arDelta_1 = egin{bmatrix} k_1 - 1 & -1 & \dots & -1 \ -1 & k_1 - 1 & & -1 \ \vdots & & \ddots & \vdots \ \vdots & & \ddots & \vdots \ -1 & -1 & \dots & k_1 - 1 \end{bmatrix} = k_1^{k_1 - 2}$$

(one has to check that these formulas give  $\Delta_i = 1$  if  $k_i = 0$  (i = 2, ..., m) and  $\Delta_1 = 1$  if  $k_1 = 1$ ). Since

$$\Delta = \Delta_1 \dots \Delta_n$$

the assertion is proved.

18. Since the number of rooted trees on n (labelled) points is  $n^{n-1}$  by Cayley's formula and at most (n-1)! of these are isomorphic, we have

$$W_n \ge \frac{n^{n-1}}{(n-1)!} = \frac{n^n}{n!} > 2^n \quad (n \ge 6).$$

Each rooted tree can be drawn in the plane. Join its root to a new point of degree 1 to get a plane tree on n+1 points rooted at an endpoint. This correspondence shows that the number of non-isomorphic rooted trees is not larger than the number of essentially different plane trees rooted at an endpoint, which is, by 4.13b,

$$\frac{1}{n}\binom{2n-2}{n-1}<4^n$$

19. (a) Following the hint, we write the right-hand side in the form

$$x(1 + x + x^{2} + \ldots)^{W_{1}} (1 + x^{2} + x^{4} + \ldots)^{W_{2}} \ldots =$$

$$= x \prod_{T} (1 + x^{|V(T)|} + x^{2|V(T)|} + \ldots),$$

where T ranges over all rooted trees. The coefficient of  $x^n$  in this product is the number of representations of the number n in the form

(1) 
$$n = 1 + \sum_{T} v_T |V(T)|, \qquad v_T \ge 0.$$

Given such a representation, take  $v_T$  copies of T for each rooted tree T and join their roots to a new root. In this way we obtain an n-point rooted tree and conversely, every n-point rooted tree yields a solution of (1). Thus the coefficient of  $x^n$  in the product is  $W_n$ , which proves the identity. (Convergence questions can be settled easily on the basis of 4.18, and will be ignored.)

(b) The two identities are equivalent by 3.7. (For  $x \le 1$  the series on the right-hand sides are convergent because then  $w(x^n) = 0(x^n)$ . The first one follows easily from (a):

$$\log \frac{w(x)}{x} = -\sum_{n=0}^{\infty} W_n \log (1 - x^n) = \sum_{n=0}^{\infty} W_n \sum_{k=1}^{\infty} \frac{x^{kn}}{k} = \sum_{k=1}^{\infty} \frac{1}{k} w(x^k),$$

which proves the identity.

To get the second identity directly by the Pólya-Redfield method, set  $D = \{1, \ldots, d\}$ ,  $R = \{\text{isomorphism types of rooted trees}\}$ , w(T) = |V(T)| for  $T \in R$ , and let  $\Gamma$  be the symmetric group on D. A rooted n-point tree is a d-tuple of "branches", where d is the degree of the root. The branches are themselves rooted trees with total weight n-1. Thus the number  $W_n^{(d)}$  of rooted n-point trees in which the root has degree d is equal to the number of essentially different mappings of D into R with total weight n-1. Hence by 3.29,

$$\sum_{n=1}^{\infty} W_n^{(d)} x^n = x p_d(w(x), w(x^2), \ldots).$$

Summing over all d, we obtain the identity in the problem [A. Cayley].

20. (a) The fact that the radius of convergence of w(x) satisfies  $0 < \tau < 1$  is trivial by 4.18. It follows that

$$\varphi(z) = z \exp\left(\frac{w(z^2)}{2} + \frac{w(z^3)}{3} + \ldots\right)$$

is analytic in the circle of radius  $\sqrt{\tau} > \tau$ . Write the first identity of 4.19a as

$$(1) w(z) e^{-w(z)} = \varphi(z).$$

By 4.7 this can also be written as

(2) 
$$w(z) = t(\varphi(z)) = \sum_{n=1}^{\infty} \frac{n^n}{n!} (\varphi(z))^n.$$

Consider first the values  $0 \le x < \tau$ . Since  $\varphi$  has positive coefficients, it follows by (1) that

 $0 < \varphi'(x) = w'(x) (1 - w(x)) e^{-w(x)}, \quad w(x) < 1.$ 

Again by (1),

$$\varphi(x) < \frac{1}{e}$$
.

By the positivity of the coefficients of  $\varphi$ , we must have  $|\varphi(z)| < \frac{1}{e}$  for  $|z| \le \tau$ 

except for  $z = \tau$ . We must have  $\varphi(\tau) = 1/e$ , otherwise (2) would define w analitically in a larger circle. Hence w is continuous at  $\tau$  and  $w(\tau) = 1$ . Moreover, (2) defines w analytically along the boundary except for  $z = \tau$ .

Investigating now the behavior in the neighborhood of  $\tau$  for  $0 < x < \tau$ , set

 $y = \tau - x$ , u = 1 - w. Then (1) can be written in the form

$$\frac{u^2}{2} + \frac{u^3}{3} + \ldots + \frac{u^n}{n(n-2)!} + \ldots = 1 - e\varphi(x) = a_1 y + a_2 y^2 + \ldots,$$

where

$$a_1 = e\varphi'(\tau) > 0.$$

Hence

(3) 
$$u\sqrt{\frac{1}{2} + \frac{u}{3} + \dots} = \sqrt{y} \sqrt{a_1 + a_2 y + \dots}$$

(the square-roots are to be taken positive if y, u are small positive numbers). The left-hand side is analytic for sufficiently small u and its derivative is non-zero at 0, therefore u can be expressed as an analytic function of the right hand side in a sufficiently small neighborhood of 0. Consequently u is an analytic function of  $\sqrt{y}$  for a sufficiently small y. Thus

(4) 
$$w(x) = \sum_{k=0}^{\infty} b_k (\tau - x)^{k/2},$$

where  $b_0 = 1$  and by (3),  $b_1 = -\sqrt{2a_1} < 0$ .

(b) By (4) above, we have for  $|z| < \tau$ 

(5) 
$$w''(z) = \sqrt{\frac{a_1}{8}} (\tau - z)^{-3/2} + \frac{3b_3}{4} (\tau - z)^{-1/2} + h(z),$$

where h(z) has the following expansion in a neighborhood of  $\tau$ :

(6) 
$$h(z) = \sum_{k=4}^{\infty} \frac{k(k-2)}{4} b_k(\tau-z)^{\frac{k-4}{2}}.$$

(5) implies that h(z) is analytic in the open disc  $|z| < \tau$  and even at all points of the boundary except  $z = \tau$ . (6) implies h(z) is continuous even here. Therefore if we expand h(z) around 0, we have

$$h(z) = \sum_{n=0}^{\infty} c_n z^n,$$

where

$$c_n=rac{1}{2\pi\,i} igoplus_{|z|= au} rac{h(z)}{z^{n+1}}\,dz=O( au^{-n}).$$

Hence by (5), the coefficient of  $x^n$  in w''(x) is

$$\sqrt{\frac{a_1}{8}} \binom{-\frac{3}{2}}{n} (-1)^n \tau^{-n-\frac{3}{2}} + \frac{3}{4} b_3 \binom{-\frac{1}{2}}{n} (-1)^n \tau^{-n-\frac{1}{2}} + O(\tau^{-n}).$$

Here the second term is  $0(\tau^{-n})$  as well, while the first term is asymptotic to

$$c\sqrt{n} \tau^{-n}$$

by elementary calculations using the Stirling formula. This proves the assertion of the problem.

**21.** Let  $a_{1, \pi(1)} a_{2, \pi(2)} \ldots a_{n, \pi(n)}$  be an expansion term of per A. Observe that the number of 1-factors consisting of a  $(u_1, v_{\pi(1)})$ -edge, a  $(u_2, v_{\pi(2)})$ -edge, . . . , a  $(u_n, v_{\pi(n)})$ -edge is exactly

$$a_{1, \pi(1)} a_{2, \pi(2)} \dots a_{n, \pi(n)}$$

because we can choose a  $(u_1, v_{\pi(1)})$ -edge, a  $(u_2, v_{\pi(2)})$ -edge, etc. independently. This has to be summed over all permutations  $\pi$ , i.e. over all expansion terms. This proves the assertion.

22. Let  $a_n$  denote the number required. We have

$$a_0 = 1, \ a_1 = 1.$$

Let F be any 1-factor of the ladder with n steps. There are two possibilities. If F starts with a vertical edge at the left end, then there are  $a_{n-1}$  possible ways to continue. If F starts with horizontal edges, then it can be one of  $a_{n-2}$  1-factors. Hence

$$(2) a_n = a_{n-1} + a_{n-2}.$$

- (1) and (2) show that  $a_n$  is the  $n^{th}$  Fibonacci number (cf. 1.27).
- **23.** The number of 1-factors in  $K_{n,n}$  is, obviously, n! If we have n edges removed from  $K_{n,n}$ , then we do not want to count those 1-factors of  $K_{n,n}$  containing a removed edge. The number of 1-factors containing j given edges is (n-j)!. Hence, by inclusion-exclusion, the number under consideration is

$$\sum_{j=0}^{n} (-1)^{j} \binom{n}{j} (n-j)! = n! \sum_{j=0}^{n} \frac{(-1)^{j}}{j!}.$$

The sum here is a partial sum of the series for  $\frac{1}{e}$  the remainder being less than  $\frac{1}{(n+1)!}$ . Hence

$$\left| n! \sum_{j=0}^{n} \frac{(-1)^{j}}{j!} - \frac{n!}{e} \right| < \frac{1}{n+1}.$$

Thus, our number is the integer next to  $\frac{n!}{e}$ .

24. We may assume that n is even, otherwise the assertion is trivial. Observe that each non-zero term in Pf B corresponds to a 1-factor of G and conversely. Thus, the number of 1-factors of G is  $\leq |Pf B|$  with equality iff all terms in Pf B have the same sign. Let

$$arepsilon\,b_{i_1\,i_2}\,b_{i_3\,i_4}\ldots\,b_{i_{n-1}\,i_n}, \ \delta\,b_{j_1\,j_2}\,b_{j_3\,j_4}\ldots\,b_{j_{n-1}\,j_n}\;(arepsilon,\,\delta=\pm 1)$$

be two terms in Pf B. If they are non-zero, we must have  $(v_{i_{2l-1}}, v_{i_{2l}}) \in E(G)$ ,  $(v_{j_{2l-1}}, v_{j_{2l}}) \in E(G)$ . Let

$$F = (v_{i_1}, v_{i_2}), \dots, (v_{i_{n-1}}, v_{i_n}),$$
  
 $F' = (v_{j_1}, v_{j_2}), \dots, (v_{j_{n-1}}, v_{j_n})$ 

be the two 1-factors of G corresponding to the expansion terms above.  $F \cup F'$  decomposes into alternating circuits  $C_1, \ldots, C_r$  and some edges belonging to both. Let us orient each  $C_i$  in an arbitrary way and let  $\pi$  denote the permutation corresponding to the resulting cycles. Since we have freedom in the ordering of the pairs  $(i_1, i_2), \ldots, (i_{n-1}, i_n)$ , as well as in the choice of order within the pairs we may assume  $(v_{i_{2l-1}}, v_{i_{2l}})$ ,  $(v_{j_{2l-1}}, v_{j_{2l}})$  are either identical or consecutive edges on the same (directed) cycle  $C_i$ .

Now the ratio, or product, of the two expansion terms of the Pfaffian is

(1) 
$$(\varepsilon \delta) b_{i_1 i_2} b_{j_1 j_2} b_{i_3 i_4} b_{j_3 j_4} \dots b_{i_{n-1} i_n} b_{j_{n-1} j_n}.$$

Here  $\varepsilon \delta$  is the sign of the permutation

$$egin{pmatrix} inom{i_1\,i_2\,\ldots\,i_n} \ j_1\,j_2\,\ldots\,j_n \end{pmatrix} = \pi,$$

i.e.

$$\varepsilon\delta=(-1)^r$$
.

On the other hand, the product  $b_{i_1 i_2} \dots b_{j_{n-1} j_n}$  contains a -1 for every edge of  $F \cup F'$  whose orientation in G does not agree with the direction on the  $C_i$  containing it (for the edges of  $F \cap F'$  the corresponding two factors  $b_{i_2 i_1 i_2 i_3} b_{j_2 i_3 i_4} b_{j_2 i_3 i_4} b_{j_3 i_4 i_5} b_{j_3 i_5 i_5} c$  cancel each other out. Since the number of such edges is odd in each  $C_i$ , (1) is equal to

 $(-1)^r(-1)^r = 1$  [P.W. Kasteleyn; see B].

**25.** I. If all circuits C of G which alternate with respect to a given 1-factor  $F_0$  have the property that an odd number of their edges is oriented in a given direction, then  $|\dot{\mathbf{Pf}}| B|$  is the number of 1-factors of G. In fact, let

$$F_0 = \{(v_{i_1}, v_{i_2}), \ldots, (v_{i_{n-1}}v_{i_n})\}$$

and let

$$(1) b_{i_1 i_2} \dots b_{i_{n-1} i_n}$$

be the expansion term of Pf B corresponding to  $F_0$ , then the preceding solution shows that any other non-zero expansion term has the same sign. Hence |Pf B| counts the number of 1-factors.

II. If |Pf B| is equal to the number of 1-factors, then every circuit C which alternates with respect to any 1-factor has the property that an odd number of edges is oriented in a given direction.

In fact, let C be a circuit which alternates with respect to a 1-factor F,

and let

$$F' = F \triangle E(C)$$
.

Assume indirectly that C contains an even number of edges going in one direction. Then the same computation as used before shows that the expansion terms corresponding to F and F' have opposite signs. Since the non-zero expansion terms of Pf B correspond to 1-factors of G and two of these cancel out, the number of 1-factors is greater than |Pf B|, which is a contradiction [ibid.].

**26.** Let 
$$B = (b_{ij})_{i,j=1}^n$$
, then

$$\det B = \sum \varepsilon(\pi) \, b_{1,\,\pi(1)} \dots b_{n,\,\pi(n)},$$

where  $\pi$  ranges over all permutations of  $\{1, \ldots, n\}$  and  $\varepsilon(\pi)$  is the sign of  $\pi$ . Suppose the cycle decomposition of  $\pi$  contains an odd cycle  $(i_1i_2\ldots i_{2k+1})$ ,  $k\geq 1$ . Let  $\pi'$  be the permutation obtained by replacing  $(i_1i_2\ldots i_{2k+1})$  by  $(i_{2k+1}i_{2k}\ldots i_1)$ . Then the expansion terms corresponding to  $\pi$  and  $\pi'$  cancel out as B is skew symmetric.

Also, if  $\pi$  has a fixed element, then the corresponding expansion term is

0 because B has 0's in the diagonal. Thus

$$\det B = \sum \varepsilon(\pi) b_{1, \pi(1)} \dots b_{n, \pi(n)},$$

where it suffices to consider only those permutations  $\pi$  which decompose into even cycles (and these cycles partition  $\{1, \ldots, n\}$ ). This proves the assertion if n is odd, since then det  $B = (\operatorname{Pf} B)^2 = 0$ . Call two permutations equivalent if their cycle decompositions differ in the direction of cycles only. Then all terms corresponding to equivalent permutations are equal. For a permutation  $\pi$  with even cycles that partition  $\{1, \ldots, n\}$ , let  $r(\pi)$  be the number of cycles and  $s(\pi)$  the number of 2-cycles. Then the sum of all terms equivalent to  $\pi$  is

$$(-1)^{r(\pi)} 2^{r(\pi)-s(\pi)} b_{1,\pi(1)} \dots b_{n,\pi(n)}$$

and hence

$$\det B = \sum (-1)^{r(n)} 2^{r(n)-s(n)} b_{1, \pi(1)} \dots b_{n, \pi(n)},$$

where  $\pi$  now ranges over a system of representatives of equivalence classes. Consider (Pf B)<sup>2</sup>. If we expand this, each term will have the form

$$\varepsilon \delta b_{i_1 i_2} \ldots b_{i_{n-1} i_n} b_{j_1 j_2} \ldots b_{j_{n-1} j_n},$$

where  $\{\{i_1, i_2\}, \ldots, \{i_{n-1}, i_n\}\}, \{\{j_1, j_2\}, \ldots, \{j_{n-1}, j_n\}\}$  are partitions and

 $\varepsilon$ ,  $\delta$  are the corresponding signs.

Consider the graph on  $\{1, \ldots, n\}$  whose edges are  $(i_{2l-1}, i_{2l})$  and  $(j_{2l-1}, j_{2l})$   $(l=1,\ldots,n)$ . This graph consists of s edges which arise in both forms and r-s circuits whose edges are alternatingly  $(i_{2l-1}, i_{2l})$  and  $(j_{2l-1}, j_{2l})$ . We may assume  $(i_{2l-1}, i_{2l})$  and  $(j_{2l-1}, j_{2l})$  are either identical or consecutive edges on these circuits. Then set

$$\varrho = \begin{pmatrix} i_1 i_2 \dots i_n \\ j_1 j_2 \dots j_n \end{pmatrix}.$$

 $\varrho$  is determined up to equivalence by the given term of (Pf B)<sup>2</sup>.

Also,  $\varepsilon \delta$  is the sign of  $\varrho$ . Hence, (2) is equal to

$$(-1)^{r(\varrho)} b_{1,\,\varrho(1)} \dots b_{n,\,\varrho(n)}$$

Moreover, we get equivalent permutations for  $2^{r(\varrho)-s(\varrho)}$  terms of  $(Pf B)^2$ . Thus

$$(\operatorname{Pf} B)^{2} = \sum_{\varrho} (-1)^{r(\varrho)} 2^{r(\varrho) - s(\varrho)} b_{1, \, \varrho(1)} \dots b_{n, \, \varrho(n)} = \det B.$$

27. First we prove the assertion of the hint by induction. If the graph is a tree, there is no bounded face and the assertion is trivial.

Suppose G is not a tree. Remove an edge of G which is on the boundary of the infinite face and which also belongs to a circuit and orient the rest as required. Put this edge back, then all bounded faces but the one containing this edge

hence

and

have the property that going around their boundary in the positive sense, we find an odd number of edges oriented in the direction of our walk. A suitable orientation of e will guarantee this for the extra bounded face too.

Now we prove that any circuit C which alternates with respect to the given 1-factor F contains an odd number of edges oriented in one way (or the other) around the circuit. G contains an even number 2p of points inside the circuit C because F matches these points. Let |V(C)| = 2k, and denote by  $A_1, \ldots, A_f$  the faces inside C; let  $q_i$  be the number of edges passed in the right direction, when going around  $A_i$  in the positive sense.

Consider  $q_1 + \ldots + q_f$ . Note that here the edges inside C are counted exactly once; the edges on C are counted iff they are oriented according to the positive

orientation of C. Thus we are interested in the number

$$q = q_1 + \ldots + q_f - m \equiv f - m \pmod{2},$$

where m is the number of edges inside C.

Now consider the graph G' formed by C and those edges and points of G inside C. G' has f+1 faces, m+2k edges and 2p+2k points. By the Euler polyhedron theorem (cf. also 5.24),

$$f + 1 + 2p + 2k = m + 2k,$$
  

$$m \equiv f + 1 \pmod{2}$$
  

$$q \equiv f - m \equiv 1 \pmod{2}$$

as stated [P.W. Kasteleyn; see B].

28. If we consider the orientation of the ladder shown in Fig. 13 we find that the boundary of any bounded face contains an odd number (3) of edges going in the positive direction. Thus, by the preceding solution, any circuit which alternates with respect to a 1-factor contains an odd number of edges oriented one given way around the circuit.

The corresponding matrix B defined as in 4.24 is

where 
$$A = \begin{pmatrix} A & I \\ -I & -A \end{pmatrix}, \\ A = \begin{pmatrix} 0 & 1 & & 0 \\ -1 & 0 & & \\ & & \ddots & \\ & & & 0 & 1 \\ 0 & & -1 & 0 \end{pmatrix}, \quad I = \begin{pmatrix} 1 & & 0 \\ & \ddots & \\ & & & 0 \\ 0 & & 1 \end{pmatrix}.$$

The number of 1-factors of the ladder is given by

$$\det \begin{pmatrix} A & I \\ -I & -A \end{pmatrix}^{1/2} = \det \begin{pmatrix} A & I - A^2 \\ -I & 0 \end{pmatrix}^{1/2} = \left(\det (I - A^2)\right)^{1/2} =$$

$$= \left(\det (I - A) \det (I + A)\right)^{1/2} = \det (I + A),$$

because

$$I - A = (I + A)^T.$$

It is quite easy to verify by induction that this is the same as the  $n^{th}$  Fibonacci number.

**29.** (a) What we have to enumerate is the number of 1-factors of a  $(2n) \times (2n)$  "lattice", i.e. the graph shown in Fig. 26. If we consider the orientation given there, it will satisfy the condition that the boundary of any bounded face con-




Fig. 26

tains one edge going one way and three the other way. Hence by the solution of 4.27, every circuit which alternates with respect to a given 1-factor has an odd number of edges oriented in a given direction around the circuit. Hence, if we form the corresponding matrix B as in 4.24, the number of 1-factors will be equal to  $\sqrt{\det B}$ .

Now B can be written in the form

(1) 
$$B = \begin{bmatrix} A & I & & \mathbf{0} \\ -I & -A & I & & \\ & -I & \cdot & \cdot & \\ & & \cdot & A & I \\ \mathbf{0} & & & -I - A \end{bmatrix}$$

where

$$A = egin{bmatrix} 0 & 1 & & \mathbf{0} \\ -1 & 0 & & & \\ & \cdot & \cdot & \cdot & \\ & & \cdot & \cdot & 1 \\ \mathbf{0} & & -1 & & 0 \end{pmatrix}.$$

If we multiply the first column, then the 3<sup>rd</sup> and 4<sup>th</sup> row, then the 4<sup>th</sup> and 5<sup>th</sup> column, then the 7<sup>th</sup> and 8<sup>th</sup> row etc. of the partitioned matrix (1) by -1, we get the partitioned matrix

$$B' = \begin{bmatrix} -A & I & & \mathbf{0} \\ I & -A & I & & \\ & I & \cdot & \cdot & \\ & & \cdot & \cdot & \cdot \\ & & & \cdot & -A & I \\ \mathbf{0} & & & I - A \end{bmatrix}.$$

Set

then

$$\det B = \det B' = \det p_n(A).$$

By 1.29, the eigenvalues of A are  $2\cos\frac{k\pi}{2n+1}$ ,  $k=1,\ldots,2n$ . Hence

$$\det \, p_{\scriptscriptstyle n}(A) = \det \coprod_{k=1}^{2n} \left[ 2 \, \cos \frac{k\pi}{2n+1} \, I - A \right] = \coprod_{k=1}^{2n} \det \left[ 2 \, \cos \frac{k\pi}{2n+1} \, I - A \right].$$

Let

$$q_n(\lambda) = \det (A - \lambda I),$$

then, in turn,

$$\det B = \prod_{k=1}^{2n} q_n \left( 2 \cos \frac{k\pi}{2n+1} \right).$$

The eigenvalues of A can be determined in the same way as those of A' and they turn out to be  $2i\cos\frac{k\pi}{2n+1}$ ,  $k=1,\ldots,2n$ . Thus

$$\det B = \prod_{k=1}^{2n} \prod_{l=1}^{2n} \left( 2\cos\frac{k\pi}{2n+1} - 2i\cos\frac{l\pi}{2n+1} \right) =$$

$$egin{aligned} &=2^{4n^2}\prod_{k=1}^{2n}\prod_{l=1}^{2n}\left|\cosrac{k\pi}{2n+1}-i\cosrac{l\pi}{2n+1}
ight|=\ &=2^{4n^2}\prod_{k=1}^{2n}\prod_{l=1}^{2n}\left(\cos^2rac{k\pi}{2n+1}+\cos^2rac{l\pi}{2n+1}
ight)^{1/2}=\ &=2^{4n^2}\prod_{k=1}^{n}\prod_{l=1}^{n}\left(\cos^2rac{k\pi}{2n+1}+\cos^2rac{l\pi}{2n+1}
ight)^2. \end{aligned}$$

Hence,

$$egin{align} a_n &= \sqrt{\det B} = 2^{2n^2} \prod_{k=1}^n \prod_{l=1}^n \left( \cos^2 rac{k\pi}{2n+1} + \cos^2 rac{l\pi}{2n+1} 
ight) = \ &= \prod_{k=1}^n \prod_{l=1}^n \left( 4 \cos^2 rac{k\pi}{2n+1} + 4 \cos^2 rac{l\pi}{2n+1} 
ight). \end{split}$$

(b) The polynomial with roots  $4\cos^2\frac{k\pi}{2n+1}$ ,  $k=1,\ldots,n$  is

$$f(x) = x^n - {2n-1 \choose 1} x^{n-1} + {2n-2 \choose 2} x^{n-2} \dots$$

Observe that  $a_n$  is the resultant of f(x) and f(-x). By the Sylvester form of the resultant,

$$a_n = egin{bmatrix} 1 & -inom{2n-1}{1} & inom{2n-2}{2} & \dots \\ 0 & 1 & -inom{2n-1}{1} & \dots \\ 0 & 1 & -inom{2n-1}{1} & \dots \\ \vdots & \ddots & \ddots & \ddots \\ 0 & 1 & inom{2n-2}{2} & \dots \\ 0 & 1 & inom{2n-2}{2} & \dots \\ 0 & \vdots & \ddots & \dots \\ 0 & \vdots & \dots & \dots \\ 0 & \vdots$$

Add the  $(n+k)^{\text{th}}$  row to the  $k^{\text{th}}$   $(k=1,\ldots,n)$ , then divide the first k rows by 2 and subtract the  $k^{\text{th}}$  row from the  $(n-k)^{\text{th}}$ . The resulting determinant is

$$a_n = 2^n \begin{bmatrix} \binom{2n}{0} & 0 & \binom{2n-2}{2} & 0 & \dots \\ 0 & \binom{2n}{0} & 0 & \binom{2n-2}{2} & \dots \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \binom{2n-1}{1} & 0 & \binom{2n-3}{3} & \dots \\ 0 & 0 & \binom{2n-1}{1} & 0 & \dots \\ \vdots & \vdots & \ddots & \vdots \\ 2n \text{ columns} \end{bmatrix}^{n \text{ rows.}}$$

Suppose, e.g. n is even. Then the last column looks like this:

$$a_n = 2^n egin{bmatrix} 2n & 0 & 0 & 2n-2 \ 0 & 0 & 2n \ 0 & 0 & 2n-2 \ 0 & 0 & 2n-3 \ 0 & 0 & 2n-3 \ 0 & 0 & 2n-1 \ 0 & 0 & 0 \ \end{bmatrix} n ext{ rows.}$$

2n columns

Thus expanding by the first and last columns we get

$$a_n = 2^n egin{bmatrix} 2n & 0 & 1 & 2n-2 \ 0 & 2n & 0 & 2n-2 \ 0 & 2n-1 & 0 & 2n-3 \ 0 & 2n-1 & 0 & 2n-3 \ 0 & 2n-1 & 0 & 2n-3 \ 0 & 2n-3 & 2n-2 \ 0 & 2n-2 & 2n-2$$

Now this determinant is the direct product of two determinants of the form

$$\begin{pmatrix} 2n \\ 0 \end{pmatrix} & \begin{pmatrix} 2n-2 \\ 2 \end{pmatrix} & \begin{pmatrix} 2n-4 \\ 4 \end{pmatrix} \cdots \\ 0 & \begin{pmatrix} 2n \\ 0 \end{pmatrix} & \begin{pmatrix} 2n-2 \\ 2 \end{pmatrix} \cdots \\ \vdots & \vdots & \ddots \\ \vdots & \ddots & \vdots \\ 2n-1 \\ 1 \end{pmatrix} & \begin{pmatrix} 2n-3 \\ 3 \end{pmatrix} & \begin{pmatrix} 2n-5 \\ 5 \end{pmatrix} \cdots \\ 0 & \begin{pmatrix} 2n-1 \\ 1 \end{pmatrix} & \begin{pmatrix} 2n-3 \\ 3 \end{pmatrix} \cdots \\ \vdots & \vdots & \ddots \\ \vdots & \vdots & \ddots & \vdots \\ 2n-3 \\ 3 \end{pmatrix} \cdots \\ \vdots & \vdots & \vdots & \vdots \\ \frac{n}{2} \text{ rows.}$$

This proves the assertion. The case of odd n follows similarly [see E.W. Montrall, in: Applied Combinatorial Mathematics, (E.E. Beckenbach, ed.) Wiley, 1964].

(c) We have

$$\frac{\log a_n}{n^2} = \frac{1}{n^2} \sum_{k=1}^n \sum_{l=1}^n \log \left( 4 \cos^2 \frac{k\pi}{2n+1} + 4 \cos^2 \frac{l\pi}{2n+1} \right).$$

The right-hand side tends to

$$(1) \qquad \qquad rac{4}{\pi^2}\int\limits_0^{\pi/2}\int\limits_0^{\pi/2}\log\left(4\cos^2x+4\cos^2y
ight)dx\,dy=cpprox 1,2\,.$$

Thus,

$$\frac{\log a_n}{n^2} \to c \approx 1.2$$

[ibid.].

**30.** Let G denote the  $(2n-1)\times(2n-1)$  lattice; its points are (i,j)  $(0\leq i\leq n)$  $\leq 2n-2$ ,  $0 \leq j \leq 2n-2$ ). Call point (i,j) black if i and j are both even; green if one of them is odd; red if both are odd. The black points form an  $n \times n$ lattice graph H. Let T be any spanning tree of H. Let x be a black point. Then there is a well-defined first edge on the path in T connecting x to a = (2n - 2, 2n - 2) and this contains a green point x'.

Let y be a red point. In the lattice of red points, there is a unique path which connects y to the outside boundary of the lattice G and on this, there is a first edge which contains a green point. Let y' be this green point.

The pairs (x, x'), (y, y') form, as is easily verified, a 1-factor of G - a

(Fig. 27).

Conversely, let F be a 1-factor of G-a. Consider the set T of those edges of H which contain an edge of F. These form a spanning tree. In fact, the number of edges of F adjacent to black points is  $n^2 - 1$ , so T contains  $n^2 - 1 =$ = |V(H)| - 1 edges; it suffices to prove they do not form circuit. Suppose indirectly they formed a circuit C. The number of points of G inside C is odd (this easily follows, e.g. by induction on the length of C) and so, F cannot match them, a contradiction.




Fig. 27

Thus we have established a one-to-one correspondence between spanning trees of H and 1-factors of G-x [H.N.V. Temperley, in: Combinatorics, London Math. Soc. Lecture Notes Series 13 (1974) 202–204].

**31.** Orient the edge  $(u_i, v_i)$  from  $u_i$  to  $v_i$  and then identify  $u_i$  and  $v_i$   $(i=1,\ldots,n)$ ,

our graph is thereby mapped onto the transitive tournament  $T_n$ .

If M is a k-element matching in G, then M is mapped onto a set M' of edges of  $T_n$  such that at most one edge of M' enters and leaves any given point. Hence M' consists of disjoint directed paths; if we consider the points of  $T_n$  not touched by M' as one-element paths, we can observe that M corresponds to a system of n-k disjoint directed paths which cover all points of  $T_n$ . Conversely, the edges of a system of n-k disjoint directed paths covering  $V(T_n)$  corresponds to a k-element matching of G.

Now let  $P_1, \ldots, P_{n-k}$  be disjoint directed paths which cover  $V(T_n)$ . Then  $\{V(P_1), \ldots, V(P_{n-k})\}$  is a partition of  $V(T_k)$ . Each partition of  $V(T_k)$  into n-k classes arises uniquely in this way; if  $\{V_1, \ldots, V_{n-k}\}$  is a partition of  $V(T_n)$  and  $P_i$  is the unique Hamiltonian path of the (transitive) subtournament induced by  $V_i$ , then  $V(P_i) = V_i$  and  $P_1, \ldots, P_{n-k}$  are disjoint directed paths

that cover  $V(T_n)$ .


Hence, there is a one-to-one correspondence between k-element matchings of G and partitions of  $V(T_n)$  into n-k (non-empty) classes. Thus the required number is S(n, n-k).

**32.** Let  $F_n$  denote the number of such permutations. By the assumption,  $\pi(n) = n$  or  $\pi(n) = n - 1$ . The number of permutations  $\pi$  with the desired property which fix n is  $F_{n-1}$ . The number of those with  $\pi(n) = n - 1$  is  $F_{n-2}$ , because, then it necessarily exchanges n and n-1 (nothing else can be mapped onto n). Thus

$$F_n = F_{n-1} + F_{n-2}$$
.

Since  $F_0 = F_1 = 1$ ,  $F_n$  is the n<sup>th</sup> Fibonacci number.

One can formulate the problem as follows. Let  $\{v_1, \ldots, v_n\}$ ,  $\{u_1, \ldots, u_n\}$  be two disjoint sets of points. Join  $u_i$  to  $v_j$  iff  $|i-j| \leq 1$  (Fig. 28). If  $\pi$  is any



admissible permutation, then the pairs  $(i, \pi(i))$  form a 1-factor of the resulting graph G and conversely. Thus we want to know the number of 1-factors of the graph G. Since G is isomorphic to the graph in Fig. 1 (p. 32) it follows that this number is the n<sup>th</sup> Fibonacci number.

33. It is easy to see that

Expanding this by the first row we get

To get a complete recurrence relation we also expand  $b_n$ ,  $c_n$  by their first columns:

$$(2) b_n = a_{n-1} + b_{n-1};$$

but with  $c_n$  we get a new permanent:

Expand  $d_n$  too:

$$d_n = a_{n-1} + e_{n-1},$$

where

Trivially,

$$(5) e_n = a_{n-1}$$

(1)—(5) yield recurrence relations for  $a_n$ ,  $b_n$ ,  $c_n$ ,  $d_n$ ,  $e_n$ , starting with the value n=2:  $a_2=b_2=c_2=d_2=2$ ,  $e_2=1$ . Set

$$A = egin{bmatrix} 1 & 1 & 1 & 0 & 0 \ 1 & 1 & 0 & 0 & 0 \ 0 & 1 & 0 & 1 & 0 \ 1 & 0 & 0 & 0 & 0 \ 1 & 0 & 0 & 0 & 1 \ \end{pmatrix}, \qquad \mathbf{v}_n = egin{bmatrix} a_n \ b_n \ c_n \ d_n \ e_n \ \end{pmatrix},$$

then

$$\mathbf{v}_n = A\mathbf{v}_{n-1} = \ldots = A^{n-2}\mathbf{v}_2 = A^{n-1}\mathbf{v}_1 = A^n\mathbf{v}_0,$$

where

$$\mathbf{v_0} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \qquad \mathbf{v_1} = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$

by convention. Hence,

$$\sum_{n=0}^{\infty} t^n \mathbf{v}_n = \left(\sum_{n=0}^{\infty} t^n A^n\right) \mathbf{v}_0 = (I - tA)^{-1} \mathbf{v}_0.$$

We are interested in the first entry of  $(I - tA)^{-1}\mathbf{v}_0$ , i.e. the upper left entry of  $(I - tA)^{-1}$ . By Cramer's rule this is equal to

$$f(t) = \frac{\begin{vmatrix} 1-t & 0 & 0 & 0 \\ -t & 1 & -t & 0 \\ 0 & 0 & 1 & -t \\ 0 & 0 & 0 & 1 \end{vmatrix}}{\begin{vmatrix} 1-t & -t & -t & 0 & 0 \\ -t & 1-t & 0 & 0 & 0 \\ -t & 1-t & 0 & 0 & 0 \\ 0 & -t & 1 & -t & 0 \\ -t & 0 & 0 & 1 & -t \\ -t & 0 & 0 & 0 & 1 \end{vmatrix}} = \frac{1-t}{t^5 - 2t^3 - 2t + 1}$$

[D. H. Lehmer, in: Comb. Theory Appl. Coll. Math. Soc. J. Bolyai 4, Bolyai–North-Holland (1970) 755–770].

34. Expanding the permanent in the hint by its first row we get

$$u_{n,p}=p\cdot u_{n-1,p}.$$

Hence

$$u_{n,p} = pu_{n-1,p} = \ldots = pu^{n-p} u_{p,p} = p^{n-p} p!$$

35. We want to determine

$$\begin{pmatrix}
1 & \dots & 1 & 0 & \dots & 0 \\
\vdots & & & & \ddots & \vdots \\
1 & & & & & \ddots & \vdots \\
0 & & & & & & 1 \\
\vdots & & & & & & \ddots & \vdots \\
0 & \dots & & & & & & \vdots \\
0 & \dots & & & & & & & \vdots \\
0 & \dots & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & & \vdots \\
0 & \dots & & & & & & & & & & & & \vdots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots \\
0 & \dots & \dots & \dots & \dots & \dots & \dots \\
0 & \dots & \dots & \dots & \dots & \dots & \dots \\
0 & \dots & \dots & \dots & \dots & \dots & \dots \\
0 & \dots & \dots & \dots & \dots & \dots & \dots \\
0 & \dots & \dots & \dots & \dots & \dots & \dots \\
0 & \dots & \dots & \dots & \dots & \dots & \dots \\
0 & \dots & \dots & \dots & \dots & \dots & \dots \\
0 & \dots & \dots & \dots & \dots & \dots \\
0 & \dots & \dots & \dots & \dots & \dots \\
0 & \dots & \dots & \dots & \dots & \dots & \dots \\
0 & \dots & \dots & \dots & \dots & \dots & \dots \\
0 & \dots & \dots & \dots & \dots & \dots & \dots \\
0 & \dots & \dots & \dots & \dots & \dots \\$$

i.e. the number of those expansion terms of

$$\operatorname{per}\left(\begin{array}{ccc} 1 & \dots & 1 \\ \vdots & & \vdots \\ 1 & \dots & 1 \end{array}\right),$$

which do not contain any entry from the two triangles which contain 0's in (1). By inclusion-exclusion this is equal to

$$\sum_{U,V} (-1)^{|U|+|V|} K_{U\cup V},$$

where U is a subset of the entries in the triangle in the lower left corner and V is a set of entries in the triangle in the upper right corner, and  $K_{U \cup V}$  is the number of expansion terms containing  $U \cup V$ . Obviously, it suffices to consider those sets U, V in which no two entries are in a row or column.

We have

$$K_{U \cup V} = (n - |U \cup V|)!.$$

Thus our number is

$$\sum_{U,V} (-1)^{|U|+|V|} (n-|U|-|V|) ! = \sum_{k=0}^{p} \sum_{l=0}^{p} (-1)^{k+1} (n-k-l) ! \sum_{\substack{|U|=k\\|V|=l}} 1.$$

By 4.31, the number of sets U under consideration with |U| = k is S(p+1, p+1-k) and, similarly, we have S(p+1, p+1-l) sets V. Thus the result is

$$\sum_{k=0}^{p} \sum_{l=0}^{p} (-1)^{k+l} (n-k-l) ! S(p+1, p+1-l) S(p+1, p+1-k).$$

36. If an expansion term of the permanent contains k 1's from the upper left block, then it contains n-k 1's from the upper right block, n-k 1's from the lower left block and k 1's from the lower right block. If we select the n-k 1's from the upper right block and n-k 1's from the lower left block, then there are  $(k!)^2$  possible ways to select the rest.

Observe that the (n-k)-tuples of 1's, no two in a row or column, out of the upper right  $n \times n$  block correspond to k-element matchings of the graph defined in 4.31; hence their number is S(n, k). Similarly for the lower left corner. Thus the number of selections for a fixed k is

$$S(n, k)^2 (k!)^2$$
,

which proves the assertion [K. Vesztergombi, Studia Sci. Math. Hung. 9 (1974) 181–185].

## § 5. Parity and duality

1. (a) Summing the degrees of a graph, we get every edge counted twice (once for each of its endpoints). Thus, the sum of degrees is twice the number of edges.

As 3+3+3+3+5+6+6+6+6+6+6+6 is odd (there is an odd number of odd summands), this sequence is not the degree-sequence of any graph. This gives generally that the number of odd degrees is even.

(b) Let  $d_1, \ldots, d_k$  be the degrees of the "upper" points,  $r_1, \ldots, r_e$  the degrees of "lower" points of a bipartite graph. Then both  $d_1 + \ldots + d_k$  and  $r_1 + \ldots + r_e$  give the number of edges, hence

$$(*) d_1 + \ldots + d_k = r_1 + \ldots + r_e.$$

Now suppose that the degrees are as given in the problem. Let  $d_1 = 5$  (say). Then the right-hand side of (\*) is divisible by 3, while the left-hand side is not. Thus, this sequence is not the degree-sequence of any bipartite graph (it is, however, the degree sequence of a graph).

- (c) Since the point with degree 9 is adjacent to all other points, it must in particular be adjacent to both points of degree 1. Hence, the point with degree 8 cannot be adjacent to either of the two points of degree 1; but then there are only 7 more points left and so, it cannot have degree 8 [cf. problem 7.51].
- **2.** Assume there is a k-regular simple graph on n points. By the solution of 5.1a,  $k \cdot n$  is twice the number of edges, thus
- (\*)  $k \cdot n$  is even;

also trivially

$$(**) k \le n-1.$$

We show that, if (\*) and (\*\*) are satisfied, there is a k-regular simple graph on n points.




Fig. 29

Case 1. k is even. Consider the vertices of a regular n-gon and join each of them to its neighbors,  $2^{nd}$  neighbors, etc.,  $\binom{k}{2}^{th}$  neighbors. Since k < n, this gives no multiple edges (Fig. 29).

Case 2. k is odd. Then by (\*), n is even. Consider the graph of degree k-1 on n points constructed in Case 1. By (\*\*) the endpoints of the longest diagonals are not joined; thus, if we add these edges, we get a k-regular simple graph.

3. If the graph G is bipartite and C is a circuit of G, then C, as a subgraph of G, is bipartite as well and hence it is even. Now suppose G has only even circuits and let  $P_1$ ,  $P_2$  be (x, y)-paths. We prove by induction on  $V(P_1)$  that  $P_1$ ,  $P_2$ .

have the same parity. If  $P_1 \cup P_2$  is a circuit, then, since it is even,  $P_1$  and  $P_2$ 

have the same parity.

So suppose  $P_1$ ,  $P_2$  have a point in common; z. Let z split them into pieces  $P_1'$ ,  $P_1''$  and  $P_2'$ ,  $P_2''$ , respectively. The notation may be chosen so that  $P_1'$ ,  $P_2'$  are (xz)-paths and hence by the induction hypothesis, they have the same parity. Similarly,  $P_1''$  and  $P_2''$  have the same parity, which proves that so do  $P_1 = P_1' \cup P_1''$  and  $P_2 = P_2' \cup P_2''$ .

To prove 2-colorability we may assume G is connected. Let  $\pi(x,y)$  denote the

common parity of (x, y)-paths  $(\pi(x, y) = 0 \text{ or } 1)$ . Let  $x_0 \in V(G)$  and set

$$S_i = \{y : \pi(x_0, y) = i\}$$
  $(i = 0, 1).$ 

Then  $\{S_0, S_1\}$  is a good bicoloration; for let (u, v) be an edge and  $u \in S_0$  (say), and consider an  $(x_0, u)$ -path P. If  $v \notin V(P)$ , then P + (u, v) is an  $(x_0, v)$ -path, whence

$$\pi(x_0, v) = \pi(x_0, u) + 1 = 1.$$

If v is on P, and divides it into an  $(x_0, v)$ -piece  $P_1$  and a (v, u)-piece  $P_2$ , then  $P_2 + (u, v)$  is a circuit and hence, it is even. This means that  $P_2$  has odd length Now

$$0 = \pi(x_0, u) \equiv |E(P)| = |E(P_1)| + |E(P_2)| \equiv \pi(x_0, v) + 1 \pmod{2}$$

This shows that  $\pi(x_0, v) = 1$ , i.e.  $v \in S_1$ .

The statement is not true for digraphs, since the triangle with transitive orien-

tation is not bipartite and has no odd cycles (no cycles at all, in fact).

However, it remains true for strongly connected digraphs. To show this, first one proves that if G has no odd cycles and P is an (xy)-path while Q is a (y, x)-path, then P, Q have the same parity. This goes exactly like the first part of the proof above. Now let  $P_1, P_2$  be two (x, y)-paths, then choose a (y, x)-path Q (here we use the fact that G is strongly connected), then  $P_1$  and Q have the same parity. But so do  $P_2$  and Q, and therefore  $P_1$  and  $P_2$  have the same parity. The rest of the proof runs exactly as in the undirected case.

4. If there is a potential p(x), the work on any walk is the change in potential between the endpoints. In particular, it is 0 on any closed walk, so on any circuit.

Now suppose that the sum of work on any circuit is 0. We claim it is 0 on any closed walk W. We use induction on the length of W. If W is a circuit the statement is the assumption. So suppose W uses a point x twice. x splits W into two smaller closed walks,  $W_1$ ,  $W_2$ . The work on  $W_1$  and on  $W_2$  is 0 by the induction hypothesis, but then so is the work on W.

Now if  $P_1$  and  $P_2$  are any two (x, y)-walks we can form a closed walk W by going on  $P_1$  from x to y and then returning on  $P_2$ . Since W has total work 0,

 $P_1$  and  $P_2$  need the same work.

Now fix a point  $x_0$  and let p(y) be defined as the work needed to go from  $x_0$  to y. By the above, this is independent of the way we go from  $x_0$  to y. It is easy to see that this potential function satisfies the requirements.

We remark that the values of v(e) and p(x) can be from an arbitrary commutative group. Moreover, if this group is such that each element has order 2, the prior to the prior to the prior of C plays no relative to the prior to the pri

orientation of G plays no role.

5. First let C be an even cycle. We can 2-color C in such a way that each point of C is joined to a point of opposite color, simply coloring its points alternatingly. If C does not span G, we extend this 2-coloration as follows. There is a point  $x_1$  joined to C, because G is strongly connected; similarly, select points  $x_2, \ldots, x_m$  so that  $x_i$  is joined to  $C \cup \{x_1, \ldots, x_{i-1}\}$   $(i = 2, \ldots, m)$ . Finally, every point of G not on C becomes one of the  $x_i$ 's. Now if  $C \cup \{x_1, \ldots, x_{i-1}\}$  is already 2-colored, give  $x_i$  the color opposite to the color of a point of  $C \cup \{x_1, \ldots, x_{i-1}\}$  which  $x_i$  is joined to. Obviously, this coloration has the desired property.

Suppose now that there is a 2-coloration of G with the property in consideration, with color classes  $S_1$ ,  $S_2$ . Let  $G_0$  denote the graph consisting of the edges between  $S_1$  and  $S_2$ . Since  $G_0$  has positive outdegrees by the assumption, it

contains a cycle C. Since  $G_0$  is bipartite, C has even length.

**6.** Let L be a maximum closed trail in G. If L does not contain all the edges' then there is an edge e(x, y) which has a point in common with L; for otherwise L would be a connected component of G which is impossible because G is weakly connected. We may assume e has its tail x on L.

Let us start with this edge and walk along edges of G - E(L), using an edge only once as long as we can. We cannot get stuck at any point  $z \neq x$ ; for there are as many edges of G - E(L) leaving it as running into it and any time we

use an edge leaving z we had to use one running into it before.

Thus we get stuck at x; now let us go on to traverse L starting with x. This gives us a trail longer than L.

- 7. (a) The indegree and outdegree of any point are both equal to n. G is strongly connected. For  $(a_1, \ldots, a_k)$ ,  $(a_2, \ldots, a_k, b_1)$ ,  $(a_3, \ldots, b_2)$ ...  $(a_k, b_1, \ldots, b_{k-1})$ ,  $(b_1, \ldots, b_k)$  is a directed walk from  $(a_1, \ldots, a_k)$  to  $(b_1, \ldots, b_k)$ . Thus G is Eulerian.
- (b) The case k=2 is trivial. Let  $k\geq 3$ . If (a,b) is an edge of  $G_{k-1,n}$ , then  $a=(x_1,\ldots,x_{k-1}),\,b=(x_2,\ldots,x_k)$  for some  $1\leq x_i\leq n$  by definition. Let us associate the point  $(x_1,\ldots,x_k)$  of  $G_{k,n}$  with the edge (a,b). It is immediate that this yields an isomorphism between  $L(G_{k-1,n})$  and  $G_{k,n}$ . An Euler trail of  $G_{k-1,n}$  yields a Hamiltonian circuit of  $L(G_{k-1,n})$ .
- 8. Let  $(a_1, \ldots, a_{2^k})$  be a Hamiltonian cycle in  $G_{k,2}$ . Then, by the definition of adjacency,  $a_1 = (x_1, \ldots, x_k)$ ,  $a_2 = (x_2, \ldots, x_k, x_{k+1}), \ldots, a_{2^k} = (x_{2^k}, x_1, \ldots, x_{k-1})$ . Let the numbers  $x_1, \ldots, x_{2^k}$  be associated with the points of the cycle. Then the arcs of length k yield the 01-sequences  $a_1, \ldots, a_{2^k}$ . Since a Hamiltonian cycle contains each point exactly once, the arcs yield each 01-sequence exactly once.
- 9. If the pieces  $L_1, \ldots, L_d$  described in the hint are given, any permutation of them determines an Euler trail, and two of these Euler trails are the same iff the two permutations are the same as cyclic permutations. Thus there are (d-1)! Euler trails obtained from  $L_1, \ldots, L_d$ . This proves that the total number of Euler trails is divisible by (d-1)! which is even because  $d \geq 3$ .
- 10. We cannot get stuck anywhere except at  $x_0$ ; this follows just as in 5.6. Assume now that there are edges of G not on the trail L we have traversed. L uses every edge incident to  $x_0$ ; this is obvious for the edges leaving  $x_0$  and follows.

lows for the incoming edges by  $d^+(x_0) = d^-(x_0)$ . If  $(x, y) \notin L$ , then the (unique) edge of T starting from x does not belong to L either, by (\*). Thus, there are edges of T not on L. Let us consider one whose head z is nearest to  $x_0$  (the distance measured on T). Since L does not use all edges running into z, it returns to z less than  $d^+(z) = d^-(z)$  times, hence by (\*), it does not use the (unique) edge (z, u) of T leaving z. However, u is nearer to  $x_0$  than is z (on T), which is a contradiction.

11. We consider all Euler trails as starting from  $x_0$  along  $e_0$ . Let L be an Euler trail, and T the set of those edges (x, y)  $(x \neq x_0)$ , which satisfy

(\*\*) every edge starting from x is used earlier than (x, y).

Then T is a spanning arborescence. For we first observe that there is exactly one edge of T starting from every  $x \neq x_0$ . Moreover, T is acyclic; for if

$$C = (x_1, e_1, x_2, e_2, \dots, e_n, x_1)$$

is a cycle in T, then  $e_2$  is the last edge incident with  $x_2$  on L and hence  $e_1$  anticipates  $e_2$ . Similarly,  $e_2$  anticipates  $e_3, \ldots, e_n$  anticipates  $e_1$ , a contradiction. The definition of T assures that L arises from T by the construction of

the preceding problem.

If T is given, then we can characterize any Euler trail obtained from it by specifying an ordering of edges of G - E(T), starting from x, for any  $x \neq x_0$ , and an ordering of the edges  $\neq e_0$  starting from  $x_0$ ; the orderings indicate in which order the Euler trail has to use the edges. Thus there are exactly  $(d_0 - 1)! \dots (d_{n-1} - 1)!$  such Euler trails. Hence, the number of Euler trails is  $(d_0 - 1)! \dots (d_{n-1} - 1)! \times (\text{number of spanning arborescences rooted at } x_0)$ . [Aarden e-Ehrenfest, de Bruijn; see B].

12. We cannot get stuck at  $x \neq x_0$ ; this follows just as in 5.6.

 $x_0$  is a "good" point. For if there is an edge  $(x_0, y)$  we have not passed from  $x_0$  to y, then we are not stuck; but if every edge  $(x_0, y)$  has been used in this direction, then we left  $x_0$   $d(x_0)$  times, hence we entered it  $d(x_0)$  times, i.e. all edges have been used to enter  $x_0$ .

Assume we meet a "bad" point on our walk and let x be the first one. We enter x on an edge (y, x) and this being the first time we meet x, this will be the edge marked. x is "bad" means we enter x less than d(x) times and hence, by (\*\*) we do not use (x, y) to leave x. However, this contradicts the fact that

y is "good".

Thus, each point we meet is a "good" point. By the definition of "good" points, their neighbors are reached during the walk and are, consequently, "good" points as well. Thus, the good points form a connected component of G. Since G is connected, every point is "good". This completes the proof. [Tarry algorithm; see B].

13. To show the statement of the hint, let  $C_1$  be a circuit of G (G is not a forest as a forest which has an edge has a point of degree 1). Let  $G_1$  be the graph obtained by removing the isolated points of  $G - E(C_1)$ . Then  $G_1$  has even degrees (each degree is reduced by 2 or 0), thus

$$G_1 = C_2 \cup \ldots \cup C_k$$

by induction, and

$$G = C_1 \cup \ldots \cup C_k$$
.

Now assume that G is a graph with even degrees. Then

$$G = C_1 \cup \ldots \cup C_k$$

(we may forget about isolated points). Orienting each  $C_i$  cyclically, we get a desired orientation of G.

- 14. (a) If G has an Euler trail it is, obviously, connected and has even degrees. Conversely, if it is connected and has even degrees, then it can be oriented in such a way that each point has equal indegree and outdegree, by 5.13. The resulting graph has an Euler trail by 5.6. This Euler trail gives an Euler trail of G.
- (b) If G' is the graph obtained by adding a new point and joining it to all points of G of odd degree, then G' has even degrees and is connected, thus it has an Euler trail. Removing the new point, this trail decomposes into kedge-disjoint trails covering G.

15. We may assume G has all degrees at least 4.

Let  $x \in V(G)$  and let  $A_1, \ldots, A_{2d}$  be short beginning segments of the edges incident with x, in this cyclic order. We may assume that if  $G = (A_1 \cup \ldots \cup A_{2d})$ is disconnected (this is the case, e.g. if there is a loop at x), then  $A_1$ ,  $A_2$ go to separate components of it. Remove  $A_1 \cup A_2$  but connect their endpoints different from x by a new arc A (Fig. 30). The resulting map G' is connected:




Fig. 30

this follows from the observation that each component of  $G - (A_1 \cup \ldots)$ ...  $\bigcup A_{2d}$ ) is connected to x by at least two  $A_i$ 's because G is Eulerian. Also, G' has fewer edges than G. Thus by induction, G' has an appropriate Euler trail. Replacing A by  $A_1 \cup A_2$  again, we obtain an appropriate Euler trail of G.

16. The statement of the hint is easily verified. As for the assertion of the problem, it is trivial for a tree  $(2^{(n-1)-n+1} = 1$ , in accordance with the fact that the only "good" subgraph of a tree G is  $(V(G), \emptyset)$ ). Suppose G is connected and has M "good" subgraphs. Those "good"

subgraphs of G + e which do not contain e are exactly the "good" subgraphs

of G and thus their number is M. All we have to do is to show that the number of "good" subgraphs containing e is the same. Let C be a circuit of G+e containing e (which exists because G is connected). Then  $G_1$  is a good subgraph of G+e containing e iff  $(V(G), E(G_1) \triangle E(C))$  is a good subgraph not containing e. This correspondence is one-to-one and proves the assertion.

17. (a) We may restrict ourselves to simple graphs as the removal of two parallel edges does not alter the assertion.

If every point has even degree, we take  $V_1 = V(G)$ ,  $V_2 = \emptyset$ . Suppose  $\alpha$  is

a point of odd degree. Let S be the set of its neighbors. Define  $G_1$  by

$$\begin{split} V(G_1) &= V(G) - \{a\}, \\ (x,y) &\in E(G_1) &\quad \text{iff} \begin{cases} (x\ y) \notin E(G) & \text{if} \quad x,y \in S, \\ (x\ y) \in E(G) & \text{otherwise.} \end{cases} \end{split}$$

By induction on n, we may assume  $V(G_1) = W_1 \cup W_2$ , where  $W_1$  and  $W_2$  span subgraphs of  $G_1$  with even degrees  $G_1$ . Since

$$|S \cap W_1| + |S \cap W_2| = |S| \equiv 1 \pmod{2},$$

we may assume  $|S \cap W_1|$  is even and  $|S \cap W_2|$  is odd. Set

$$V_1 = W_1 \cup \{a\}, \quad V_2 = W_2.$$

Then  $V_1$ ,  $V_2$  span subgraphs of G with even degrees. First, let  $x \in V_1$ . If  $x \notin S$ , then its degree is even in  $G[V_1]$  obviously. Let  $x \in S$ . Then

$$\begin{split} d_{G[V_1]}(x) &= d_{G_1[W_1]}(x) - d_{G_1[W_1 \cap S]}(x) + d_{G[W_1 \cap S]}(x) + 1 = \\ &= d_{G_1[W_1]}(x) - d_{G_1[W_1 \cap S]}(x) + (|W_1 \cap S| - 1 - d_{G_1[W_1 \cap S]}(x)) + 1 \\ &= d_{G_1[W_1]}(x) - 2d_{G_1[W_1 \cap S]}(x) + |W_1 \cap S|, \end{split}$$

and here each term is even. It follows similarly for  $x \in V_2$  that it has even degree in  $G[V_2]$ .

- (b) Let v be a new point, joined to all points of V(G), and let  $G_1$  be the resulting graph. By the preceding exercise,  $V(G_1) = U_1 \cup U_2$ , where  $U_1, U_2$  span subgraphs with even degrees. Now if  $v \in U_2$  (say), then  $V_1 = U_1, V_2 = U_2 \{v\}$  is a desired partition of V(G) [T. Gallai, unpublished; W.K. Chen,  $SIAM\ J.\ Appl.\ Math.\ 20\ (1971)\ 526-529$ ; the proof given here is due to L. Pósa].
- 18. (a) Let S be the set of r-element matchings of the complete graph on V(G). Let  $e_1, \ldots, e_m$  be the edges of  $\overline{G}$  and  $A_i$  the set of those elements of S which contain  $e_i$ . We want to determine

$$m_r(G) = \left| S - \bigcup_{i=1}^m A_i \right|.$$

By the sieve formula (2.2),

$$m_r(G) = |S| - \sum_{1 \le i \le m} |A_i| + \sum_{1 \le i < j \le m} |A_i \cap A_j| - \dots +$$
  
  $+ (-1)^r \sum_{1 \le i_1 < \dots < i_r \le m} |A_{i_1} \cap \dots \cap A_{i_r}|;$ 

here  $|A_{i_1} \cap \ldots \cap A_{i_{\nu}}|$  is the number of those elements of S which contain  $e_{i_1}, \ldots, e_{i_{\nu}}$ . This is non-zero only if  $e_{i_1}, \ldots, e_{i_{\nu}}$  are independent and then it is  $\binom{n-2\nu}{2r-2\nu}$   $(2r-2\nu-1)!!$ . Hence,

$$m_r(G) = \sum_{\nu=0}^r (-1)^{\nu} m_{\nu}(\overline{G}) \binom{n-2\nu}{2r-2\nu} (2r-2\nu-1)!! \qquad ((-1)!!=1).$$

(b) Let n = 2k. By (a),

$$m_k(G) = \sum_{\nu=0}^k (-1)^{\nu} {2k-2\nu \choose 2k-2\nu} (2k-2\nu-1) ! ! m_{\nu}(\bar{G}) \equiv \sum_{\nu=0}^k m_{\nu}(\bar{G}) \pmod{2},$$

thus

$$m_k(G) \neq 0$$
.

- (c) Let A be the adjacency matrix of G and consider det A. Each expansion term here which is non-zero and symmetric with respect to the main diagonal corresponds to a 1-factor and vice versa. The other expansion terms correspond to each other in pairs under reflexion in the main diagonal. Thus the number of 1-factors of G has the same parity as det A, and is even iff det A vanishes over GF(2). This occurs iff the rows of A are linearly dependent over GF(2), i.e. there are elements  $g_1, \ldots, g_n$  of GF(2) such that not all of them are 0 and multiplying the rows of A in order by  $g_1, \ldots, g_n$ , the sum of rows will be 0. Let S be the set of points for which the corresponding  $g_i$  is 1, then S has the property that each point is adjacent to an even number of elements of it. Conversely, each such set S yields appropriate coefficients  $g_1, \ldots, g_n$ .
- 19. Let S be the set of all Hamiltonian paths of the complete digraph on V(G) and let  $e_1, \ldots, e_m$  be the edges of  $\overline{G}$ . Let  $A_i$  denote the set of elements of S containing  $e_i$ , then we have

$$h(G) = \sum_{\nu=0}^{n-1} (-1)^{\nu} \sum_{1 \leq i_1 < \dots < i\nu \leq m} |A_{i_1} \cap \dots \cap A_{i\nu}|.$$

(the term with v=0 is |S|). Now,  $|A_{i_1}\cap\ldots\cap A_{i_{\nu}}|$  is the number of Hamiltonian paths of the complete digraph containing  $e_{i_1},\ldots,e_{i_{\nu}}$ . This is 0 unless  $e_{i_1},\ldots,e_{i_{\nu}}$  form disjoint paths; in the letter case it is (n-v)!, since the graph  $(V(G);\{e_{i_1},\ldots,e_{i_{\nu}}\})$  has n-v components and any Hamiltonian path through  $e_{i_1},\ldots,e_{i_{\nu}}$  defines an ordering of these components and conversely. Hence,

$$|A_{i_1}\cap\ldots\cap A_{i_{\pmb{v}}}|$$
 is  $\left\{egin{array}{ll} 1 & ext{if } \ \pmb{v}=n-1 \ ext{and} \ e_{i_1},\ldots,e_{i_{\pmb{v}}} \ ext{form a} \ ext{Hamiltonian path,} \end{array}
ight.$  even otherwise.

This proves that  $h(G) \equiv h(\overline{G}) \pmod{2}$ .

If G is an undirected graph, then, defining S,  $e_i$ ,  $A_i$  similarly, we have the same formula as (\*), but now we have

$$|A_{i_1}\cap\ldots\cap A_{i_{m{
u}}}|=rac{(n-{m{
u}})\,!}{2}\,2^{\mu},$$

if  $e_{i_1}, \ldots, e_{i_{\nu}}$  form  $\mu$  vertex-disjoint proper paths. If  $n \geq 4$ , then this is also even except when  $\nu = n - 1$  and  $e_{i_1}, \ldots, e_{i_{\nu}}$  form a Hamiltonian path. Hence, we can conclude as above. [T. Szele; see B].

20. If the statement of the hint is true, the assertion of the problem follows easily: by reversing edges we get a transitive tournament, which has one Hamiltonian path. As the parity remained the same, the original tournament had an odd number of Hamiltonian paths.

Thus, it suffices to show that, if T is a tournament, and T' is the tourna-

ment obtained from it by reversing an edge E, we have

$$h(T') \equiv h(T) \tag{mod 2}.$$

Let  $G_1$  and  $G_2$  be the digraphs obtained from T by removing e and adding the inverse of e, respectively. Then a simple computation shows that

$$h(G_1) + h(G_2) = h(T) + h(T').$$

Moreover,  $\bar{G}_1$  arises from  $G_2$  simply by reversing all edges, whence

$$h(\bar{G}_1) = h(G_2).$$

Finally,

$$h(G_1) \equiv h(\overline{G}_1) \tag{mod 2}$$

by 5.19. Hence

$$h(T) + h(T') = h(G_1) + h(G_2) = h(G_1) + h(\overline{G}_1) \equiv 0 \pmod{2},$$

which proves the assertion [L. Rédei; see B].

21. If  $F_1$ ,  $F_2$  are disjoint 1-factors of G, then  $E(G)-F_1-F_2$  is a 1-factor. Conversely, if we have a partition of E(G) into three 1-factors, then we can choose as  $F_1$  the 1-factor containing e and any one of the two others as  $F_2$ . Hence, the number m of pairs  $(F_1, F_2)$  of 1-factors with  $F_1 \cap F_2 = \emptyset$ ,  $e \in F_1$ ,  $e \notin F_2$  is twice the number of partitions of E(G) into 1-factors. Thus, m is even. Now consider  $F_1 \cup F_2$ ; this consists of even circuits covering V(G). If, conversely,  $H \subseteq E(G)$  is the set of edges of a system of k disjoint even circuits covering V(G) and containing e, then H can be decomposed into  $H = F_1 \cup F_2$  ( $F_1$ ,  $F_2$  1-factors,  $e \in F_1$ ,  $e \notin F_2$ ) in exactly  $2^{k-1}$  ways. Thus, if  $m_k$  denotes the number of systems of k disjoint even circuits covering V(G) and containing e, then

$$m = m_1 + 2m_2 + \ldots + 2^{k-1}m_k + \ldots$$

Note here that  $m_1$  is the number of Hamiltonian circuits through e and therefore

$$m_1 \equiv m \equiv 0 \pmod{2}$$
.

[C.A.B. Smith; see B.]

22. We prove the assertion by induction on |V(G)|. If |V(G)| = 4, it is

obvious. Also, we may assume that G is connected.

If there is a double edge in G, then all Hamiltonian circuits must use one of the two and hence the Hamiltonian circuits occur in pairs, differing only in which of the two parallel edges they use. Thus the total number of Hamiltonian circuits is even again.




Fig. 31

So suppose G is simple. Let (x, y),  $(x, u_1)$ ,  $(x, u_2)$ ,  $(y, v_1)$ ,  $(y, v_2) \in E(G)$ . (see Fig. 14). Remove x and y and join  $u_1$  to  $v_1$ ,  $u_2$  to  $v_2$  to get G'; also, join

 $u_1$  to  $v_2$  and  $u_2$  to  $v_1$  to get G''.


The Hamiltonian circuits of G may or may not contain (x, y). There are four kinds of Hamiltonian circuits through (x, y), containing  $(u_1xyv_1)$ ,  $(u_1xyv_2)$ ,  $(u_2xyv_1)$ ,  $(u_2xyv_2)$ , respectively (Fig. 31). Let their numbers be  $h_1$ ,  $h_2$ ,  $h_3$ ,  $h_4$ . Also, those Hamiltonian circuits not containing (x, y) may go like  $(\dots u_1 x u_2 \dots$  $\dots v_1 y v_2 \dots$ ) or  $(\dots u_1 x u_2 \dots v_2 y v_1 \dots)$ . Let the numbers of such Hamil-

tonian circuits be  $h_5$  and  $h_6$ .

The  $h_1$  Hamiltonian circuits defined above are in a one-to-one correspondence with those Hamiltonian circuits of G' containing  $(u_1, v_1)$  but not  $(u_2, v_2)$ . Similarly,  $h_2$ ,  $h_3$ ,  $h_4$  are equal to the numbers of Hamiltonian circuits of G''and G' containing  $(u_1, v_2)$ ,  $(u_2, v_1)$ ,  $(u_2, v_2)$  but not the other new edge. The  $h_5$ Hamiltonian circuits of G correspond to those Hamiltonian circuits of G' containing both  $(u_1, v_1)$  and  $(u_2, v_2)$  and going through them like  $(\dots u_1 v_1 \dots$  $\dots u_2 v_2 \dots$ ). Similarly, the  $h_6$  Hamiltonian circuits correspond to those Hamiltonian circuits of G'' going through  $(u_1, v_2)$  and  $(u_2, v_1)$  like  $(\dots u_1 v_2 \dots$  $\dots u_2v_1\dots$ ).

The Hamiltonian circuits of G' not considered so far are those going through  $(u_1, v_1)$  and  $(u_2, v_2)$  like  $(\dots u_1 v_1 \dots v_2 u_2 \dots)$  and those not containing the

new edges. Let  $h_7$  and  $h_8$  be the numbers of such Hamiltonian circuits.



In G'', we have not considered so far the Hamiltonian circuits of form  $(\ldots u_1v_2\ldots v_1u_2\ldots)$  and those not containing any of the new edges. The numbers of such Hamiltonian circuits are, as is easily seen,  $h_7$  and  $h_8$ , respectively (Fig. 32, see p. 257).

Now the number of Hamiltonian circuits of G is  $h_1 + h_2 + h_3 + h_4 + h_5 + h_6 \equiv (h_1 + h_4 + h_5 + h_7 + h_8) + (h_2 + h_3 + h_6 + h_7 + h_8) \equiv 0 \pmod{2}$ , since here in the brackets we have the numbers of Hamiltonian circuits of

G' and G'', respectively [J. Bosák; see B].

23. To show the statement of the hint, let  $F^*$  be the graph on  $V(G^*)$  formed by those edges of  $G^*$  which are not crossed by the edges of F.  $F^*$  does not contain a cycle, for then the points of F inside this cycle would not be connected to those outside. Moreover,  $F^*$  is connected. For if  $F^* = F_1 \cup F_2$ ,  $F_1 \cap F_2 = \emptyset$ , then let U be the union of faces of G with "capital" in  $F^*$ . Then U is not the whole plane (no point of  $F_2$  belongs to it) and therefore, its boundary B is non-empty. Now B consists of certain edges of F, each having a face of U on one side and a face not in U on the other. Therefore, there is no point of degree 1 on the boundary, i.e. it contains a circuit. This is a contradiction.

Hence  $F \to F^*$  gives a one-to-one correspondence between the spanning

trees of G and  $G^*$ . This proves the assertion of the problem.

Remark: Our argument uses some facts of plane topology like the Jordan curve theorem (each simple closed curve divides the plane into two parts) and other similar statements. As our goal is to illuminate the combinatorial content of the problems we assume these facts without proof.

**24.** Consider a spanning tree F of G and  $F^*$  as in 5.24. Then

$$|E(F)| = |V(G)| - 1,$$
 $|E(F^*)| = |V(G^*)| - 1$ 

and by the definition of  $F^*$ ,

 $|E(F)| + |E(F^*)| = |E(G)|.$ 

Hence

$$|V(G^*)| = |E(G)| + 2 - |V(G)|.$$

25. Since each face has at least three [four] edges on its boundary and each edge is on the boundary of exactly two faces, we get by Euler's formula


$$2m \ge 3(m-n+2)$$
  $[2m \ge 4(m-n+2)]$ 

or, equivalently,

$$m \le 3n - 6 \qquad [m \le 2n - 4].$$


**26.** We may assume there are no loops. We use induction on |E(G)|.

Let F be a face bounded by the edges  $e_1, \ldots, e_m$ . We claim that every point x is incident with an even number of them. For let k "corners" in x belong to F, then two of these "corners" cannot have a common edge e because this edge would be a cut-edge (see Fig. 33) and hence a component of G - e would contain exactly one point of odd degree. Thus x is incident with 2k edges of the boundary of F.



Now remove the edges  $e_1, \ldots, e_m$ . The remaining map G' has even degrees, thus its faces can be 2-colored. The faces of G' are the union of F and its neighbors, as well as all the other faces of G. Interchanging the color of F but keeping it everywhere else, we get a 2-coloration of the faces of G.

- 27. First, 2-color the faces according to 5.26 with red and blue, say. Then, orient each edge so that the red face incident with a given edge e is on its left-hand side. This orientation satisfies the requirement.
- 28. (a) Suppose there were a planar map with even degrees, and all faces triangular except one which is pentagonal. 2-color the faces (5.26) with red and blue, and suppose the pentagon is red (say). Count the number of edges. Each blue face has 3 edges on its boundary, and this counts each edge exactly once; hence the number of edges is divisible by 3. On the other hand, the red faces have altogether 3k + 5 edges on their boundaries (k is the number of triangular red faces), which should give the same total number of edges, a contradiction.
- (b) The graph G' constructed in the hint has even degrees at every point except possibly z, but then, by 5.1a, z has an even degree as well. Let  $a_1 + 1$ ,



 $a_2+1,\ldots,a_{2s}+1$  be the sizes of faces incident with z in a cyclic order. 2-color the faces (Fig. 34 see p. 259). The color containing the face of size  $a_1+1$  has  $(a_1+1)+(a_3+1)+\ldots+(a_{2s-1}+1)+3k$  edges on its boundary, the other one has  $(a_2+1)+\ldots+(a_{2s}+1)+3N$  edges. In both cases we get the total number of edges of G. Hence,

$$a_1 + a_3 + \ldots + a_{2s-1} \equiv a_2 + a_4 + \ldots + a_{2s} \pmod{3}$$
.

Moreover, we have  $\sum_{i=1}^{2s} (a_i - 1) = 5$ . It is easy to verify that the only solution of

these equations is

$$a_1 = 2, \quad a_2 = 5$$

(or conversely). Thus, exactly two, neighboring points of the pentagon have odd degrees [T. Gallai].

29. Let the three colors be red, blue, green. Each triangle whose points get different colors has one red-blue edge; any other triangle has 0 or two. Moreover, every red-blue edge is counted twice, in both triangles adjacent to it. So this adds up to an even number, i.e. the number of triangles with 3 different colors is even.

Remark: The statement is a special case of Sperner's lemma in algebraic topology, which is equivalent to a similar statement for n-dimensional triangulations [see e.g. L. S. Pontryagin, Grundzüge der Kombinatorischen Topologie, Berlin, 1956, p. 73].

**30.** Consider the coloration defined in the hint. If there is a face with points x, y, z, of color red, blue, green (in this order) and  $z \in V$ , (say), then z plus the (x, a)-path in  $V_1$  yields a (z, a)-path in  $V_1$ , whence z should have been colored red.

Now if there is no (a, c)-path in  $V_1$  and no (b, d)-path in  $V_2$ , then c, d are green. Hence, if we add the edge (a, c), we get a triangulation with only the face abc meeting three colors. This contradicts the preceding exercise.

- **31.** (a) True; in fact, A is regular iff det  $A \neq 0$ , a fact whose standard proof uses considerations valid in any field.
  - (b) True; for

$$\det A^T A = \det A^T \det A = (\det A)^2 \neq 0.$$

(c) False; for we may have a vector  $\mathbf{u} \neq \mathbf{0}$  with  $\mathbf{u}^T \mathbf{u} = 0$  (e.g. if F = GF(2),  $\mathbf{u} = \mathbf{e}_1 + \mathbf{e}_2$  or F is the complex field and  $\mathbf{u} = (1, i)^T$ ), and then the transformation A defined by

$$A\mathbf{e}_1 = \mathbf{u}, A\mathbf{e}_2 = \ldots = A\mathbf{e}_n = \mathbf{0}$$
 (for some basis  $\mathbf{e}_1, \ldots, \mathbf{e}_n$ )

satisfies  $A \neq 0$ ,  $\mathbf{y}A^T A \mathbf{x} = (A\mathbf{y})_{ij}^{T} A \mathbf{x} = 0$  for any  $\mathbf{x}$ ,  $\mathbf{y}$ , thus  $A^T A = 0$ .

- (d) False; if  $\mathbf{u}^T \mathbf{u} = 0$ ,  $\mathbf{u} \neq \mathbf{0}$ , then let  $M = \langle \mathbf{u} \rangle$  (the subspace generated by  $\mathbf{u}$ ); then  $M \subseteq M^{\perp}$ .
- (e) False; for the preceding M, we have  $\langle M, M^{\perp} \rangle = M^{\perp} \neq V$ , as, e.g. one of  $\mathbf{e}_1, \ldots, \mathbf{e}_n$  is surely non-orthogonal to M.

(f) True; let A be as in the hint. Then

$$\mathbf{x} \in M^{\perp} \Leftrightarrow \mathbf{x}^{T} \mathbf{v}_{i} = 0 \quad (i = 1, \dots, k) \Leftrightarrow \mathbf{x}^{T} A \mathbf{e}_{i} = 0 \quad (i = 1, \dots, k),$$

$$\Leftrightarrow \mathbf{e}_{i}^{T} A^{T} \mathbf{x} = 0 \quad (i = 1, \dots, k) \Leftrightarrow A^{T} \mathbf{x} \in \langle \mathbf{e}_{k+1}, \dots, \mathbf{e}_{n} \rangle,$$

$$\Leftrightarrow \mathbf{x} \in (A^{T})^{-1} \langle \mathbf{e}_{k+1}, \dots, \mathbf{e}_{n} \rangle.$$

Hence

$$M^{\perp} = (A^T)^{-1} \langle \mathbf{e}_{k+1}, \ldots, \mathbf{e}_n \rangle,$$

which shows that dim  $M^{\perp} = n - k$ .

(g) True; for let  $\mathbf{u} \in M$ , then  $\mathbf{u}$  is orthogonal to every element of  $M^{\perp}$ , i.e.  $\mathbf{u} \in (M^{\perp})^{\perp}$ . Thus

$$M\subseteq (M^\perp)^\perp.$$

On the other hand,

$$\dim (M^{\perp})^{\perp} = n - \dim M^{\perp} = n - (n - \dim M) = \dim M,$$

which proves the assertion.

**32.**  $\mathbf{x} \in \langle M, M^{\perp} \rangle^{\perp}$  iff  $\mathbf{x}$  is orthogonal to both M and  $M^{\perp}$ , i.e.  $\mathbf{x} \in M^{\perp} \cap (M^{\perp})^{\perp} = M \cap M^{\perp}$ . Hence

$$\langle M, M^{\perp} \rangle = (M \cap M^{\perp})^{\perp}.$$

Now, let  $\mathbf{u} \in M \cap M^{\perp}$ . Then  $\mathbf{u} \in M$  and  $\mathbf{u} \in M^{\perp}$ , thus  $\mathbf{u}^T \mathbf{u} = 0$ . This means, however, that the number of 1's in  $\mathbf{u}$  is even, which is equivalent to  $\mathbf{u}^T \mathbf{j} = 0$ . Thus  $\mathbf{j} \in (M \cap M^{\perp})^{\perp} = \langle M, M^{\perp} \rangle$ . [T. Gallai, W. K. Chen, loc. cit. on p. 254.]

**33.** (a) We show  $U_G$  consists of the cuts. Cuts form a subspace; for the sum of cuts determined by the sets  $S_1$  and  $S_2$ , respectively, is the cut determined by  $S_1 \triangle S_2$ . Each star is a cut. Conversely, the cut determined by  $S_1 \triangle S_2$  is the sum of stars determined by the points of  $S_1 \triangle S_2$ .

The star of x is orthogonal to a set A of edges iff the subgraph determined by A has even degree at x. Hence,  $W_G$  consists of the sets of edges of subgraphs with even degrees. By the solution of 5.13,  $W_G$  is generated by the circuits

of G.

If all circuits are even, then  $\mathbf{j} = (1, \dots, 1)$  is orthogonal to every circuit, hence  $\mathbf{j} \in (U_G^{\perp})^{\perp} = U_G$ . This says by (a) that G is bipartite. Hence 5.3 follows.

To show 5.16, we have to determine dim  $W_G$ . It is enough to determine dim  $U_G$  by 5.21f. Let  $A_1, \ldots, A_n$  be all the stars. Since

$$A_n = A_1 + \ldots + A_{n-1},$$

we have  $U_G = \langle A_1, \ldots, A_{n-1} \rangle$ . We show that  $A_1, \ldots, A_{n-1}$  are linearly independent. Any linear dependence between them would be of the form

$$A_{i_1}+\ldots+A_{i_k}=0,$$

as we are working over GF(2). However,  $A_{i_1} + \ldots + A_{i_k}$  is a cut determined by a non-empty proper subset of V(G), which is non-empty because G is connected.

Thus dim  $U_G = n - 1$ , dim  $W_G = m - n + 1$ , q.e.d. 5.17 follows from 5.32 immediately.

(b) The assertion is trivial for disconnected graphs, so suppose G is connected. The number of decompositions, i.e. the number of solutions of  $\mathbf{a} + \mathbf{b} = \mathbf{j}$ ,  $\mathbf{a} \in U_G$ ,  $\mathbf{b} \in W_G$ , is clearly equal to  $|U_G \cap W_G|$ . This is the same as the number of vectors  $\mathbf{u} \in U_G$  orthogonal to every element of  $U_G$ .

Let  $A_G$  be the point-edge incidence matrix of G and  $A_0$  the matrix arising from it by cancelling a row. Then the range of  $A_0^T$  is  $U_G$ . We also know  $A_0^T$  is one-to-one. Thus, we are interested in the number of vectors  $\mathbf{x}$  such that

$$(A_0 \mathbf{x})^T (A_0 \mathbf{y}) = \mathbf{x}^T A_0 A_0^T \mathbf{y} = 0$$

for every y. But this holds for every y iff  $A_0 A_0^T \mathbf{x} = 0$ . This equation has a unique solution iff det  $A_0 A_0^T \neq 0 \pmod{2}$ . By 4.9 this means the number of spanning trees is odd.

**34.** Let  $C_1, \ldots, C_{f-1}$  be the finite faces, considered as circuits of G, i.e. elements of  $W_G$ . We show that every circuit of G is the sum of some  $C_i$ 's. Let C be a circuit of G. Each face  $C_i$  lies either inside or outside C. Let, say,  $C_1, \ldots, C_r$  lie inside C. Then

$$C = C_1 + \ldots + C_r;$$

in fact, if e is an edge of C, then exactly one of the two faces incident with e lies inside C, hence  $e \in E(C_1 + \ldots + C_r)$ . If e is inside (outside) C, then both (none) of these two faces are inside C, hence e does not occur in  $C_1 + \ldots + C_r$ . On the other hand, if  $C_1, \ldots, C_{f-1}$  were dependent, say

$$C_1+\ldots+C_r=0,$$

then draw a continuous line from an inner point x of  $C_1$  to the infinity, avoiding the vertices. This line leaves the union of faces  $C_1, \ldots, C_r$  at a point which belongs to an edge e of G. Then exactly one of the two faces adjacent to e belongs to  $C_1, \ldots, C_r$ , hence

$$e \in C_1 + \ldots + C_r$$

a contradiction.

Remark: Since the dimension of  $W_G$  is m-n+1 by the previous problem, we get

$$m = n + 1 = f - 1$$
,

i.e. we also obtain Euler's formula at least for 2-connected graphs. Conversely, the use of Euler's formula would have made one half of the proof superfluous.

**35.** (a) Let  $e \in \sum_{i \in I \cup J} C_i$ , and say  $e \in C_\mu$ ,  $\mu \in I \cup J$ . We claim  $\mu \in I$ . This is clear if  $\mu \notin J$ . Suppose  $\mu \in J$ , then since no other  $C_\nu$ ,  $\nu \in I \cup J$  contains e, we have

$$e \in \sum_{i \in J} C_i \subseteq \bigcup_{i \in I} C_i,$$

whence  $\mu \in I$ . Thus,  $e \in \sum_{i \in I} C_i = K$ , i.e.  $\sum_{i \in I \cup J} C_i \subseteq K$ . Since K is a circuit and  $\sum_{i \in I \cup J} C_i \neq \emptyset$ , we have

If we have 
$$K = \sum_{i \in I \cup J} C_i$$
, while  $I = \emptyset$  we have the first  $I = \emptyset$ 

Since K has a unique decomposition into the sum of  $C_i$ 's, we get

$$I \cup J = I$$
,  $J \subseteq I$ .

(b) If  $C_1, \ldots, C_f$  are the only circuits, then it is easy to see that they are precisely the blocks of G (not counting cutting edges).

Let  $K \neq C_1, \ldots, C_f$  be a circuit such that its representation as the sum of  $C_i$ 's has a minimum number of terms. Let, say,

$$K = C_1 + \ldots + C_r$$
  $(r \ge 2), I = \{1, \ldots, r\}$ 

and, e.g.  $C_1 \cap C_2 \neq \emptyset$ . Then  $C_1 + C_2$  has even degrees and therefore

$$(1) C_1 + C_2 = K_1 + \ldots + K_s,$$

where  $K_1, \ldots, K_s$  are edge-disjoint circuits. At least one of them, say  $K_1$ , must be different from  $C_1, \ldots, C_f$ ; otherwise (1) would show that  $C_1, \ldots, C_f$  are linearly dependent. Let

at k to but a complete of  $K_1 = \sum_{i \in J} C_i$  .

By (a),  $J \subseteq I$  and hence by the minimality of I, we have J = I. Thus  $K = K_1$  and so

$$(2) K \subseteq C_1 + C_2.$$

We claim that  $K=C_1+C_2$ . Suppose there is an edge of  $C_1$  not in  $K\cup C_2$  (say). This edge belongs to a  $C_k$ ,  $3\leq k\leq r$ . (2) holds for  $C_k$  as well, i.e.

$$(3) K \subseteq C_1 + C_k.$$

But (2) and (3) imply that an edge of K not in  $C_1$  must belong to both of  $C_2$  and  $C_k$  and thus it cannot occur in  $\sum_{i=1}^{r} C_i$ , a contradiction. Thus  $C_1 + C_2$  is a circuit.

(c) We use induction on the number of edges. Let, say,  $C_1 + C_2 = C$  be a circuit. Remove the edges common to  $C_1$  and  $C_2$  and the isolated points that possibly arise. Let G' be the resulting graph.

Consider the system  $C, C_3, \ldots, C_f$ . It is obvious from the construction that each edge of G' is contained in at most two of them. Moreover, if A is a sub-

graph of G' with even degrees, then

$$A = \sum_{i \in I} C_i$$

 $(I \subseteq \{1,\ldots,f\})$ , and here either both  $C_1$  and  $C_2$  occur or neither of them, whence A is a linear combination of  $C,C_3,\ldots,C_f$ . Moreover,  $C,C_3,\ldots,C_f$  are, obviously, linearly independent. Hence, by the induction hypothesis, G' can be embedded in the plane so that  $C,C_3,\ldots,C_f$  are boundaries of faces. To get G we have to put back  $C_1\cap C_2$ , which is a path joining two points of C and thus, it can be done and  $C_1,C_2$  will become the new boundaries.

(d) It is easy to see that a graph is planar iff each block of it is planar. A similar assertion holds for the property that  $W_G$  has a basis such that each edge belongs to at most two elements of it. Therefore to prove the equivalence of these two properties, we may restrict ourselves to 2-connected graphs.

For these, the necessity of MacLane's condition of planarity follows by 5.34. We remark that we may require that the elements of the basis should be circuits. For suppose there exists a basis  $A_1, \ldots, A_f$  of  $W_G$  such that each edge belongs to at most two of them.  $A_1$  is the union od egde-disjoint circuits. One of these must clearly be linearly independent of  $A_2, \ldots, A_n$ . Replacing  $A_1$  with this circuit we obtain a basis with the same property. Going on similarly we can replace  $A_2, \ldots, A_f$  by circuits.

Now the sufficiency of MacLane's condition follows by (c). [S. MacLane;

see W.]

**36.** Suppose G is a planar map and let  $G^*$  be its dual graph. Then if  $\varphi$  associates with each  $e \in E(G)$  the edge of  $G^*$  crossing it, will satisfy the requirements of the statement by 5.23.

Conversely, suppose  $G^*$  and  $\varphi$  exist, we show G is planar.

First, we prove that the edges of a star in  $G^*$  correspond to the edges of an element of  $W_G$ . For let  $X \subseteq V(G)$  and suppose  $\varphi(X)$  is the star of a point  $x \in V(G^*)$ . Let  $B_1, \ldots, B_s$  be the branches of  $G^*$  relative to x and let  $A_i$  be the set of  $(x, B_i)$ -edges. Then  $A_i$  is a minimal set which meets every spanning tree of  $G^*$ . Therefore,  $\varphi^{-1}(A_i)$  is a minimal set which is not contained in any spanning tree of  $G^*$ , i.e.  $\varphi^{-1}(A_i)$  is a circuit. Hence

$$X = \sum_{i=1}^{s} \varphi^{-1}(A_i) \in W_{G^{\bullet}}$$


Now, let  $C_1, \ldots, C_{n^*}$  be the elements of  $W_G$  corresponding to the stars of points in  $G^*$ . Then trivially each edge of G is contained in exactly two  $C_i$ 's. Moreover if we set  $f = n^* - 1, C_1, \ldots, C_f$  will form a basis of  $W_G$ . In fact,

$$\dim W_G = e - n + 1 = n^* - 1 = f$$

(from the definition of  $G^*$ ), and  $C_1, \ldots, C_f$  are linearly independent over GF(2), because so are the corresponding stars in  $G^*$ .

Thus G is planar by MacLane's criterion. [H. Whitney; see, e.g. Wi.]

- 37. (a) It is trivial that G is 2-connected. Suppose indirectly that  $G = G_1 \cup G_2$  with  $V(G_1) \cap V(G_2) = \{x, y\}$ ,  $|V(G_i)| \geq 3$ . Let  $P_i$  be an (x, y)-path in  $G_i$  and  $H_i = G_i + P_{3-i}$ . Then  $H_i$  is planar; embed  $H_i$  into the plane so that the path  $P_{3-i}$  lies on the boundary of the unbounded domain (this can be achieved by an inversion). Then identify the two points x and y and delete the paths  $P_i$  (Fig. 35). This results in an embedding of G in the plane, a contradiction.
- (b) Let  $(x_0, \ldots, x_m)$  be a longest path in G.  $x_0$  has degree at least 3, and it is adjacent to no point outside this path because of the maximality of it. Hence it has two neighbors  $x_i$ ,  $x_j$  with 1 < i < j. Then  $(x_0, \ldots, x_j)$  is a circuit with a chord (cf. also 6.35).
- (c) Let (x, y) be a chord of a circuit C and choose C so that embedding G-(x,y) in the plane the number of faces inside C is as large as possible. Observe first that there is no point outside C. In fact, let  $G_0$  be a component of G-V(G) and suppose indirectly that  $G_0$  lies outside C. Since G is 3-connected, there must be three points of C adjacent to  $G_0$  and at least two of these, u and v say, are not separated by x and y. Then replacing the (u, v)-arc of e



not containing x, y by a (u, v)-path through  $G_0$ , we obtain a circuit C' with chord (x, y) and more faces inside.

The same reasoning shows that all chords of C running outside connect inner

points of the two (x, y)-arcs of C.

Now let us consider the bridges of C inside C. Call such a bridge flappable if its endpoints do not separate the endpoints of any outside chord of C. It is clear that we can "flap" all these bridges to the outside of C. Among the remaining bridges there must be one containing inner points of both (x, y)-arcs of C, otherwise x and y could be connected inside C and G would be planar. So there is a bridge G inside G and a chord G outside G such that the endpoints of G separate G from G and G from G and G and G separate G from G and G from G and G and G separate G from G and G and G separate G from G and G from G and G separate G from G and G from G and G separate G from G and G separate G from G and G from G and G separate G from G and G from G and G separate G from G and G separate G from G and G from G and G separate G from G and G separate G from G and G from G from G from G and G from G

- (a) B contains inner points of the (x, a)-arc and (y, v)-arc (or, symmetrically, of (x, v) and (y, a));
- (b) B contains v, an inner point of (x, a) and moreover a point of (y, a) different from a (or any symmetrical situation);
  - (c) B contains x, y, a, v.

Take a path P connecting two of the mentioned endpoints of B. In case b) take a path connecting P to the third one. In case (c) take two paths connect-




FIG. 36




Fig. 37

ing P to the other two endpoints. If these two paths meet each other, then let them have a common initial piece. So we get from case (c) two subcases according to whether the mentioned paths in B form an H or an H or


In case (a), we see a subdivision of  $K_{3,3}$ ; by the minimality of G, there cannot be other edges or any subdividing points, i.e.  $G \simeq K_{3,3}$ . In cases (b) and (c1) the graph properly contains a subdivision of  $K_{3,3}$ ; this is impossible since each proper subgraph of G is supposed to be planar. In case (c2) we see a subdivision of  $K_5$  and hence  $G \simeq K_5$ .

(d) Suppose G is planar. Then, obviously, G cannot contain a subdivision of  $K_5$  or  $K_{3,3}$ . Conversely, suppose G is not planar. Then G contains a minimal non-planar graph  $G_0$ . If we get rid of the points of degree 2 from  $G_0$  (removing them and joining their two neighbors successively), we get another minimal non-planar graph, this time with degrees at least 3. This graph is  $K_5$  or  $K_{3,3}$  by (c), thus  $G_0$  is a subdivision of  $K_5$  or  $K_{3,3}$  [see S II].

38. We use induction on the number of points. If this is at most 3, the assertion is trivial.

First we show that if G is any planar graph we can introduce new edges to turn all faces into triangles without getting parallel edges. For let us draw new edges as long as we can without getting parallel ones. The graph G has no cutpoints; for if  $G = G_1 \cup G_2$  with  $V(G_1 \cap G_2) = \{x\}$ , then take a point  $x_i$  of  $G_i - x$  on the boundary of the face which meets both  $G_1 - x$  and  $G_2 - x$ ;  $x_1, x_2$  could be connected by a further edge.

So G is 2-connected and, therefore, each face is a circuit. Suppose C is the boundary of a face with at least 4 vertices and let a, b, c, d be four consecutive points on C. One of the two edges (a, c), (b, d) must be missing; since they both ought to run outside C and therefore, they ought to cross. Suppose a and c



are non-adjacent; then they can be connected by an edge inside C. Hence all faces of G are triangles.

Thus it suffices to prove the assertion for triangulations. Now we find an edge (x, y) which is contained in two triangles only. For let x be a point which is contained inside some triangle T (any point not on the outermost triangle has this property) and choose x, T so that the number of faces inside T is minimal. Let y be any neighbor of x. Now if (x, y) were an edge of three triangles  $(x, y, z_1)$ ,  $(x, y, z_2)$ ,  $(x, y, z_3)$ , then all these triangles would be properly contained in T and, say,  $(x, y, z_1)$  would contain  $z_3$  inside it, contrary to the minimality of T.

So choose an edge (x, y) such that the only two triangles having (x, y) as an edge are the two triangular faces  $(x, y, z_1)$ ,  $(x, y, z_2)$  incident with it. Contract (x, y) to a point p and remove one edge from both arising pairs of parallel edges. This way we get a new simple triangulation  $G_0$  and by induction hypothesis, there is a triangulation  $G_0$  with straight edges such that the faces

of  $G_0$  and  $G'_0$  correspond to each other.

Now consider the edges of G' corresponding to  $(p, z_1)$  and  $(p, z_2)$ . They split the angle around p into two angles; one of these contains the edges whose pre-images in G are adjacent to x, the other one those whose pre-images are adjacent to y. Therefore, we can "pull x, y apart" and get an appropriate representation of G with straight edges (Fig. 38) [K. Wagner–I. Fáry; see S].

## § 6. Connectivity

1. We use induction on |E(G)|. If |E(G)| = 0, G consists of isolated points and hence, c(G) = |V(G)|. Let  $e \in E(G)$ . Then

$$c(G) \ge c(G-e) - 1,$$

because e connects either two points of the same component of G - e, in which case G - e has the same connectivity classes, or two points in different components, in wich case c(G) drops by 1. By induction,

$$c(G-e)+|E(G-e)|\geq |V(G-e)|=|V(G)|$$
 , hence  $c(G)+|E(G)|\geq c(G-e)-1+|E(G-e)|+1=$   $=c(G-e)+|E(G-e)|\geq |V(G)|.$ 

6.2

2. (a) Let H be a component of the graph  $G^*$  constructed in the hint. Define

$$\widetilde{H} = \bigcup_{(s_i, t_i) \in E(H)} (S_i \cap T_i).$$

Suppose  $(x, y) \in E(G_1 \cup G_2)$ ,  $x \in \widetilde{H}$ . Let  $(x, y) \in E(G_1)$  and  $x \in S_i \cap T_j$  (say). Since  $S_i$  is a component of  $G_1, y \in S_i$ . Let  $y \in T_j$ . Now  $s_i \in V(H)$ ,  $(s_i, t_j) \in E(H)$ , hence  $S_i \cap T_j \subseteq \widetilde{H}$ ,  $y \in \widetilde{H}$ .

Hence,  $\widetilde{H}$  consists of one or more components of  $G_1 \cup G_2$ , whence

$$c(G_1 \cup G_2) \ge c(G^*).$$

If  $S_i \cap T_j \neq \emptyset$ , then it consists of one or more components of  $G_1 \cap G_2$ , hence  $c(G_1 \cap G_2) \geq |E(G^*)|$ .

By the previous exercise,

$$c(G_1 \cup G_2) + c(G_1 \cap G_2) \ge |V(G^*)| = c(G_1) + c(C_2).$$

(b) Set  $V = V(G_1) \cup V(G_2)$ , and add the points of  $V - V(G_i)$  to  $G_i$  as isolated points. Let  $G_i$  be the resulting graph. Then

$$egin{aligned} V(G_1') &= V(G_2') &= V, \ c(G_1') &= c(G_1) + |V| - |V(G_1)|, \ c(G_1' \cap G_2') &= c(G_1 \cap G_2) + |V| - |V(G_1 \cap G_2)|, \ c(G_1' \cup G_2') &= c(G_1 \cup G_2). \end{aligned}$$

By (a),

$$c(G_1' \cup G_2') + c(G_1' \cap G_2') \ge c(G_1') + c(G_2').$$

So

$$\begin{split} c(G_1 \cup G_2) + c(G_1 \cap G_2) &= c(G_1' \cup G_2') + c(G_1' \cap G_2') - \\ &- |V| + |V(G_1 \cap G_2)| \geq c(G_1') + c(G_2') - |V| + |V(G_1 \cap G_2)| = \\ &= c(G_1) + c(G_2) + |V| - |V(G_1)| - |V(G_2)| + |V(G_1 \cap G_2)| = c(G_1) + c(G_2). \end{split}$$

3. Assume indirectly that G is disconnected and let  $G_1$  be a component not containing  $x_n$ . Let  $|G_1|=k$ , and  $x_{i_1},\ldots,x_{i_k}$  its points  $(1 \le i_1 < \ldots < i_k < n)$ . Since the component containing  $x_n$  has at least  $d_n+1$  points, we have  $k \le n-d_n-1$ . Moreover,

$$d_k \leq d_{i_k} \leq k-1,$$

a contradiction [A. Bondy; see B].

**4.** Suppose first that  $G_1$  contains an odd circuit C and that  $G_1$ ,  $G_2$  are connected. To follow the hint, let  $x, y \in V(G_1)$ . If there is an (x, y)-walk in  $G_1$  of length k, then there is one of length k+2 (since we can go on an edge back and forth). Therefore it suffices to show there are both odd and even (x, y)-walks. There is a walk touching C because  $G_1$  is connected; if we add a tour around C to this walk, we get a walk of opposite parity. This proves the statement of the hint. Now let (x, u) and (y, v) be two points of  $G_1 \times G_2$ .  $G_2$  contains a walk (u = x)

 $=u_0,u_1,\ldots,u_k=v$ ). We may assume this walk is very long. Then by the above,  $G_1$  contains a walk  $(x=x_0,x_1,\ldots,x_k=y)$  of the same length. Then  $((x_0,u_0),(x_1,u_1),\ldots(x_k,u_k))$  is a walk in  $G_1\times G_2$  connecting (x,u) to (y,v). Conversely, assume that  $G_1\times G_2$  is connected. Trivially,  $G_1,G_2$  are connected. Suppose both of  $G_1$  and  $G_2$  are bipartite, and let  $V(G_1)=A_1\cup B_1,\ V(G_2)=A_2\cup B_2$  be 2-colorations of them. Then no edge of  $G_1\times G_2$  connects  $(A_1\times A_2)\cup (B_1\times B_2)$  to  $(A_1\times B_2)\cup (B_1\times A_2)$ , a contradiction.

**5.** Suppose  $G = G_1 \cup G_2$ ,  $V(G_1) \cap V(G_2) = \{x\}$ ,  $|V(G_i)| \ge 2$ . Then  $1 \le d_{G_1}(x) \le k-1$ , on the other hand,  $d_{G_1}(y) = k$  for  $y \in V(G_1) - \{x\}$ . Let  $u_1, \ldots, u_r, x; v_1 \ldots v_s$  be the points in the two color classes of  $G_1$ . Then

$$|E(G_1)| = d_{G_1}(u_1) + \ldots + d_{G_1}(u_r) + d_{G_1}(x) = d_{G_1}(v_1) + \ldots + d_{G_1}(v_s)$$

whence

$$k \cdot r + d_{G_1}(x) = k \cdot s, \qquad k \mid d_{G_1}(x),$$

a contradiction.

**6.** (a) Let  $P = (x_0, x_1, \ldots, x_m)$  be a maximum path in G. Suppose  $G - x_0$  is not connected. Let  $G_1$  be a component of  $G - x_0$  which does not contain  $P - x_0$ . Since G is connected, there must be an edge  $(y, x_0)$  joining  $G_1$  to  $x_0$ . Now  $(y, x_0, x_1, \ldots, x_m)$  is a longer path than P.

A cycle shows that the assertion does not hold for strongly connected

digraphs.

(b) Let  $P = (x_0, x_1, \ldots, x_m)$  be a longest path in G. If  $G - x_0 - x_1$  is connected we are finished. Assume not, then there is a point y separated from  $x_2, \ldots, x_m$  by  $x_0$  and  $x_1$ . Let Q be a (y, P)-path in G. Then Q hits P at  $x_0$  or  $x_1$ . However, by the maximality of P, it must hit at  $x_1$ . Also, since  $Q \cup (P - x_0)$  is a path and has length  $\leq |E(P)|$  by the maximality of P, Q consists of a single edge, joining y to  $x_1$ . No other edge can leave y, since it cannot go to a point outside P (by the maximality of P) and cannot go to a point of P since  $x_0, x_1$  disconnect P and Y. Thus, Y is of degree 1.

Now observe that  $(y, x_1, \ldots, x_m)$  is a maximum path and therefore the same argument yields a point  $z \neq y$  of degree 1 connected to  $x_1$ . This contradicts

the assumption of the theorem.

- 7. (a) Let  $e \in E(T_1) E(T_2)$ . Then  $T_1 e$  has two components and  $T_2$  has an edge f connecting these.  $T_1 e + f$  is a spanning tree having more edges in common with  $T_2$  than  $T_1$ . Repeating this we can transform  $T_1$  into  $T_2$ .
- (b) Let W be a largest common subtree of  $T_1$  and  $T_2$ . We prove the assertion by induction on |V(G)| |V(W)|. If  $|V(W)| \ge |V(G)| 1$  the assertion is trivial. Suppose  $|V(W)| \le |V(G)| 2$ , and let  $e_i = (x_i, y_i)$  be an edge of  $T_i$  with  $x_i \in V(W)$ ,  $y_i \notin V(W)$ .
- Case 1.  $y_1 \neq y_2$ . Let  $T_3$  be a spanning tree of G containing  $W + e_1 + e_2$ . Then by the induction hypothesis  $T_1$  can be transformed into  $T_3$  and  $T_3$  can be transformed into  $T_2$  in the required way, which proves the assertion.
- Case 2.  $y_1 = y_2$ . Let  $e_3 = (x_3, y_3)$  be any edge of  $G y_1$  such that  $x_3 \in V(W)$ ,  $y_3 \notin V(W)$  (here we use the fact that G is 2-connected). Let  $T_3$  and  $T_4$  be spanning trees of G containing  $W + e_1 + e_3$  and  $W + e_2 + e_3$ , respectively.

Then by the induction hypothesis, we can transform  $T_1$  to  $T_3$  to  $T_4$  to  $T_2$ ,

which proves the assertion.

Note that this procedure has the additional property that if  $T_1$  and  $T_2$  have a common subtree W this remains unchanged during the sequence of transformations, and if W consists of a single point this never plays the role of the endpoint x.

8. (a) Let  $e = (x_1, x_2)$  be an edge of G and let  $T_i$  consist of e and a spanning tree of  $G - x_i$ . Then by 6.7b,  $T_1$  can be transformed into  $T_2$  through a sequence of intermediate trees in the given way, and by the remark after the solution of 6.7b, we may assume all these trees contain e. The branch of these trees relative to e containing  $x_1$  has one point in  $T_1$  and n-1 points in  $T_2$  and its size changes by at most one at each step. Hence it will have exactly  $n_1$  points in some intermediate spanning tree T. Now the two branches of T relative to e give the partition we wanted.

(b) The assertion is clearly equivalent to the fact that a 2-connected non-bipartite graph G has a spanning tree T such that the (unique) 2-coloration

of T has equal color-classes.

Let  $C = (x_0, \ldots, x_{2k})$  be an odd circuit in G. Consider a spanning tree W of  $G - x_0$  containing  $C - x_0$  and the two spanning trees  $T_1, T_2$  of G obtained from W by adding the edges  $(x_0, x_1)$  and  $(x_{2k}, x_0)$ , respectively. By 6.7b and the remark after its solution, we can obtain the tree  $T_2$  from  $T_1$  through a sequence of spanning trees of G, each of which arises from the preceding one by the following operation: we remove the edge adjacent to an endpoint  $x \neq x_0$  and connect x to the rest by another edge of G. Let us 2-color each of these trees with red and blue in such a way that x is red. Then if  $T_1$  has k red points,  $T_2$  has 2m - k + 1 red points, because clearly every point except x has different colors in each of them. Moreover, the number of red points changes by at most one at each step. Hence there is an intermediate tree with exactly m red points. [A. Bondy-L. Lovász; see L. Lovász, Acta Math. Acad. Sci. Hung. 30 (1977) 241–251.]

**9.** If G is strongly connected between a and b, let  $\emptyset \neq X \subset V(G)$ ,  $a \in X$ ,  $b \in V(G) - X$ . Walking along a (directed) (a, b)-path, at some point we have to leave x. The next edge of the (a, b)-path connects the set X to V(G) - X (in this direction).

Conversely, suppose there is no (a, b)-path. Let X be the set of those points which are accessible from a along a directed path. Then  $a \in X$ ,  $b \notin X$  and there is no edge (x, y) with  $x \in X$ ,  $y \notin X$  since any (a, x)-path plus this edge would give an (a, y)-path, which, however, does not exist because  $y \notin X$ .

10. Let  $G_0$  be the digraph obtained by contracting the red edges and removing the green ones.

Suppose first there is no directed (x, y)-path in  $G_0$ . Then there is a set  $S_0$ ,  $x \in S_0$ ,  $y \notin S_0$  such that no black edge goes from  $S_0$  to  $V(G_0) - S_0$ . Let  $S_0$  be the co-image of  $S_0$  under the contraction of red edges; then no red edge connects S and V(G) - S and no black edge goes from S to V(G) - S, i.e. (ii) is satisfied.

Trivially one obtains that if there is an (x, y)-path in  $G_0$ , then (i) is satisfied.

Finally, (i) and (ii) cannot be satisfied simultaneously. Suppose indirectly that there is a path P as in (i) and a set S as in (ii). Then P has a first edge f joining S to V(G) - S. f cannot be green since P has no green edges and it cannot be red since no red edge joins S to V(G) - S.

Thus f is black. But f cannot be oriented from S to V(G) - S by (ii), neither can it be oriented conversely since it lies on P. [G. J. Minty; see B, ch. 1.]

11. (a) Let F be a minimum set of edges such that G-F is not strongly connected. Then by 6.8, there is a set  $\emptyset \subset X \subset V(G)$  with  $\delta_{G-F}(X)=0$ . Any edge e in F must join a point of X to a point of V(G)-X; otherwise e could be placed back and we would still have  $\delta_{G-F+e}(X)=0$ . Therefore, if we put back the edges of F inversely the resulting digraph G' has  $\delta_{G'}(X)=0$ , i.e. G' is not strongly connected.

(b) Let F be a minimum set of edges such that G/F is strongly connected, and let  $G_0$  be obtained from G by reversing the edges of F. We claim that  $G_0$ 

is strongly connected.

First we consider the case  $F = \{f\}$ . G is not strongly connected by the minimality of F; thus there is a set  $X \subset V(G)$ ,  $X \neq \emptyset$  with  $\delta_G(X) = 0$ . Since G/f is strongly connected, f must join a point  $y \in V(G) - X$  to a point  $x \in X$ . Invert f, and assume indirectly that the resulting digraph  $G_0$  is not strongly connected either. Thus there is a set  $Y \subset V(G)$ ,  $Y \neq \emptyset$  such that  $\delta_G(Y) = 0$ . Again since G/f is strongly connected,  $y \in Y$  but  $x \in V(G) - Y$ .

Now if  $X \cap Y \neq \emptyset$ , then  $\delta_{G/f}(X \cap Y) = 0$  which is impossible. Thus  $X \cap Y = \emptyset$ . Similarly we obtain  $X \cup Y = V(G)$ . But then  $\delta_G(X) = \delta_G(Y) = 0$  implies that f is an isthmus, a contradiction. This settles the case |F| = 1.

The general case follows by an easy induction on |F|. Let  $f \in F$  and denote by H the digraph obtained from G by reversing f. The digraph  $G/(F-\{f\})$  clearly contains no isthmuses, is not strongly connected (by the minimality of F) and contracting f in it we obtain a strongly connected digraph. Hence if we reverse f the resulting graph  $H/F-\{f\}$  will be strongly connected. Moreover,  $F-\{f\}$  is a minimal set with this property. For if  $H/F_0$  were strongly connected for some  $F_0 \subset (F-\{f\})$ , then so would be  $(H/F_0)/f = G/F_0 \cup \{f\}$  which would contradict the minimality of F as  $F_0 \cup \{f\} \subset F$ . Thus we may apply the induction hypothesis and conclude that the graph obtained from H by reversing the edges of  $F-\{f\}$  is strongly connected. But this digraph is just  $G_0$  [A. Frank].

(c) Suppose H is a digraph which contains no cycles. Then H has a point x with outdegree 0; otherwise, starting from any point and walking along edges in their direction, we would never get stuck and sooner or later we would complete a cycle. Using induction on V(H), we may assume that the points of H-x can be ordered in such a way that every edge has larger head than tail. Putting x on the top, we obtain such an ordering of V(H).

Now let  $\overline{F}$  be a minimal set of edges such that H - F is acyclic. Then V(H) has an ordering such that every edge of H - F has larger head than tail. If we put back any edge of F the ordering loses this property, because the graph is no longer acyclic. Thus all edges of F go "downward". But then inverting them they will have a larger head than tail, i.e. the graph obtained from G by inverting the edges of F is acyclic. [E. J. Grinberg, J. Dambit,

Latv. Mat. E. 2 (1966) 65–70; T. Gallai, Theory of Graphs (P. Erdős–G. O. H. Katona, eds.) Akadémiai Kiadó, (1968) 115–118.]

12. If the tournament T has a Hamiltonian cycle, then it is obviously strongly connected. Now suppose that T is strongly connected. Let  $C = (y_1, \ldots, y_k)$  be a maximal cycle in T (this exists, as the acyclic tournament is not strongly connected). Suppose indirectly that C is not a Hamiltonian cycle.

Let x be a point not on C. Suppose, e.g.  $(y_1, x) \in E(T)$ . If  $(x, y_2) \in E(T)$ , then  $(y_1, x, y_2, \dots, y_k)$  is a longer cycle. So  $(y_2, x) \in E(T)$ . Similarly  $(y_i, x) \in E(T)$ .

 $\in E(T)$  for  $i=1,2,\ldots,k$ .

Now, let X be the set of all points x such that  $(y_1, x) \in E(T)$ . Then  $(y_i, x) \in E(T)$  for every  $x \in X$  as above. Let  $(x, z) \in E(T)$  be an edge with  $x \in X$ ,  $z \notin X$  (which exists by  $\delta_T(X) > 0$ ). Then  $z \notin C$ , and thus,  $z \notin X$  implies  $(z, y_1) \in E(T)$ . Now  $(x, z, y_1, \ldots, y_k)$  is a cycle longer than C [P. Camion; see B].

13. Let  $C = (y_1, \ldots, y_k)$  be a longest cycle which is not a Hamiltonian cycle; this exists as  $n \ge 4$ . If C has length |V(T)| - 1, it misses exactly one point x

and so T - x is strongly connected.

Assume indirectly that C has length at most |V(T)| - 2. The same argument as in the previous solution yields that, for each  $x \in V(T) - V(G)$ , either  $(y_i, x) \in E(T)$  for each  $1 \le i \le k$  or  $(x, y_i) \in E(T)$  for each  $1 \le i \le k$ . Let X denote the set of points with the first property. Let (x, z) be an edge leaving X. Then

$$(x, z, y_1, \ldots, y_{k-1})$$

is a cycle longer than C but still missing a point  $(y_k, \text{ in fact})$ .

To show that there are at least two such points let  $y_1$  have this property.  $y_1$  is contained in some circuit shorter than n; for let X denote the set of points z with  $(y_1, z) \in E(T)$ , then there is an edge (z, u) leaving X and  $(y_1, z, u)$  is a 3-cycle.

Consider now a longest cycle  $(y_1, \ldots, y_k)$  with k < n; we find as before that k = n - 1 and this yields a point  $x \neq y_1$  such that T - x is strongly connected.

14. Invert all edges of F. The resulting graph G' has no circuit, therefore the beginning point of a longest path is the tail of every edge incident with it. Inverting the edges of F again, we get that this point satisfies the requirements.

15. We use induction on |V(T)|. For |V(T)| = 1, 2 the assertion is trivial.

Let  $|V(T)| \geq 3$ .

If  $\varphi$  is one-to-one then it is an automorphism. Denoting by T' the subtree formed by the inner points,  $\varphi$  maps T' onto itself. If  $\varphi$  is not one-to-one, then  $\varphi(T)$  is a proper subtree mapped into itself by  $\varphi$ .

In both cases we are finished by induction.

Remark: The assertion is a degenerate case of the Lefschetz Fixed Point Theorem (see, e.g. E. Spanier, Algebraic Topology, McGraw-Hill, 1966).

16. The path connecting two points of the intersection is unique and is contained, therefore, in each of the given subtrees. Thus, it belongs to their intersection.

17. (a) Let  $x_1, x_2$  be the endpoints of  $Q, x_i \in P_i$ .  $x_i$  divides  $P_i'$  into two pieces  $P_i', P_i''$ . We may assume  $P_i'$  is at least as long as  $P_{i,i}''$  and that  $P_1'$  is at least as long as  $P_2'$ . Then

$$|E(P_1' \cup Q \cup P_2')| > |E(P_1')| + |E(P_2')| \ge 2 |E(P_2')| \ge |E(P_2')| + |E(P_2'')| = |E(P_2)|,$$

i.e. the path  $P_1' \cup Q \cup P_2'$  is longer than  $P_2$ , a contradiction.

(b) Let  $P_1$ ,  $P_2$  be two maximum paths. Their intersection Q is a path by 6.15, hence  $P_1 \cup P_2$  has the following form: there are two paths  $P_1'$ ,  $P_2'$  starting from one endpoint of Q, and two others:  $P_1''$ ,  $P_2''$  from the other endpoint, so that

$$P_1 = P_1' \cup Q \cup P_1'', \quad P_2 = P_2' \cup Q \cup P_2''.$$

We have  $|E\left(P_{1}'\right)|=|E\left(P_{2}'\right)|$ , because, e.g.  $|E(P_{1}')|>|E(P_{2}')|$  would imply that the path  $P_{1}'\cup Q\cup P_{2}''$  is longer than  $P_{2}$ . Similarly,  $|E(P_{1}'')|=|E(P_{2}'')|$ . Since  $P_{1}'\cup P_{2}'$  is a path, we have  $|E(P_{1}')|\leq 1/2\;|E\left(P_{1}\right)|$  and similarly,  $|E(P_{1}'')|\leq 1/2\;|E(P_{1})|$ .

This shows that the middle point (points) of  $P_1$  belongs to Q and, hence, to  $P_2$ . Since this holds for any  $P_2$ , this point (points) is in the intersection of all

maximum paths.

Remark: An example of H. J. Walther [WV; Fig. 39] shows that this is not true for every connected graph, as was conjectured by Gallai.




Fig. 39

18. First solution. For k=2 the statement is obvious. Using induction on k, we may assume  $G_1\cap\ldots\cap G_{k-1}=G_0\neq\emptyset$ . Suppose  $G_0\cap G_k=\emptyset$ . Let P be a  $(G_0,G_k)$ -path. Consider an edge (x,y) of P (x nearer  $G_k$ ). Then obviously  $x\notin V(G_0)$ , thus  $x\notin V(G_i)$  for some  $1\leq i\leq k-1$ . Hence  $(x,y)\notin E(G_i)$ . Now G-(x,y) is disconnected, and x and y are in distinct components of it. Obviously,  $G_k$  is in the component containing x. Since  $G_i$  is a connected subgraph of G-(x,y) and meets  $G_k$ , it also lies in the same component. But this is impossible because  $G_0\subseteq G_i$  and  $G_0$  has points in the other component.

Second solution. Let x be a point of degree 1 of G, joined to y. Suppose the assertion holds true for G - x.

6.19

If none of  $G_1, \ldots, G_k$  is the one-point graph with the point x, then  $G_1 - x, \ldots$ ,  $G_k - x$  intersect mutually; for if x is a common point of  $G_i$  and  $G_j$ , then so is y. Thus by the induction hypothesis  $G_1 - x, \ldots, G_k - x$  have a point in common, hence so do  $G_1, \ldots, G_k$ . If  $G_1$  (say) has only one point x, then x is a common point of  $G_1, \ldots, G_k$ .

19. To show  $d(x, y) + d(y, z) \ge d(x, z)$ , consider an (x, y)-path P of length d(x, y) and a (y, z)-path of length d(y, z).  $P \cup Q$  contains an (x, z)-path, which obviously has length at most d(x, y) + d(y, z). Thus d(x, z), the minimum length of all such paths, is also  $\le d(x, y) + d(y, z)$ .

To show  $D(x,y) + D(y,z) \ge D(x,z)$ , consider an (x,z)-path P of length D(x,z). Since G is connected, we have a (y,P)-path Q; let t be the endpoint of Q on P. t splits P into two paths  $P_1$ ,  $P_2$ , of lengths  $l_1$  and  $l_2$ , say. Let Q have length k. Then  $Q \cup P_1$  is an (x,y)-path, hence

$$D(x, y) \geq k + l_1$$
.

Similarly,

$$D(y,z) \geq k + l_2$$

whence

$$D(x, y) + D(y, z) \ge 2k + l_1 + l_2 = 2k + D(x, z) \ge D(x, z).$$

**20.** In the formulation given in the hint, let us see how many times a given edge e is counted on each of the two sides. G - e consists of two components; let  $\alpha$  of the points  $p_i$  and  $\beta$  of the points  $q_i$  be in one of them. Then e is contained in

$$lpha(n-lpha)$$
 of the  $(p_i,p_j)$ -paths,  $eta(n+1-eta)$  of the  $(q_i,q_j)$ -paths,  $lpha(n+1-eta)+eta(n-lpha)$  of the  $(p_i,q_j)$ -paths.

It suffices to show that

$$\alpha(n-\alpha)+\beta(n+1-\beta)\leq \alpha(n+1-\beta)+\beta(n-\alpha)$$

or, equivalently,

$$\beta - \alpha \leq (\alpha - \beta)^2$$
,

which is true because  $\alpha$ ,  $\beta$  are integers. [J.B. Kelly, in: Comb. Structures and their Appl. Gordon and Breach (1969) 201–208; cf. 13.16.]

21. (a) First we prove the assertion of the hint. Since all points at distance  $\tilde{d}(x)$  from x are, obviously, of degree 1,  $\tilde{d}(x)$  decreases at every point. On the other hand, only one point was omitted from the longest path starting from x; hence  $\tilde{d}(x)$  decreases by 1.

Now suppose  $|V(G)| \ge 2$ . Let Z be the set of points, where  $\tilde{d}(x)$  is minimal. If Z contains no points of degree 1, remove all points of degree 1 from G. By the assertion of the hint, this does not alter Z and hence the statement follows by induction. If a point x of degree 1 belongs to Z, look at the neighbor y of x. Obviously, y is strictly nearer than x to any other point. Hence,  $\tilde{d}(x)$  can be

minimal only if  $\tilde{d}(x) = 1$ , i.e. G is the tree with two points. In this case, the statement is obvious.

Note that the center (bicenter) is contained in every maximum path (cf. 6.16b).

(b) Let P be a path of length  $\tilde{d}(x)$  starting from x. If P does not contain y or z, then obviously,

$$\tilde{d}(y) = \tilde{d}(x) + 1, \quad \tilde{d}(z) = \tilde{d}(x) + 1$$

and the assertion follows. If P contains y (say), then

$$\tilde{d}(y) \ge \tilde{d}(x) - 1, \quad \tilde{d}(z) = \tilde{d}(x) + 1,$$

which proves the statement. [See, e.g. K.]

22. (a) Let  $k_1$ ,  $k_2$  be the numbers of points in the components of G - x containing y and z, respectively (obviously, these are different components). Moving from x to y,  $k_1$  points approach but  $n - k_1$  depart, hence

$$s(x) = s(y) + k_1 - (n - k_1) = s(y) + 2k_1 - n.$$

Similarly,

$$s(x) = s(z) + 2k_2 - n.$$

Summing,

$$2s(x) = s(y) + s(z) + 2(k_1 + k_2 - n) \le s(y) + s(z) - 2.$$

(b) Assume indirectly that there are two non-adjacent points x, y such that s(x) = s(y) is minimal. Let

$$(x_1 = x, x_2, \ldots, x_p, x_{p+1} = y)$$

be the (x, y)-path. Then

$$s(x_2) \geq s(x)$$
.

By (a),

$$s(x) + s(x_3) > 2s(x_2) \ge s(x_2) + s(x), \ s(x_3) > s(x_2) \ge s(x),$$
  
 $s(x_2) + s(x_4) > 2s(x_3) > s(x) + s(x_2), \ s(x_4) > s(x),$ 

and so on, finally we get

$$s(x_{p+1}) = s(y) > s(x_p),$$

a contradiction [K].

(c) Let  $x_1, \ldots, x_p$  be the points of a path and  $y_1, \ldots, y_q$  other points connected to  $x_1$ . Then, if p is even, the center of the tree is  $x_p$ ; on the other hand, the baricenter is  $x_1$ , if  $q = \binom{p}{2}$ . Thus, the distance of the baricenter and center is  $\frac{p}{2} - 1$ , which is arbitrarily large indeed.

23. In

$$\sum_{x,y\in V(G)}d(x,y),$$

there are exactly n-1 terms equal to 1; if G is the star, all the other terms are 2; otherwise, they are not less than 2 with at least one of them 3. Hence, the sum is minimal for the star.

To prove the statement given in the hint for the second part of the problem, note that, for an endpoint x of a path P we have

$$s(x) = 1 + 2 + \ldots + (n-1).$$

If G is any tree and x an endpoint of it, and d(x) = d, we have at least one point of distance 1, 2, ..., d from x, therefore the sum defining s(x) looks like

(2) 
$$s(x) = 1 + 2 + \ldots + d + d_1 + \ldots + d_{n-1-d},$$

where  $d_1, \ldots, d_{n-1-d} \leq d$ . Obviously,  $(2) \leq (1)$  and equality holds only if d = n - 1, i.e. G is a path and x its endpoint.

Now we prove by induction on n that

$$\sum_{x,y\in V(G)}d(x,y)$$

is maximal (G ranging over all n-point trees) if G is a path. Let a be a point of degree 1 in G. Then

$$\sum_{x, y \in V(G)} d(x, y) = s(a) + \sum_{x, y \in V(G-a)} d(x, y).$$

Here the first term is maximal if G is a path and a is an endpoint of it. Fortunately, in this case the second term is also maximal by the induction hypothesis.

**24.** Let  $P = (x_1, \ldots, x_{2k-2})$  be a path of length 2k-3 in G. With any point y not on P, associate a path  $P_y$  in the following way. Let Q be the path connecting y to P and having no other point in common with P. If Q is longer than k-1, let  $P_y$  be the segment of Q of length k, starting from y. Otherwise, let  $P_y$  consist of Q and a subpath of P. Such a choice is possible since P has length 2k-3, hence one of the pieces of P incident with the endpoint of Q is of length k-1. Furthermore, let  $P_{x_i}$  be a subpath of P incident with  $x_i$  for  $k \leq i \leq 2k-2$ .

It is easy to see that the  $P_{\nu}$ 's are distinct and that their number is n-k.

25. The resulting graph is obviously a spanning tree. Let H be an optimal spanning tree with the maximum number of edges in common with G and let  $e_i = (x, y)$  be an edge of G not in H, selected at the  $i^{\text{th}}$  step. Let P be the path in H connecting x to y; then there is an edge f of P connecting  $G_i$  to a point outside it. Since we preferred to choose  $e_i$  rather than f,  $v(e_i) \leq v(f)$ . On the other hand,  $H - f + e_i$  is a spanning tree and has no more expense but more edges in common with G than H, a contradiction.

Remark: If  $G_i$  is the component containing a given point at every step, the graph formed by the edges selected in the first i steps is always connected. [J.B. Kruskal; see S.]

**26.** Let G, G' be two optimal trees. Let e be an edge of G not in G'. Consider the path in G' connecting the endpoints of e. Some edge of it, f say, connects the two components of G - e. Now either G - e + f or G' - f + e has less expense than G.

27. If e, f are contained in the same circuits, then f is contained in no circuit of G - e, hence G - e - f is disconnected.

Conversely, if there is a circuit containing f but not e (say), then f is on a circuit of G - e, whence G - e - f is connected. This proves the statement of the hint.

Now (a), (b) follow immediately. To show (c), let e be an edge not in the removed equivalence class P, and let  $f \in P$ . e is not a cut edge of G - f, hence there is circuit C containing e but not f. C obviously contains no other edge of P, whence C is a circuit of G - P. Obviously C lies in the connected component of G - P containing e. Hence e is not a cut edge of this component.

To show (d), we show that each component  $G_0$  of G-P is incident with two edges of P. Let (x, y) be an edge of P incident with  $G_0$  ( $x \in V(G_0), y \notin V(G_0)$ ). Let C be a circuit containing (x, y). Let us walk along C, starting from x through (x, y). Upon reaching a point of  $G_0$  again, let us go back to x within  $G_0$ . The circuit defined this way contains exactly two edges of P incident with  $G_0$ ; on the other hand, it contains all edges of P; hence there are exactly two edges of P incident with  $G_0$ .

Thus, contracting the components of G-P, we get a connected regular graph of degree 2. This is a circuit.

- 28. We can select a circuit  $G_1$  of G. Assume  $G_1, \ldots, G_j$  are already selected in such a way that  $G_{i+1}$   $(1 \leq i < j)$  is either a path having its endpoints in common with  $G_1 \cup \ldots \cup G_i$  or a circuit having one point in common with  $G_1 \cup \ldots \cup G_i$ . If  $G_1 \cup \ldots \cup G_j = G$  we are finished. Otherwise, there is an edge e = (x, y) not in  $G_1 \cup \ldots \cup G_j$  but having a point x in common with  $G_1 \cup \ldots \cup G_j$ . Let G be a circuit containing G; start from G0 walking through G1 and then along G2 until we meet G1 G1 again; let G2 again; let G3 again; let G4 be the subgraph consisting of those edges and points we touched. Then, obviously, G4 is either a path with two endpoints in G4 G5 or a circuit G6 meeting G5 in the point G6 where G7 in the point G6 being finite, sooner or later we have to have decomposed the whole G5.
- **29.** First solution. If we orient G in some way and the resulting G is strongly connected, let (x, y) be an edge of G, directed from x to y. There is a path in G connecting y to x; hence (x, y) is contained in a circuit of G (even of G). Now assume that G is 2-edge-connected. We use induction on the number of edges. Remove an equivalence-class P defined in 6.27. The components of the remaining graph are 2-connected and thus they can be oriented so that they are strongly connected. The edges of P lie on a circuit G; let us orient them in accordance with a cyclic orientation of G. It is easy to see that the resulting digraph is strongly connected.

Second solution. (for the second part): Let  $G = G_1 \cup \ldots \cup G_r$  as in 6.28, then we can define an orientation of G by considering an orientation which makes an oriented path or cycle out of each  $G_i$ .


Let  $a \in V(G_1)$ . Then it easily follows by induction on j that any  $b \in V(G_j)$  is accessible from a on a directed path and conversely, a is accessible from any other point. By the transitivity of accessibility this proves  $\vec{G}$  is strongly connected. [H.E. Robbins; see B, Wi.]

**30.** By induction on k, we can find a circuit  $C_0$  such that  $e_1, \ldots, e_{k-1}$  are well fitted with  $C_0$ . If  $e_k$  is well fitted with  $C_0$ , we are finished. If both endpoints of  $e_k$  are on  $C_0$ ,  $e_k$  and one of the arcs bounded by its endpoints give the desired circuit. Thus we may assume e has one endpoint x on  $C_0$  and one endpoint y not on  $C_0$ .

Contract  $C_0$ . The resulting graph G' is obviously 2-edge-connected and by induction, it has a circuit  $C_1$  with which all the edges  $e_i$  not on  $C_0$  are well-fitted. Let  $C'_1$  be the subgraph of G mapped onto  $C_1$  by the contraction. There are

three possibilities.

- (a)  $C'_1$  is a circuit disjoint from  $C_0$ . Then  $C'_1$  is the desired circuit.
- (b)  $C'_1$  is a circuit having one point in common with  $C_0$ . Then it must contain  $e_k$  (since  $e_k$  is well fitted with  $C_1$ ), hence it does not meet the  $e_i$ 's on  $C_0$ , and has the desired property again.
- (c)  $C'_1$  is a path connecting two points of  $C_0 \cdot e_k$  lies again on  $C'_1$ . Now  $C'_1$  and one of the arcs of  $C_0$  bounded by its endpoints gives a circuit with the desired properties (Fig. 40).



31. The only non-trivial statement to prove is that mentioned in the hint. Walk from  $e_3$  in both directions on  $C_2$  till we meet  $C_1$ ; let x and y be the two points reached (it may happen that x and/or y are endpoints of  $e_3$ ). x and y are distinct, so they bound two arcs of  $C_1$ , one of which contains  $e_1$ ; this arc and the arc of  $C_2$  bounded by x, y and containing  $e_3$ , give a circuit containing  $e_1$  and  $e_3$ .

32. (i)  $\Rightarrow$  (iii): First let e and f be two edges with a common endpoint x. Let y and z be their other endpoints. Since G - x is connected, there is a path

in it connecting y to z; this forms with e and f a circuit.

Using 6.31, we get that any two edges are on a circuit; in fact, an equivalenceclass of the relation "being on a circuit" contains every edge which is adjacent to its edges; therefore consists of the edges of a component. But G is connected, hence this equivalence-class is the whole graph.

- (iii) ⇒ (ii): Given two points, consider one edge incident with each and a circuit through these two edges.
- (ii)  $\Rightarrow$  (i): Assume that the removal of a point x disconnects G. Let a, b lie in different components of G x. Then no circuit can contain both a and b.
- **33.** First solution. We show that  $G = G_1 \cup \ldots \cup G_r$ , where  $G_1$  is a circuit and  $G_{i+1}$  is a path having exactly its endpoints in common with  $G_1 \cup \ldots \cup G_i$ . Let  $G_1$  be a circuit and choose  $G_2, \ldots, G_i$  satisfying the requirement above. If  $G_1 \cup \ldots \cup G_i \neq G$ , we can find an edge (x, y) in  $G E(G_1 \cup \ldots \cup G_i)$  such that  $x \in V(G_1 \cup \ldots \cup G_i)$ . Since G x is connected, we can find a  $(y, G_1 \cup \ldots \cup G_i x)$ -path Q in G x. Then Q and (x, y) give a  $G_{i+1}$ . Therefore, sooner or later we have  $G = G_1 \cup \ldots \cup G_r$ .

Now  $G_1 \cup \ldots \cup G_{r-1}$  is 2-connected, which can be seen very easily. Hence

 $P_{\bullet} = G_r$  satisfies the requirements.

Second solution. Let  $P_1, P_2, P_3$  be three paths connecting a and b  $(a, b \in V(G))$  and assume  $a, b, P_1, P_2, P_3$  are chosen so that  $P_1$  is of minimum length. First we show  $P_1$  has inner points of degree 2 only. Assume (x, y) would be an edge incident with an inner point x of  $P_1$  but not belonging to  $P_1$ . Since G-x is connected there is a  $(y, P_1 \cup P_2 \cup P_3 - x)$ -path Q in G-x. The other endpoint of Q may be on  $P_1$  or on  $P_2 \cup P_3$ ; in either case we get two points connected by three independent paths, one of them being shorter than  $P_1$ , a contradiction with the choice of  $P_1$  (Fig. 41).




Fig. 41

Thus the inner points of  $P_1$  are of degree 2. Remove the inner points of  $P_1$ . The resulting graph G' is 2-connected. In fact, consider two points a,b of G'-c for some c. There is an (a,b)-path P in G-c. If P does not go through  $P_1$ , it is an (a,b)-path in G'-c. If  $P_1 \subseteq P$ , one of  $P_2$ ,  $P_3$  does not contain c and we can "avoid"  $P_1$  using it. Thus G'-c is connected.

This proves that  $P_1$  satisfies our requirements. [H. Whitney, Amer. J. Math. 55 (1933) 236-244; also see G. A. Dirac, J. Reine Angew. Math. 228 (1967)

204-216.]

**34.** Let  $G = G_1 \cup \ldots \cup G_r$ , where  $G_1$  is a (p,q)-path and  $G_{i+1}$  is a path which has exactly its endpoints in common with  $G_1 \cup \ldots \cup G_i$ . Such a decomposition can be found just as in the first solution of the preceding exercise.

Now order the points of  $G_1$  according to their position on  $G_1$ ; insert the points of  $G_2$  between them, so that they are ordered in accordance with their positions

on  $G_2$  (and thus, lie between the two endpoints of  $G_2$ ) etc. The ordering obtained finally has the property that each point is adjacent to both earlier and later

points.

Now orient any edge, so that its tail anticipates its head in this ordering. Then starting from any edge (x, y), we can find edges  $(y, y_1)$ ,  $(y_1, y_2)$  etc. by the properties of the ordering the last of which must end, of course, in q. Similarly, there are edges  $(x_1, x)$ ,  $(x_2, x_1)$  etc. the last starting from p. These edges yield a (p, q) path of G containing (x, y). [A. Ádám, A problem of the Schweitzer Competition, Mat. Lapok, 22 (1971) 34.]


35. The assertion given in the hint follows from the second solution of problem 6.33; it proves immediately that (i) implies (ii). Conversely, if (ii) holds, let e = (x, y) be an edge such that G - e is 2-connected. Then x and y lie on a circuit of G - e which has e as a chord, a contradiction. [M. D. Plummer, Trans. Amer. Math. Soc. 134 (1968) 85-94.]

**36.** If G = C we have nothing to prove. Otherwise, there is a (C, C)-path in G. For let u be a point of C of degree at least 3, (u, v) an edge not on C, and  $P_0$  a (v, C - u)-path in G - u. Then  $P_0 + (u, v)$  is a (C, C)-path.

Now let P be a  $(C, \overline{C})$ -path selected as in the hint. R has at least 2 edges by 6.35. Assume z has degree at least 3. Then it follows as above that z is the

endpoint of a (C, C)-path Q. Let w be the other endpoint of Q.

Observe that if P and Q intersect or w is outside R, we can find a circuit such that (x, z) is a chord of it, which contradicts 6.35 (see Fig. 42). However, this means that the arc connecting z and w (in R) is a proper sub-arc of R, which contradicts the choice of P. [M. D. Plummer, ibid.]



**37.** G' is a forest by 6.36. By the argument in the preceding proof, there is a (G', G')-path  $Q_1$ . If G' is connected, then it contains a path  $Q_2$  connecting the endpoints of  $Q_1$ .  $C = Q_1 \cup Q_2$  is a circuit whose points of degree 2 form an arc of it (these are the inner points of  $Q_2$ , in fact).

Now let P be a (C, C)-path; one of the arcs connecting its endpoints has no points of degree 2. On the other hand, this arc contains a minimal arc of this type and this has a point of degree 2 by the solution of the preceding exercise.

[M. D. Plummer, ibid.]

38. Let  $G_1$ ,  $G_2$  be two isomorphic trees such that every endpoint of them is at distance at least 1000 from their centers, and which have no points of degree 2. Identify the corresponding endpoints of  $G_1$  and  $G_2$ . The resulting graph

G is 2-connected; for take any two points x, y of it. Let x', y' be the corresponding points in the other tree (maybe x = x' or y = y'). We show that even x, x', y, y' lie on a circuit. We may assume  $x, y \in G_1$   $x', y' \in G_2$ . Let  $P_1$  be the path in  $G_1$  connecting x to y. We can lengthen  $P_1$  to a path P which connects two endpoints of  $G_1$ . Let P' be the path corresponding to P; then  $P \cup P'$  is a circuit which contains x, x', y, y'.

On the other hand, removing an  $e \in E(G_1)$  (say) the resulting graph will not even be 2-edge connected, because the edge  $e' \in E(G_2)$  corresponding to e

will be a cutting edge of it.

**39.** (a) If G has k edge-disjoint (a, b)-paths, it is obviously k-edge-connected between a and b. To prove the other part, remove edges till the removal of any further edge will destroy k-edge-connectivity between a and b. Then obviously, there will be no edge with head at a or tail at b. Assume first there is an edge  $e_1$  incident neither to a nor to b. Since  $G - e_1$  no longer satisfies the conditions, it has a (k-1)-element (a,b)-cut C'. Then,  $C = C' \cup \{e_1\} = \{e_1, \ldots, e_k\}$  is a k-element (a,b)-cut and by the choice of  $e_1$ , the set S' determining C satisfies  $|S| \geq 2$ ,  $|V(G) - S| \geq 2$ .

Let  $G_1, G_2$  be the graphs obtained by contracting S and V(G) - S, respectively;

let a' and b' be the images of a in  $G_1$  and b in  $G_2$ , respectively.

Obviously,  $G_1$  is k-connected between a' and b, and thus by the induction hypothesis, there are k edge-disjoint (a', b)-paths  $P_1, \ldots, P_k$ . Since the edges going out from a' are only  $e_1, \ldots, e_k$ , we may assume  $e_i \in P_i$ . Similarly, there are k edge-disjoint (a, b')-paths  $Q_1, \ldots, Q_k$  in  $G_2$ ,  $e_i \in Q_i$ . Then  $P_1 \cup Q_1, \ldots, P_k \cup Q_k$  form k-edge-disjoint (a, b)-paths in G.

What is left is the case, when each edge has tail at a or head at b.

If there is an (a, b)-edge, we can remove it and proceed by induction on k,

thus we may assume there is no such edge.

For any  $x \neq a$ , b, let k(x) be the minimum of the numbers of (a, x)-edges and (x, b)-edges. Then obviously, there are  $\sum_{x \neq a, b} k(x)$  edge-disjoint (a, b)-paths.

On the other hand, let S be the set of all points x which are connected to b by k(x) edges. Then, the cut determined by  $\{a\} \cup S$  has exactly  $\sum_{x \neq a, b} k(x)$  edges. Hence

$$\sum_{x \neq a, b} k(x) = k,$$

which proves the assertion.

- (b) As suggested in the hint, consider a graph G', which has points a, b and two points  $x_1, x_2$  for each  $x \in V(G)_1$ ,  $x \neq a, b$ . Put  $a_1 = a_2 = a$  and  $b_1 = b_2 = b$ . For any edge  $e = (x, y) \in E(G)$ , G' has the edge  $e' = (x_2, y_1)$ , moreover, for each  $x \in V(G)$ ,  $x \neq a, b$ , the edge  $(x_1, x_2)$ . Now
- (i) G' is k-edge-connected between a and b iff G is k-connected between a and b;

(ii) G' has k-edge-disjoint (a, b)-paths iff G has k vertex-disjoint (a, b)-paths.

To show (i), consider an (a, b)-cut C in G'. Let A consist of all points x such that  $(x_1, x_2) \in C$  and all other edges of C. Then |A| = |C| and A separates a and b in G; for if P is any (a, b)-path in G, then the edges of G' corresponding

to edges and inner points of P, form an (a, b)-path P' in G and since P' contains an edge of C, P contains an edge or point of A.

Conversely, if A is a set of edges and points separating a and b in G, then the construction above associates a set C of edges of G' with it, and C will be an

(a, b)-cut with |C| = |A|. This proves (i).

Now consider k edge-disjoint (a, b)-paths  $P_1, \ldots, P_k$  in G'. If  $x_i \in P_j$  then, obviously,  $(x_1, x_2)$  is an edge of  $P_j$  and  $x_{3-i}$  is also on  $P_j$ . Hence,  $P_1, \ldots, P_k$  are vertex-disjoint and even contracting the edges  $(x_1, x_2)$  to get G back, we obtain k vertex-disjoint (a, b)-paths of G.

Conversely, if there are k vertex-disjoint (a, b)-paths in G, then the (a, b)-paths of G' associated with them in the natural way are vertex-disjoint, and

hence edge-disjoint. This proves (ii).

(i) and (ii) prove the statement of (b), by (a).

(c) The directed graph  $\bar{G}$  obtained as in the hint has the same connectivity

and edge-connectivity between any two points as G.

Moreover, the maximum number of edge-disjoint (vertex-disjoint) (a, b)-paths of G and  $\overline{G}$  are the same. For edge[vertex]disjoint (a, b)-paths of G yield such paths in  $\overline{G}$ . Conversely, if we have some edge-disjoint (a, b)-paths of  $\overline{G}$ , we may assume that they do not use both (x, y) and (y, x); for in this case one can easily find another system of the same number of (a, b)-paths, which contain neither (x, y) nor (y, x) (in the vertex-connectivity case, this difficulty does not even arise). These paths yield edge-disjoint (vertex-disjoint) (a, b)-paths in G [K. Menger; see any textbook on graph theory.]

- **40**. The graph G' constructed in the hint is k-connected between a and b; since a and b are non-adjacent, it suffices to show that the (a, b)-paths cannot be covered by k-1 points; but this is the assumption. Therefore by Menger's theorem, there are k independent (a, b)-paths in G'; removing a and b from them, we get k (A, B)-paths as desired.
- 41. To prove the statement formulated in the hint, let us substitute w(x) distinct points for each x, and connect two new points if the original points are connected. Denoting by G' the resulting graph and by A' and B' the sets of points substituted for A and B, respectively, these sets will satisfy the conditions of 6.40. For let X be a set which meets all (A', B')-paths of G'. Choose X to be minimal. If X contains a point  $x_1$  substituted for x, then it contains every other point  $x_2$  substituted for x; since if  $X x_1$  fails to meet some path P, then  $P \cap X = \{x_1\}$  and hence if  $x_2 \notin X$ , replacing  $x_1$  by  $x_2$  in P we get a path disjoint from X. Let  $X_0$  be the set of all points  $x \in V(G)$  such that X contains the corresponding points of G', then

$$|X| = \sum_{x \in X_0} w(x),$$

on the other hand X meets all (A, B)-paths of G, hence  $|X| \geq k$ .

There are, thus, k(A', B')-paths in G' which have no point in common. Contracting the points substituted for x onto x for each  $x \in V(G)$ , we get k(A, B)-paths in G as required.

Menger's theorem is obtained by putting w(a) = w(b) = k and w(x) = 1 elsewhere; 6.40 follows by putting w(x) = 1 for every x. A third special case

of interest is when  $A = \{a\}$  and w(a) = k, w(x) = 1 for  $x \neq a$ . Then we get: if any set which does not contain a but meets every (a, B)-path is of cardinality at least k, then there are k (a, B)-paths which have no point but a in common. [G. A. Dirac, J. London Math. Soc. 38 (1962) 148–163.]

**42.** If  $R_1, \ldots, R_k$  are chosen in accordance with the hint, we claim that, if the beginning edge of a  $Q_j$   $(1 \le j \le k+r)$  is not the beginning edge of any  $R_i$ , then  $Q_j$  is independent of every  $R_i$ . For assume  $Q_j$  meets some of them and let x be the first point of  $Q_j$  in common with an  $R_i$ . Let  $R'_i$  be the path consisting of the (a, x)-part of  $Q_j$  and the (x, B)-part of  $R_i$ . Then

$$|E(R_1 \cup \ldots \cup R'_i \cup \ldots \cup R_k) - E(Q_1 \cup \ldots \cup Q_{k+r})| < |E(R_1 \cup \ldots \cup R_k) - E(Q_1 \cup \ldots \cup Q_{k+r})|,$$

a contradiction.

Now there are at least r of  $Q_1, \ldots, Q_{k+r}$  whose beginning edge is not a beginning edge of  $R_1, \ldots, R_k$ . [H. Perfect, J. Math. Anal. Appl. 22 (1968) 96–111.]

- **43.** With  $B = V(P_0)$ , the conditions of 6.42 are satisfied: we have k independent (a, B)-paths  $P_1, \ldots, P_k$  and, on the other hand, suitable beginning sections of  $Q_0, \ldots, Q_k$  give k+1 independent (a, B)-paths (no matter where they end, i.e. where they meet  $P_0$ ). Therefore, we have k+1 independent (a, B)-paths  $R_1, \ldots, R_k$  and R such that  $R_1, \ldots, R_k$  end at  $b, R \cup P_0$  contains a  $(k+1)^{\text{st}}$  (a, b)-path, independent of  $R_1, \ldots, R_k$ . [For this and the next problem see G. Hajós, Theory of Gr. Int. Symp. Rome, Dunod, Paris-Gordon and Breach, New York (1967) 147.]
- **44.** Let  $c \neq a, b$ . Assume indirectly that c is not connected to anyone of a and b. The addition of an (a, c)-edge produces k + 1 independent (a, b)-paths; removing this edge again we get k independent (a, b)-paths and a (b, c)-path independent of them. Similarly we get k independent (a, b)-paths and an (a, c)-path independent of them. By 6.43, there are k + 1 independent (a, b)-paths in G, which is a contradiction.

We show by a similar argument that, if x is joined to a but not to b and y is joined to b but not to a, then x and y are not adjacent. Assume indirectly that there was an edge e connecting them. Subdivide e by a point c. Then, adding, the edges (a, y) and (b, x) and applying 6.43, we get a contradiction just

as above.


This statement means the points connected to both a and b separate them. Hence their number is at least k. On the other hand, through each of these points there goes an (a, b)-path of length 2 and these paths are independent. This proves that the number of points joined to both a and b is k.

Now if  $G_1$  is spanned by a and its neighbors,  $G_2$  by b and its neighbors, then joining two points of  $G_1$  or  $G_2$  cannot produce k+1 independent (a,b)-paths as a and b will still be separated by k points. Thus  $G_1$ ,  $G_2$  are complete and we

are finished.

To deduce Menger's theorem, consider an undirected graph G and  $a, b \in V(G)$ . We may assume a, b are independent, since the removal of an (a, b)-edge decreases both the maximum number of independent (a, b)-paths and the con-

nectivity between a and b by 1. Let k denote the maximum number of independent (a, b)-paths; we have to show a and b can be separated by k points. Add edges till k remains the same; the graph we finally get is described above and has a k-element set separating a and b. But the same set separates a and b in G.



**45.** To disprove (i), consider the graph shown in Fig. 43. Any (a, b)-path starts through  $e_1$ , whence any two of them have an edge in common, similarly for (b, a)-paths. Taking an (a, b)-path P and a (b, a)-path Q, they connect two opposite corners of the broken rectangle; hence they intersect somewhere. Now each intersection point is a point of the graph and has degree 3, therefore they have an edge in common.

There are two vertex-disjoint paths connecting c to d; hence, if an edge is contained in each (a, b)-path it is either  $e_1$  or  $f_2$ . Since, similarly, an edge contained in each (b, a)-path is  $e_2$  or  $f_1$ , no edge is contained in all (a, b)-paths

and (b, a)-paths.

This example is not the simplest one. It was given here because contracting  $e_1$ ,  $e_2$ ,  $f_1$ ,  $f_2$ , it gives a graph G which is 2-connected between the image a' of a and the image b' of b, but any (a, b)-path has an edge in common with any (b, a)-path. This disproves (iii).

Finally, (ii) is obviously false for a graph consisting of k vertex disjoint

(a, b)-paths.

**46.** Let C be an (a, b)-cut, determined by S. Then

$$|C| \ge |C| - |C^*| = \sum_{x \in S} (d_G^+(x) - d_G^-(x)) = k,$$

which proves the statement by Menger's theorem.

- 47. (i) If there are neither (a, b)-paths, nor (b, a)-paths we have nothing to prove. If P is an (a, b)-path say, then by 6.46, G E(P) has a (b, a)-path Q which is edge-disjoint from P; a contradiction.
- (ii), (iii). Assume G is k-edge-connected between a and b. Then there are k edge-disjoint (a, b)-paths by 6.39. Remove the edges of these (a, b)-paths,

then the remaining graph has k edge-disjoint (b, a)-paths by 6.46. This proves both (ii) and (iii) [A. Kotzig, Wiss. Z. Martin-Luther Univ. Halle-Wittenberg 10 (1961-62) 118-125].

- **48.** (a) Every edge joining  $X \cap Y$  to V(G) X Y is counted twice on both sides. Those edges joining X Y to Y X are counted on the right-hand side only. Any other edge is counted exactly once on each side.
- (b) This follows from (a) by substituting V(G) Y for Y, and taking into account the fact that

$$\delta_G(Y) = \delta_G(V(G) - Y).$$

(c) It is clear that

1° if an edge connects two of X - Y - Z, Y - X - Z, Z - X - Y,  $X \cap Y \cap Z$ , then it is counted twice on the left-hand side and twice on the right-hand side,

 $2^{\circ}$  if an edge connects one of X-Y-Z, Y-X-Z, Z-X-Y,  $X \cap Y \cap Z$  to a point not in these sets, it is counted once on the left-hand side and at least once on the right-hand side,

3° no other edge is counted on the left-hand side.

**49.** Let  $\delta_G(X) = k$ , |X| minimal,  $x \in X$ . If all edges incident with x go to V(G) - X, then their number is  $\leq \delta_G(X) = k$ , so the degree of x is  $\leq k$ .

By the k-edge-connectivity, we have equality here.

Suppose that x is adjacent to  $y \in X$ . Removing (x, y), the edge-connectivity decreases, hence there is an  $\emptyset \neq Z \subset V(G)$  with  $\delta_{G-(x,y)}(Z) \leq k-1$ . This can only be if  $\delta_G(Z) = k$  and Z separates x and y, say  $x \in Z$ ,  $y \in V(G) - Z$ . We may assume  $X \cup Z \neq V(G)$ , otherwise we could consider V(G) - Z instead of Z. By 6.48a,

$$\delta_G(X \cap Z) + \delta_G(X \cup Z) \leq \delta_G(X) + \delta_G(Z).$$

Here  $\delta_G(X \cap Z) \geq k$ ,  $\delta_G(X \cup Z) \geq k$ ,  $\delta_G(X) = \delta_G(Z) = k$ . Therefore, we must have equality throughout, in particular  $\delta_G(X \cap Z) = k$ . This contradicts the minimality of  $X \in W$ . Mader, Math. Ann. 191 (1971) 21–28].

**50.** (a) We have to show that if we contract  $G - (F_1 \cup \ldots \cup F_m)$  we get a graph G' with not more than 2m points. Suppose we will have p points. These points have degree at least k and thus

$$|E(G')| \ge \frac{k \cdot p}{2}.$$

On the other hand, each edge of G' belongs to one of the cuts  $F_1, \ldots, F_m$ , otherwise it should be contracted. Thus

$$|E(G')| \leq \sum_{i=1}^{m} |F_i| = m \cdot k,$$

whence

$$p \leq 2m$$

as stated.

**51.** Let y, z be two neighbors of x. Remove two edges (x, y) and (x, z) and create a new edge (y, z). If the resulting graph G' is not k-edge-connected between two points of V(G) - x, then there is a set  $\emptyset \neq U \subset V(G) - \{x\}$  such that

$$\delta_{G}(U) < k$$
.

Since G and G' are Eulerian, all cuts in them are even; so k is even and  $\delta_{G'}(U)$  as well, whence

$$\delta_{G'}(U) \leq k - 2.$$

Obviously,

$$\delta_{G'}(U) \geq \delta_{G}(U) - 2 \geq k - 2,$$

whence

$$\delta_G(U) = k$$
,  $\delta_{G'}(U) = \delta_G(U) - 2$ .

The latter equality means  $y, z \in U$ .

Now let y be any neighbor of x and U a maximum set such that  $U \subset V(G) - \{x\}$ ,  $y \in U$ ,  $\delta_G(U) = k$ . Since

$$\delta_G(U \cup \{x\}) \ge k = \delta_G(U),$$

it follows that U cannot contain all neighbors of x; let  $z \notin U$  be a neighbor of x. Now if the pair (y, z) does not satisfy the requirements of our problem, there is a set  $V \subset V(G) - \{x\}$  with  $y, z \in V$  and  $\delta_G(V) = k$ . Now by 6.48a,

$$\delta_G(U \cup V) + \delta_G(U \cap V) \leq \delta_G(U) + \delta_G(V) = 2k$$
,

and since  $U \cap V \neq \emptyset$ ,

$$\delta_G(U \cap V) \geq k$$
.

This implies that

$$\delta_G(U \cup V) \leq k$$
.

By the maximality of U, this can only occur if

$$U \cup V = V(G) - \{x\}.$$

Now observe that

$$\delta_G(U-V)+\delta_G(V-U)<\delta_G(U)+\delta_G(V),$$

because in the counting in the solution of 6.48a the edge (x, y) is only counted on the right-hand side. Here

$$\delta_G(U) = \delta_G(V) = k, \quad \delta_G(U - V), \quad \delta_G(V - U) \ge k,$$

a contradiction.

52. We use induction on |V(G)|; if |V(G)| = 2 then we have the initial graph I. Now, suppose  $|V(G)| \ge 3$  and  $x \in V(G)$ . By the preceding result, we can find two points  $y_1, z_1$  adjacent to x such that if we remove  $(x, y_1)$  and  $(x, z_1)$  but connect  $y_1$  to  $z_1$ , then the resulting graph  $G_1$  is still (2k)-edge-connected between any two points of  $V(G) - \{x\}$ . If x is still not isolated, we can find two points  $y_2, z_2$  adjacent to x in  $G_1$  such that  $G_1 - (x, y_2) - (x, z_2) + (y_2, z_2)$  is again 2k-edge connected between any two points of  $V(G) - \{x\}$ , etc. Here

loops can never arise, because a point with loops could be separated from the other points by less than k edges. Thus we get a (2k)-edge-connected, (2k)regular graph G' which arises from G by removing the edges  $(x, y_1), (x, z_1), \ldots$  $(x, y_k)$   $(x, z_k)$  (and then the isolated point x) and creating k new edges  $(y_1, z_1)$ , ...,  $(y_k, z_k)$ . Now G arises from G' by construction II. [A. Kotzig, Doctoral dissertation, Bratislava, 1959; cf. also G. J. Simmons, Infinite and Finite Sets, Coll. Math. Soc. J. Bolyai 10, Bolyai-North-Holland (1974) 1277-1349.

53. We are going to follow the solution of 6.51 as long as we can. For each pair (y, z) of neighbors of x we find a set U such that  $\emptyset \neq U \subset V(G) - \{x\}$ ,  $y, z \in U$ , and

$$\delta_G(U) \leq k+1$$

(because here we do not gain 1 as we did in (1) in the solution of 6.51).

Now let y be a fixed neighbor of x and consider the set H of all sets U like above for all points z. If the assertion is not true, then these cover all neighbors of x. Let  $U_1, \ldots, U_l$  be a minimum number of sets of H which cover all neighbors of x. Since

$$\delta_G(U_i \cup \{x\}) \geq k \geq \delta_G(U_i) - 1,$$

it follows that at most  $\frac{1}{2}d_G(x)$  edges connect x to  $U_i$ . Since x is joined to y

which belongs to all  $U_i$ , it follows that two  $U_i$ 's cannot contain all neighbors

of x. Hence  $l \geq 3$ .

Consider  $U_1$ ,  $U_2$ ,  $U_3$ . Since  $U_1$  could not be omitted from H, there is a neighbor of x which is covered by  $U_1$  but not by  $U_2$  or  $U_3$ . Similarly we find that  $U_2 - U_3 - U_1 \neq \emptyset$  and  $U_3 - U_1 - U_2 \neq 0$ .

Now use a slight sharpening of 6.48c.

$$\begin{split} \delta_{\mathcal{G}}(\boldsymbol{U}_1 - \boldsymbol{U}_2 - \boldsymbol{U}_3) + \delta_{\mathcal{G}}(\boldsymbol{U}_2 - \boldsymbol{U}_3 - \boldsymbol{U}_1) + \delta_{\mathcal{G}}(\boldsymbol{U}_3 - \boldsymbol{U}_1 - \boldsymbol{U}_2) + \\ + \delta_{\mathcal{G}}(\boldsymbol{U}_1 \cap \boldsymbol{U}_2 \cap \boldsymbol{U}_3) &\leq \delta_{\mathcal{G}}(\boldsymbol{U}_1) + \delta_{\mathcal{G}}(\boldsymbol{U}_2) + \delta_{\mathcal{G}}(\boldsymbol{U}_3) - 2. \end{split}$$

(The -2 comes from the observation that the edge (x, y) is counted 3 times on the right-hand side but only once on the left.) Here all terms on the lefthand side are  $\geq k$ , but all terms on the right are  $\leq k+1$ . Hence


$$4k \leq 3(k+1)-2, \quad k \leq 1,$$

a contradiction.

**54.** (a) Let  $\emptyset \neq X \subset V(G)$ . Then

$$\delta_G(X) = \delta_G^{\pm}(X) + \delta_G^{\pm}(V(G) - X) \ge k + k = 2k.$$

(b) We prove the assertion by induction on |V(G)|. If |V(G)| = 2 it is trivial. Let  $|V(G)| \geq 3$ . We may assume G is critically 2k-edge-connected; then by 6.49, there is a point x with  $d_G(x)=2k$ . As in the solution of 6.52, we find a 2k-edge-connected graph G' and k edges  $(y_1, z_1), \ldots, (y_k, z_k) \in E(G')$ such that if we subdivide each of these k edges by a point and identify these new points to a point x, we get G (note that now 6.53 must be applied rather than 6.51). Let  $\vec{G}'$  be an orientation of G' which is k-edge-connected. This induces an orientation  $\vec{G}$  of G: if  $(y_i, z_i)$  is directed from  $y_i$  to  $z_i$  (say), then we direct  $(y_i, x)$  from  $y_i$  to x and  $(x, z_i)$  from x to  $z_i$ . It is easy to see that  $\vec{G}$  is k-edge-connected. [In a stronger form this was proved by Nash-Williams; see B.]



55. The necessity of the conditions given is obvious. Consider the graph shown in Fig. 44. Here the conditions are satisfied since a quadrilateral is 2-edge-connected. On the other hand any (a, a')-path has an edge in common with any (b, b')-path.

56. (a) Take two new points u, u'. Connect u to a and b by  $\alpha$  and  $\beta$  edges, respectively and, similarly, connect u' to a' and b' by  $\alpha$  and  $\beta$  edges, respectively. Condition (\*) now implies that the resulting graph G' is  $(\alpha + \beta)$ -connected between u and u'. Hence, it contains  $\alpha + \beta$  edge-disjoint (u, u')-paths and thus G contains  $\alpha + \beta$  edge-disjoint  $(\{a, b\}, \{a', b'\})$ -paths,  $\alpha$  of them starting at  $\alpha$  and the same number of them ending at a', as stated in the hint. Let  $P_1, \ldots, P_{\alpha+\beta}$  be these paths.

Now  $G - E(P_1) - \dots - E(P_{\alpha+\beta})$  has even degrees, so by 5.13 it has an orientation with equal out-, and indegrees. Orienting the paths  $P_i$  from their endpoint in  $\{a,b\}$  toward their endpoint in  $\{a',b'\}$ , we get a desired orientation of G.

(b) Let  $\vec{G}$  be an orientation of G as in part (a). First, we show that there are  $\alpha$  edge-disjoint (a, a')-paths in  $\vec{G}$ . Let  $X \subseteq V(\vec{G}) - \{a'\}$ ,  $a \in X$ . If  $b, b' \notin X$ , then

$$\delta^+_{\vec{G}}(X) \geq \delta^+_{\vec{G}}(X) - \delta^-_{\vec{G}}(X) = \delta^+_{\vec{G}}(a) - \delta^-_{\vec{G}}(a) = \alpha.$$

We argue similarly, if  $b, b' \in X$  or  $b \in X$ ,  $b' \notin X$ . Suppose  $b' \in X$ ,  $b \notin X$ . Then we have

$$(1) \qquad \delta_{\vec{G}}^{+}(X) - \delta_{\vec{G}}^{-}(X) = \left(\delta_{\vec{G}}^{+}(a) - \delta_{\vec{G}}^{-}(a)\right) + \left(\delta_{\vec{G}}^{+}(b') - \delta_{\vec{G}}^{-}(b')\right) = \alpha - \beta.$$

On the other hand

(2) 
$$\delta_{\vec{G}}^{+}(X) + \delta_{\vec{G}}^{-}(X) = \delta_{G}(X) \ge \alpha + \beta.$$

Summing (1) and (2) we obtain

$$2\delta_{G}^{\pm}(X) \geq 2\alpha$$

which proves that there are  $\alpha$  edge-disjoint (a, a')-paths  $P_1, \ldots, P_{\alpha}$  in  $\vec{G}$ .


Now consider  $G' = \overline{G} - E(P_1) - \ldots - E(P_{\alpha})$ . In this graph  $\delta_{G'}^+(b) - \delta_{\overline{G'}}^-(b) = \beta$ ,  $\delta_{G'}^+(b') - \delta_{\overline{G'}}^-(b') = -\beta$ ,  $\delta_{G'}^+(x) = \delta_{\overline{G'}}(x)$  for  $x \neq b, b'$ .

Therefore by 6.46, G' contains  $\beta$  edge-disjoint (b, b')-paths  $Q_1, \ldots, Q_{\beta}$ . Now  $P_1, \ldots, P_{\alpha}, Q_1, \ldots, Q_{\beta}$  (more exactly, the undirected paths of G corresponding to these directed paths) form a required set of paths. [T.C. Hu, *Integer Programming and Network Flows*, Addison-Wesley, 1969; this proof is due to C. St. J. A. Nash-Williams.]

57. Let G be a 5-connected planar map with a quadrangular face F; let a, c, b, d be the vertices of F in this cyclic order. Then, obviously, any (a, b)-path intersects any (c, d)-path.

Thus it suffices to find a 5-connected planar graph with a quadrangular face.

A little experimentation yields, e.g., Fig. 45.



F1G. 40

Remark: It can be shown [P. Mani–H. A. Jung, see Jung, in: Comb. Structures Appl. Gordon and Breach, 1970 189–191] that all examples are planar. In particular, there are no 6-connected examples. More generally, there exists a function g(n) such that given any 2n points  $a_1, \ldots, a_n, b_1, \ldots, b_n$  in a g(n)-connected graph, there are n vertex-disjoint paths  $P_1, \ldots, P_n, P_i$  connecting  $a_i$  to  $b_i$ .

**58.** First solution. C separates a and b; for any (a, b)-path intersects  $A \cup B$  (even A) and the first point of it belonging to  $A \cup B$  is a point of C.

Similarly, D defined in the hint is a separating set. Hence,  $|C| \ge k$ ,  $|D| \ge k$ . Now

$$C \cap D \subseteq A \cap B$$
;

for if  $x \in C \cap D$ , then there is an  $(a, A \cup B)$ -path  $P_1$  connecting x to a and a  $(b, A \cup B)$ -path connecting x to b. Then  $P_1 \cup P_2$  is a path, which meets  $A \cup B$  in x

only; since each (a, b)-path must meet both A and B, x must belong to both A and B. Thus

$$|A \cap B| \ge |C \cap D| = |C| + |D| - |C \cup D| \ge k + k - |A \cup B| =$$
  
=  $|A| + |B| - |A \cup B| = |A \cap B|$ ,

which shows that we have equality everywhere, in particular |C| = |D| = k.

Second solution. Let  $P_1, \ldots, P_k$  be independent (a, b)-paths (these exist by Menger's theorem), then A(B) contains exactly one point  $a_i(b_i)$  from  $P_i$ .

Let  $c_i$  be the one of  $a_i$  and  $b_i$  nearer a on  $P_i$ .

Now  $C = \{c_1, \ldots, c_k\}$ . Obviously,  $c_i \in C$ . On the other hand, if  $c \in C$ , then  $c \in A \cup B$  and thus  $c \in P_i$  for some i. Assume  $c \neq c_i$ , then c is on the  $(c_i, b)$ -piece of  $P_i$ . By definition, there is an  $(a, c_i)$ -path not meeting A, which gives with the  $(c_i, b)$ -piece of P an (a, b)-path not meeting A, a contradiction.

Thus, |C| = k. The fact that it separates a and b follows easily as above

[cf. L. Lovász, Acta Math. Acad. Sci. Hung. 2 (1970) 365-368].

**59.** Let T be as in the hint. It is enough to show that if T separates G - (x, y), then, in fact, it separates x and y; for then, obviously,  $T \cap V(H)$  separates x and y in H, i.e. H - (x, y) has a (k - 1)-element separating set.

Now if, to the contrary, x, y are in the same component of G - (x, y) - T, then, adding (x, y), we see that G - T is still disconnected and this is impossible.

**60.** (a) 1° In G-C, there are no edges between  $G_1\cap G_3$  and the remainder of the graph, which is non-empty. Hence C is a separating set and since  $G_1\cap G_3$  is a proper subgraph of  $G_1$  (as  $B\cap V(G_1)\neq\emptyset$ ), C cannot be a k-element separating set.

 $2^{\circ}$  Let  $x \in C$ . Then there is an  $(a, A \cup B)$ -path P connecting a to x. If  $x \notin A \cap B$ , then  $x \notin A$  or  $x \notin B$ . If  $x \notin A$  (say), then P does not intersect A, i.e. x is in the same component  $G_1$  of G - A as a.

 $3^{\circ}$  If  $b \in G_2 \cap G_4$ , then by 6.58 C has cardinality k.

40

$$|V(G_1)| > |B \cap V(G_1)| + |V(G_1 \cap G_4)|,$$

while

$$|V(G_4)| = |A \cap V(G_4)| + |V(G_1 \cap G_4)|.$$

Now since

$$|B \cap V(G_1)| + |A \cap B| + |A \cap V(G_3)| \ge |C| > k$$

and

$$|A \cap V(G_4)| + |A \cap B| + |A \cap V(G_3)| = |A| = k$$

it follows that

$$|B \cap V(G_1)| > |A \cap V(G_4)|,$$

whence

$$|V(G_1)| > |V(G_4)|,$$

which contradicts the minimality of  $G_1$  (Fig. 46).




Fig. 46

To show the second statement, assume first that  $V(G_2) \subseteq B$ . Since  $|V(G_1)| \leq |V(G_2)|$ , this immediately implies that  $|V(G_1)| \leq \tilde{k}/2$ . Secondly, assume that  $V(G_2) \subseteq B$ . Let, e.g.  $G_2 \cap G_3 \neq \emptyset$ . Since  $(A \cap V(G_3)) \cup$  $\bigcup (A \cap B) \bigcup (B \cap V(G_2))$  separates G, it follows that

$$|A \cap V(G_3)| + |A \cap B| + |B \cap V(G_2)| \ge k = |B| =$$
  
=  $|V(G_1)| + |A \cap B| + |B \cap V(G_3)|,$ 

whence

$$|V(G_1)| \leq |A \cap V(G_3)|.$$

If  $G_2 \cap G_4 \neq \emptyset$  then similar argument yields that

$$|V(G_1)| \le |A \cap V(G_4)|,$$

and hence

$$|V(G_1)| \le \frac{1}{2}|A| = \frac{k}{2}.$$

If  $G_2 \cap G_4 = \emptyset$  then similar argument yields that

$$|\mathit{V}(G_4)| \leq |\mathit{B} \cap \mathit{V}(G_2)|,$$

and so

$$|V(G_1)| \le |V(G_4)| \le |B \cap V(G_2)| \le |B| - |V(G_1)| = k - |V(G_1)|,$$

whence the assertion follows again. [W. Mader, Arch. Math. 22 (1971) 333-336; M. E. Watkins, J. Comb. Th. 8 (1970) 23-29.

(b) Obviously, G has no multiple edges. We may assume G is not a (k+1)clique. Then  $G_1$  of (a) exists. Assume  $G_1$  has at least two points, then  $G_1$  has an edge e = (x, y). G - e has a (k - 1)-element separating set T. Then G - T is




Fig. 47

connected but G-T-e is not, whence G-T-e has two components connected by e. Suppose the component containing x has more than one point, then  $T \cup \{x\}$  is a k-element separating set of G, having a point x in common with  $G_1$  but not containing another point, y of it. This is impossible by (a). Thus  $G_1 = \{x\}$  and it follows that x has degree k. [R. Halin, J. Comb. Theory 7 (1969) 150–154.]

- **61.** (a) Let  $\{x, y\}$  be a separating set of G e. Since  $G \{x, y\}$  is connected, e connects different components of  $G e \{x, y\}$ , hence a and b are separated by x, y in G e.
- (b) Let  $\{u, v\}$  be a separating set of G/e. We claim one of u, v is the point e is contracted onto. For otherwise  $G/e \{u, v\}$  would arise by contraction of an edge from the connected graph  $G \{u, v\}$  and would be connected. Now if v is this point, then  $\{a, b, u\}$  is a separating set. To show it separates x and y, consider a component G' of  $G \{a, b, u\}$ . Since G is 3-connected, this component is connected to each of a, b, u and, thus there is an (a, b)-path  $P_1$  with inner points in G'.  $P_1$  contains one of x and y, x say. Considering another component G'' of  $G \{a, b, u\}$ , the same argument shows that y must be in G''.
- (c) Assume the component  $G_1$  of  $G-e-\{x,y\}$  containing a has a point  $c\neq a$ . Since G is 3-connected, there is a (c,x)-path P in  $G-\{a,y\}$ . P does not go through b and u, because b, u and c are separated by a, x, y (even by e, x, and y); hence P is a (c,x)-path in  $G-\{a,b,u\}$ . Similarly, there is a (c,y)-path Q in  $G-\{a,b,u\}$ ; however,  $P\cup Q$  then gives an (x,y)-path in  $G-\{a,b,u\}$ , which contradicts (b) (Fig. 47).

The degree of a is now 3, a contradiction.

**62.** First we prove the assertion of the hint. Assume G-T is disconnected, where  $|T| \leq k-1$ . If T does not contain either of the endpoints x, y of e,

then (G-T)/e = G/e - T is disconnected, which is a contradiction, because G/e is k-connected. If  $x, y \in T$ , then G/e can be cut even by the (k-2)-element image of T. Therefore, we may assume that  $x \in T$ ,  $y \notin T$  (say).

G/e - T can be obtained from G - T by removing y. G/e - T is connected while G - T is not, which means y was a component of G - T. Hence T contains all the neighbors of y, which is impossible, because y has degree at least k.

Now the result follows easily. G/e is 3-connected by the preceding problem. Suppose it is not minimal, then G/e - f = (G - f)/e is still 3-connected for some  $f \in E(G)$ . Since the endpoints of e have degree at least 3 in G - f, it follows by the above that G - f is 3-connected, contrary to the assumption.

**63.** Let C be the circuit under consideration. If |V(C)| = 2 the statement is void as an edge-critically 3-connected graph cannot have multiple edges.

Assume  $|V(C)| \geq 3$ . If C has at most one point of degree 3, it has an edge e whose endpoints have degree at least 4. Then, by 6.62, G/e is a critically 3-connected graph, and the circuit C is mapped by the contraction of e onto a shorter circuit C' which still has at most one point of degree 3. This is a contradiction.

This result — with a more complicated proof — remains valid for critically k-connected graphs. [See W. Mader, Arch. Math. 23 (1972) 219–224.]

**64.** The statement of the hint follows just like (b) in the solution of 6.61. Take a circuit C through  $b_1$  and  $b_2$  in G-a (which is 2-connected so that such a circuit exists by 6.32), then one of the  $(b_1, b_2)$ -arcs of C contains  $u_3$ , the other one contains  $b_3$ . Thus C is a circuit through  $b_1$  and  $b_3$  and hence by an argument similar to the one above, it also contains  $u_2$ . Thus each circuit of G-a containing two of  $b_1, b_2, b_3$  also contains the third one as well as  $u_1, u_2$ , and  $u_3$ . Further  $u_i$  separates  $b_{i-1}$  and  $b_{i+1}$ .

It follows that each  $(u_1, u_2)$ -path must contain either  $b_3$  or both  $b_1$  and  $b_2$ ; otherwise we could replace an arc of C by it and get a circuit violating the above assertion. Thus  $b_3$  and  $b_2$  separate  $u_1$  and  $u_2$  in G - a, but then they also separate  $u_1$ ,  $u_2$  in G, a contradiction (Fig. 48). [Cf. W.T. Tutte, Connectivity in

Graphs, Toronto University Press, Toronto, 1966.]




Fig. 48

- **65.** Construct G as in the hint. I. G is 3-connected. For let us remove 2 points of it. If both are inner points of T, then, since the points of T are of degree at least 3, each component of the remainder of T meets C and this "connects" them. The same argument works if a point of T and a point of C are removed. Finally, if we remove two points of C, the remainder of T is connected and meets all pieces of the remainder of C, thus the resulting graph is connected again.
- II. First suppose  $e \in E(C)$ , then G-e is at most 2-connected, because the endpoints of e have degree 2 in it. Secondly suppose  $e \in E(T)$ . Then T-e has exactly two components  $T_1, T_2$ . The endpoints of  $T_i$  form an arc  $A_i$  of C (the precise proof of this fact is left to the reader). Thus, there are two edges  $f_1, f_2$  connecting  $A_1$  and  $A_2$ .  $G-e-f_1-f_2$  is disconnected, showing that G-e is not even 3-edge-connected.
- III. Choose the tree T such that some point x has distance 1000 from all endpoints and all inner points have degree 1000.
- **66.** We use induction on k. Let C be a circuit passing through  $x_1, \ldots, x_{k-1}$ . If  $x_k \in V(C)$  we are finished; assume  $x_k \notin V(C)$ . Now we distinguish two cases.
- (a) Assume there are k ( $x_k$ , C)-paths which have only  $x_k$  in common. The points  $x_1, \ldots, x_{k-1}$  divide C into k-1 arcs, thus one of these arcs (including its endpoints) contains two endpoints of the paths  $P_1, \ldots, P_k$ . Now adding these two paths to C, but removing the arc between their endpoints (which does not contain any  $x_i$ ,  $1 \le i \le k-1$ ), we get a circuit through  $x_1, \ldots, x_k$ .
- (b) If there are not k ( $x_k$ , C)-paths which have no point in common except  $x_k$ , then by 6.41 there is a set X, |X| = k 1,  $x_k \notin X$ , which meets each ( $x_k$ , C)-path. Since G X is connected, each point of C X can be connected to  $x_k$  in G X by a path. This is a contradiction except when  $X = V(C) = \{x_1, \ldots, x_{k-1}\}$ . In this latter case an argument similar to the one above shows that there are k 1 ( $x_k$ , C)-paths, having only  $x_k$  in common. These paths connect  $x_k$  to  $x_1, \ldots, x_{k-1}$ . Adding two of them, which connect  $x_k$  to neighboring points of C, but removing the edge between their endpoints we get a desired circuit again. [G. Dirac; see B.]
- **67.** Suppose G is a 3-connected graph and  $e_1$ ,  $e_2$ ,  $e_3$  are edges of G not on a circuit. We show first that there is a circuit C containing two of them, which does not meet the third one.

Let  $C_1$  be a circuit through  $e_1$  and  $e_2$ . If  $e_3 = (x, y)$  has no endpoint on  $C_1$ , we have the desired circuit. If  $x \in V(C_1)$ , but  $y \notin V(C_1)$ , consider an  $(y, C_1 - x)$ -path P of G - x; let this path P end at  $z \in V(C_1)$ . Then  $e_3$ , P and one of the (x, z)-arcs of  $C_1$  give a desired circuit. Finally, if  $x, y \in V(C_1)$ , then  $e_3$  and one of the (x, y)-arcs of  $C_1$  yield C. Now  $C - e_1 - e_2 = P_1 \cup P_2$ , where  $P_1, P_2$  are disjoint paths. Consider all  $(\{x, y\}, P_1)$  paths of  $G - V(P_2)$ . No two of them are disjoint, since this would yield a circuit containing  $e_1, e_2, e_3$ . Thus, there is a point u meeting each of them. Similarly, there is a point meeting each  $(\{x, y\}, P_2)$ -path of  $G - V(P_1)$ . Then  $\{u, v\}$  meets all  $(\{x, y\}, C)$ -paths. Since G is 3-connected, this is only possible if  $\{u, v\} = \{x, y\}$ , say u = x, v = y. Thus, any  $(\{x, y\}, C)$ -path which ends on  $P_1$   $(P_2)$  starts from x (y).


If we show that  $G - \{e_1, e_2, e_3\}$  does not contain any (x, y)-path we are

through. Assume indirectly that there is such a path T.

Assume first that T does not meet C. Let Q be a (T,C)-path of  $G-\{x,y\}$  ending in  $P_1$  (say); then Q and a suitable piece of T would give an  $(\{x,y\},C)$ -path starting from y, which is impossible. We reach a similar conclusion when T meets  $P_1$  only. Thus the only case we have to deal with is when T meets both  $P_1$  and  $P_2$ . Then it contains a  $(P_1,P_2)$ -path R, different from  $e_1$  and  $e_2$ . Let  $a \in P_1$ ,  $b \in P_2$  be the endpoints of R. Since  $G-\{a,b\}$  is connected, there is an  $(\{x,y\},C-\{a,b\})$ -path  $Q_1$  in it. Let the endpoints of  $Q_1$  be e.g. x and  $z_1 \in P_1$ . Let  $Q_2$  be a (y,C)-path in G-x, then obviously, the endpoint  $z_2 \neq y$  of  $Q_2$  is on  $P_2$ .  $Q_1$  and  $Q_2$  are disjoint for otherwise we would get on (x,y)-path of  $G-\{e_1,e_2,e_3\}$  disjoint from C.

One of the  $(z_1, z_2)$ -arcs of C does not contain a. This arc forms with  $Q_1, Q_2$  and  $e_3$  a circuit  $C_0$ , containing  $e_3$  and one of  $e_1$  and  $e_2$ ,  $e_2$  say. Now there are  $(e_1, C_0)$ -paths connecting the same endpoint of  $e_1$  to both components of  $C_0 - \{e_2, e_3\}$ ,

a contradiction (Fig. 49).



**68.** Using induction on k, we may assume there is an (a, b)-path P which passes through  $x_1, \ldots, x_{k-1}$  and assume  $x_k \notin P$ . It is impossible to cover all  $(x_k, P)$ -paths by k points different from  $x_k$  since, if X were a k-element set meeting all of them  $(x_k \notin X)$ , then  $x_k$  and  $P - X \neq \emptyset$  would belong to different components of G - X. Hence by 6.41, there are k + 1  $(x_k, P)$ -paths  $Q_1, \ldots, Q_{k+1}$ , having only  $x_k$  in common. Since  $x_1, \ldots, x_{k-1}$  divide P into k pieces, one of these meets two  $Q_j$ 's. Then removing the subpath connecting these two endpoints but adding the two  $Q_j$ 's, we get an (a, b)-path through  $x_1, \ldots, x_k$  [M.D. Plummer, in:  $Proc.\ 2^{nd}\ Louisiana\ Conf.\ on\ Comb.\ (Univ.\ of\ Manitoba\ Press,\ Winnipeg, 1971)\ 458-472$ ].

**69.** Let C be the boundary circuit of a face F. Let B be any bridge of C. We show, following the hint, that B contains all points of C. Assume  $x \in V(C)$ ,  $x \notin V(B)$ . Let y, z be the two points of  $B \cap C$  next to x on C. Then the other (y, z)-arc of C has an inner point; this is trivial, if B is a chord and follows from the fact that B has at least 3 points on C, if B has an inner point. Since  $G - \{y, z\}$  is connected, it has a path P connecting the two (y, z)-arcs of C. Let Q be a (y, z)-path in B. Both P and Q run outside F (as F is a face) and so, they inter-




Fig. 50

sect by planarity. This means, however, that P belongs to B, a contradiction to the definition of y and z (Fig. 50).

The statement just shown implies that C has no chords. Suppose  $B_1$ ,  $B_2$  are two bridges. Then  $C + B_1$  divides the exterior of F into regions having at most 2 boundary points on C and  $B_2$  lies inside one of these (as it does not meet  $B_1$ ).

But  $B_2$  has at least 3 points on C, a contradiction.

Since the result of this problem characterizes the boundaries of faces of a planar 3-connected map independently of the embedding, it follows that every isomorphism between two planar 3-connected maps preserves faces. By the Schoenflies Theorem in plane topology it follows that every isomorphism between two 3-connected maps on the sphere is induced by a homeomorphism of the sphere onto itself [H. Whitney, Trans. Amer. Math. Soc. 34 (1932) 339-362].

70. Let C be defined as in the hint; we show it has no bridge other than B. As in the previous solution, we show first that each point of C is on B. Suppose  $x \in V(C), x \notin V(B)$ . Let y, z be the two points of  $\overline{B}$  next to x on C. As in the previous solution, we find a path P joining the two (y, z)-arcs of C in G - y - z. Let a, b be the endpoints of P and let R be an (a, b)-arc of C. Then  $P \cup R$  is a circuit such that the bridge of  $P \cup R$  containing f properly contains B, a contradiction.

So every point of C is in B. Suppose  $B_1$  is another bridge of C. Let c, d be two points of  $B_1$  on C. Then a (c, d)-arc of C has an inner point (since  $B_1$  is either a chord or has at least 3 points on C). Denote by R the other (c, d)-arc of C and denote by P a (c, d)-path of  $B_1$ , then  $R \cup P$  is a circuit through c, dand the bridge of  $R \cup P$  containing f properly contains B. A contradiction again [W. T. Tutte, Proc. London Math. Soc. 13 (1963) 743-767].

71. With the notation of the hint, let  $P_0$  be an (a, b)-path, formed by some edges of  $e_1, \ldots, e_k$ . Since  $\{e_1, \ldots, e_{k-1}\}$  does not contain any (a, b)-path,

there is an (a, b)-cut  $C_0$  such that  $C_0$  contains none of  $e_1, \ldots, e_{k-1}$ . Hence,  $C_0 \cap P_0 = \{e_k\}$ . Moreover, by definition,

$$\min_{e \in P_{\bullet}} v(e) = v(e_k), \max_{e \in C_{\bullet}} v(e) = v(e_k).$$

On the other hand, let C be any (a, b)-cut, then  $C \cap P_0 \neq \emptyset$ , say  $e_i \in C \cap P_0$ ,  $i \leq k$ . Then

$$\max_{e \in C} v(e) \ge v(e_i) \ge v(e_k)$$

and similarly

$$\min_{e \in P} v(e) \leq v(e_k)$$

for any (a, b)-path P, whence

$$\max_{P} \min_{e \in P} v(e) = v(e_k) = \min_{C} \max_{e \in C} v(e)$$

[cf. J. Edmonds, D.R. Fulkerson, J. Comb. Theory 8 (1970) 299].

**72.** I. Suppose  $\varphi$  is a function satisfying (\*) and  $P = (a, x_1, \ldots, x_k, b)$  is any (a, b)-path. Then

$$egin{aligned} u(P) &= v(a,x_1) + v(x_1,x_2) + \ldots + v(x_k,b) \geq \ &\geq arphi(x_1) + igl(arphi(x_2) - arphi(x_1)igr) + \ldots + igl(arphi(b) - arphi(x_k)igr) = arphi(b). \end{aligned}$$

II. Thus it suffices to construct an (a, b)-path P and a function  $\varphi$ , satisfying (\*), such that  $u(P) = \varphi(x)$ . We define  $\varphi$  step by step. Put  $\varphi(a) = 0$ . Suppose  $\varphi$  is already defined for a set  $S \subset V(G)$ ,  $a \in S$  and has the property that for each point  $x \in S$  there is an (a, x)-path in S of value  $\varphi(x)$ . Consider all edges (x, y) with  $x \in S$ ,  $y \notin S$  and choose one for which  $\varphi(x) + v(x, y)$  is minimal. Set  $\varphi(y) = \varphi(x) + v(x, y)$ . This extends the domain of  $\varphi$ . We show that the property (\*) is preserved. Let (z, u) be any edge spanned by  $S \cup \{y\}$ , we have to show

$$\varphi(u) - \varphi(z) \leq v(z, u).$$

If  $u, z \in S$  this is obvious. If  $z \in S$ , u = y then, by the choice of (x, y),

$$\varphi(y) = \varphi(x) + v(x, y) \le \varphi(z) + v(z, u).$$

So let  $u \in S$ , z = y. In this case we claim  $\varphi(u) = \varphi(y)$ , which will prove the assertion. For this we show, by induction, that a point chosen later has no smaller  $\varphi$ -value than a point chosen earlier. Suppose indirectly that  $\varphi(y) < \varphi(u)$ . Then  $\varphi(x) \leq \varphi(y) < \varphi(u)$  and hence, by induction, x was chosen earlier than u. But then instead of u one should have chosen y. So  $\varphi$  satisfies (\*). Moreover, by induction, there is an (a, x)-path with value  $\varphi(x)$  and adding (x, y) to this path we get an (a, y)-path with value  $\varphi(y)$ .

In particular, we have an (a, b)-path with value  $\varphi(b)$  when  $\varphi$  is completely defined. This proves the assertion [For most problems of the rest of this para-

graph see FF or Hu].

**73.** Since f is an (a, b)-flow, the only non-zero term in the expression given in the hint is

$$\sum_{e=(a,y)} f(e) - \sum_{e=(y,a)} f(e),$$

which is the value of f. On the other hand, each edge e spanned by S occurs twice with opposite signs, the edges of C occur with positive sign, the edges of  $C^*$  with negative sign. Thus, the value of f is

$$\sum_{e \in C} f(e) - \sum_{e \in C^*} f(e).$$

74. We follow the hint. Assume there is an (a, b)-path P in  $G_1$ . Let

$$\epsilon = \min \sum_{e \in P} v_0(e) > 0$$
.

Now set for  $e \in E(G)$ ,

$$f_1(e) = \begin{cases} f(e) \, + \, \varepsilon & \text{if} \quad e \in E(P), \\ f(e) \, - \, \varepsilon & \text{if} \quad e' \in E(P), \\ f(e) & \text{otherwise.} \end{cases}$$

Then, obviously,  $f_1(e)$  is an (a, b)-flow of value  $w(f) + \varepsilon$ , a contradiction. Thus,  $G_1$  is not connected between a and b. Hence, by 6.9 there is an  $S \subseteq V(E)$  such that  $a \in S$ ,  $b \notin S$  and the cut C of G determined by S satisfies

$$f(e) = v(e)$$
 if  $e \in C$ ,  
 $f(e) = 0$  if  $e \in C^*$ .

Hence, by 6.73,

$$w(f) = \sum_{e \in C} f(e) - \sum_{e \in C^*} f(e) = \sum_{e \in C} v(e).$$

Thus

$$\max_{f} w(f) \ge \min_{e} \sum_{e \in C} v(e).$$

On the other hand, if f is an arbitrary (a, b)-flow with  $f \leq v$ , and C is any (a, b)-cut, we have

$$w(f) = \sum_{e \in C} f(e) - \sum_{e \in C^*} f(e) \le \sum_{e \in C} f(e) \le \sum_{e \in C} v(e),$$

which proves the assertion. [C.R. Ford-D.R. Fulkerson; FF.]

75. (a) Consider the network in Fig. 51, where  $\alpha$  is the positive root of  $\alpha^3 + \alpha - 1 = 0$ , all capacities are 1 and the values of a flow  $f_0$  are indicated.




Fig. 51

We start with the flow in Fig. 51, and use the paths (atwvub), (atuvwb), (avutwb), (avutub), and repeat this cyclically. We consider the change of values of  $f_k$  on the quadrilateral (uvwt), since the other edges have a large enough capacity not to interfere.

We claim that after 4m steps we will have the flow shown in Fig. 52, and the next 4 increases will be  $\alpha^{4m+1}$ ,  $\alpha^{4m+2}$ ,  $\alpha^{4m+3}$ ,  $\alpha^{4m+4}$ . Using induction on m, it suffices to consider four consecutive steps; so Fig. 53 proves the assertion; the flow values on the edge (a, t) will be partial sums of

$$\alpha - \alpha^2 + \alpha^4 - \alpha^6 + \alpha^6 - \alpha^{10} + \dots$$




Fig. 52




Fig. 53

so they will be strictly between 0 and 1; similarly for the other edges at a and b. Now the value of these flows converges to

$$lpha^2+(lpha+lpha^2+\ldots+lpha^k+\ldots)=lpha^2+rac{lpha}{1-lpha}=1+lpha+2lpha^2<4.$$

On the other hand, the maximum flow clearly has value 4.

(b) Let us form a new digraph  $G_0$  as follows: we orient all edges of  $P_k$  and  $P_{k+1}$  toward b and take the union of the resulting directed paths  $P_k$  and  $P_{k+1}$ ; but we remove those pairs (x, y) (y, x) of edges where  $(x, y) \in P_k$  and  $(y, x) \in P_{k+1}$ . Those common edges of  $P_k$  and  $P_{k+1}$ , which are passed in the same sense are taken, however, twice.

Now  $G_0$  has equal outdegrees and indegrees except at a and b. Thus by 6.46 there are two edge-disjoint (directed) (a, b)-paths  $Q'_1$  and  $Q'_2$  in  $G_0$ . Let  $Q_1$ 

and  $Q_2$  be the two corresponding undirected (a, b)-paths in G.

Now observe that  $Q_1$  and  $Q_2$  have the property that, after the k<sup>th</sup> step, they come into consideration together with  $P_k$ , i.e. they have the property that their edges e directed toward a satisfy  $f_k(e) > 0$  and those directed toward b satisfy  $f_k(e) < v(e)$ . This follows immediately from the observation that any edge of  $Q_i$  is an edge of  $P_k$  or  $P_{k+1}$  and  $Q_i$  passes it in the same sense.

Thus by the minimality of  $P_k$ ,

$$|E(Q_i)| \geq |E(P_k)|$$
.

Since

$$|E(Q_1)| + |E(Q_2)| \le |E(P_k)| + |E(P_{k+1})|$$
,

it follows that

$$|E(P_k)| \leq |E(P_{k+1})|.$$

(c) Suppose that contrary to the assertion formulated in the hint,  $|E(P_{k+l})| = |E(P_k)|$  ( $\geq$  follows by (b)). Let l be minimal. Then by (b)

$$|E(P_{k+1})| = \ldots = |E(P_{k+l-1})| = |E(P_k)|$$

and the paths  $P_{k+1}, \ldots, P_{k+l-1}$  do not use any edge of  $P_k$  or  $P_{k+l}$  oppositely to  $P_k$  or  $P_{k+l}$  by the minimality of l. Hence it follows that those edges e of  $P_{k+l}$  not on  $P_k$  already satisfy the requirement at the k<sup>th</sup> step:  $f_k(e) > 0$  if e is directed toward e (on e) and e0 of e1 is directed toward e2.

Now we get, by an argument similar to the one in (b), two (a, b)-paths  $Q_1, Q_2$  such that  $Q_1 \cup Q_2 \subset P_k \cup P_{k+l}, Q_1 \cap Q_2 \subseteq P_k \cap P_{k+l}$ , and  $Q_1, Q_2$  avoid those edges which are passed by  $P_k$  and  $P_{k+l}$  oppositely; moreover,  $Q_1, Q_2$  pass each edge in the same direction as  $P_k$  or  $P_{k+l}$ . By the above,  $Q_1, Q_2$  were candidates for  $P_k$  and hence,

$$|E(Q_i)| \geq |E(P_k)|$$
.

Moreover, since at least one edge is used by  $P_k$  and  $P_{k+1}$  in opposite directions,

$$|E(Q_1)| + |E(Q_2)| \le |E(P_k)| + |E(P_{k+l})| - 2.$$

Thus

$$|E(P_{k+l})| \ge |E(P_k)| + 2.$$

This proves the assertion of the hint. To get (c) observe that if  $P_k, P_{k+1}, \ldots, P_{k+l}$  do not pass any edge in opposite directions, then the number of those edges e in  $P_1 \cup \ldots \cup P_{k+l}$  which satisfy  $f_{k+i}(e) = 0$  and are directed toward a (on the path  $P_{k+j}$  containing them), or  $f_{k+i}(e) = v(e)$  and are directed toward b, increases at each step. Hence

$$l \le |E(G)| < n^2.$$

Thus, in every  $n^2$  steps the length of  $P_k$  increases by at least two. Since it is less than n, the number of steps does not exceed

$$n^2 \cdot \frac{n}{2} = \frac{n^3}{2}.$$

[J. Edmonds and R.M. Karp; see Hu, Ch. 8.]

76. If there is a function as required, we have

$$\sum_{x} u(x) = \sum_{x} \left\{ \sum_{y} f(x, y) - \sum_{y} f(y, x) \right\}$$

and here each edge occurs twice with opposite signs. This gives a necessary condition

$$\sum_{x} u(x) = 0.$$

Moreover, define

$$f(a_0, x) = u(x),$$
  
 $f(x, b_0) = -u(x).$ 

Then we obtain an  $(a_0, b_0)$ -flow, satisfying  $f(e) \leq v(e)$  and of value

$$V = \sum_{u(x)>0} u(x) = \sum_{u(x)<0} (-u(x)).$$

Since such an  $(a_0, b_0)$ -flow exists, we must have

$$\sum_{e \in C_1} v(e) \ge V$$

for any  $(a_0, b_0)$ -cut  $C_1$  of  $G_1$ . Now let  $C_1$  be determined by  $S \cup \{a_0\}$ ,  $S \subseteq V(G)$  and let C be the cut of G determined by S, then

$$\sum_{e \in C_1} v(e) = \sum_{e \in C} v(e) + \sum_{\substack{x \in S \\ u(x) < 0}} \left( -u(x) \right) + \sum_{\substack{x \in V(G) - S \\ u(x) > 0}} u(x).$$

Thus.

(2) 
$$\sum_{e \in C} v(e) \ge \sum_{x \in S} u(x).$$

Conversely, if (1) and (2) are satisfied, then  $G_1$  has a flow  $f(e) \leq v(e)$  of value V by 6.77. This flow has, necessarily, f(e) = v(e) for the edges adjacent to  $a_0$  or  $b_0$ , and consequently gives the required function, when restricted to G. [D. Gale; see FF.]

77. (a) Let us substitute v(e) parallel edges for each edge e and let  $G_1$  be the resulting graph. Let C be an (a, b)-cut of G and  $C_1$  the (a, b)-cut of  $G_1$ , determined by the same set. Then

$$|C_1| = \sum_{e \in C} v(e).$$

Thus, putting

$$V = \min_{C} \sum_{e \in C} v(e),$$

we get  $|C_1| \ge V$  and hence, by Menger's theorem, we find V edge-disjoint (a, b)-paths in  $G_1$ . Let f(e) be the number of edges parallel to e used by these paths, then f(e) is an (a, b)-flow of value V and with integral entries.

We remark that the proof of 6.77 constructs an integer valued flow whenever

v is integer valued.

(b) If v(e) is an integer for each e, (a) solves the problem. If v(e) is rational for each e, consider  $s \cdot v(e)$ , where s is the least common multiple of denominators of the v(e)'s, this reduces the problem to the integral case. Finally, if v(e) is arbitrary, take a sequence  $v_i \to v$  with rational values  $v_i(e)$ . Since there are only finitely many (a, b)-cuts, we may assume the minimum

$$m_i = \min_{C} \sum_{e \in C} v_i(e)$$

is attained for the same cut C. For each i, there is an (a, b)-flow  $f_i$  with

$$0 \le f_i(e) \le v_i(e); \qquad w(f_i) = m_i.$$

As the graph is finite, we may assume  $f_i(e)$  converges for any e. The limit

$$f(e) = \lim_{i \to \infty} f_i(e)$$

is an (a, b)-flow of value

$$\lim_{i\to\infty}m_i=\sum_{e\in C}v(e).$$

This proves the non-trivial part of the Max-Flow-Min-Cut Theorem.

78. Let f' be as in the hint. We show  $w(f') = w_1$ . Assume  $w(f') > w_1$ . Let G' be the subgraph of G determined by those edges e of G for which f'(e) > 0. G' is connected between a and b; otherwise we could find an (a, b)-cut G of G which would contain no edge of G', i.e. we would have

$$\sum_{e \in C} f'(e) = 0.$$

On the other hand, the solution of 6.74 yields

$$\sum_{e \in C} f'(e) \ge w(f') > w_1 \ge 0,$$

a contradiction. Thus, we have an (a, b)-path P with f'(e) > 0 for each  $e \in P$ . Subtract 1 from the flow-value of each edge. Then w(f') is decreased but is still  $\geq w_1$ , and f' is an (a, b)-flow  $\leq f$ . So  $w(f') = w_1$ .

Now let  $f_1=f', f_2=f-f'.$  Then by the above,  $f_1, f_2$  are (a,b)-flows and  $0 \le f_1 \le f, \qquad 0 \le f_2 \le f,$   $w(f_1)=w_1, \ w(f_2)=w(f)-w(f_1)=w_2.$ 

## § 7. Factors of graphs

1. First, let  $F_1$  be a system of  $\varrho(G)$  edges which cover all points. Since removing any edge from  $F_1$  yields an uncovered point, each edge in  $F_1$  has an endpoint which has degree one in  $\langle V(G), F_1 \rangle$ . This implies that  $F_1$  consists of disjoint stars.

The number of components of  $\langle V(G), F_1 \rangle$  is, obviously,  $|V(G)| - |F_1| = |V(G)| - \varrho(G)$ . Select an edge from each component of  $\langle V(G), F_1 \rangle$ . Then we have  $|V(G)| - \varrho(G)$  independent edges, whence

(1) 
$$\nu(G) \ge |V(G)| - \varrho(G), \quad \nu(G) + \varrho(G) \ge |V(G)|.$$

On the other hand, let  $F_2$  be a system of  $\nu(G)$  independent edges. Select one edge from each of the  $|V(G)| - 2\nu(G)$  points not covered by  $F_2$ . Together with  $F_2$ , these edges cover all points of G and their number is

$$v(G) + (|V(G)| - 2v(G)) = |V(G)| - v(G),$$

whence

(2) 
$$\varrho(G) \le |V(G)| - \nu(G), \quad \nu(G) + \varrho(G) \le |V(G)|.$$

- (1) and (2) yield the desired equality. [T. Gallai, Ann. Univ. Sci. R. Eötvös, Sectio Math. 2 (1959) 133–138.]
- **2.** First solution. Let  $\{A, B\}$  be a bicoloration of G, i.e. suppose  $V(G) = A \cup B$ ,  $A \cap B = \emptyset$  and each edge of G joins A to B. Consider two new points a, b. Join a to all points of A and b to all points of B. Let G' be the resulting graph. Now observe
- (i) a set  $S \subseteq V(G)$  separates a and b in G' if and only if S covers all edges of G (ii) the maximum number of independent (a, b)-paths is v(G); for any v(G) independent edges of G yield v(G) independent (a, b)-paths and conversely, any k independent (a, b)-paths contain k independent edges of G.

Since, by Menger's theorem, the minimum cardinality of sets separating a and b is equal to the maximum number of independent (a, b)-paths, (i) and (ii) prove  $\nu(G) = \tau(G)$ . Since by 7.1,  $\nu(G) + \varrho(G) = |V(G)|$  and, obviously,  $\alpha(G) + \tau(G) = |V(G)|$ , the other equation of this problem follows immediately.

Second solution. We show  $\nu(G) = \tau(G)$ . Since obviously,  $\nu(G) \le \tau(G)$ , it suffices to show that  $\nu(G) \ge \tau(G)$ .

Let G' be a minimal subgraph of G such that  $\tau(G') = \tau(G)$ . We show G' consists of disjoint edges. This will be sufficient since the number of these edges must be  $\tau(G') = \tau(G)$ , thus  $\nu(G) \geq \tau(G)$ .

Suppose indirectly that there is a point x adjacent to two edges  $e_1 = (x, y_1)$ ,  $e_2 = (x, y_2)$  of G'. By the minimality of G', there is a set  $S_i$  with  $|S_i| = \tau(G) - 1$ , which covers all edges of  $G' - e_i$  (i = 1, 2). Obviously,  $x, y_i \notin S_i$ .

but  $y_1 \in S_2$ ,  $y_2 \in S_1$ . Set  $R = S_1 \cap S_2$ , |R| = p,  $\tau(G) = t$ , and let  $G_1$  denote the subgraph of G' induced by  $(S_1 - R) \cup (S_2 - R) \cup \{x\}$ .

Since  $G_1$  is a subgraph of G, it is bipartite. Let  $\{A, B\}$  be a bicoloration of  $G_1$ 

7.3

and, say,  $|A| \leq |B|$ . Then

$$|A| \leq \left[\frac{|(S_1 - R) \, \cup \, (S_2 - R) \, \cup \, \{x\}|}{2}\right] = \left[\frac{2(t - 1 - p) + 1}{2}\right] = t - p - 1.$$

We claim that  $A \cup R$  covers all edges of G'. In fact, let  $e \in E(G')$ . If e has an endpoint in R, we are done. We show that, if e is not covered by R, then it belongs to  $G_1$ . This is obvious if  $e = e_i$ , since  $x, y_i \in V(G_1)$ . If  $e \neq e_i$ , then both  $S_1 - R$  and  $S_2 - R$  have to cover e and hence, e has to join  $S_1 - R$  to  $S_2 - R$ . Thus e has both endpoints in  $V(G_1)$  again. Since A covers all edges of  $G_1$ , it covers e.

Thus,  $\tau(G') \leq |A \cup R| = t - p - 1 + p = t - 1$ , which contradicts the definition of G' [D. König; see any textbook on graph theory].

3. I. Suppose some point x of F is joined to a point  $y \in B_1$ . Obviously,  $x \in A$ . By (\*\*), there is a path P in F connecting x to a point  $a \in A_1$ . This path starts from a on an edge not in M. Thus by (\*), its second edge belongs to M. Its third edge is therefore not in M while by (\*) its fourth edge is in M, etc. Hence, it is a path alternating relative to M and its last edge belongs to M. Exchange the edges of  $E(P) \cap M$  and E(P) - M, then a matching M' arises to which (x, y) can be added. Thus M is not a maximum matching.

II. On the other hand, suppose no point of F is adjacent to any point in  $B_1$ . Let X = A - V(F) and  $Y = V(F) \cap B$ . We claim  $|X \cup Y| = |M|$  and  $X \cup Y$  covers all edges.

All points of  $X \cup Y$  are covered by M, for suppose not. If  $y \in Y = V(F) \cap B$ , then by (\*) y is covered by M. If  $x \in X = A - V(F)$  and  $x \in A_1$ , then the forest  $F' = F \cup \{x\}$  is larger than F, contradicting the assumption that F was maximal relative to M.

Moreover, no edge of M covers two points of  $X \cup Y$ , for if  $(x, y) \in M$  and  $y \in V(F) \cap B$ , then  $x \in V(F)$  by the maximality of F. Hence  $|X \cup Y| = |M|$ . Secondly, let (u, v) be any edge not covered by  $X \cup Y$  and suppose  $u \in A$ ,  $v \in B$ . Then  $u \in V(F)$  and  $v \notin V(F)$ . Moreover, by the assumption,  $v \notin B_1$ . Thus there is an edge  $(w, v) \in M$ , and  $w \neq u$  since  $|X \cup Y| = |M|$ . Here  $w \in V(F)$ , for otherwise it would follow as in part I, that w would be connected in F to  $A_1$  by a path starting from w through (w, v), whence  $v \in V(F)$ , a contradiction. But, if  $w \in V(F)$ , then the edges (u, v) and (w, v) can be added to F, contradicting the maximality of F.

So  $\tau(G) \leq |M|$ . Since, obviously,  $|M| \leq \nu(G) \leq \tau(G)$ , it follows that

$$|M| = \nu(G)$$
 and  $\tau(G) = \nu(G)$ .

The first inequality shows that M is a maximum matching. The second proves

König's theorem.

The desired algorithm could now be the following: Start with a matching M. Form a maximal forest with properties (\*) and (\*\*). Then either we can enlarge M or conclude that it is a maximum matching. In the latter case we also obtain a minimum cover.

We cannot go into details but remark that this algorithm does not need more time than a polynomial of the number of points.

**4.** (a) First solution. What we want to show is that  $\nu(G) = |A|$ . It is enough to prove, by 7.2, that  $\tau(G) = |A|$ . Since A covers all edges,  $\tau(G) \leq |A|$ . Thus, it suffices to show that any set S covering all edges has at least |A| elements. Observe now that if S covers all edges of G, then

$$\Gamma(A-S) \subseteq S \cap B$$

because, if  $x \in A - S$  is joined to  $y \in B$ , then to cover the edge (x, y), S has to contain y. Hence by the assumption of the theorem,

$$|S \cap B| \ge |\Gamma(A - S)| \ge |A - S|$$

and thus,

$$|S| = |S \cap A| + |S \cap B| \ge |S \cap A| + |A - S| = |A|.$$

Second solution. We use induction on |A|. Let us distinguish two cases:

Case 1. There is a set  $A_1 \subset A$ ,  $A_1 \neq \emptyset$ , such that  $|\Gamma(A_1)| = |A_1|$ . Let  $G_1$  be the subgraph of G induced by  $A_1 \cup \Gamma(A_1)$  and let  $G_2 = G - A_1 - \Gamma(A_1)$ . We claim both  $G_1$  and  $G_2$  satisfy the condition of the problem.

Let  $X \subseteq A_1$ . Then  $\Gamma(X) \subseteq \Gamma(A_1)$  and so,

$$\Gamma_{G_1}(X) = \Gamma(X), \ |\Gamma_{G_1}(X)| = |\Gamma(X)| \ge |X|.$$

On the other hand, suppose  $X \subseteq A - A_1$ . Then

$$\begin{aligned} |\Gamma_{G_2}(X)| &= |\Gamma(X \cup A_1)| - |\Gamma(A_1)| \ge |X \cup A_1| - |\Gamma(A_1)| = \\ &= |X \cup A_1| - |A_1| = |X|. \end{aligned}$$

Thus, by the induction hypothesis, there exists a matching  $F_1$  in  $G_1$  which matches  $A_1$  with  $\Gamma(A_1)$  and a matching  $F_2$  in  $G_2$  which matches  $A-A_1$  with (certain points of)  $B-\Gamma(A_1)$ .  $F_1\cup F_2$  is a desired matching.

Case 2.  $|\Gamma(X)| > |X|$  for every  $X \subset A$ ,  $X \neq \emptyset$ . Let  $x \in A$ ,  $y \in B$  be adjacent points. We claim  $G_1 = G - x - y$  satisfies the conditions of the problem. Let  $X \subseteq A - \{x\}$ ; if  $X = \emptyset$ , then  $|\Gamma(X)| = 0 = |X|$ , so we may assume  $X \neq \emptyset$ . Also  $X \neq A$ , thus by the assumption,  $|\Gamma(X)| > |X|$ . So

$$|\Gamma_{G_1}(X)| \geq |\Gamma(X)| - 1 \geq |X|.$$

Hence, by the induction hypothesis, there is a matching  $F_1$  in  $G_1$  which covers all points of  $A - \{x\}$ . Now  $F_1 \cup \{(x, y)\}$  is a matching of the type desired.

(b) If G has a 1-factor, then obviously,

(1) 
$$|\Gamma(X)| \ge |X|$$
 for each  $X \subseteq A$  and  $|A| = |B|$ .

Conversely, if (1) holds, then (a) implies G has a 1-factor. [P. Hall; B, O, S.]

5. By 7.2, it suffices to show that

$$\tau(G) = |A| - \delta.$$

First let  $X \subseteq A$  be a set such that

$$\delta = |X| - |\Gamma(X)|.$$

Set  $S = \Gamma(X) \cup (A - X)$ . Then S covers a ledges of G, and hence

$$\tau(G) \leq |S| = |\Gamma(X)| + |A| - |X| = |A| - \delta.$$

Now let S be a minimum point-cover and X = A - S. Then, as before,

$$\Gamma(X) \subseteq S \cap B$$

and hence,

$$|\Gamma(X)| \le |S \cap B| = |S| - |S \cap A| = |S| - |A| + |A - S| = \tau(G) - |A| + |X|,$$

whence

$$\delta \geq |X| - |\Gamma(X)| \geq |A| - \tau(G)$$
, and thus  $\tau(G) \geq |A| - \delta$ . [O. Ore; O, B]

6. (a) By the observation in the hint,

$$\begin{split} |\varGamma(X_1 \cup X_2)| + |\varGamma(X_1 \cap X_2)| &\leq |\varGamma(X_1) \cup \varGamma(X_2)| + |\varGamma(X_1) \cap \varGamma(X_2)| = \\ &= |\varGamma(X_1)| + |\varGamma(X_2)| = |X_1| + |X_2| + 2k. \end{split}$$

On the other hand,

$$|\Gamma(X_1 \cup X_2)| \ge |X_1 \cup X_2| + k$$
 and  $|\Gamma(X_1 \cap X_2)| \ge |X_1 \cap X_2| + k$ ,

because of (\*) and since  $X_1 \cap X_2 \neq \emptyset$ . Thus

$$\begin{split} |\varGamma(X_1 \cup X_2)| + |\varGamma(X_1 \cap X_2)| &\geq |X_1 \cup X_2| + |X_1 \cap X_2| + 2k = \\ &= |X_1| + |X_2| + 2k. \end{split}$$

Hence equality must hold throughout, in particular

$$|\Gamma(X_1 \cap X_2)| = |X_1 \cap X_2| + k.$$

(b) Let  $G_1$  be defined as in the hint. Let  $x \in A$ , and let S be the intersection of all sets X with

$$|\Gamma_G(X)| = |X| + k$$
 and  $x \in X$ .

Then, by the repeated application of (a), we see that S itself satisfies

$$|\Gamma_{G_1}(S)| = |S| + k \text{ and } x \in S.$$

(If the assertion is true,  $S = \{x\}$ , but we do not know this as yet.) Let y be any point adjacent to x. Since  $G_1 - (x, y)$  does not have property (2), we find a set  $X \subseteq A$ ,  $X \neq \emptyset$ , such that

$$|\Gamma_{G_1-(x,y)}(X)| < |X| + k$$
.

Since, on the other hand,  $|\Gamma_{G_1-(x,y)}(X)|$  can differ from  $|\Gamma_{G_1}(X)|$  only by 1, we must have

$$|\Gamma_{G_1}(X)| = |X| + k, \quad x \in X, \quad y \notin \Gamma_{G_1-(x,y)}(X).$$

Thus, by the definition of  $S, S \subseteq X$  and also

$$(4) y \notin \Gamma_{G_i}(S - \{x\}),$$

because

$$\Gamma_{G_1}(S - \{x\}) = \Gamma_{G_1 - (x, y)}(S - \{x\}) \subseteq \Gamma_{G_1 - (x, y)}(X).$$

Now since (4) holds for each neighbor y of x, we get

$$\Gamma_{G_1}(x) \cap \Gamma_{G_1}(S - \{x\}) = \emptyset.$$

Hence

$$|\varGamma_{G_1}(S)| = |\varGamma_{G_1}(x) \cup \varGamma_{G_1}(S - \{x\})| = |\varGamma_{G_1}(x)| + |\varGamma_{G_1}(S - \{x\})|.$$

Here

$$|\Gamma_{G_i}(x)| \ge k + 1$$

and

$$|\Gamma_{G_1}(S - \{x\})|$$
  $\begin{cases} = 0 = |S - \{x\}| & \text{if } S - \{x\} = \emptyset, \\ \geq |S - \{x\}| + k \geq |S - \{x\}| & \text{if } S - \{x\} \neq \emptyset. \end{cases}$ 

Consequently,

$$|\Gamma_{G_1}(S)| \ge k + 1 + |S - \{x\}| = |S| + k.$$

Comparing this with (1), we find that equality must hold everywhere, in particular in (5). [L. Lovász, Acta Math. Acad. Sci. Hung. 21 (1970) 443-446.]

7. (i)  $\Rightarrow$  (iii). G has a 1-factor, hence |A| = |B|. Let  $\emptyset \neq X \subset A$  and assume indirectly that  $|\Gamma(X)| \leq |X|$ . We must then have equality as G has a 1-factor. Now  $X \cup \Gamma(X)$  is joined by an edge e to the rest of the graph, because G is connected. Obviously, e joins  $\Gamma(X)$  to A-X. Now this edge e is not contained in any 1-factor; for if F is a 1-factor, then it contains |X| edges leaving X. But these edges cover all of  $\Gamma(X)$  so that e cannot belong to F.

(iii)  $\Rightarrow$  (ii). We have to show G-x-y has a 1-factor if  $x \in A$ ,  $y \in B$ . Since  $|A-\{x\}|=|B-\{y\}|$ , it suffices to show that there is a matching in G-x-y covering  $A-\{x\}$ . But this is equivalent to saying that, for each  $X \subseteq A-\{x\}$ ,

$$|\Gamma_{G-x-y}(X)| \ge |X|.$$

This is obvious if  $X = \emptyset$ . Otherwise,

$$|\varGamma_{G-\mathbf{x}-\mathbf{y}}(X)| \ge |\varGamma(X)| - 1 \ge |X|.$$

(ii)  $\Rightarrow$  (i). Suppose G is disconnected. Then we can find a component  $G_1$  such that  $|V(G_1) \cap A| \leq |V(G_1) \cap B|$ . Let  $x \in V(G_1) \cap A$ ,  $y \in B - V(G_1)$ . Then G - x - y cannot have 1-factor, because  $V(G_1) \cap B$  is adjacent to  $|V(G_1) \cap A| - 1 < |V(G_1) \cap B|$  points only. Thus G is connected.

Now let e = (x, y) be any edge of G. Then G - x - y has a 1-factor, which extends to a 1-factor of G containing e. Thus each edge of G is contained in a

1-factor. [G. Hetvei, Pécsi Tan. Főisk. Közl. (1964) 151–168.]

## 8. I. First we show that the graphs of form

$$G = G_0 \cup P_1 \cup \ldots \cup P_k$$

where  $P_1, \ldots, P_k$  are chosen as described in the problem, are elementary. They are obviously bipartite and connected.

We use induction on k. Suppose that

$$G' = G_0 \cup P_1 \cup \ldots \cup P_{k-1}$$

is elementary and let x, y be the two endpoints of  $P_k$ . Observe first that every 1-factor of G' extends to a 1-factor of G, simply by adding every second edge of  $P_k$ , starting with the second edge. Thus, all edges of G' as well as every second edge of  $P_k$  belongs to some 1-factor of G.

On the other hand, G' - x - y has a 1-factor  $F_0$ . Add to  $F_0$  every second edge of  $P_k$ , starting with the first edge. Then we get a 1-factor of G which includes

the edges of  $P_k$  not used before.

Thus we have shown that G is connected and each edge of it belongs to a 1-factor. This proves by the preceding result that it is elementary.

II. Now suppose that G is any elementary graph. Let F be a 1-factor of G and let  $G_0$  be the graph formed by an edge of F. We choose  $P_1, P_2, \ldots, P_k$  one by one so that they have the properties required in the problem and in addition that (\*) they are alternating paths with respect to F, starting with an edge not in F (we allow  $P_i$  to be a single edge not in F).

Suppose  $P_1, \ldots, P_i$  are chosen, and  $G' = G_0 \cup P_1 \cup \ldots \cup P_i \neq G$ . Let e be an edge not in G' but having at least one endpoint x in G'. Since G is connected,

such an edge exists. Let  $F_1$  be a 1-factor of G containing e.

Observe that by the additional requirement (\*), G' has the property that  $F \cap E(G')$  is a 1-factor of G', and  $F_0 = F - E(G')$  does not meet G' at all. Thus, if we consider  $F_1 \cup F_0$ , one component of it will be a path ending with e. This path  $P_{i+1}$  has both endpoints in G' (one of them is x) and has no other point in common with G'. Its end-edges belong to  $F_1$ , and hence by reasons of parity,  $P_{i+1}$  has odd length and connects two points in different classes of G'. By its definition, it alternates with respect to F in the required manner. Thus, if  $G_0 \cup P_1 \cup \ldots \cup P_i \neq G$ , we were able to extend this union by  $P_{i+1}$ . Sooner or later G will be exhausted, which proves the assertion. [G. Hetyei, ibid.]

## **9.** Represent G in the form

$$G_0 \cup P_1 \cup \ldots \cup P_k$$

as in the preceding problem. Here  $P_k$  cannot be a single edge, because

$$G_0 \cup P_1 \cup \ldots \cup P_{k-1}$$

is elementary and therefore cannot be obtained from G by removing a single edge. Thus  $P_k$  has an inner point (at least two, in fact) and these have degree 2 [G. Hetvei, ibid.].

The stronger statement that every edge would have an endpoint of degree 2 is false. The graph shown in Fig. 54 arises from the construction of 7.8, so it is elementary. It is also minimal. To see this we remark that an elementary graph has degrees at least 2 by 7.7. Removing any edge except e and f, the remain-




Fig. 54

ing graph will have a point of degree 1, and so will not be elementary. If e or f is removed, the three points marked by x or O will be joined to three points only, showing by 7.7 that the remaining graph is not elementary again. However, both endpoints of e have degree 3 [J. Csima].

10. Suppose first that G is r-regular. We show G has a 1-factor. Obviously,

$$|E(G)| = r|A| = r|B|,$$

whence |A| = |B|.

Thus, it suffices to show that G contains a matching which covers A or by 7.4, that for each  $X \subseteq A$ ,

$$|\Gamma(X)| \geq |X|$$
.

Suppose  $X \subseteq A$  and let us count the edges leaving X. There are, by regularity, r|X| such edges. On the other hand, each of them goes to a point of  $\Gamma(X)$  and any point of  $\Gamma(X)$  meets at most r of them. Hence,

$$r|X| \le r|\Gamma(X)|, |X| \le |\Gamma(X)|.$$

Thus G has a 1-factor.

Let F be a 1-factor of G. G - F is an (r - 1)-regular bipartite graph and, by induction, it decomposes into the union of r - 1 1-factors. Adding F to them

we obtain a desired decomposition.

Now let G be any bipartite graph with maximum degree r. Add new (isolated) points to make the two color-classes of G equal. Then add new edges as long as you can without increasing the maximum degree. The graph  $G_1$  obtained this way must be r-regular; for if there was a point x with degree  $d_{G_1}(x) < r$ , then the opposite color class would contain such a point y as well and we could add an (x, y)-edge.

Thus,  $G_1$  is an r-regular graph containing G. Hence we may split  $G_1$  into r 1-factors, and thereby produce a decomposition of G into r matchings.

11. Let  $G_1$  denote the bipartite graph obtained as described in the hint. Then

$$d_{G_1}(x) \le k \qquad \left(x \in V(G_1)\right)$$

and hence, by the preceding problem,  $E(G_1)$  is the union of k matchings  $F_1, \ldots, F_k$ .

Now identify those points of  $G_1$  which correspond to the same point of  $G_1$ , then  $F_1, \ldots, F_k$  are mapped onto certain subgraphs  $F'_1, \ldots, F'_k$ . Since

 $F_i$  is a matching, there is at most one  $F_i$ -edge incident with any of the  $\left[\frac{d(x)}{dx}\right]^*$  points of  $G_1$  corresponding to  $x \in V(G)$ . Hence,

7.12

$$d_{F_i'}(x) \leq \left[\frac{d(x)}{k}\right]^*.$$

On the other hand, there are  $\left[\frac{d(x)}{k}\right]$  points of  $G_1$  corresponding to x which have degree k. There must be an  $F_i$ -edge starting from each of these, whence

$$d_{F_i'}(x) \ge \left[\frac{d(x)}{k}\right]$$

[D. de Werra; B].

12. By 7.11, E(G) has a partition  $G_1 \cup \ldots \cup G_r$  where

$$\left\lceil \frac{d(x)}{r} \right\rceil \leq d_{G_i}(x) \leq \left\lceil \frac{d(x)}{r} \right\rceil^*$$

for any point x. Since  $d(x) \ge r$  by the definition of r, this implies that

$$d_{G_i}(x) \geq 1$$
,

i.e.  $G_i$  is an edge-cover [R.P. Gupta, Theory of Gr. Int. Symp. Rome, Dunod, Paris–Gordon and Breach, New York, (1967) 135–138].

13. Suppose  $G_1$  as defined in the hint has no 1-factor. Let  $\{A, B\}$  be a bicoloration of G, obviously |A| = |B|. Thus, if  $G_1$  has no 1-factor, 7.4b implies that there exists an  $X \subseteq A$  such that

$$|\Gamma_{G_i}(X)| < |X|.$$

We count the edges of G leaving X by two different methods. Firstly, the number of these edges is  $r \cdot |X|$ . Secondly, there are at most  $|X| \cdot (n - |\Gamma_{G_1}(X)|)$  edges joining X to  $B - \Gamma_{G_1}(X)$  (since these edges are simple edges) and at most  $r \cdot |\Gamma_{G_1}(X)|$  edges joining X to  $\Gamma_{G_1}(X)$ , since any point of  $\Gamma_{G_1}(X)$  has degree r. Thus

$$r \cdot |X| \leq |X| \cdot (n - |\Gamma_{G_1}(X)|) + r |\Gamma_{G_1}(X)|$$

or, equivalently,

(2) 
$$(r-n) |X| \le (r-|X|) |\Gamma_{G_1}(X)|.$$

Now obviously, we are interested in the case r > n, otherwise not even the existence of multiple edges follows. Then |X| < r and (1), (2) imply that

$$(r-n)|X| \le (r-|X|)(|X|-1)$$

or, equivalently,

$$r \leq (n-|X|+1) \cdot |X| \leq \left\lceil \frac{n+1}{2} \right\rceil \cdot \left\lceil \frac{n+2}{2} \right\rceil$$

Thus we have shown that  $r > \left[\frac{n+1}{2}\right] \cdot \left[\frac{n+2}{2}\right]$  implies the existence of a 1-factor in  $G_1$ .

Conversely, let |A| = |B| = n,  $A' \subset A$ ,  $|A'| = \left[\frac{n+1}{2}\right]$ ,  $B' \subset B$  and  $|B'| = \left[\frac{n-1}{2}\right]$ . Join each point of A' to each point of B' by  $\left[\frac{n+2}{2}\right]$  edges, each point of A - A' to each point of B - B' by  $\left[\frac{n+1}{2}\right]$  edges; and each point of A' to each point of B - B' by one edge. The resulting graph G is  $\left[\frac{n+1}{2}\right] \cdot \left[\frac{n+2}{2}\right]$ -regular. For any  $r \leq \left[\frac{n+1}{2}\right] \cdot \left[\frac{n+2}{2}\right]$ , G contains an r-regular spanning subgraph  $G_r$  (see 7.10). Observe that G and, consequently,  $G_r$  has no 1-factor whose edges have a parallel in  $G_r$ . This proves that  $r(n) = \left[\frac{n+1}{2}\right] \cdot \left[\frac{n+2}{2}\right] + 1$ .

14. I. We show we cannot get stuck in any situation other than  $x = a_1$  and

 $a_1$  is joined to some  $b_i$  by r parallel edges.

Suppose we get stuck when  $x = b_i$ . The reason why we cannot continue is that we have come from  $a_{\mu}$  ( $1 \le \mu \le n$ ) and  $b_i$  has no neighbor other than  $a_{\mu}$ . Since  $b_i$  has degree r-1, when  $x = b_i$ , we must have r-1 ( $a_{\mu}$ ,  $b_i$ )-edges. So at the previous step when  $x = a_{\mu}$ , there were r ( $a_{\mu}$ ,  $b_i$ )-edges. Suppose x came to  $a_{\mu}$  from  $b_j \ne b_i$ . Then the edge ( $a_{\mu}$ ,  $b_j$ ) was doubled, i.e. the degree of  $a_{\mu}$  was at least r+2, which is not allowed.

Similarly it follows that if we get stuck when  $x = a_i$ , then i = 1 and we

have the situation described.

II. So it remains to show that the procedure cannot run infinitely. Suppose indirectly that it does. Note that all edges in all graphs arising are parallel to edges of G. Consider those edges of G which x passes infinitely many times. These edges must be passed infinitely often in both directions, otherwise their multiplicities would tend to  $+\infty$  or  $-\infty$ , both impossible.

Consider a point  $b_i$ , and suppose 3 edges  $(a_{\mu}, b_i)$ ,  $(a_{\nu}, b_i)$ ,  $(a_{\varrho}, b_i)$  are used infinitely often  $(\mu < \nu < \varrho \le n)$ . Then  $(a_{\varrho}, b_i)$  could never be passed from  $b_i$  to  $a_{\varrho}$ , since  $(a_{\mu}, b_i)$  or  $(a_{\nu}, b_i)$  is always a choice with  $\mu$ ,  $\nu < \varrho$ . But this is impossible. Similarly it follows that each  $a_{\mu}$  is adjacent to at most two in-

finitely used edges.

Now consider a stage when all the finitely used edges have already been used up. Since each point is adjacent to at most two infinitely used edges, x can only run around a circuit. But then the edges of this circuit are always passed

in the same direction, a contradiction.

The procedure thus results in an r-regular graph which is obtained from a subgraph of G by multiplying edges and in which the pair  $\{a_1, b_i\}$  spans a component. Repeating it, we finally obtain a graph which arises from a 1-factor of G by taking each edge r times. Thus we obtain a 1-factor of G. Note that no data must be stored; the procedure can be carried out on a blackboard with an eraser and one piece of chalk [J. Csima].

15. (a) We use simultaneous induction on |V(G)| and k.

Suppose first G is elementary. Let  $x \in A$ ,  $y \in B$  be adjacent. Then the 1-factors of G containing (x, y) correspond to the 1-factors of G - x - y. Since (x, y) is contained in some 1-factor, G - x - y has a 1-factor. Moreover, every point of A - x has degree  $\geq k - 1$  in G - x - y, so G - x - y has at least (k - 1)! 1-factors. Thus, each edge of G is contained in at least (k - 1)! 1-factors.

Fix an  $x \in A$ . There are at least k edges incident with x and each of them is contained in at least (k-1)! 1-factors, This gives k(k-1)! = k! 1-factors which are obviously different.

Now suppose that G is not elementary. Then by 7.7, there is an  $\emptyset \neq X \subset A$ 

such that

$$|\Gamma(X)| = |X|.$$

Let F be any 1-factor of G and let  $F_1$  be the set of its edges starting from A-X. Then, obviously, the other edges of F cover  $\Gamma(X)$ , so  $F_1$  matches

up A - X with  $B - \Gamma(X)$ .

Observe that the subgraph  $G_1$  induced by  $X \cup \Gamma(X)$  satisfies the same requirements as G: every point of X has degree at least k in  $G_1$ , and it has a 1-factor (e.g.  $F - F_1$ ). Therefore,  $G_1$  has at least k! 1-factors by the induction hypothesis. Adding  $F_1$  to each of these 1-factors, we obtain k! 1-factors of G [M. Hall, Bull. Amer. Math. Soc. 54 (1948) 922–926].

(b) We use induction on m - n + 2. If this = 2, then G is a single circuit and the assertion is trivial. Suppose m > n. By 7.8,

$$G = G_1 \cup P$$
,

where  $G_1$  is an elementary bipartite graph and P is an odd path, which has only its endpoints x, y in common with  $G_1$  and x, y belong to different color classes of  $G_1$ . Clearly

$$|E(G_1)| - |V(G_1)| + 2 = m - n + 1 < m - n + 2$$

and thus, by the induction hypothesis,  $G_1$  contains at least m-n+1 1-factors. Each of these can be extended to a 1-factor of G. Moreover by 7.7,  $G_1-x-y$  contains a 1-factor which can also be extended to a 1-factor of G using every second edge of P. Thus we indeed have found (m-n+1)+1=m-n+2 1-factors in G [cf. L. Lovász-M.D. Plummer, in: Infinite and Finite Sets, Coll. Math. Soc. J. Bolyai 10, Bolyai-North-Holland (1975) 1051-1079].

**16.** I. Suppose G has an f-factor F. Then, obviously,

$$\sum_{x \in A} f(x) = |F| = \sum_{x \in B} f(x),$$

which proves (i). Let  $X \subset A$  and  $Y \subseteq B$ . Then there are at most m(X, Y) edges of F joining X to Y and at most  $\sum_{x \in B} f(y)$  edges of F joining X to B - Y.

Since there are exactly  $\sum_{x \in X} f(x)$  edges of F joining X to B, (ii) follows.

II. Now suppose that (i) and (ii) are satisfied. Direct all edges from A to B. Take two points a, b and join a to each point of A and each point of B to b by directed edges. Define the capacity c(e) of an edge e of the resulting digraph  $G_0$  by

 $c(e) = \begin{cases} f(x) & \text{if } e = (a, x) \text{ or } e = (x, b), \\ 1 & \text{if } e = (x, y), \ x \in A, \ y \in B. \end{cases}$ 

Observe that  $G_0$  has an integral (a, b)-flow with value  $\sum_{x \in A} f(x) = \sum_{y \in B} f(y)$ , if and only if G has an f-factor. So what we have to verify is, by 6.74 and 6.77, that each (a, b)-cut of  $G_0$  has capacity at least  $\sum f(x)$ .

Let S determine an (a, b)-cut  $(a \in S \subset V(G_0))$ . Set  $X = S \cap A$  and Y = B - S. Then the capacity of the cut determined by S is

$$\sum_{x \in A - X} f(x) + \sum_{y \in B - Y} f(y) + m(X, Y) \ge \sum_{x \in A - X} f(x) + \sum_{x \in X} f(x) = \sum_{x \in A} f(x)$$

by (ii). This completes the proof [O. Ore, D. Gale-H. J. Ryser; see O].

17. Suppose G is embedded into a d-regular bipartite graph G' with color classes A, B. Let  $A_1 = A \cap V(G)$ ,  $B_1 = B \cap V(G)$ ,  $A_2 = A - A_1$ ,  $B_2 = B - B_1$ . Since there are exactly d - d(x) edges of G' joining x to  $B_2$  for any  $x \in A_1$ , we have

 $\sum_{x \in A_1} (d - d(x)) \le d \cdot |B_2|$ 

or, equivalently,

$$d \cdot |A_1| - m \le d \cdot |B_2|.$$

Similarly,

$$d \cdot |B_1| - m \le d \cdot |A_2|.$$

Summing, we get

$$dn - 2m \le d(|A_2| + |B_2|) = d(|V(G')| - n),$$

thus

$$|\mathit{V}(G')| \geq 2n - rac{2m}{d}$$

and, since |V(G')| is an even integer,

$$|V(G')| \ge 2n - 2\left[\frac{m}{d}\right].$$

Next we show that G can be embedded as an induced subgraph into a d-regular bipartite graph G' with exactly  $2n-2\left[\frac{m}{d}\right]$  points. Let  $\{A,B\}$  be a bicoloration of G. Since  $m \leq d \cdot |A|$  and  $m \leq d \cdot |B|$ , we have

$$|B| = n - |A| \le n - \left\lceil \frac{m}{d} \right\rceil$$
 and  $|A| \le n - \left\lceil \frac{m}{d} \right\rceil$ .

So we can construct two disjoint sets  $A' \supset A$  and  $B' \supset B$  with

$$|A'| = |B'| = n - \left\lceil \frac{m}{d} \right\rceil.$$

We add now (A, B' - B)-edges and (B, A' - A)-edges so that all the degrees in  $A \cup B$  become equal to d. This is possible because

$$\sum_{x \in A} (d - d(x)) = |A| \cdot d - m = nd - |B| \cdot d - m \le$$

$$\le nd - |B| \cdot d - \left\lceil \frac{m}{d} \right\rceil \cdot d = |B'| \cdot d - |B| \cdot d = |B' - B| \cdot d$$

and similarly,

$$\sum_{x \in B} (d - d(x)) \le |A' - A| \cdot d$$

(and, because we do not mind multiple edges). Finally, we add (A'-A, B'-B)-edges to make all degrees d.

18. Considering the "bipartite complement"  $\widetilde{G}$  of G as defined in the hint, by 7.16 it suffices to show that whenever  $X \subseteq A$ ,  $Y \subseteq B$ , then

$$(n-2d)(n-|Y|)+m_{\tilde{G}}(X,Y)\geq (n-2d)\cdot |X|$$

or, equivalently,

$$(n-2d)(|X|+|Y|-n) \leq m_{\tilde{G}}(X, Y).$$

Since this inequality is symmetric in X, Y we may assume that  $|X| \ge |Y|$ . Moreover,

$$m_{\tilde{G}}(X, Y) = |X| \cdot |Y| - m_{G}(X, Y),$$

and so, it suffices to show that

$$(1) (n-2d)(|X|+|Y|-n) \leq |X|\cdot |Y|-m_G(X,Y).$$

Here

$$m_G(X, Y) \leq d \cdot |Y|$$

and so, (1) would follow from

$$(|X| + d - n)(|Y| + 2d - n) \ge d(2d - n).$$

If  $|Y| \le n - 2d$ , then  $|Y| + 2d - n \le 0$  and so,

$$(|X|+d-n)(|Y|+2d-n) \ge (n+d-n)(|Y|+2d-n) \ge d(2d-n).$$

If  $n-2d \le |Y| \le |X| \le n-d$  and  $|X| \ge d$ , then  $|X|+d-n \le 0$ , thus  $(|X|+d-n) \, (|Y|+2d-n) \ge (|X|+d-n)d \ge (2d-n)d$ .

If  $n-2d \leq |Y|$  and  $n-d \leq |X|$ , then recalling also that  $d \leq n/2$ , we have

$$(|X| + d - n) (|Y| + 2d - n) \ge 0 \ge d(2d - n).$$

Finally, if  $|X| \leq d$ , we use

$$m_G(X, Y) \leq |X| \cdot |Y|$$

and thus it suffices to show that

$$(n-2d)(|X|+|Y|-n) \leq 0,$$

which is clear, since  $|Y| \le |X| \le d \le \frac{n}{2}$ .

19. The transformation of the problem given in the hint is easily verified

(cf. 4.21).

Suppose G has no 1-factor, then by 7.4b, there is a set  $X \subseteq A = \{u_1, \dots, u_n\}$ with  $|\Gamma(X)| < |X|$ . Let, say,  $X = \{u_1, \ldots, u_k\}$  and  $B - \Gamma(\overline{X}) = \{v_1, \ldots, v_l\}$ . Thus k + l > n, and  $a_{ij} = 0$  for  $1 \le i \le k$ ,  $1 \le j \le l$ . Now

$$\sum_{i=1}^n a_{ij} = 1,$$

because A is doubly stochastic, hence

$$\sum_{j=1}^{l}\sum_{i=1}^{n}a_{ij}=l,$$

On the other hand,

$$\sum_{j=1}^{l} \sum_{i=1}^{n} a_{ij} = \sum_{i=k+1}^{n} \sum_{j=1}^{l} a_{ij} \le \sum_{i=k+1}^{n} \sum_{j=1}^{n} a_{ij} = n - k < l,$$

a contradiction.

- 20. (a) It is clear from the remark in the hint.
- (b) Suppose G has no 1-factor. Then  $\det(a_{ij}) = 0$ . Let, say, the first k+1 columns form a minimal linearly dependent set of columns. Since the rank of the matrix A' formed by these k+1 columns is k, it contains a  $k \times k$ non-singular submatrix. We may assume that the first k rows and columns form this submatrix.

Since the first k+1 columns are linearly dependent, we have numbers  $\lambda_1, \ldots, \lambda_{k+1}$  such that

(1) 
$$\sum_{j=1}^{k+1} \lambda_j a_{ij} = 0 \qquad (i = 1, \dots, n).$$

Since  $\lambda_{\mu} \neq 0$  by the minimality of k, we may divide by any  $\lambda_{\mu}$  and determine the numbers  $\lambda_{\nu}/\lambda_{\mu}$  by Cramer's rule from the first k rows. Thus  $\lambda_{\nu}/\lambda_{\mu}$  belongs to the field generated by  $\{a_{ij}: 1 \leq i \leq k, \ 1 \leq j \leq k+1\}$ . We claim that  $a_{ij}=0$  if  $k+1 \leq i \leq n$  and  $1 \leq j \leq k+1$ . We show, e.g. that  $a_{k+1,1}=0$ . We have

$$a_{k+1,1} = -\frac{1}{\lambda_1} \sum_{j=2}^{k+1} \lambda_j a_{k+1,j}$$

from (1). Here the right-hand side is in the field generated by  $\{a_{ij}; (i,j) \neq$  $\neq (k+1,1)$ . By the algebraic independence of the  $a_{ij}$ 's this is only possible if  $a_{k+1,1} = 0$ .

Setting  $X = \{a_{k+1}, \ldots, a_n\}$ , we have

$$\Gamma(X) \subseteq \{b_{k+2}, \ldots, b_n\}, |\Gamma(X)| < |X|,$$

which proves the non-trivial part of 7.4b [J. Edmonds, J. Res. Natl. B. Standards 71B (1967) 241-245].

21. If the conditions imposed upon y are linearly independent the answer is clearly m-n, where n=|V(G)|; if n-1 of them are linearly independent but the  $n^{\text{th}}$  one is not, the dimension is m-n+1. We claim the first case occurs, if G is not bipartite and the second one, if it is.

To prove this, consider a non-trivial linear relation among the left-hand sides of the constraints (1). We want to show that this can exist only, if G is

bipartite and that in this case it is essentially unique. So let

$$\sum_{v \in V(G)} \lambda_v \sum_{e_i \ni v} y_i$$

be identically 0, i.e.

$$\lambda_n + \lambda_n = 0$$

for every edge (u, v). Hence using the fact that G is connected we conclude that  $|\lambda_v| = \text{const} = \lambda$  and points with  $\lambda_v = \lambda$  are only connected to points with  $\lambda_v = -\lambda$  and vice versa. Hence G is bipartite. Since the 2-coloration of a bipartite graph is unique, we also see that the linear relation among the left-hand sides of the constraints (1) is unique up to a scalar factor. Thus in the bipartite case there is the unique relation

$$\sum_{v \in A} \sum_{e_i \ni v} y_i = \sum_{v \in B} \sum_{e_i \ni v} y_i,$$

where  $\{A, B\}$  is the bicoloration of G [H. Sachs, Wiss. Z. TH Ilmenau 12 (1966) 7–12].

22. Following the hint, let  $(x, y) \in F$ . If there are at least 3 edges joining  $\{x, y\}$  to  $\{u, v\}$ , then there are two of them which are independent, e.g. (x, u) and (y, v). Now  $F = \{(x, y)\} \cup \{(x, u), (y, v)\}$  is a bigger matching than F, a contradiction. So each edge of F is joined to  $\{u, v\}$  by at most two edges. Since u and v are not joined to each other or to any point not on the edges of F (otherwise F would not be maximal), this implies that

$$d(u) + d(v) \le 2|F| \le 2(n-1).$$

This, however, contradicts the hypothesis that d(u) and d(v) are at least n (cf. also problem 10.19).

- 23. Let X be the set of points covered by  $F_0$ . Following the hint, let F be a maximum matching of G containing as many edges of  $F_0$  as possible. We claim F covers X. Suppose indirectly that there exists an  $x \in X$  not covered by F. Let (x, y) be the edge of  $F_0$  incident with x. Since  $F \cup \{(x, y)\}$  is not a matching by the maximality of F, there is an  $e \in F$  incident with e. Set e0 be a maximum matching. Moreover, it has more edges in common with e1 than e2, a contradiction.
- **24.** (a) For k=1, a single edge satisfies the requirements. Suppose  $G_{k-1}$  is a graph with a unique 1-factor and with degrees at least k-1. Let  $G'_{k-1}$  be another disjoint copy of it, and also consider two new points x, y. Join x to all points of  $G_{k-1}$ , y to all points of  $G'_{k-1}$ , and x to y. Let  $G_k$  be the resulting graph (Fig. 55).




Fig. 55

Every point of  $V(G_{k-1}) \cup V(G'_{k-1})$  has greater degree than in  $G_{k-1}$  or  $G'_{k-1}$ , hence it has degree at least k. Obviously, this holds for x and y. Moreover,  $G_k$  has a unique 1-factor. For let F be any 1-factor of  $G_k$ . By reasons of parity it must contain the cut-edge (x, y). Then, however, the rest of it forms a 1-factor of  $G_{k-1}$  and a 1-factor of  $G'_{k-1}$ . Since these are unique, F is uniquely determined. So  $G_k$  has at most one 1-factor. But the one described (composed of the 1-factors of  $G_{k-1}$ ,  $G'_{k-1}$  and of (x, y)) is indeed a 1-factor of  $G_k$ . Thus  $G_k$  has a unique 1-factor.

(b) First solution. Suppose G is a minimum counterexample. Let F be its unique 1-factor and let  $e_1 \in E(G) - F$ . Consider the equivalence-class defined in 6.27 containing  $e_1$ . If it consists only of  $e_1$ , then  $G - e_1$  is 2-edge-connected and we get a contradiction of the minimality of G. Suppose it has other edges  $e_2, \ldots, e_k, k \geq 2$ .


Let  $G_1$  be any component of  $G - \{e_1, \ldots, e_k\}$ . By 6.27, it is incident with exactly two edges of  $\{e_1, \ldots, e_k\}$ . If these two edges do not belong to F, we get by induction that  $G_1$  has two distinct 1-factors which, together with  $F - E(G_1)$ , yield two distinct 1-factors of G. Thus at least one of the two edges of  $\{e_1, \ldots, e_k\}$  incident with  $G_1$  belongs to F. Since this is true for any component of  $G - \{e_1, \ldots, e_k\}$ , at least half of the edges  $\{e_1, \ldots, e_k\}$  belong to F. Since, in turn, this holds for any equivalence class, at least half of the edges of G belong to F. Hence  $|E(G)| \leq |V(G)|$ . Since G is 2-edge-connected, this is possible only, if G is a circuit. However, an (even) circuit has two 1-factors.

Second solution. Suppose G is 2-edge-connected. Let F be a 1-factor of G. Let C be a circuit such that any edge of F either lies on C or is completely disjoint from C. Such a circuit exists by 6.30. Now C must be alternating relative to F (since F is a 1-factor). Thus, "switching" on C (i.e. replacing the edges of  $E(C) \cap F$  by the edges of E(C) - F, we get another 1-factor [A. Kotzig,  $Mat.\ Fyz.\ \check{C}asopis\ 9\ (1959)\ 73-91$ ; L. Beineke—M. D. Plummer,  $J.\ Comb.\ Th.\ 2\ (1967)\ 285-289.$ ]

(c) The proof is by induction on |E(G)|.

The sharpening of (b) formulated in the hint follows from (b) trivially. It suffices to prove this for connected graphs, since if G has a unique 1-factor, so do its components. Double all edges of G not in F. This cannot produce any new 1-factors, so the resulting graph G' has a cut-edge by (b). This edge must belong to F, since it is not doubled.

Let e = (x, y) be a cut-edge of G, belonging to the unique 1-factor F of G, then G - x - y has a unique 1-factor. By the induction hypothesis,



 $E(G-x-y) \le (n-1)^2$ . Moreover, since (x, y) is a cut-edge, no point is joined to both x and y. Hence, at most 1 + (2n-2) = 2n - 1 edges are adjacent to one of x and y. Thus

$$|E(G)| \leq (n-1)^2 + 2n - 1 = n^2.$$

The result is sharp, as shown by graphs of structure similar to those in Fig. 56 [G. Hetyei, loc. cit. p. 307].

**25.** Let  $x_1, \ldots, x_{\delta}$  be the points not covered by a maximum matching F,  $\delta = \delta(G) = |V(G)| - 2\nu(G)$ . Let  $F = \{e_1, \ldots, e_i\}$ , and let  $f_i$  be an edge adjacent to  $x_i$   $(i = 1, \ldots, \delta)$ . Then each  $f_i$  joins  $x_i$  to a point covered by F. Moreover, no two  $f_i$ 's go to different endpoints of the same  $e_j$ , since this would make it possible to enlarge F. Therefore, we can select an endpoint  $u_j$  of  $e_j$   $(j = 1, \ldots, \nu)$  so that  $\{u_1, \ldots, u_{\nu}\}$  covers all  $f_1, \ldots, f_{\delta}$ . Since  $d(u_j) \leq d$ , we have

$$\delta \leq (d-1)v$$
 or  $n-2v \leq (d-1)v$ ,

and hence

$$v\geq rac{n}{d+1}$$
 .

**26.** Following the hint, observe that if x and y are adjacent, then  $v(G - x - y) < \langle v(G) \rangle$ , because any matching of G - x - y can be extended using (x, y). Let x, y be two points at distance k > 1, and let z be any inner point

Let x, y be two points at distance k > 1, and let z be any inner point of a minimum (x, y)-path. Suppose indirectly that v(G - x - y) = v(G). Let F be any maximum matching in G - x - y. Also, consider a maximum matching  $F_0$  in G - z; then  $|F| = |F_0| = v(G)$  (v for brevity). Observe that

- (a) F must cover z, otherwise, it would be a  $\nu$ -element matching of G-x-z, which contradicts the induction hypothesis  $\nu(G-x-z) < \nu$ , and
  - (b) similarly,  $F_0$  must cover x and y.

Now  $F \cup F_0$  is the union of some disjoint alternating paths, circuits and common edges of F and  $F_0$ . There is an alternating path  $P_1$  starting from x and such a path  $P_2$  starting from y. If  $P_1 = P_2$ , then replace the edges of  $F \cap E(P_1)$  by the edges of  $F_0 \cap E(P_1)$  in F; this way we get a (v+1)-element matching (since both end-edges of  $P_1$  belong to  $F_0$ ), which is a contradiction. Thus,  $P_1 \neq P_2$ . At least one of  $P_1$  and  $P_2$  does not contain z, say  $z \notin V(P_1)$ . Now replace the edges of  $F \cap E(P_1)$  by the edges of  $F_0 \cap E(P_1)$  in F. The resulting matching F' has  $|F'| \geq |F|$  (because  $P_1$  starts from x with an  $F_0$ -edge and so, if the last edge of  $P_1$  is in  $F_0$ , then |F'| > |F|, otherwise

|F'| = |F|). Moreover, z is still not covered by F' (since  $z \notin V(P_1)$ ) and thus F' is a matching of cardinality  $\geq v$  in G - y - z, again a contradiction.

Thus, v(G-x-y) < v(G) for every x, y. Let  $\overline{F}$  be any maximum matching of G. Then there is at most one point not covered by F, since if there were two such points x, y, then we would have

$$v(G - x - y) \ge |F| = v(G),$$

a contradiction. Moreover, F cannot be a 1-factor for if it were, then

$$|v(G-x)| \le \frac{1}{2} |V(G-x)| < \frac{1}{2} |V(G)| = |F| = |v(G)|,$$

which contradicts one of the hypotheses. Hence,

$$\nu(G) = |F| = \frac{1}{2}(|V(G)| - 1),$$

which proves the assertion [T. Gallai, MTA Mat. Kut. Int. Közl. 8 (1963) 135–139].

27. (a) Let  $X \subseteq V(G)$  and F a maximum matching of G. Let  $G_1, \ldots, G_k$  be the odd components of G - X. Let, say,  $G_1, \ldots, G_i$  be those containing a point not covered by F. From every  $G_j$ ,  $i < j \le k$ , there must be an edge of F going to X by parity. Since these edges have to go to different points of X, we have

$$k-i \leq |X|$$
.

On the other hand, each of  $G_1, \ldots, G_i$  contains an uncovered point, so

(1) 
$$\delta(G) \ge i \ge k - |X| = c_1(G - X) - |X|.$$

It remains to show — and this is the crux of the argument — that there exists a set X for which equality holds in (1). We show this by induction on |V(G)|.

Suppose first that there exists an  $x \in V(G)$  such that  $\nu(G - x) < \nu(G)$ . Obviously,  $\nu(G - x) = \nu(G) - 1$  and hence

$$\delta(G-x) = |V(G-x)| - 2\nu(G-x) = |V(G)| - 2\nu(G) + 1 = \delta(G) + 1.$$

Thus by the induction hypothesis, there exists an  $X' \subseteq V(G-x)$  such that

$$\delta(G) + 1 = c_1(G - x - X') - |X'|.$$

Now set  $X = X' \cup \{x\}$ . Then

$$\delta(G) = c_1(G - X) - |X|.$$

Secondly, suppose v(G - x) = v(G) for every  $x \in V(G)$ . Let  $G_1, \ldots, G_k$  be the components of G, and  $x \in V(G_i)$ . Then

$$v(G) = v(G_1) + \ldots + v(G_k) = v(G - x) =$$
  
=  $v(G_1) + \ldots + v(G_i - x) + \ldots + v(G_k),$ 

hence  $\nu(G_i - x) = \nu(G_i)$ . By the preceding problem, this implies that  $|V(G_i)|$  is odd and  $\delta(G_i) = 1$ . Thus

$$\delta(G) = \sum_{i=1}^k \delta(G_i) = k = c_1(G),$$

and  $X = \emptyset$  satisfies the requirement [C. Berge; see B].

(b) G has a 1-factor if and only if  $\delta(G) = 0$ , i.e. by (a), if and only if

(2) 
$$\max_{X} \left\{ c_1(G - X) - |X| \right\} = 0.$$

Since  $c_1(G) - |\emptyset| \ge 0$ , the maximum is always  $\ge 0$  and so, (2) is equivalent to

$$c_1(G-X)-|X|\leq 0$$

for every  $X \subseteq V(G)$  as stated [W. T. Tutte; see B].

**28.** (a) Let  $V_1$  be the set of those points of G joined to every other point and  $V_2 = V(G) - V_1$ . We want to show that if  $a, b, c \in V_2$ , and b is adjacent to both a and c, then a and c are adjacent.

Suppose this is not the case. Since  $b \in V_2$ , there is a fourth point d which is not adjacent to b. By the maximality assumption on G, G + (a, c) has a 1-factor  $F_1$  and G + (b, d) has a 1-factor  $F_2$ . Obviously,  $(a, c) \in F_1$ ,  $(b, d) \in F_2$ ,  $(b, d) \notin F_1$  and  $(a, c) \notin F_2$ .

Consider  $F_1 \cup F_2$ . This consists of disjoint alternating circuits and common edges of  $F_1$  and  $F_2$ . (a, c) is on an alternating circuit  $C_1$  and similarly, (b, d) is on an alternating circuit  $C_2$ .

Case 1.  $C_1 \neq C_2$ . Then replace the edges of  $E(C_1) \cap F_1$  by the edges of  $E(C_1) \cap F_2$  in  $F_1$  and get a 1-factor of G, a contradiction.

Case 2.  $C_1 = C_2$ . Starting from b through (b, d) and walking along  $C_1$ , sooner or later we must hit (a, c). Without loss of generality we may suppose we hit a before c.

So we have a (b, a)-path starting with the  $F_2$ -edge (b, d) and ending with another  $F_2$ -edge, since  $(a, c) \in F_1$ . Thus, K = P + (a, b) is a circuit alternating relative to  $F_2$ . Replacing the edges of  $E(K) \cap F_2$  by the edges of  $E(K) - F_2$  we get a 1-factor of G from  $F_2$ . This is again a contradiction.

Thus we have shown that adjacency is an equivalence relation on  $V_2$ , i.e.  $V_2$  decomposes into disjoint complete graphs.

(b) We only prove the non-trivial part of 7.27b, which says that if G has no 1-factor, then there is an  $X \subseteq V(G)$  with

$$c_1(G-X)>|X|.$$

If |V(G)| is odd,  $X = \emptyset$  has this property; so we may assume |V(G)| is eventet us add edges to G as long as we do not produce any 1-factor. When we have to stop we will have a graph G' on the same point-set as G with the property that G' has no 1-factor, but joining any two non-adjacent points of G' produces a graph with a 1-factor.

By (a), G' has the following form: if  $V_1$  is the set of those points connected to every other point, then  $G' - V_1$  consists of disjoint complete subgraphs

 $G_1,\ldots,G_k$ 

We claim there are at least  $|V_1|+1$  odd complete graphs among  $G_1,\ldots,G_k$ . Suppose not. Select a 1-factor of each even  $G_i$ , select a maximum matching of each odd  $G_i$ , this misses exactly one point. Select independent edges matching these points with some points in  $V_1$ . Since |V(G)| is even, there are an even number of points of  $V_1$  still uncovered. But we may match them arbitrarily with each other, because  $V_1$ , obviously, spans a complete subgraph of G'. So we get a 1-factor, and hence a contradiction.

Thus, there are at least  $|V_1| + 1$  odd components of  $G' - V_1$ . When the edges of E(G') - E(G) are removed, these odd components may fall apart, but each of them yields at least one odd component of  $G - V_1$ . Hence

$$c(G - V_1) > |V_1|,$$

i.e.  $X = V_1$  is an appropriate choice for X.

Remark: The reader may easily verify the fact (which was not needed here) that  $G_1, \ldots, G_k$  are all odd and  $k = |V_1| + 2$  [G. Hetyei, Pécsi Tanárképző Főiskola Közl. 1974; L. Lovász, J. Comb. Th. 19 (1975) 269–271].

29. (a) By Tutte's theorem, we have to show that

$$c_1(G-X) \leq |X|$$

for every  $X \subseteq V(G)$ .

Let  $G_1$  be any odd component of G-X. Since G is 2-connected, there are at least 2 edges joining  $G_1$  to X. There cannot be exactly two such edges, however, for this would imply the sum of the degrees of  $G_1$  is  $3|V(G_1)|-2$ , an odd number, which is impossible. So there are at least 3 edges joining  $G_1$  to X. Thus, there are at least  $3c_1(G-X)$  edges coming from the odd components of G-X to X. But each point of X is adjacent to at most 3 of them, so their number is at most 3|X|. Thus


$$3c_1(G-X) \leq 3|X|, \ c_1(G-X) \leq |X|$$

as required.

(Note that only 2-edge-connectivity of G was used. It is trivial to see though that, if a 3-regular graph is 2-edge-connected then it is also 2-connected.) [J. Petersen; for a proof not using Tutte's theorem, see K, S.]

(b) If e is a cut-edge of a 3-regular graph, then e is contained in every 1-factor (if any). In fact, the two components of G-e are odd (they contain exactly one point of even degree). Thus, G-e has no 1-factor, so each 1-factor, if any, contains e.

It therefore suffices to construct a simple cubic graph with three separating edges meeting at some point: any 1-factor of such a graph must contain all three of them, which is impossible. Figure 57 shows such a construction (the reader may wish to verify that this is the smallest such example).



30. By Tutte's theorem (7.27b), we only have to show that

$$c_1(G'-X) \leq |X|$$

for any  $X \subseteq V(G)$ . Suppose this does not hold for some X. Since |V(G)| is even and so

$$c_1(G'-X) \equiv |V(G')-X| \equiv |X| \pmod{2}$$

we have

$$c_1(G' - X) \ge |X| + 2.$$

Let  $G_1, \ldots, G_m$  be the odd components,  $G_{m+1}, \ldots, G_M$  the even components of G' - X. For  $1 \le i \le M$ , let  $\alpha_i$   $(\beta_i)$  be the number of edges of E(G) - E(G') joining  $G_i$  to X (to the other  $G_j$ 's), and let  $\gamma_i$  denote the number of edges of G' joining  $G_i$  to X. Then  $\alpha_i + \beta_i + \gamma_i$  is the total number of edges of G leaving  $G_i$ , which is at least k-1.

Moreover, if  $G_i$  is an odd component, then  $\alpha_i + \beta_i + \gamma_i > k - 1$ ; for  $\alpha_i + \beta_i + \gamma_i = k - 1$  would mean that the sum of degrees of the subgraph of G induced by  $V(G_i)$  is  $k|V(G_i)| - k + 1 = k(|V(G_i)| - 1) + 1 \equiv 1 \pmod{2}$ . Hence

(2) 
$$\sum_{i=1}^{m} \alpha_i + \sum_{i=1}^{m} \beta_i + \sum_{i=1}^{m} \gamma_i \ge k \cdot m.$$

The number of removed edges between two of  $G_1, \ldots, G_M$  and X is  $\sum_{i=1}^{M} \alpha_i + \frac{1}{2} \sum_{i=1}^{M} \beta_i$ , whence

$$2 \sum_{i=1}^{M} \alpha_i + \sum_{i=1}^{M} \beta_i \leq 2(k-1).$$

The total number of edges entering X is  $\sum_{i=1}^{M} \alpha_i + \sum_{i=1}^{M} \gamma_i$ , whence

$$\sum_{i=1}^{M} \alpha_i + \sum_{i=1}^{M} \gamma_i \leq k \cdot |X|.$$

Adding the last two inequalities,

$$3\sum_{i=1}^{M}\alpha_{i} + \sum_{i=1}^{M}\beta_{i} + \sum_{i=1}^{M}\gamma_{i} \leq k \cdot (|X| + 2) - 2.$$

Comparing this with inequality (2), we get

$$km \le \sum_{i=1}^{M} \alpha_i + \sum_{i=1}^{M} \beta_i + \sum_{i=1}^{M} \gamma_i \le 3 \sum_{i=1}^{M} \alpha_i + \sum_{i=1}^{M} \beta_i + \sum_{i=1}^{M} \gamma_i \le$$
  
 $\le k \cdot (|X| + 2) - 2 < k(|X| + 2)$ 

or, equivalently,


$$m < |X| + 2$$

which contradicts (1) [J. Plesnik, Mat. Čas. Slov. Akad. Vied. 22 (1972) 310–318].

**31.** Let F be a maximum matching which misses a given point x. Let y be a point adjacent to x and let  $F_0$  be a 1-factor of G-y.  $F \cup F_0$  consists of disjoint alternating circuits, common edges of F and  $F_0$  and an alternating (x,y)-path P. Now  $P_0=P+(x,y)$  is an odd circuit which contains  $\frac{1}{2}(|V(P_0)|-1)$  edges of F.

Now suppose that  $P_0, \ldots, P_i$  have been chosen so that for all  $j < i, P_{j+1}$  is an odd  $(P_0 \cup \ldots \cup P_j, P_0 \cup \ldots \cup P_j)$ -path or an odd circuit having exactly one point in common with  $P_0 \cup \ldots \cup P_j$  and  $P_{j+1}$  alternates relative to F (starting and ending with an edge not in F). If  $P_{j+1}$  happens to be an odd circuit, then we adopt the convention that it starts and ends at its common point with  $P_0 \cup \ldots \cup P_j$ . Also suppose  $P_0 \cup \ldots \cup P_i \neq G$ . Let (a, b) be any edge not in  $P_0 \cup \ldots \cup P_i$  such that  $a \in V(P_0 \cup \ldots \cup P_i)$  (such an edge exists since G is connected).

If  $b \in V(P_0 \cup \ldots \cup P_i)$ , we may take  $P_{i+1} = (a, b)$ . So suppose  $b \notin V(P_0 \cup \ldots \cup P_i)$ . Let  $F_i$  be a 1-factor of G - b.  $F \cup F_i$  consists of disjoint alternating circuits, common edges of F and  $F_i$ , and an alternating (b, x)-path Q. Traverse Q from b to the first point c of  $P_0 \cup \ldots \cup P_i$ . The last edge traversed



21\*

then belongs to  $F_i$ , because each F-edge meeting  $P_0 \cup \ldots \cup P_i$  belongs to this union by the construction. Let Q' be the piece of Q between b and c, and  $P_{i+1} = Q' + (a, b)$ . Then  $P_{i+1}$  satisfies the requirements (Fig. 58).

Thus if  $P_0 \cup \ldots \cup P_i \neq G$ , we can choose a  $P_{i+1}$ . Sooner or later we exhaust G, and the required decomposition is obtained [L. Lovász, *Studia Sci. Math. Hung.* 7 (1972) 279–280; also cf. the Edmonds Matching Aglorithm 7.34].

**32.** (a) Set  $A = A_G$ ,  $C = C_G$ , ...;  $A' = A_{G-x}$ ,  $C' = C_{G-x}$ , ... It suffices to prove that D' = D, since then C is the set of points not adjacent to D = D' and so is the same as C'.

Observe that since G-x contains no maximum matching of G, v(G-x)=v(G)-1. Hence if M is any maximum matching of G, M-x is a maximum matching of G-x. Since for each  $y \in D$  there is a maximum matching of G avoiding G and this yields such a maximum matching of G and G is the follows that

 $u \in D'$ , i.e.  $D \subset D'$ .

Conversely, let  $y \notin D$  and suppose indirectly that  $y \in D'$ . Then there is a maximum matching M' of G-x avoiding y. Let M be a maximum matching of G avoiding a neighbor  $z \in D$  of x. Consider  $M \cup M'$ . Since y is covered by M, but not by M', the component of  $M \cup M'$  containing y is a path P starting from y with an edge of M. P cannot end with an edge of M', since then replacing the edges of  $M \cap E(P)$  by the edges of  $M' \cap E(P)$  we would get a maximum matching of G avoiding G. Thus G ends with an edge of G. Now replace the edges of G on G with the edges of G on G avoiding G avoiding G avoiding G avoiding G of G avoiding G and G is new matching cannot lie in G or G. This implies that G ends at G are now G of G avoiding G and G is a maximum matching avoiding G a contradiction again.

(b) Let  $G_1, \ldots, G_t$  be those components of G - A contained in D. Since no edge connects D to C, these components partition D. Let H be the subgraph

induced by C.

If we remove the points of A one by one, (a) implies that the sets C, D do not change. Hence every maximum matching of G-A will cover all points of H, but each point of D will not be covered by some maximum matching of G-A. The maximum matchings of G-A consist of a maximum matching  $M_0$  of H and maximum matchings  $M_i$  of  $G_i$ ,  $i=1,\ldots,t$ . Since every point of H must be covered,  $M_0$  is a 1-factor of H. Moreover, since each point of  $G_i$  is avoided by some maximum matching of  $G_i$ ,  $G_i$  is factor-critical by 7.26.

(c) If M is any maximum matching of G then by the argument at the beginning of the proof, M-A is a maximum matching of G-A. In part (b), we have already seen what M-A must look like.

(d) By (b),

$$\nu(G-A) = \frac{1}{2} \big( |V(G)| - |A| - t \big).$$

By the remark above concerning removal of points of A one at a time,

$$\nu(G-A)=\nu(G)-|A|.$$


Hence the equality follows.

(e) Suppose that  $c_1(G-X) \ge |X|$  holds for every X. In particular,  $c_1(G-A) \ge |A|$ . On the other hand  $c_1(G-A) = t$ . Hence by (d),

$$v(G) \ge \frac{1}{2} |V(G)|,$$

i.e. G has a 1-factor. (The other direction of Tutte's theorem is trivial.) [T. Gallai, MTA Mat. Kut. Int. Közl. 8 (1963) 373–395; J. Edmonds, Canad. J. Math. 17 (1965) 449–467.]

(f) See Fig. 59.



**33.** I. Suppose M' is not a maximum matching in G'. Then G' has a matching  $M_0$  with  $|M_0| > |M'|$ .  $M_0$  is a matching in G which contains at most one point of C. Hence we can add k edges of C to  $M_0$  to get a matching  $M_1$  with

$$|M_1| = |M_0| + k > |M'| + k = |M|,$$

a contradiction.

II. Now suppose M' is a maximum matching of G'. Consider the sets D', A', C' defined for G' as in 7.32. Let c be the image of C in G'. Since M' avoids c,  $c \in D'$ . Hence if we "blow up" c, the component of G' - A' containing c becomes a component of G - A' containing C, also odd. Hence

$$c_1(G - A') = c_1(G' - A')$$


and

$$\begin{split} &v(G) \leq \frac{1}{2} \left\{ |V(G)| - c_1(G - A') + |A'| \right\} = \\ &= \frac{1}{2} \left\{ |V(G)| - c_1(G' - A') + |A'| \right\} = \\ &= \frac{1}{2} \left\{ |V(G')| - c_1(G' - A') + |A'| \right\} + k = |M'| + k = |M|, \end{split}$$

showing that M is a maximum matching in G.

**34.** (a) I. A forest with properties (\*)-(\*\*) certainly exists; e.g. let the forest consist of the points not covered by M and no edges. Let F be a maximal such forest. Then no outer point can be adjacent to V(G)-V(F), for by the properties of F,  $(G-V(F))\cap M$  is a 1-factor of G-V(F) (since all the uncovered points must be roots in F) and so, if  $a\in V(G)-V(F)$  is adjacent to the outer point b of F and  $(a,c)\in M$ , then (a,b) and (a,c) could be added to F.

II. If two outer points, x and y say, in distinct components of F are adjacent, then consider the paths P and Q connecting a and y to the roots of their components in F. Then  $R = P \cup Q \cup \{(a, y)\}$  is a path alternating relative to M and if we replace  $M \cap E(R)$  by E(R) - M, we get a matching larger than M (Fig. 60).



III. Now suppose that two outer points u, v of the same component of F are adjacent. Then they form with certain edges of F an odd circuit C. Let w be the point of C nearest the root r of this component of F (w = r may occur). "Switching" on the alternating path connecting w to r, we get a new matching

 $M_0$  such that  $M_0$  contains  $\frac{1}{2}(|V(C)|-1)$  edges of C and no other edge of

 $M_0$  meets C. By the preceding result, if we contract C, then  $M_0 = M_0 - E(C)$  will be a maximum matching in the resulting graph G' iff  $M_0$  (or, equivalently, M) is a maximum matching of G. Moreover, if we find in G' a matching larger than  $M_0$  we can easily extend it to a matching of G larger than M.

IV. Finally, suppose that the outer points of G are independent. Let A be the set of inner points. Then the outer points will be isolated in G - A; their number is

$$c_1(G-A) = |M \cap E(F)| + (|V(G)| - 2|M|),$$

while

$$|A| = |M \cap E(F)|.$$

Hence

$$|M| = \frac{1}{2} (|V(G)| + |A| - c_1(G - A)).$$

But by 7.27 the right-hand side is an upper bound for  $\nu(G)$ . Thus M is a maximum matching.

The desired algorithm is now clear by summarizing the above results. At each step, we have a matching M of G and a forest  $F \subset G$  satisfying (\*) and (\*\*). We consider the edges adjacent to outer points of F.

- 1° If there is an edge connecting an outer point of F to a point of V(G) V(F), we enlarge F as in part I.
- $2^{\circ}$  If there is an edge connecting two outer points in different components of F, we enlarge M as in part II.
- $3^{\circ}$  If there is an edge connecting two outer points in the *same* component of F, we contract an odd circuit as in part III and try to enlarge the resulting matching of the resulting graph. We can either conclude that M is maximal or enlarge it.
- $4^{\circ}$  If all edges adjacent to outer points connect them to inner points, we conclude that M is a maximum matching.

Remarks. 1. The above algorithm is efficient in the sense that it performs in  $O(n^4)$  steps. In fact, let f(n) denote the least upper bound on the number of steps the algorithm needs to check whether a given matching M of an n-point




Fig. 61

graph is maximum and replace it by a larger one if not. It takes  $O(n^2)$  steps to find a maximal forest with properties (\*) and (\*\*). (We carry a list of certain outer points. At each step we check whether or not the first point on the list is adjacent to any point ouside F. If so, we enlarge F, and put the new outer point on the end of the list. If not, we cancel this outer point from the list. One such step needs the inspection of O(n) adjacencies. At each step either V(F) or the number of cancelled outer points increases, so the number of steps is O(n).) Then we have to carry out one of the steps  $2^{\circ}$ ,  $3^{\circ}$ ,  $4^{\circ}$ . Since this takes, in the worst case when step  $3^{\circ}$  must be carried out, at most  $f(n-2) + O(n^2)$  steps, we get that

$$f(n) = f(n-2) + O(n^2),$$

whence  $f(n) = O(n^3)$ . Since the size of the matching is increased at most O(n)

times, the whole algorithm finishes in no more than  $O(n^4)$  steps.

- 2. The following observation is very significant: If we carry out step  $3^{\circ}$ , the forest F is mapped onto a forest of the same kind for resulting graph. Using this one can improve the upper bound on the length of the algorithm to  $O(n^3)$ . The same observation and a more careful analysis of the algorithm would enable us to avoid the use of 7.33 (and through this, the use of the Gallai–Edmonds Structure Theorem). We also remark that the sets  $A_G$ ,  $C_G$ ,  $D_G$  can be determined and several previous results (e.g. 7.26, 7.27, 7.31, 7.32) can be deduced from the algorithm. The reader may find it interesting and instructive to work out the details.
- (b) See' Fig. 61 (cf. also 7.32f) [J. Edmonds, Canad. J. Math. 17 (1965) 449–467].
- **35.** Suppose G has no 1-factors. Consider the sets A, C, D as in 7.32. Let  $G_1, \ldots, G_t$  be components of G[D], then by 7.32 they are factor-critical. 7.31 implies (it is also easy to see directly) that if  $|V(G_i)| > 1$ , then  $G_i$  con-

tains an odd circuit. Hence, at most one of  $G_1, \ldots, G_t$  has more than one point. Let, say,

$$|V(G_1)| \ge 1$$
,  $|V(G_2)| = |V(G_3)| = \ldots = |V(G_t)| = 1$ .

Now there are exactly k edges joining  $G_i$  to A for  $i = 2, \ldots, t$ . Since each point in A is adjacent to at most k of these, we get

$$k(t-1) \le k \cdot |A|$$

or, equivalently,

$$t \leq |A| + 1$$
.

Now since

$$t = c_1(G-A) \equiv |V(G-A)| \equiv |A| \pmod{2},$$

this implies that

$$t \leq |A|, \ \ v(G) = \frac{1}{2} \left\{ |V(G)| - t + |A| \right\} \geq \frac{1}{2} |V(G)|,$$

whence G has a 1-factor, a contradiction [D.R. Fulkerson, A.J. Hoffman, M.H. McAndrew, Canad. J. Math. 17 (1965) 166-177].

**36.** (a) Let x be as in the hint, and (x, y) any edge incident with x. y must be covered by some edge e of F, otherwise F + (x, y) would be a larger matching. Now F - e + (x, y) is a maximum matching containing (x, y).

(b) We use induction on |E(G)|. By 7.24b we know that G has two distinct 1-factors  $F_1$ ,  $F_2$ . Let C be a circuit in  $F_1 \cup F_2$ . If some point of C is of degree 2 we are done, since then the two edges adjacent to it belong to the 1-factors


 $F_1$  and  $F_2$ , respectively.

So suppose each point of C has degree at least 3. We claim that G-e is 2-connected for some  $e \in E(C)$ . This will settle the problem; for G-e has a 1-factor (e belongs to exactly one of  $F_1$  and  $F_2$ ) and so by the induction hypothesis, there is a point x such that each edge of G-e adjacent to x, belongs to some 1-factor of G-e. Since e belongs to a 1-factor, x will have the same property in G.

Suppose, indirectly, that G - e has a cut-point  $x_e$  for each  $e \in E(C)$ . Clearly  $x_e \in V(C)$ . Let e be chosen in such a way that  $x_e$  is as close to e on C as possible.

Obviously,  $x_e$  is not an endpoint of e.

Let f be the edge of C adjacent to e on the shorter  $(e, x_e)$ -arc of C (Fig. 62).



Then by the choice of e,  $x_f$  must lie on the other  $(e, x_e)$ -arc. Now the common point y of e and f is adjacent to a third edge (y, z). Since G - y is connected, z is joined to some point of C - y by a path P. Now it is seen that no matter where P hits C, we get a contradiction to the fact that both  $\{e, x_e\}$  and  $\{f, x_f\}$  separate G. [J. Zaks, Combin. Structures Appl. Gordon and Breach (1970) 481-488.]

**37.** I. Let w be any 2-matching and choose an independent set  $X \subseteq V(G)$ . Set

$$\overline{w}(x) = \sum_{\substack{e \in E(G) \ e \ni X}} w(e), \quad \delta_X = \sum_{x \in X} (2 - \overline{w}(x)).$$

The total w-weight of edges going from X to  $\Gamma(X)$  is

$$2|X|-\delta_X$$

on the other hand, it is  $\leq 2 |\Gamma(X)|$ , whence

$$\delta_X \ge 2 |X| - 2 |\Gamma(X)|.$$

Thus,

$$2 |w| = \sum_{x \in V(G)} \overline{w}(x) = 2 |V(G)| - \sum_{x \in V(G)} (2 - \overline{w}(x)) \le$$

$$\le 2 |V(G)| - \sum_{x \in X} (2 - \overline{w}(x)) = 2 |V(G)| - \delta_X \le 2 (|V(G)| - |X| + |\Gamma(X)|).$$

This proves that the maximum size of a 2-matching is

$$\leq |V(G)| - \max_{X \text{ indep}} (|X| - |\Gamma(X)|).$$

II. First we remark that

$$\max_{X \text{ indep}} \left\{ |X| - |\Gamma(X)| \right\} = \max_{\text{all } X} \left\{ |X| - |\Gamma(X)| \right\}.$$

In fact, for an arbitrary  $X \subseteq V(G)$  define  $X' = X - \Gamma(X)$ . Then X' is independent, and  $\Gamma(X') \subseteq \Gamma(X) - X$ . Hence

$$|X| - |\Gamma(X)| = |X'| - |\Gamma(X) - X| \le |X'| - |\Gamma(X')|$$

showing that the maximum of  $|X| - |\Gamma(X)|$  is attained by some independent set X.

Let  $G_0$  be the bipartite graph defined in the hint. Observe that for  $X \subseteq V(G)$ ,  $|\Gamma_{G_0}(X)| = |\Gamma(X)|$ . Therefore, we have

$$\delta = \max_{X \subseteq V(G)} \{|X| - |\Gamma_{G_0}(X)|\} = \max_{X \subseteq V(G)} \{|X| - |\Gamma(X)|\}.$$

By 7.5,  $G_0$  has a set F of  $|V(G)| - \delta$  independent edges. For  $e = (u, v) \in E(G)$  let  $w_0(e)$  denote the number of edges of F among (v', u') and (u', v'); so  $w_0(e) = 0$  1 or 2. Moreover, for any fixed v, there is at most one edge of F leaving v and at most another one leaving v'. Hence

$$\overline{w}_0(v) = \sum_{\substack{e \in E(G) \ e \ni v}} w_0(e) \le 2,$$

i.e.  $w_0$  is a 2-matching. Moreover,

$$|w_0| = \sum_{e \in E(G)} w_0(e) = |F| = |V(G)| - \delta,$$

whence

$$\max_{w \text{ a 2-matching}} |w| \geq |w_0| = |V(G)| - \delta = |V(G)| - \max_{X \subseteq V(G)} \left\{ |X| - |\Gamma(X)| \right\}$$

[cf. W. T. Tutte, Proc. Amer. Math. Soc. 4 (1953) 922-931].

**38.** I. Observe that a 2-matching of size |V(G)| is necessarily maximum. Therefore, if G has a 1-factor F, then taking the edges of F with weight 2, we get a maximum 2-matching containing the maximum 1-matching F.

II. Now let G be factor-critical and suppose |V(G)| > 1. Let F be any maximum matching of G. The solution of 7.31 shows that there exists an odd circuit

 $P_0$  such that F has  $\frac{1}{2}|V(P_0)|-1$ ) edges on  $P_0$ ; i.e.  $F-E(P_0)$  is a 1-factor

of 
$$G - V(P_0)$$
. Set 
$$w(e) = \begin{cases} 1 & \text{if } e \in E(P_0), \\ 2 & \text{if } e \in F - E(P_0), \\ 0 & \text{otherwise} \end{cases}$$

then w is a 2-matching of size |V(G)|, containing F.

III. Now suppose G is not factor-critical and has no 1-factor. Consider the Gallai-Edmonds decomposition of G (7.32). Let  $G_1, \ldots, G_t$  be the components of  $G[D_G]$ ; let, say,

$$|V(G_1)| = \ldots = |V(G_k)| = 1 < |V(G_{k+1})|, \ldots, |V(G_t)|,$$
  
 $V(G_i) = \{y_i\}, \quad (i = 1, \ldots, k), \quad Y = \{y_1, \ldots, y_k\}.$ 

Also set, as before

$$\delta = \max_{X \subseteq A(G)} \{ |X| - |\Gamma(X)| \}.$$

Since, therefore,

$$|\Gamma(X)| \ge |X| - \delta$$

holds for every  $X \subseteq Y$ , we get from 7.5, applied with the bipartite graph formed by the edges incident with Y, that there are  $|Y| - \delta = k - \delta$  independent edges joining Y to  $\Gamma(Y) \subseteq A$ . We may assume  $y_1, \ldots, y_{k-\delta}$  can be matched with  $k - \delta$  points of A.

By 7.23, there exists a maximum matching F of G covering  $y_1, \ldots, y_{k-\delta}$ . We define a maximum 2-matching w containing F as follows: let  $G_{k+1}, \ldots G_m$  be joined to A by some edge of F ( $m \le t$ ). By 7.32,  $F \cap E(G_i)$  is a maximum matching in  $G_i$  ( $i = m + 1, \ldots, t$ ), and hence, as in part II of the proof, we can define a 2-matching  $w_i$  of  $G_i$  of size  $|V(G_i)|$  containing  $F \cap E(G_i)$ , for  $i = m + 1, \ldots, t$ . Now, for each  $e \in E(G)$ , let

$$w(e) = \begin{cases} w_i(e) & \text{if} \quad e \in E(G_{m+1}) \cup \ldots \cup E(G_t), \\ 2 & \text{if} \quad e \in F - E(G_{m+1}) - \ldots - E(G_t), \\ 0 & \text{otherwise.} \end{cases}$$

Then, obviously, w is a 2-matching which contains F. We have to show w is maximal. Observe that exactly two edges of w contain each point except  $y_{k-\delta+1}, \ldots, y_k$ , which are contained in no edge of w. Thus

$$w = |V(G)| - \delta,$$

which shows by 7.37 that w is a maximum 2-matching.

**39.** Let L be an Euler trail of G. Consider every second edge of L. Since the total number of edges is even, this is possible. Then the considered edges form a d-factor.

**40.** Let  $\vec{G}$  be a pseudosymmetric orientation of G (see 5.13), set  $V(G) = \{v_1, \ldots, v_n\}$  and define a bipartite graph  $G_0$  by setting

$$\begin{split} V(G_0) &= \{v_1, \ldots, v_n, \ v_1', \ldots, v_n'\}, \\ E(G_0) &= \{(v_i, v_j') : (v_i, v_j) \in E(\vec{G})\}. \end{split}$$

Observe that  $G_0$  is a d-regular bipartite graph and G arises from it by identifying v, and v' for each  $1 \le i \le n$ 

ing  $v_i$  and  $v_i'$  for each  $1 \le i \le n$ . Let us color the edges of  $G_0$  by d colors so that each color forms a 1-factor (this is possible by 7.10). Identifying  $v_i$  and  $v_i'$ , this yields a d-coloration of the edges of G such that each color forms a 2-factor. (Note the difference between this and the construction in the proof of 7.37; there two edges of the bipartite graph were mapped onto the same edge of G, therefore we could only conclude that a matching of the bipartite graph yields a 2-matching of the graph; here no two edges are mapped onto the same edge, therefore a 1-factor of the bipartite graph yields a 2-factor of the graph.)

7.41

41. First suppose all degrees are even. If the number of edges is even, the same proof as in 7.39 gives a desired subgraph; i.e. consider an Euler trail and take every second edge of it. On the other hand, if such a subgraph F exists, then

$$2 |E(F)| = \sum_{x \in V(G)} d_F(x) = \sum_{x \in V(G)} \frac{d_G(x)}{2} = |E(G)|,$$

whence |E(G)| is even.

Now suppose there are points with odd degree. Let v be a new point and join v to all points of G of odd degree. Adding a loop at v if necessary, we thereby construct a graph G' with even degrees and with an even number of edges. Thus again by 7.39, G' has a spanning subgraph F' with degrees

$$d_{F'}(x) = \frac{1}{2} d_{G'}(x) \qquad (x \in V(G')).$$

Now set F = F' - v. Then, for any  $x \in V(G)$ ,

$$d_F(x) = d_{F'}(x) = \frac{1}{2} d_{G'}(x) = \frac{1}{2} d(x)$$
 if  $d(x)$  is even,

$$d_F(x)=d_{F'}(x)=rac{1}{2}\,d_{G'}(x)=rac{d(x)+1}{2} ext{ if } d(x) ext{ is odd and } (x,v) 
otin E(F),$$

$$d_F(x) = d_{F'}(x) - 1 = \frac{d(x) - 1}{2} \qquad \qquad \text{if $d(x)$ is odd and $(x, v) \in E(F)$.}$$

So all connected graphs have this property except Eulerian graphs with an odd number of edges.

**42.** (a) Since a graph with odd degrees has an even number of points, G must have an even number of points.

Conversely, suppose |V(G)| is even, say  $V(G) = \{x_1, \ldots, x_{2m}\}$ . Let  $P_i$  be a path connecting  $x_i$  to  $x_{i+m}$   $(i = 1, \ldots, m)$ , and set

$$F = \sum_{i=1}^{m} E(P_i) \tag{mod 2}$$

Then for each  $x \in V(G)$ ,

$$d_F(x) \equiv \sum_{i=1}^m d_{P_i}(x) \equiv 1 \pmod{2},$$

and F is a spanning subgraph of the type desired.

(b) Let  $x \in V(G)$ ,  $d_G(x) \ge 4$ . Let us consider two edges (x, y), (x, z) of G such that G - (x, y) - (x, z) is connected. (There exist two such edges, for let y, z lie in distinct components of G - x if G - x is not connected. Then G - (x, y) - (x, z) is connected, if G - x is connected, because  $d_G(x) \ge 3$ 

and also if G - x is disconnected since each component of G - x is joined to x by at least two edges.) Take a new point x' and connect it to x, y and z. Contracting the edge (x, x') we recover G.

Repeating the above procedure we get, in a finite number of steps, a 3-regu-

lar graph G', for the sum

$$\sum_{x \in V(G)} \left( d_G(x) - 3 \right)$$

decreases at each step. G is a contraction of G' and G' is 2-edge-connected

(and thus, as remarked in 7.29a, 2-connected).

Hence G' contains a 1-factor F by 7.29a. Then F' = E(G') - F is a 2-factor. Contracting the new edges to get G from G', F' is mapped onto a subgraph of G with positive even degrees.

- **43.** (a) Replace each point x by a set  $M_x$  of f(x) independent points. If  $(x, y) \in E(G)$ , then each point of  $M_x$  is joined to each point of  $M_y$ , otherwise no edge connects  $M_x$  to  $M_y$ . It is straightforward to check that the resulting graph has a 1-factor iff G has an f-matching.
- **44.** Let  $\vec{G}'$  be the digraph defined in the hint; i.e. set  $V' = \{x' : x \in V(G)\}$ ,  $V'' = \{x'' : x \in V(G)\}$ ,  $V(\vec{G}') = V' \cup V''$  and  $E(\vec{G}') = \{(x', y'') : (x, y) \in E(G)\}$ . Let G' arise by forgetting about the orientations. Define

$$h(y) = \begin{cases} g(x) & \text{if} \quad y = x', \\ f(x) & \text{if} \quad y = x'' \end{cases} \quad \big(x \in V(G)\big).$$

Obviously, G has a desired factor if and only if G' has an h-factor. By 7.16, this is so if and only if

 $\sum_{x \in V'} h(x) = \sum_{y \in V''} h(y)$ 

and for every  $X \subseteq V'$ ,  $Y \subseteq V''$ ,

$$\sum_{x \in X} h(x) \leq \sum_{y \in Y} h(y) + m_G, (X, V'' - Y).$$

In other words, if and only if

$$\sum_{x \in V(G)} f(x) = \sum_{x \in V(G)} g(x)$$

and for all  $X, Y \subseteq V(G)$ ,

$$\sum_{x \in X} g(x) \leq \sum_{y \in Y} f(y) + m_G(X, V(G) - Y),$$

where  $m_G(X, Z)$  denotes the number of edges with tail in X and head in Z.

**45.** Following the hint, let  $G_1$  be the graph obtained by subdividing the edges of G and extend f to  $V(G_1)$  as described. Observe that G has a desired orientation if and only if  $G_1$  has an f-factor.

 $G_1$  is, obviously, bipartite with color classes V(G) and  $V(G_1) - V(G)$ . Thus

by 7.16,  $G_1$  has an f-factor if and only if

(1) 
$$\sum_{x \in V(G)} f(x) = |E(G)|$$

and

(2) for each  $X \subseteq V(G)$ ,  $Y \subseteq E(G)$  the number of incidences between X and Y is not less than

$$\sum_{x \in X} f(x) - |E(G) - Y|.$$

Obviously, it suffices to require this for the special case when there is no incidence between X and Y, but Y contains all edges not incident with X. Then (2') says that

- (2') at least  $\sum_{x \in X} f(x)$  edges are incident with X, for every  $X \subseteq V(G)$ .
- (1) and (2') are the necessary and sufficient conditions for G being orientable in the desired way.
- **46.** (a) We use induction on the number of points. Suppose first that G is a circuit,  $V(G) = \{x_1, \ldots, x_n\}$  and  $E(G) = \{(x_1, x_2), \ldots, (x_n, x_1)\}$ . If g(x) > 0 for every  $x_i \in V(G)$ , we can take  $F = (V(G), \emptyset)$ . So suppose, e.g.  $g(x_1) = 0$ . Then put  $(x_1, x_2)$  into F. Going along the cycle, put  $(x_i, x_{i+1})$  into F whenever  $(x_{i-1}, x_i) \in F$  and  $g(x_i) = 1$  or  $(x_{i-1}, x_i) \notin F$  and  $g(x_i) = 0$ . Deciding this way about every edge we end up with 1 or 2 F-edges at  $x_1$ , thus  $d_F(x_1) \neq g(x_1) = 0$ . The other points satisfy the requirement, because F has been constructed that way.

Now suppose that G is not a single circuit. Then we find a point  $x \in V(G)$  such that  $G_1 = G - x$  is connected and has a circuit. Define  $g_1$  on  $V(G_1)$  as follows: if g(x) > 0, we set  $g_1 = g$ ; if g(x) = 0, we select a neighbor g(x) = 0 of g(x) = 0.

$$g_1(z) = \begin{cases} g(z) & \text{if } z \neq y, \\ g(y) - 1 & \text{if } z = y. \end{cases}$$

By the induction hypothesis,  $G_1$  has a spanning subgraph  $F_1$  with degree different from  $g_1$ . In case g(x) > 0,  $F = F_1$  is a desired subgraph of G. If g(x) = 0,  $F = F_1 \cup \{(x, y)\}$  has the required property.

- (b) I. Suppose both (i) and (ii) hold; let F be a spanning subgraph with degrees  $\neq g(x)$  and  $\vec{G}$  an orientation with indegrees g(x). Reverse the orientation of the edges of F. Since the resulting digraph is acyclic, it has a point x with indegree 0. However, reversing the edges of F again, x will have indegree g(x), whence it is adjacent to g(x) edges of F, a contradiction.
- II. Let G be any tree, we show either (i) or (ii) is satisfied. Let x be a point of degree 1. If g(x) < 0 or g(x) > 1, then (ii) is satisfied. For add a loop to x, apply (a), and then remove the loop. We get a spanning subgraph having degree  $\neq g(y)$  at each  $y \neq x$  and of course, its degree is different from g(x) at x.

Thus we may assume g(x) = 0 or g(x) = 1. Suppose, e.g. g(x) = 1 (the other case follows similarly). If G - x has a spanning subgraph F with degrees  $\neq g(y)$  for each  $y \in V(G) - \{x\}$ , the same subgraph (with x as an isolated point) satisfies the requirements for G; thus (b) holds again. Suppose, therefore, that no such F exists. Then, by the induction hypothesis, G - x has an orientation with indegrees g(y) for each  $y \in V(G) - x$ . Orient the edge adjacent to x toward x. Then the resulting orientation shows that (i) holds [L. Lovász,  $Periodica\ Math.\ Hung.\ 4\ (1973)\ 121-123$ ].

**47.** The necessity is obvious as a tree on n points has n-1 edges. Now suppose that  $d_1+\ldots+d_n=2n-2$ . We use induction on n. The case  $n\leq 2$  is trivial, so suppose  $n\geq 3$ . Then  $d_1=1$ , since  $d_1\geq 2$  would imply  $d_1+\ldots+d_n\geq 2n>2n-2$ . Also  $d_n>1$  as  $d_n=1$  would imply  $d_1+\ldots+d_n=n<<2n-2$ . Since

$$d_{2} + \ldots + d_{n-1} + (d_{n} - 1) = 2n - 4 = 2(n-1) - 2,$$

there is a tree T with degrees  $d_2, \ldots, d_{n-1}, d_n - 1$ . Add a new point and join it to the point of T with degree  $d_n - 1$ ; then the resulting tree has the desired degrees [see B].

48. Again the necessity is obvious. We prove the sufficiency using induction on  $\sum_{i=1}^{n} d_i$ . We distinguish two cases.

I.  $d_{n-2} < d_n$ . Then  $d_n - 1$  is a largest element in  $d_1, d_2, \ldots, d_{n-2}, d_{n-1} - 1$ ,  $d_n - 1$  and so what we have to verify is

$$(1') d_1 + \ldots + d_{n-2} + (d_{n-1} - 1) + (d_n - 1) \equiv 0 \pmod{2},$$

$$(2') d_1 + \ldots + d_{n-2} + d_{n-1} - 1 \ge d_n - 1,$$

which are obvious from (1) and (2).

II.  $d_{n-2} = d_n$ . Then  $d_{n-1} = d_n$ . Obviously, (1') holds and

$$(2'') d_1 + \ldots + d_{n-3} + (d_{n-1} - 1) + (d_n - 1) \ge d_{n-2},$$

provided  $d_{n-2} \geq 2$ . If  $d_{n-2} = 1$ , then the left-hand side of (2") is odd by (1), and is therefore at least 1 trivially.

Thus  $d_1, \ldots, d_{n-2}, d_{n-1}-1, d_n-1$  satisfy the assumptions of the problem and by induction, there exists a graph without loops on n points realizing this sequence. Joining the points with degree  $d_{n-1}-1$  and  $d_n-1$  by a new edge, we get a graph with degrees  $d_1, \ldots, d_n$  [see B].

**49.** Let K be the simple digraph without loops defined by  $V(K) = \{v_1, \ldots, v_n\}$ ,  $E(K) = \{(v_i, v_j) : i \neq j\}$ . The existence of a digraph with the desired properties is equivalent to the existence of a subgraph of K with given outdegrees and indegrees. By 7.44, this is equivalent to (1) and (2), since the number of edges of K connecting  $\{v_i; i \in I\}$  to  $\{v_j; j \notin J\}$  is |I|(n-|J|)-|I-J|. (Remember here that K has no loops!)

Now if  $f_1 \leq \ldots \leq f_n$  and  $g_1 \leq \ldots \leq g_n$ , then for every I, J with |I| = k, |J| = l, we have

$$\sum_{I} f_{i} \leq \sum_{i=n-k+1}^{n} f_{i},$$

as well as

$$\sum_{j \in J} g_j + k(n-l) - |I - J| \ge \sum_{j=1}^l g_j + k(n-l) - \min(k, n-l).$$

Thus (2) can be replaced by

(2') 
$$\sum_{i=n-k+1}^{n} f_i \leq \sum_{j=1}^{l} g_j + (n-l)k - \min(k, n-l).$$

For (2') is, of course, a special case of (2) (with  $I = \{n - k + 1, \ldots, n\}$  and  $J = \{1, \ldots, l\}$ ), so it is necessary; and moreover as we have seen (2') implies (2) under the assumption  $f_1 \leq \ldots \leq f_n, g_1 \leq \ldots \leq g_n$  [see B].

**50.** Suppose first that there exists a simple graph G with degrees  $d_1, \ldots, d_n$  and let  $v_i$  be the point with degree  $d_i$ . Also set  $p=n-d_n$ . We claim G can be chosen so that  $v_n$  is adjacent to each of  $v_1, \ldots, v_{n-1}$  (but to no other point). For let us choose G among all graphs with  $d_G(v_i) = d_i$  such that  $v_n$  is adjacent to as many points in  $\{v_p, \ldots, v_{n-1}\}$  as possible. Suppose  $v_n$  is not adjacent to  $v_k$ , for some  $p \leq k \leq n-1$ . Then  $v_n$  must be adjacent to some  $v_v$ ,  $1 \leq v \leq 2n-1$ , since  $d_n = n-p$ . Since  $d_v \leq d_k$ , there must be a point  $v_m, m \neq k, v$ , adjacent to  $v_k$  but not to  $v_v$ . Now remove the edges  $(v_k, v_m)$  and  $(v_v, v_n)$ , but add  $(v_k, v_n)$  and  $(v_v, v_m)$ . This way a graph with the same degrees is obtained in which  $v_n$  is adjacent to more points in  $\{v_p, \ldots, v_{n-1}\}$  than in G, a contradiction.

Thus we have a graph G with degrees  $d_1, \ldots, d_n$  in which all of  $v_p, \ldots, v_{n-1}$  are adjacent to  $v_n$ . Removing  $v_n$  we obtain a graph with degrees  $d_1, \ldots, d_{p-1}, d_p-1, d_{p+1}-1, \ldots, d_{n-1}-1$ .

Conversely, suppose the numbers  $d'_k$  are degrees of a simple graph G'. We can then add  $v_n$  trivially [V. Havel, Cas. Pest. Mat. 80 (1955) 477-480].

51. (a) Following the hint, choose H with the given out- and indegrees such that the number of 2-cycles in H is maximal. Let  $\tilde{H}$  denote the subgraph formed by those edges  $(x, y) \in E(H)$  for which  $(y, x) \notin E(H)$ .

Now  $\widetilde{H}$  contains no even cycle. For suppose  $(x_1, \ldots, x_{2k})$  is an even cycle of  $\widetilde{H}'$ . Then remove  $(x_2, x_3), \ldots, (x_{2k}, x_1)$ ; but add  $(x_2, x_1), (x_4, x_3), \ldots, (x_{2k}, x_{2k-1})$ . We thereby get a graph with the same out- and indegrees

but with more 2-cycles.

The same argument shows that  $\widetilde{H}$  cannot have any closed directed trail with an even number of edges. Since  $\widetilde{H}$  has equal indegrees and outdegrees, it is the union of edge-disjoint cycles (cf. 5.6). By the above, these are odd and no two can have a point in common, because they would then form a trail with an even number of edges.

Suppose there are two cycles  $C_1, C_2$  in the above cycle-decomposition of  $\widetilde{H}$ ,  $C_1 = (x_0, \ldots, x_{2k})$  and  $C_2 = (y_0, \ldots, y_{2l})$ . If  $(x_0, y_0) \in E(H)$ , then  $(x_0, y_0) \notin E(\widetilde{H})$  by the above, and so  $(y_0, x_0) \in E(H)$  and  $C_1 \cup C_2 \cup \{(x_0, y_0), (y_0, x_0)\}$  is a trail with an even number edges, whence we get a contradiction as above. So  $(x_0, y_0), (y_0, x_0) \notin E(H)$ . Then remove  $(x_0, x_1), (x_2, x_3), \ldots, (x_{2k}, x_0), (y_0, y_1), (y_2, y_3), \ldots, (y_{2l}, y_0)$ , but add  $(x_2, x_1), \ldots, (x_{2k}, x_{2k-1}), (y_2, y_1), \ldots$ 

...,  $(y_{2l}, y_{2l-1})$ ,  $(x_0, y_0)$ ,  $(y_0, x_0)$ . We again get a simple digraph with the same degrees and more 2-cycles.

Thus  $\widetilde{H}$  contains at most one cycle. If it contains exactly one, then this is odd and  $\sum_{i=1}^{n} d_i = |E(H)|$  is odd, a contradiction of the assumption. Thus  $\widetilde{H}$  has no edge; i.e. H consists of pairs of opposite edges. Replacing each such edge by a single undirected edge we get a simple graph with degrees  $d_1, \ldots, d_n$  [cf. D.R. Fulkerson, A.J. Hoffman, M.H. McAndrew, Canad. J. Math. 17 (1965) 166–177].

(b) By (a), the existence of a simple graph with degrees  $d_1, \ldots, d_n$  is equivalent to the existence of a simple digraph H on  $\{v_1, \ldots, v_n\}$  such that  $d_H^+(v_i) = d_H^-(v_i) = d_i$   $(i = 1, \ldots, n)$  (under the necessary assumption that  $d_1 + \ldots + d_n$  is even). By 7.49, such a digraph exists iff for every  $k, l \in \{1, \ldots, n\}$ ,

(2') 
$$\sum_{i=n-k+1}^{n} d_i \leq \sum_{j=1}^{l} d_j + k(n-l) - \min(k, n-l).$$

It suffices to require (2') for the case  $k+l \leq n$ . In fact, if k+l > n, then (2') is equivalent to

(2") 
$$\sum_{i=l+1}^{n} d_i = \sum_{j=1}^{n-k} d_j + k(n-l) - \min(k, n-l),$$

which can also be derived from (2') with n-l and n-k playing the roles of k and l, respectively.

Now if  $k+l \leq n$ , then

$$\sum_{j=1}^{l} d_j + k(n-l) \ge \sum_{j=1}^{n-k} \min(d_j, k) + k^2.$$

So if

(2) 
$$\sum_{i=n-k+1}^{n} d_i \leq \sum_{j=1}^{n-k} \min(d_j, k) + k^2 - k,$$

then (2') is satisfied. On the other hand, (2) is a special case of (2') when we take  $m = \max\{j: d_j \leq k\}$  and  $l = \min\{n-k, m\}$ . So (2) is a necessary condition [P. Erdős, T. Gallai; see B].

**52.** Suppose first that  $d_1, \ldots, d_n$  are the degrees of a connected graph. Then 7.51 implies (1), while (2) and (3) are trivial.

Suppose, conversely, that  $d_1, \ldots, d_n$  satisfy (1), (2) and (3). By (1), there is a simple graph G with degrees  $d_1, \ldots, d_n$ . Choose G so that it has the least

possible number of components.

We claim that G is connected. Suppose not, then by (3), one of its components contains a circuit. Let  $G_1$  be this component and (x, y) an edge of  $G_1$  on a circuit. Let  $G_2$  be any other component and (u, v) any edge of  $G_2$   $(G_2$  has an edge since it is not an isolated point by (2)). Then  $G = \{(x, y), (u, v)\}$ 

 $+\{(x, u), (y, v)\}\$  is a graph on the same set, with the same degrees and with fewer components, a contradiction [P. Erdős, T. Gallai; B].

**53.** Let G be a graph with a 1-factor F and with points  $v_1, \ldots, v_n$  such that  $d(v_i) = d_i$ . Then, obviously G - F has degree sequence  $d_1 - 1, \ldots, d_n - 1$ . To prove the converse, let G be any simple graph on  $V = \{v_1, \ldots, v_n\}$  with  $d_G(v_i) = d_i$  and G' another simple graph on V with  $d_{G'}(v_i) = d_i - 1$ . Moreover, among all such pairs of graphs, choose a pair with |E(G) - E(G')| minimal. We claim  $G' \subseteq G$ . Suppose not, then there are points incident with edges of E(G') - E(G). Let X be one adjacent to the maximum number X of edges of X of X of X be one adjacent in X of X degree of X of X degree of X deg

Since  $d_G(z) > d_{G'}(z)$ , there is a point v such that  $(z, v) \in E(G) - E(G')$ . Choose an  $1 \le i \le k+1$  such that  $y_i \ne v$ . Then  $(v, y_i) \in E(G)$ . For if  $(v, y_i) \notin E(G)$ , then removing  $(x, y_i)$  and (z, v) from G, but adding (x, z) and  $(v, y_i)$ , we get a graph with the same degrees as G, but with more edges in common with G'. Similarly,  $(v, y_i) \notin E(G')$ , for if  $(v, y_i) \in E(G')$  we could remove  $(v, y_i)$  and (x, z) from G', but add  $(x, y_i)$  and (z, v) to G' and get a graph with the same degrees as G', but having more edges in common with G. Thus,  $(v, y_i) \in E(G')$ 

 $\in E(G) - E(G').$ 

Now, if  $v \neq y_1, \ldots, y_{k+1}$ , then v is adjacent to at least k+2 edges of E(G)-E(G') ( $(v,z), (v,y_1), \ldots, (v,y_{k+1})$ ) so by hypothesis it is adjacent to at least k+1 edges of E(G')-E(G), a contradiction of the choice of x. If  $v=y_1$  (say) then, again, it is adjacent to at least k+2 edges of E(G)-E(G') (in this case ( $(v,z), (v,x), (v,y_2), \ldots, (v,y_{k+1})$ ) are these edges), and we get a contradiction as before [S. Kundu, Discrete Math. 6 (1973) 367-376; L. Lovász, Periodica Math. Hung. 5 (1974) 149-151].

## § 8. Independent sets of points

1. Let S be a maximal independent set. Then any point  $x \in V(G) - S$  is joined to a point in S. Since a point of S can only be joined to at most d points of V(G) - S, we have

$$|V(G)| - |S| = |V(G) - S| \le d|S|,$$

i.e.

$$|S| \ge \frac{1}{d+1} |V(G)|.$$

2. Define  $S_1, S_2, \ldots$  as in the hint. Then

$$|S_i| \leq \alpha(G[S_{i+1} \cup S_i]) \leq \alpha(G - S_1 - \ldots - S_{i-1}) = |S_i|,$$

whence

$$\tau(G[S_{i+1} \cup S_i]) = |S_{i+1}|.$$

Hence by König's theorem 7.2, we have  $|S_{i+1}|$  independent edges in  $G[S_i \cup S_{i+1}]$ , which is a matching  $F_i$  of  $S_{i+1}$  into  $S_i$ .

Now  $F_1 \cup F_2 \cup \ldots$  consists of  $|S_2|$  disjoint paths, which cover all points except  $|S_1| - |S_2|$  of the points of  $S_1$ . Taking these as one-point paths, we get  $|S_1|$  paths covering V(G).

3. If G consists of independent points, the assertion is obvious. So suppose

this is not the case. We use induction on  $\alpha(G)$ .

Let P be a longest path in G and let x be an endpoint of P. If x has degree 1, remove x and its neighbor y. Observe that  $\alpha(G-x-y)<\alpha(G)$ , since any independent set of G-x-y can be completed with x. Hence, G-x-y can be covered by  $\alpha(G)-1$  disjoint circuits, edges and points, and adding

the edge xy, we get a desired cover of G.

So suppose x has degree at least 2. Each point adjacent to x is on P, because P is maximal. Let y be the farthest point on P adjacent to x and let C be the circuit formed by the (x, y)-piece of P and the edge (x, y). Observe again that  $\alpha(G - V(C)) < \alpha(G)$ , because C contains all neighbors of x. Hence G - V(C) can be covered by  $\alpha(G) - 1$  disjoint circuits, edges and points, and adding C, we get a desired cover of G again [L. Pósa, MTA Mat. Kut. Int. Közl. 8 (1963) 355-361].

**4.** We use induction on |V(G)|. For |V(G)| = 1 the assertion is trivial.

Consider a set S such that there are disjoint paths  $P_1, \ldots, P_{|S|}$  starting from points of S and covering all points; suppose, moreover, that |S| is minimal. What we have to show is  $|S| \leq \alpha(G)$ . Suppose this is not the case, then, obviously, S is not independent, i.e. there are  $x_1, x_2 \in S$  with  $(x_1, x_2) \in E(G)$ . Let, say,  $P_i$  be the path starting from  $x_i$ . Then  $P_1$  is not a single point, since then the paths

 $P_2' = P_2 + (x_1, x_2), P_3, \dots, P_{|S|}$ 

would cover every point, which is impossible, because they are disjoint and start from the points of  $S - \{x_1\}$ . Thus,  $P_1$  has a second point z. Consider now  $G' = G - x_1$ ,  $S' = S - x_1 \cup \{z\}$  and the paths

$$P_1' = P_1 - x_1, P_2, P_3, \dots, P_S.$$

They start from the points of S', cover V(G'), and they are disjoint. Hence, by the induction hypothesis, there is an  $S_0 \subseteq S'$  with  $|S_0| \le \alpha(G') \le \alpha(G) < |S|$  such that certain disjoint paths, starting from the points of  $S_0$ , cover G. Now if  $z \in S_0$ , then adding  $x_1$  to the path starting from z, we get  $|S_0| < |S|$  disjoint paths starting from a proper subset of S and covering G, which is a contradiction. A similar argument works if  $z \notin S_0$  but  $x_2 \in S_0$ . Finally, suppose  $x_2, z \notin S_0$ . Then  $|S_0| \le |S| - 2$ , thus we may add  $\{x_1\}$  as a path to the system to get  $|S_0| + 1 < |S|$  disjoint paths, starting from a subset of S and covering all points of G. This is a contradiction again [T. Gallai; see B].

- 5. (a) If S is a maximal independent set, then each  $x \notin S$  must be adjacent to a  $y \in S$ . By the symmetry assumption, this means that both (x, y) and (y, x) are edges of the digraph.
- (b) Let x be a point with indegree 0 (the existence of such a point follows from the fact that the digraph G is acyclic); let T be the set of neighbors of x. Obviously,  $G_0 = G x T$  is acyclic, so by induction on the number of points it has a (unique) kernel  $S_0$ . We claim that  $S = \{x\} \cup S_0$  is a kernel of G. It is independent since  $G_0$  contains no neighbor of x; moreover, any  $y \in V(G) S$

is either  $\in V(G_0) - S_0$  in which case it can be reached from  $S_0$ , or  $\in T$ , in which case it can be reached from x on an edge of G.

Now if S' is any other kernel of G, then  $x \in S'$  (since x cannot be reached from ony point) and hence,  $T \cap S' = \emptyset$ . Let  $S'_0 = S' \cap V(G_0)$ , then  $S'_0$  is a kernel af  $G_0$  and hence by the uniqueness of  $S_0$ ,  $S'_0 = S_0$ . Thus

$$S' = S'_0 \cup \{x\} = S_0 \cup \{x\} = S_{\bullet}$$

(c) Let  $G_0$  be a strongly connected component of G such that no edge of G enters  $G_0$ . By 5.3,  $G_0$  is bipartite; let  $S_0 \neq \emptyset$  be one of its color classes, and let T be the set of all neighbors of  $S_0$  in  $V(G) - V(G_0)$ . Using induction on |V(G)| we may assume  $G' = G - V(G_0) - T$  has a kernel  $S_1$ . Set

$$S = S_1 \cup S_0$$

Then S is, obviously, independent. Further let  $x \in V(G) - S$ . If  $x \in V(G') - S_1$ , then it can be reached from  $S_1$  through an edge, because  $S_1$  is a kernel of G'. If  $x \in T$ , it is adjacent to a  $y \in S_0$  by the definition of T and the edge joining them goes from y to x by the definition of  $G_0$ . Finally, let  $x \in V(G_0) - S_0$ . Since  $G_0$  is strongly connected and  $S_0 \neq \emptyset$ , we have an edge (x, z) of  $G_0$  leaving x. Since  $S_0$  has been one of the color classes of  $G_0$ ,  $z \in S_0$ . This proves that S is a kernel of G [see B].

**6.** Let x have the highest outdegree among the points of T. We claim any other point y can be reached from x on a path of length at most 2.

Suppose y is a point, which cannot be reached. Then

(a) the edge joining x and y has tail in y,

(b) whenever (x, z) is an edge, (y, z) is an edge (otherwise, (xzy) would be

an (x, y)-path of length at most 2).

Hence, for each edge leaving x, there is an edge leaving y and (a) yields another edge leaving y. This means that the outdegree of y is larger than the outdegree of x, a contradiction.

7. Let  $x \in V(G)$  and let T be the set of all points, which can be reached from x on an edge. Set G' = G - T - x and suppose, by the induction hypothesis, that G' contains an independent set S' such that every point of G' - S' can be reached on a path of length at most 2 from S'. Now, we distinguish two cases:

Case 1.  $S' \cup \{x\}$  is independent. Then set  $S = S' \cup \{x\}$  and observe that a  $y \in V(G') - S'$  can be reached from S' on a path of length at most 2, while a  $y \in T$  can be reached from x even on an edge.

Case 2.  $S' \cup \{x\}$  is not independent, i.e. there is a  $z \in S'$  adjacent to x. Since  $z \notin T$ ,  $(z, x) \in E(G)$ . Now set S = S'. If  $y \in V(G') - S'$ , it can be reached from S' in at most 2 steps by the definition of S'. If  $y \in T$ , it can be reached on zxy. [V. Chvátal—L. Lovász, in: Hypergraph Seminar, Lecture Notes in Math. 411, Springer (1974) p. 175.]

8. (a) To be precise, we define a *winning move* as a point such that, if a player occupies it, he can win (no matter what his opponent does). E.g. a point with outdegree 0 is a winning move, since there is no response at all. Now it follows from the definition that

- (1) winning moves are independent points; for if x is a winning move and  $(x, y) \in E(G)$  then, by definition, if I occupy x, I can win; this happens, in particular, if my opponent occupies y at the next step, so y is not a winning move;
- (2) any point x, which is not a winning move is joined to a winning move. Since by definition, if my opponent occupies x, I can still win, i.e. I have a response y which is a winning move.

Thus, the set of winning moves is the "dual kernel", i.e. the kernel of the graph obtained by inverting the arrows. So A can win by choosing a point of dual kernel at each step.

- (b) Assume A cannot win if he starts with  $x_0$ ; this means that B has a response  $x_1$  such that A cannot win after this move. In this case, let A imagine he is the second player and B has started with  $x_0$ ; so respond with  $x_1$  and follow B's previous strategy. Since  $x_0$  has indegree 0, it will never happen that B makes a move which was illegal in the previous game, since the only such move would be occupying  $x_0$ . Thus A wins.
- (c) Assume first that for each strong component  $G_i$  of G (i = 1, ..., k), the second player has a winning strategy. Then he can follow this rule: whenever A moves in a component  $G_i$ , he responds in the same component  $G_i$  with the move determined by his winning strategy for this particular component. So sooner or later A has no move in this component  $G_i$ , and he either loses or has to move to another component  $G_j$ . Clearly, they have not played in  $G_j$  earlier and therefore B can consider the move of A as an opening move of the game in  $G_i$ , and respond according to his strategy in  $G_i$  etc.

Conversely, assume that there are strong components of G, where the beginner has a winning strategy. Let  $G_i$  be such a component with the property that no such component can be reached from  $G_i$  on a path (if for each such component there is another one accessible on a path we can make a closed walk through more than one component, contradicting the definition of component). Let A start in  $G_i$  according to the winning strategy in  $G_i$ . Then he can force his opponent to make a move outside  $G_i$  first. However, this move is a component, where the second player has a winning strategy and so A can force B to leave this component first (or to lose), etc.

**9.** Suppose first that G has a 1-factor  $\{e_1, \ldots, e_r\}$ . Then the second player B can win by following this rule: whenever A occupies an endpoint of an  $e_i$ , his response is to occupy the other endpoint of the same edge. So A has to choose from a new edge of the one-factor at each step and B's response is always legitimate.

Now suppose G has no 1-factor. Let  $\{e_1, \ldots, e_v\}$  be a maximum matching of G and  $x_1$  a point not covered by  $e_1, \ldots, e_v$ . We claim that, if A opens with x and follows the strategy that whenever B occupies an endpoint of an  $e_i$  he

responds by occupying the other one, then he wins.

What we have to show is that B can never make a move to occupy a point, which is not an endpoint of an edge  $e_i$ . Suppose that at the  $j^{\text{th}}$  move, B first has a move  $y_j$ , which is not an endpoint of  $e_1, \ldots, e_v$ . Let  $x, x_1, \ldots, x_{j-1}$  be the previous moves of  $A, y_1, \ldots, y_{j-1}$  the previous moves of B. Then  $(x_i, y_i)$  is an edge of the maximum matching  $\{e_1, \ldots, e_v\}$ ; say  $(x_i, y_i) = e_i$   $(i = 1, \ldots, j-1)$ .

Now observe that  $\{(x, y_1), \ldots, (x_{j-1}, y_j), e_j, \ldots, e_v\}$  is a larger matching, a contradiction [cf. W. N. Anderson, J. Comb .Theory B 17 (1972) 234–239].

10. For k=1 the statement is evident. Suppose

$$|T_1 \cup \ldots \cup T_{k-1}| + |T_1 \cap \ldots \cap T_{k-1}| \geq 2\alpha(G)$$
.

In fact, we show that

 $|T_1 \cup \ldots \cup T_k| + |T_1 \cap \ldots \cap T_k| \ge |T_1 \cup \ldots \cup T_{k-1}| + |T_1 \cap \ldots \cap T_{k-1}|$  or, equivalently

$$|T_1 \cup \ldots \cup T_k - T_1 \cup \ldots \cup T_{k-1}| \ge |T_1 \cap \ldots \cap T_{k-1} - T_1 \cap \ldots \cap T_k|$$

Now set

$$A = T_1 \cup \ldots \cup T_k - T_1 \cup \ldots \cup T_{k-1},$$
 $B = T_1 \cap \ldots \cap T_{k-1} \cap \cdots - T_1 \cap T_k,$ 
 $C = T_k - A = T_k \cap (T_1 \cup \ldots \cup T_{k-1}).$ 

Then  $|A|+|C|=|T_k|=\alpha(G)$ . On the hand,  $B\cup C$  is independent. In fact, suppose  $x,y\in B\cup C$ . Obviously, B,C are independent, so we may assume  $x\in B,\ y\in C$ . Then  $y\in T_1\cup\ldots\cup T_{k-1}$ , say  $y\in T_1$ . But also  $x\in T_1$  and hence, x and y are non-adjacent. Thus  $B\cup C$  is independent and hence,

$$|B \cup C| \leq \alpha(G) = |A \cup C|,$$

i.e.

$$|B| \leq |A|$$
.

[A. Hajnal; see B.]

11. It suffices to consider the case k = 1, since the general case is trivial by induction. Observe that

$$\Gamma(S) \subseteq \Gamma(X) - T_1.$$

Hence

$$\begin{split} |\varGamma(X)| - |\varGamma(S)| &\geq |\varGamma(X) \cap T_1| = |T_1| - |S| - |T_1 - X - \varGamma(X)| = \\ &= \alpha(G) - |S| + |X| - |T_1 \cup X - \varGamma(X)|. \end{split}$$

Here  $T_1 \cup X - \Gamma(X)$  is, obviously, independent, and has, therefore, at most  $\alpha(G)$  points. Thus  $|\Gamma(X)| - |\Gamma(S)| > |X| - |S|$ ,

which proves the assertion.

12. Let T be an  $(\alpha(G) + 1)$ -element independent set in G - (x, y). Since T is not independent in G, we have  $x, y \in T$ . Now T - y is an  $\alpha(G)$ -element independent set in G containing x, and T - x is another one missing it.

If x, y do not form a component of G, we can find a z joined to one of them, say to x. Then G - (x, z) contains an  $(\alpha(G) + 1)$ -element independent set T. Obviously,  $x, z \in T$ . If  $y \in T$ , then T - x contains y but not x; if  $y \notin T$ , then T - z contains x but not y. Finally, if x, y are adjacent, then  $y \notin T$  and so T - x misses both x and y.

13. Let G' be the graph obtained from G by replacing x by  $x_1$  and  $x_2$  as in the hint. Since  $G' - x_2 \cong G$ , we have  $\alpha(G') \geq \alpha(G)$ . On the other hand, no independent set of G contains both  $x_1$  and  $x_2$  and so, it yields an independent set of G of the same size. Hence,  $\alpha(G') = \alpha(G)$ .

Now let  $e \in E(G')$ . Suppose first that  $e = (x_1, x_2)$ . Let T be a maximum independent set of G containing x. Then  $T - \{x\} \cup \{x_1, x_2\}$  is an  $(\alpha(G) + 1)$ -

element independent set in G - e.

Suppose  $e = (x_1, y)$   $(y \neq x_2)$ . Let T be an  $(\alpha(G) + 1)$ -element independent set in G - (x, y). Then  $T - \{x\} \cup \{x_1\}$  is an  $(\alpha(G) + 1)$ -element independent set in  $G - (x_1, y)$ . Similarly, if  $e = (x_2, y)$  or (z, y)  $(z, y \neq x_1, x_2)$ , we get  $\alpha(G - e) > \alpha(G)$ . [W. Wessel, Coll. Math. Soc. J. Bolyai 4 (1970) 1123–1139].

14. Suppose we have a G such that  $G_0$  is an induced subgraph of G and  $\alpha(G-e)>\alpha(G)$  for  $e\in E(G_0)$ . Remove edges of G as long as you do not increase  $\alpha(G)$ ; then you never remove any edge from  $G_0$  and so, the  $\alpha$ -critical

graph you get contains  $G_0$  as an induced subgraph.

So it suffices to construct such a G. Let  $E(G_0) = \{e_1, \ldots, e_m\}$  and  $\alpha = \alpha(G_0)$ . Let  $S_1, \ldots, S_m$  be  $(\alpha - 1)$ -element sets disjoint from each other and  $V(G_0)$ . Set  $V(G) = V(G_0) \cup S_1 \cup \ldots \cup S_m$ . Let, moreover, each  $x \in S_j$  be joined to all other points except the points of  $S_j$  and the endpoints of  $e_j$ . Thus, we get a graph G.

Now  $\alpha(G) \geq \alpha$ , because  $S_1$ , together with an endpoint of  $e_1$ , form an  $\alpha$ -element independent set. On the other hand, if T is an independent set and it contains a point of  $S_j$ , then it contains at most  $S_j$  and one endpoint of  $e_j$ ; if  $T \subseteq V(G_0)$ ,

then  $|T| \leq \alpha(G_0)$ . Hence  $\alpha(G) = \alpha$ .

Moreover,  $G - e_j$  contains the  $(\alpha + 1)$ -element independent set  $S_j \cup e_j$   $(j = 1, \ldots, m)$ .

- 15. (a) Consider a (6t + 3)-cycle. This is  $\alpha$ -critical. Substitute a  $K_{r-1}$  for every third point of it. Then the resulting graph is r-regular, connected and  $\alpha$ -critical by 8.13.
- (b) It is trivial that the tetrahedron is  $\alpha$ -critical and that the octahedron and cube are not.

Using the fact that the automorphism group of the icosahedron is vertexand edge-transitive, it is easy to verify that its independence number is 3 but this increases to 4 if any edge is removed. Thus the icosahedron is  $\alpha$ -critical.

Note that two opposite vertices of the icosahedron, though non-adjacent, never occur in the same maximum independent set. This shows that the

assertion of problem 8.12 could not be extended in this direction.

Finally, to show that the dodecahedron is not α-critical note that it contains 8 independent vertices (Fig. 63). Remove an edge and consider an independent set in the remaining graph. This contains at most three vertices of the two faces adjacent to the removed edge and at most 2 vertices of any other face. Each element of the independent set belongs to three faces. Therefore the cardinality of this independent set is no larger than

$$\frac{2 \cdot 3 + 10 \cdot 2}{3} = 8.66,$$

showing that the independence number is still 8.




Fig. 63

**16.** Let  $S_1$ ,  $S_2$  be as in the hint. Obviously  $x \in S_1 \cap S_2$ ,  $y_i \in S_i$ ,  $y_i \notin S_{3-i}$ . Since  $S_1 \triangle S_2 \cup \{x\}$  induces a bipartite graph, it contains an independent set T with

$$|T| \ge \frac{1}{2} |S_1 \triangle S_2 \cup \{x\}| = |S_1 - S_2| + \frac{1}{2},$$

i.e.  $|T| \ge |S_1 - S_2| + 1$ . Now note that  $T \cup (S_1 \cap S_2) - \{x\}$  is independent; for let  $u, v \in T \cup ((S_1 \cap S_2) - \{x\})$ ; we show they are non-adjacent. If  $u, v \in T$  or  $u, v \in S_1 \cap S_2$ , we are done. Let, say,  $u \in T$ ,  $v \in S_1 \cap S_2$ . Since  $T \subseteq S_1 \cup S_2$ , we have, e.g.  $u \in S_1$ . So u, v are in  $S_1$  and  $v \ne x, y_1$ , hence they are non-adjacent. But

$$|T \cup ((S_1 \cap S_2) - \{x\})| = |T| + |S_1 \cap S_2| - 1 \ge |S_1 - S_2| + 1 + |S_1 \cap S_2| - 1 = |S_1| > \alpha(G),$$

a contradiction.

17. Let x be a point of the  $\alpha$ -critical graph G, adjacent to  $y_1$  and  $y_2$ . Let  $S_i$  be an  $(\alpha(G)+1)$ -element independent set in  $G-(x,y_i)$ . As before,  $x,y_i\in S_i$ ,  $y_i\notin S_{3-i}$ . Let  $G'=G[S_1\triangle S_2\cup \{x\}]$ . If G' is bipartite, we get a contradiction exactly as above. So G' contains an odd circuit. A minimal odd circuit G is chordless. Since, obviously,  $G'-(x,y_1)$  is bipartite with bipartition  $\{S_1-S_2,\ S_2-S_1\cup \{x\}\},\ C$  must go through  $(x,y_1)$ . Similarly, G goes through  $(x,y_2)$  [L. Beineke, F. Harary, M.D. Plummer, C. Berge; see B].

18. Following the hint, let  $(x, y_1)$ ,  $(x, y_2)$  be two edges, and suppose  $y_1, y_2$  are in different components of G - S. Then G contains a chordless (odd) circuit through  $(x, y_1)$  and  $(x, y_2)$ . This has another point z of S because S separates  $y_1$  and  $y_2$  and z is joined to x because S spans a complete graph. Thus C is not chordless, a contradiction.

19. (a) Let T be an independent set in G. If T contains at most one of  $x_1, x_2$ , then  $T \cap (V(G_1) - \{x\})$  and  $T \cap V(G_2)$  are independent sets in  $G_1$  and  $G_2$ , respectively, whence  $|T| \leq \alpha(G_1) + \alpha(G_2)$ . If  $x_1, x_2 \in T$ , then  $T \cap V(G_1)$ 

 $-\{x\}$   $\cup$   $\{x\}$  and  $T \cap V(G_2) - \{y_1\}$  are independent sets in  $G_1$  and  $G_2$ , respec-

tively, and we reach the same conclusion.

On the other hand, let  $T_1$  be a maximum independent set of  $G_1$  containing x and  $T_2$  an  $(\alpha(G_2) + 1)$ -element independent set of  $G_2 - (y_1, y_2)$  (thus containing  $y_1$  and  $y_2$ ). Then  $T_1 - \{x\} \cup T_2$  is an  $(\alpha(G_1) + \alpha(G_2))$ -element independent set of G. Hence  $\alpha(G) = \alpha(G_1) + \alpha(G_2)$ .

Let  $e \in E(G_1)$ . Let  $T_1$  be an  $(\alpha(G_1) + 1)$ -element independent set in  $G_1 - e$ . If  $x \in T_1$ , let  $T_2$  be an  $(\alpha(G_2) + 1)$ -element independent set of  $G_2 - (y_1, y_2)$  and  $T = T_1 - \{x\} \cup T_2$ . If  $x \notin T_1$ , let  $T_2$  be a maximum independent set of  $G_2$  missing  $y_1$  and  $y_2$  and  $T = T_1 \cup T_2$ . In both cases, T is an  $(\alpha(G) + 1)$ -

element independent set in G - e.

Let  $e \in E(\hat{G}_2)$ ,  $e \neq (y_1, y_2)$ . Let  $T_2$  be an  $(\alpha(G_2) + 1)$ -element independent set in  $G_2 - e$ . If  $y_1, y_2 \notin T_2$ , let  $T_1$  be a maximum independent set of  $G_1$  missing x and  $T = T_1 \cup T_2$ . If  $y_1 \in T_2$  (say), let z be a neighbor of  $x_2$  in  $V(G_1)$ , let  $T_1$  be an  $(\alpha(G_1) + 1)$ -element independent set in  $G_1 - (x, z)$  and  $T = T_1 - \{x\} \cup T_2$ . In both cases T is an  $(\alpha(G) + 1)$ -element independent set in G - e. This proves G is  $\alpha$ -critical. [T. Gallai, M. D. Plummer, W. Wessel; see W. Wessel, Manuscripta Math. 2 (1970) 309–334.]

(b) Let G be a connected  $\alpha$ -critical graph,  $\alpha(G) = \alpha$ . By the preceding exercise, G is 2-connected. Suppose G is not 3-connected, then  $G = G_1 \cup G_2$  with  $V(G_1) \cap V(G_2) = \{x, y\}, \ |V(G_i)| \geq 3$ . By the preceding problem again, x and y are non-adjacent.

For each  $X \subseteq \{x, y\}$  and i = 1, 2, denote the maximum size of an independent set  $T \subseteq V(G_i)$  with  $T \cap \{x, y\} = X$  by  $a_X^i$ . Then clearly

(1) 
$$a_X^1 + a_X^2 \le \alpha + |X|; \quad a_X^i \le a_0^i + |X| = a^i + |X|.$$

Let  $(x, z) \in E(G)$ ,  $z \in V(G_1)$ . Then G - (x, z) contains an  $(\alpha + 1)$ -element independent set T. Clearly,  $x, z \in T$ . If  $y \notin T$ , we get from this

(2) 
$$\begin{cases} a_x^1 + a_x^2 \ge \alpha + 1, \\ a^1 + a_x^2 \ge \alpha + 1. \end{cases}$$

Similarly if  $y \in T$ , we obtain

(3) 
$$\begin{cases} a_{xy}^1 + a_{xy}^2 \ge \alpha + 2, \\ a_y^1 + a_{xy}^2 \ge \alpha + 2. \end{cases}$$

Thus either (2) or (3) holds. Similarly we obtain that one of the following systems of inequalities holds:

(4) 
$$\begin{cases} a_x^1 + a_x^2 \ge \alpha + 1, \\ a_x^1 + a^2 \ge \alpha + 1, \end{cases}$$
 (5) 
$$\begin{cases} a_{xy}^1 + a_{xy}^2 \ge \alpha + 2, \\ a_{xy}^1 + a_y^2 \ge \alpha + 2. \end{cases}$$

Now (2) and (4) cannot hold simultaneously since (1) implies

$$(a^1 + a_x^2) + (a_x^1 + a^2) = (a^1 + a^2) + (a_x^1 + a_x^2) \le 2\alpha + 1.$$

Similarly, (3) and (5) cannot hold simultaneously. Thus, choosing the indices appropriately we get that (2) and (5) hold. Interchanging the role of x and y we get that either

(6) 
$$\begin{cases} a_y^1 + a_y^2 \ge \alpha + 1, \\ a^1 + a_y^2 \ge \alpha + 1, \end{cases} \text{ and } (7) \begin{cases} a_{xy}^1 + a_{xy} \ge \alpha + 2, \\ a_{xy}^1 + a_{xy}^2 \ge \alpha + 2, \end{cases}$$

or

(8) 
$$\begin{cases} a_y^1 + a_y^2 \ge \alpha + 1, \\ a_y^1 + a^2 \ge \alpha + 1, \end{cases} \text{ and } (9) \begin{cases} a_{xy}^1 + a_{xy}^2 \ge \alpha_y^2 + 2 \\ a_x^1 + a_{xy}^2 \ge \alpha + 2 \end{cases}$$

hold. But (2), (5) and (8), (9) cannot hold simultaneously; in fact, they would imply

$$(a^1 + a_x^2) + (a_y^1 + a^2) + (a_{xy}^1 + a_y^2) + (a_x^1 + a_{xy}^2) \ge 4\alpha + 6;$$

but the left-hand side is equal to

$$(a^1 + a^2) + (a_x^1 + a_x^2) + (a_y^1 + a_y^2) + (a_{xy}^1 + a_{xy}^2) \le 4\alpha + 4$$

by (1), a contradiction. Thus (2), (5), (6), (7) hold but (3), (4), (8), (9) do not. Since the first inequalities in (3), (4), (8), (9) also occur in (2), (5), (6), (7), their second inequalities must fail to hold, i.e.

(10) 
$$\begin{cases} a_{x}^{1} + a^{2} \leq \alpha & a_{x}^{1} + a_{xy}^{2} \leq \alpha + 1, \\ a_{y}^{1} + a^{2} \leq \alpha & a_{y}^{1} + a_{xy}^{2} \leq \alpha + 1. \end{cases}$$

Since by (1) and (6)

$$\alpha \ge a^1 + a^2 \ge a^1 + a_v^2 - 1 \ge \alpha$$

we have

$$a^1 + a^2 = \alpha$$
,  $a_v^2 = a^2 + 1$ .

But, then by (6) and (10),

$$\alpha + 1 \le a_{\nu}^1 + a_{\nu}^2 = a_{\nu}^1 + a^2 + 1 \le \alpha + 1$$

whence

$$a_{\nu}^{1} = \alpha - a^{2} = a^{1}$$
.

Similarly we deduce that

$$a_x^1 = a^1$$
,  $a_x = a^2 + 1$ ,  $a_{xy}^1 = a^1 + 1$ ,  $a_{xy} = a^2 + 1$ .

Now denote by  $\tilde{G}_1$ ,  $\tilde{G}_2$  the graphs obtained from  $G_1$  by connecting x to y and from  $G_2$  by identifying x and y, respectively. We claim these two graphs are  $\alpha$ -critical. Since G arises from them by the operation in (a), this will prove the assertion.

We have

$$\begin{split} &\alpha(G_1) \,=\, \max{(a^1,\,a_x^1,\quad a_y^1)} = a^1, \\ &\alpha(G_2) = \max{(a_2,\,a_{xy}^2 - 1)} = a^2. \end{split}$$

Let  $e \in E(G_1)$ . If e = (x, y), then  $\alpha(G_1 - e) \ge a_{xy}^1 > a^1$ ; if  $e \ne (x, y)$ , then  $e \in E(G)$ . Let T be an  $(\alpha + 1)$ -element independent set in G - e,  $T \cap \{x, y\} = X$ . Then

$$T \cap V(G_2) \leq a_X^2$$

and thus by (1)

$$T \cap V(G_1) \geq \alpha + 1 - a_X^2 + |X| = a_X^1 + 1.$$

Now, if  $X \neq \{x, y\}$ , then  $T \cap V(G_1)$  is an independent set in  $G_1 - e$  of size  $a_X^1 + 1 > a^1$ ; if  $X = \{x, y\}$ , then  $T - \{x\}$  is such a set of size  $a_{xy}^1 > a^1$ . This proves  $G_1$  is  $\alpha$ -critical. For  $G_2$  this follows similarly. [T. Gallai, W. Wessel; ibid.]

**20.** Let  $T_1, \ldots, T_k$  be all maximum independent sets of G. Then, by 8.12,

$$T_1 \cap \ldots \cap T_k = \emptyset, T_1 \cup \ldots \cup T_k = V(G).$$

Thus by 8.10,

$$|V(G)| = |T_1 \cup \ldots \cup T_k| = |T_1 \cup \ldots \cup T_k| + |T_1 \cap \ldots \cap T_k| \ge 2\alpha(G)$$

[P. Erdős, T. Gallai; B].

**21.** Let  $T_1, \ldots, T_k$  be the maximum independent sets of G. Then, as we have seen,

$$T_1 \cap \ldots \cap T_k = \emptyset$$
.

Thus setting  $S = T_1 \cap \ldots \cap T_k \cap X = \emptyset$ , we get by 8.11

$$|\Gamma(X)| - |X| \ge |\Gamma(S)| - |S| = 0$$

[A. Hajnal; see B.]

**22.** Let  $S = X \cap T_1 \cap \ldots \cap T_k$ , where  $T_1, \ldots, T_k$  are as in the hint. Then  $x \in S$ . Let  $S' = S - \{x\}$ . We claim that  $\Gamma(S') \cap \Gamma(x) = \emptyset$ . Suppose there is a v adjacent to x and to a point u in S'. Let T be an  $(\alpha(G) + 1)$ -element independent set in G - (x, v), then  $x, v \in T$  and hence  $u \notin T$ . Moreover,  $T - \{v\}$  is a maximum independent set of G containing x and so,  $T - \{v\} = T_i$  for some i. But then  $u \notin T_i$ , hence  $u \notin S$ , a contradiction.

Thus  $|\Gamma(S)| = |\Gamma(S')| + |\Gamma(x)|$ ; by 8.11 we have

$$|\Gamma(S)| - |S| \le |\Gamma(X)| - |X|$$

whence

$$\begin{split} d_{\mathcal{G}}(x) &= |\varGamma(x)| = |\varGamma(S)| - |\varGamma(S')| = |\varGamma(S)| - |S| - (|\varGamma(S')| - |S'|) + 1 \leq \\ &\leq |\varGamma(X)| - |X| + 1 \end{split}$$

[L. Lovász, L. Surányi].

23. Let  $x \in V(G)$  and let T be an independent set of size  $\alpha(G)$  containing x. By the preceding problem,

$$d_G(x) \le |\Gamma(T)| - |T| + 1 = |V(G) - T| - |T| + 1 = |V(G)| - 2\alpha(G) + 1$$

[A. Hajnal; see B].

**24.** Let S, T be maximum independent sets with  $S \cap T \neq \emptyset$ . Obviously

$$\Gamma(S \cap T) \subseteq V(G) - S - T.$$

On the other hand if  $x \in S \cap T$ , we have

$$|V(G)| - 2\alpha(G) + 1 = d_G(x) \le |\Gamma(S \cap T)| - |S \cap T| + 1$$

by 8.22, so

$$|\Gamma(S \cap T)| \ge |S \cap T| + |V(G)| - 2\alpha(G) =$$
  
=  $|V(G)| + |S| + |T| - |S \cap T| = |V(G)| - |S \cup T|,$ 

whence we have  $\Gamma(S \cap T) = V(G) - S - T$ .

Now let a be an arbitrary point of V(G), and  $b_1, \ldots, b_k (k = |V(G)| - 2\alpha(G) + 1)$  be the neighbors of a. Let  $S_i$  be an  $(\alpha(G) + 1)$ -element independent set of  $G - (a, b_i)$  and set  $T_i = S_i - a$ . Then, obviously,  $T_i$  is a maximum independent set of G, and  $b_i \in T_i$ ,  $b_i \notin T_j$  for  $j \neq i$ . Thus,  $T_i \cap T_j$  contains no neighbors of a. This means by the statement proved above that  $T_i \cap T_j = \emptyset$ . Hence

$$|V(G)| \ge 1 + \sum_{i=1}^{k} |T_i| = 1 + (|V(G)| - 2\alpha(G) + 1)\alpha(G)$$

or, equivalently,

$$(\alpha(G)-1)(|V(G)|-2\alpha(G)-1)\leq 0.$$

Now if  $\alpha(G) = 1$ , we have a complete graph. If  $\alpha(G) > 1$  and  $|V(G)| \le 2\alpha(G) + 1$ , then the degree is at most 2 and so we have  $K_2$  or a cycle. The latter is  $\alpha$ -cr only if it is odd.

25. If  $|V(G)| = 2\alpha(G)$ , we have  $d_G(x) = 1$  by 8.22 and so,  $G \simeq K_2$ . If  $|V(G)| = 2\alpha(G) + 1$ , we have  $d_G(x) \leq 2$  for every point x, thus G is a circuit or a path.

It is easily seen that only odd circuits are α-critical among these.

Now consider those graphs which are  $\alpha$ -critical and satisfy  $|V(G)| = 2\alpha(G) + 2$ . They have  $d_G(x) \leq 3$  for every point x. Observe that if we subdivide an edge of them by two points we get a graph from the same class; because this operation is a special case of the operation in 8.19 when  $G_1$  is a triangle and  $G_2$  is the given graph. Conversely, if we have such a graph and a point x of degree 2 in it (no point can have degree 1, since then its neighbor would be a one-element cutset, contradicting e.g. 8.18), then the two neighbors of x form a 2-element cutset and by part b of 8.19 we get that, contracting the two edges adjacent to x, an  $\alpha$ -critical graph arises. This graph G' has degrees at most 3 as well since simple computation shows that  $|V(G')| - 2\alpha(G') = 2$ . Therefore, one of the neighbors of x in G must have been of degree 2 and so, G arises from G' by subdividing an edge by two points. Thus, it suffices to consider those graphs with degree 3.

By the previous problem, a 3-regular graph with  $|V(G)| = 2\alpha(G) + 2$  is  $K_4$ . Hence all graphs G which are connected,  $\alpha$ -critical and satisfy  $|V(G)| = 2\alpha(G) + 2$  arise from  $K_4$  by subdividing some edges by an even number of points [B. Andrásfai, in: Theory of Gr. Int. Symp. Rome, Dunod, Paris-Gordon and

Breach, New York, 1967, 9-19].

Remark: It is known [L. Lovász, in: Combinatorics, Coll. Math. Soc. J. Bolyai 15, Bolyai–North-Holland (1977)] that for each  $\delta$  there exists a finite number of  $\alpha$ -critical graphs G with  $2\alpha(G) + \delta$  points such that all the other such graphs arise from them by subdividing some edges by an even number of points.

**26.** Remove edges till finally an  $\alpha$ -critical graph G' with  $\alpha(G') = \alpha(G)$  arises. G' does not have isolated points since for such a point x we would have  $\alpha(G-x) \le \alpha(G'-x) = \alpha(G) - 1$ . Similarly, no component of G' consists of two points; since if x, y constitute a component of G', then

$$\alpha(G - \{x, y\}) \ge \alpha(G' - \{x, y\}) = \alpha(G) - 1.$$

Hence, for each component  $G_0$  of G', we have  $|V(G_0)| \ge 2\alpha(G_0) + 1$  by 8.25. Since  $|V(G)| = 2\alpha(G) + 1$ , we have only one component and so by 8.25, this is an odd circuit.

So G' is an odd circuit. We show G' = G. Suppose there is an edge of G not in G'. This edge, together with an arc of G, forms an odd circuit G' and the rest of G', an arc of odd length, has a 1-factor G'. Now  $G'' = G \cup G'$  has  $\alpha(G'') = \alpha(G')$  and we get a contradiction with the above. [V.G. Vizing, L.S. Melnikov, Diskret. Analiz 19 (1971) 11–14.]

27. As before, let G' be an  $\alpha$ -critical spanning subgraph of G with  $\alpha(G') = \alpha(G)$ . We see as before that G has no component, which is an isolated point or a single edge. Moreover, each point of G' has degree greater than 2. For suppose x is adjacent in G' to g and g only. Then  $\alpha(G' - \{x, y, z\}) < \alpha(G)$ , because we could add g to any independent set of g and g to get an independent set of g. Thus

$$\alpha(G-\{x,\,y,\,z\}) \leq \alpha(G'-\{x,\,y,\,z\}) < \alpha(G),$$

a contradiction.

By 8.23, we have  $|V(G_0)| \ge 2\alpha(G_0) + 2$  for each component  $G_0$  of G. This implies that

 $|V(G)| \ge 2\alpha(G) + 4,$ 

if G is not connected; so we may assume G is connected. By 8.25,  $G \simeq K_4$  or

$$|V(G)| \ge 2\alpha(G) + 3$$

[E. Szemerédi, Comb. Theory Appl. Coll. Math. Soc. J. Bolyai 4. Bolyai–North-Holland (1970) 1051–1053].

## § 9. Chromatic number

- 1. Let  $x \in V(G)$ . Since every point of G x has degree at most k, by using induction we can (k + 1)-color it. x is adjacent to at most k points, therefore we can find a color, which does not occur among the neighbors of x. If we give x this color, we get a good coloration of G.
- **2.** The first statement is trivial: Consider any (good) coloration of G with  $\chi(G)$  colors, let  $V_1$  be the union of  $k < \chi(G)$  color classes and  $V_2$  the rest. The  $\chi(G)$ -coloration we started with induces a k-coloration of  $G[V_1]$  and a  $(\chi(G) k)$ -coloration of  $G[V_2]$ : thus

$$\chi(G[\,V_{\,1}]) \leq k, \qquad \chi(G[\,V_{\,2}]) \leq \chi(G) \, - \, k.$$

On the other hand,  $G[V_1]$  cannot be colored by less than k colors, since such a coloration, together with the coloration of  $G[V_2]$  we already have, would give a coloration of G by less than  $\chi(G)$  colors. Thus

$$\chi(G[V_1]) = k$$

and similarly

$$\chi(G[V_2]) = \chi(G) - k.$$

Now suppose that G is not a clique, then it contains a maximal clique  $H_1$ . Let  $V_1 = V(H_1)$ ,  $V_2 = V(G) - V_1$  and  $H_2 = G[V_2]$ . We claim that

$$\chi(H_1) + \chi(H_2) > \chi(G).$$

Since  $H_1$  is a clique,  $\chi(H_1) = |V_1| = k$  (say). We have to show that  $\chi(H_2) > 0$ 

 $> \chi(G) - k$ .

Suppose indirectly that  $H_2$  admits a  $(\chi(G) - k)$ -coloration, using colors  $1, \ldots, \chi(G) - k$ . Use colors  $\chi(G) - k + 1, \ldots, \chi(G)$  to color  $H_1$ . We are going to show that we can get rid of one of these k colors, color  $\chi(G)$ , say. To this end, look at the set T of those points of color 1, which are neighbors of the point x with color  $\chi(G)$ . Since  $H_1$  is a maximal complete subgraph, each point  $t \in T$  is non-adjacent to a certain point  $t' \in V_1$ . Now re-color the graph giving to each  $t \in T$  the color of t'. This is another legitimate coloration, because the color of t' does not occur anywhere else, and the new points of this color are independent. Now observe that in this new coloration x is the only point of color  $\chi(G)$  and has no neighbor of color 1. So if we re-color x by 1, we get a coloration by  $\chi(G) - 1$  colors. This is a contradiction.

3. Let  $\alpha_i$  be a  $\chi(G_i)$ -coloration of  $G_i$  (i = 1, 2). We may assume  $V(G_1) = V(G_2)$ , since this can always be achieved by adding isolated points to the two graphs. Then

$$\alpha(x) = (\alpha_1(x), \alpha_2(x))$$

defines a coloration of  $G_1 \cup G_2$  with  $\chi(G_1) \cdot \chi(G_2)$  colors, which is easily seen to be legitimate.

- 4. Let  $\alpha$ ,  $\beta$  be as in the hint. We show that  $\beta$  uses each color at least twice. Suppose, e.g. 1 occurs only once. Let x be the point of  $\beta$ -color 1, and let S be the color-class of x with respect to  $\alpha$ . Let  $\beta'$  be obtained from  $\beta$  by re-coloring each point of S by 1. Then the "old" common color classes of  $\alpha$  and  $\beta$  remain untouched, but we get a new color class S in common, a contradiction [T. Gallai,  $Mat.\ Kut.\ Int.\ K\ddot{o}zl.\ 8\ (1964)\ 373-385$ ].
- **5.** (a) By induction on k, we may suppose that

$$\chi(G - V_k) \le n - |V_k| - (k - 1) + 1.$$

Let  $\alpha$  be a coloration of  $G - V_k$  with colors  $1, \ldots, n - |V_k| - k + 2$ , and extend  $\alpha$  to a coloration of G by coloring the points of  $V_k$  with colors  $n - |V_k| - k + 3, \ldots, n - k + 2$ . We want to re-color the graph, so as to

get rid of one of these colors.

By the assumption, each  $V_i$  ( $1 \le i \le k-1$ ) contains a point  $y_i$ , which is non-adjacent to a certain point  $x_i$  of  $V_k$ . There must be a color which occurs in the set  $\{y_1, \ldots, y_{k-1}\}$  only, since the number of points outside this set is only n-k+1. Let, e.g. 1 be this color. Now re-color the points  $y_i$  such that  $\alpha(y_i) = 1$  with the color of  $x_i$ . It is clear that the resulting coloration is good and uses n-k+1 colors only [R.P. Gupta, in: *Theory of Graphs*, Proc. Int. Coll. Rome, Gordon and Breach, 1969].

(b) Consider a  $\chi(\bar{G})$ -coloration of  $\bar{G}$  with color classes  $V_1, \ldots, V_k$  ( $k = \chi(\bar{G})$ ). Then any two  $V_i$ 's are connected by an edge in  $\bar{G}$ , since otherwise they could be merged. Thus G satisfies the conditions of (a) and so,

$$\chi(G) \le n - k + 1 = n - \chi(\bar{G}) + 1.$$

Hence the first inequality. The second is trivial by 9.3:

$$\chi(G) \cdot \chi(\bar{G}) \geq \chi(G \cup \bar{G}) = \chi(K_n) = n.$$

[E.A. Nordhaus, J.W. Gaddum, Amer. Math. Monthly 63 (1956) 175-177].

- **6.** Let  $V(K_m) = \{x_1, \ldots, x_m\}$ . Suppose T is an independent set of  $G \oplus K_m$ . Then T contains at most one point of  $\{(y, x_1), \ldots, (y, x_m)\}\ (y \in V(G))$  since this is a complete graph. Therefore,  $|T| \leq |V(G)|$ . If equality holds, then exactly one of  $\{(y, x_1), \ldots, (y, x_m)\}$  belongs to T. Define then an m-coloration of G by giving y the color i, if  $(y, x_i) \in T$ . This is a legitimate coloration since, if y, z both have color i, then  $(y, x_i)$ ,  $(z, x_i) \in T$  and so are non-adjacent, but then so are y, z. Conversely, suppose G is m-colorable, and consider a coloration  $\alpha$  of it with colors  $1, \ldots, m$ . Then the set  $\{(y, x_{\alpha(y)}); y \in V(G)\}$  is an independent subset of  $G \oplus K_m$  of cardinality |V(G)| [V. G. Vizing; B].
- 7. (a) Let  $\alpha$  be a  $\chi(G_1)$ -coloration of  $G_1$ . Define a coloration  $\beta$  of  $G_1 \times G_2$  by  $\beta(x,y) = \alpha(x) \ (x \in V(G_1), \ y \in V(G_2))$ . This is a legitimate coloration, for if  $(x_1, y_1)$  is joined to  $(x_2, y_2)$ , then  $x_1$  is joined to  $x_2$  (and  $y_1$  is joined to  $y_2$ ), so  $\alpha(x_1) \neq \alpha(x_2)$ . Thus  $\chi(G_1 \times G_2) \leq \chi(G_1)$ . Similarly, it is  $\leq \chi(G_2)$ .
- (b) Since the points of  $\{(x, x) : x \in V(G)\}$  induce a subgraph of  $G \times G$  isomorphic to G, we have  $\chi(G \times G) \ge \chi(G)$  and by (a), we have equality.
- (c) Suppose  $k = \chi(G \times K_n) < n$ , we show that  $k \ge \chi(G)$ . Let  $\alpha$  be a kcoloration of  $G \times K_n$ . Since k < n, we have a repetition in the set  $\{\alpha(v, x)\}$ ;  $x \in V(K_n)$  for every  $v \in V(G)$ ; let  $\beta(v)$  be this more than once occurring color. We claim  $\beta$  is a good k-coloration of G. Suppose  $(v_1, v_2) \in E(G)$ . There are  $x_i, x_i'$  with  $\alpha(v_i, x_i) = \alpha(v_i, x_i') = \beta(v_i)$  (i = 1, 2). We assume  $x_1 \neq x_2$ . Then  $(v_1, x_1)$  and  $(v_2, x_2)$  are adjacent in  $G \times K_n$ , so

$$\beta(v_1) = \alpha(v_1, x_1) \neq \alpha(v_2, x_2) = \beta(v_2).$$

(d) Let  $\alpha$  be an n-coloration of  $G \times K_n$ . If, for every v, there is a color which occurs twice among  $\{\alpha(v,x):x\in V(K_n)\}$ , then we get a good n-coloration of G as in the previous solution and this contradicts  $\chi(G) > n$ . So we have a  $v \in V(G)$  such that  $\alpha(v, x), x \in V(K_n)$  are different.

We claim that the same holds for each neighbor u of v, and, in fact,  $\alpha(u, x) =$  $= \alpha(v, x)$  for each  $x \in V(K_n)$ . This follows since we have  $\alpha(u, x) \neq \alpha(v, y)$  for  $x \neq y$  and all colors but  $\alpha(v, x)$  occur among  $\alpha(v, y), y \neq x$ . So  $\alpha(u, x) = \alpha(v, x)$ and as G is connected, this will hold for any u and v. Hence, the n-coloration  $\alpha$  is the one induced by the (unique) n-coloration of  $K_n$ .

(e) If n=2, a bipartite graph G has  $2^{c(G)-1}$  2-colorations, since each component has a unique 2-coloration (up to exchanging colors) and these can be combined in  $2^{c(G)-1}$  ways.

So suppose  $n \geq 3$ . We claim that  $K_n^m$  has exactly m n-colorations (those induced by the n-colorations of the factors). We use induction on m. Let  $\alpha$  be

an *n*-coloration of  $K_n^m = K_n^{m-1} \times K_n$ . Suppose first there is a point  $v \in V(K_n^{m-1})$  such that all points  $(v, x), x \in V(K_n)$ , have the same color. Then it follows as in the previous solution that  $\alpha(v, x)$ is independent of v (as because  $K_n^{m-1}$  is connected for  $n \geq 3$  by 6.4), i.e. it is induced by an n-coloration of the last factor. So suppose that there are two points  $x_v, x_v' \in V(K_n)$  for each  $v \in V(K_n^{m-1})$  such that  $\alpha(v, x_v) = \alpha(v, x_v')$ . If  $\alpha(v, x)$  is independent of x, then it is induced by a coloration of  $K_n^{m-1}$  and we are finished by induction. So suppose there is a  $v_0 \in V(K_n^{m-1})$  and  $x_0, x_0' \in V(K_n)$  such that  $\alpha(v_0, x_0) \neq \alpha(v_0, x_0')$ . Now define

$$eta(v) = egin{cases} lpha(v_0, \, x_0) & ext{for } v = v_0, \ lpha(v, \, x_v) & ext{otherwise,} \ eta'(v) = egin{cases} lpha(v_0, \, x_0') & ext{for } v = v_0, \ eta(v) & ext{otherwise.} \end{cases}$$

Then it is easy to check that  $\beta$ ,  $\beta'$  are legitimate n-colorations of  $K_n^{m-1}$  and they differ only at  $v_0$ . But we know all n-colorations of  $K_n^{m-1}$  from the induction hypothesis and there are no two such  $\beta$ ,  $\beta'$  among them. This contradiction proves the assertion [D.L. Greenwell, L. Lovász, Acta Math. Acad. Sci. Hung. 25 (1974) 335–340].

- 8. (a) The complete graph on S.
  - (b) Take k |S| + 1 new points and join them to each other and to S.
  - (c) Take k-1 new points and join them to each other and to S.

(d) Take an odd circuit C of length  $\geq |S|$ ; join each point of C to a point of S, so that each point of S occurs. Take, furthermore, k-3 other points  $u_1, \ldots, u_{k-3}$  and join them to each other, to the points of C and to a given point  $s_0 \in S$ .

No k-coloration of the resulting graph G induces  $\{S\}$ . For if all points of S are colored the same, then  $u_1, \ldots, u_{k-3}$  are colored by k-3 other colors (being adjacent to  $s_0$ ), and we can use only two colors to color C, which is impossible. Let us consider a k-coloration of S, where at least two colors occur. Let us color  $u_1, \ldots, u_{k-3}$  by colors different from the color of  $s_0$ . Now at each point of C there are at most k-2 colors excluded (the colors of  $u_1, \ldots, u_{k-3}$  and the color of its neighbor in S) and so, there are still two legal colors. Moreover, these two colors are not the same at every point. We show that this implies that we can extend the coloration to C.

Let  $C = (x_1, \ldots, x_m)$  (we do not need in this argument the fact that C is odd). Also we may assume that the two colors permitted at  $x_1$  and  $x_m$  are not the same, i.e. there is a color  $\alpha_1$  permitted at  $x_1$  but not at  $x_m$ . Let  $\alpha_2$  be a color permitted at  $x_2$  different from  $\alpha_1$  (since two colors are permitted at  $x_2$ , there is such a color) and similarly, if  $\alpha_1, \ldots, \alpha_i$  are defined, then let  $\alpha_{i+1}$  be a color permitted at  $x_{i+1}$  different from  $\alpha_i$ . Then

$$\alpha(x_i) = \alpha_i$$
  $(i = 1, \ldots, m)$ 

defines a good coloration of C. In fact,  $x_i$ ,  $x_{i+1}$  have different colors by definition and so do  $x_1$  and  $x_m$  since the color  $\alpha_m$  is one of the colors permitted at  $x_m$  while the color of  $x_1$  is not.

(e) First solution. First, let a partition P be given. We construct a graph  $G_P$  such that  $S \subseteq V(G_P)$  and the k-colorations of  $G_P$  induce all partitions of S except P. Let  $P = \{X_1, \ldots, X_r\}$ . Let us associate a new point with each (r-1)-subset  $A = \{a_1, \ldots, a_{r-1}\}$  such that  $a_i \in A_i$   $(i=1, \ldots, r-1)$ , and connect it to  $a_1, \ldots, a_{r-1}$ . Let T be the set of these new points. Now note the following two properties of the resulting graph:

- (1) If we k-color it so that the coloration induces the partition P on S, then  $T \cup X$ , is necessarily monochromatic.
- (2) If we k-color S so that the corresponding partition is different from P, then the coloration can be extended over T so that  $T \cup X_r$  will not be monochromatic.

Now attach a graph to  $T \cup X_r$  such that a k-coloration of  $T \cup X_r$  can be extended over this graph iff  $T \cup X_r$  is not monochromatic (see part (d)). Then

the resulting graph  $G_P$  has the desired property.

If any set  $\{P_1, \ldots, P_N\}$  of partitions of S is given, construct the graph  $G_P$  for each partition  $P \neq P_1, \ldots, P_N$ . We may assume the graphs  $G_P$  have only the points of S in common. Then  $G = \bigcup G_P$  is a graph with the desired properties.

Note the additional property that S is an independent set of points.

Second solution. We only describe the construction without details. Consider  $K_k^N$ . By 9.7, this graph has exactly N k-colorations  $\alpha_1, \ldots, \alpha_N$ . Let  $\beta_i$  be a k-coloration of S inducing the partition  $P_i$  ( $i=1,\ldots,N$ ), and  $\gamma_i=\alpha_i\cup\beta_i$  (this is a k-coloration of  $S\cup V(K_k^N)$ ). Add all edges between S and  $K_k^N$  to the graph which leave the colorations  $\gamma_i$  legitimate. It is easy to verify that the k-colorations of the resulting graph are just  $\gamma_1,\ldots,\gamma_N$ , and thus it satisfies our requirements.

. Suppose first that G has no cycles. Let  $\alpha(x)$  denote the maximum length of a path starting from x. Then  $\alpha$  is a coloration with colors  $0, 1, \ldots, m-1$ . For suppose there were adjacent points x, y with the same color i; let say  $(x, y) \in E(G)$ . Let P be a path of length i starting from y; then P does not go through x, because G has no cycles. Therefore, (x, y) + P is a path of length i + 1 starting from x, hence  $\alpha(x) \geq i + 1$ , a contradiction. Observe that in this coloration, no path joins points of the same color.

Now let  $\{e_1, \ldots, e_k\}$  be a minimum set of edges whose removal destroys all cycles. Then  $G' = G - \{e_1, \ldots, e_k\}$  is acyclic and has the m-coloration constructed above. Since  $G' + e_i$  is not longer acyclic, it has a cycle through  $e_i$ . Hence, G' has a path connecting the endpoints of  $e_i$ . By our remark above, this implies that the endpoints of  $e_i$  have different colors. Thus, the m-coloration of G' is a good m-coloration of G as well [T. Gallai, B. Roy; see B].

10. Assume we have a coloration with the mentioned property. Then every circuit must go "up" and "down" equally often, i.e. has the same number of

edges oriented in each of the two directions.

Conversely, suppose the circuits have this property. Let W be any closed walk, we claim that W has the same property. We use induction on |E(W)|. If W is a circuit this is trivial; otherwise let x be a point, which occurs twice on W. Then x divides W into two shorter closed walks, and we are finished

by induction.

Define the "expense" of a walk as the number of edges used in the right direction less the number of edges used backwards. Let  $x_0$  be a point. For any point a, the expenses of all  $(x_0, a)$ -walks are the same, because otherwise, walking from  $x_0$  to a on one walk and coming back on the other the expense of this walk would be non-zero. Color a by the expense of any  $(x_0, a)$ -walk, then the coloration has the desired property.

- 11. If G is k-colorable with colors  $1, \ldots, k$ , direct (a, b) from a to b, if the color of b has a higher number than the color of a. It is easy to check that this orientation has the property stated. Conversely, assume  $\overline{G}$  is an appropriate orientation of G. Let us verify first the statements of the hint.
- (a) Observe that by the assumption on the orientation, if I walk around any circuit of G, I cannot gain. Thus, if I remove a "loop" from any (a, b)walk. I obtain an (a, b)-walk with no more expense. The number of (a, b)-paths being finite, we have one with minimal expense.
- (b) Suppose a, b are adjacent and say  $(a, b) \in E(G)$ . Let  $W_a$ ,  $W_b$  be optimal  $(x_0, a)$ - and  $(x_0, b)$ -walks with expenses  $\varepsilon(a)$  and  $\varepsilon(b)$ , respectively. Then  $W_a + (a, b)$  is an  $(x_0, b)$ -walk with expense  $\varepsilon(a) - 1$ , thus  $\varepsilon(b) \leq \varepsilon(a) - 1$ . Conversely,  $W_b + (a, b)$  is an  $(x_0, a)$ -walk with expense  $\varepsilon(b) + k - 1$ , hence  $\varepsilon(a) \leq \varepsilon(b) + k - 1$ . Thus

$$1 \le \varepsilon(a) - \varepsilon(b) \le k - 1$$

as stated.

Now fix a point  $x_0$  and define the color of a as the minimum expense of  $(x_0, a)$ walks mod  $\bar{k}$ . Thus we use colors  $0, 1, \ldots, k-1$ . This coloration is legal, because if a, b are adjacent points then by (b) above, the minimum expenses of  $(x_0, a)$  and  $(x_0, b)$ -walks belong to different residue classes mod k [G.J. Minty, Am. Math. Monthly 69 (1962) 623-624].

12. (a) Let  $x_0$  be a point with  $|C(x_0)| > d_G(x_0)$ ; suppose  $x_0, \ldots, x_i$   $(0 \le i \le i \le j \le i)$  $\leq n-1$ ) are defined, then let  $x_{i+1}$  be a point adjacent to one of them (which exists, because G is connected). So we get the sequence  $(x_0, \ldots, x_{n-1})$ .

Now color  $x_{n-1}$  with one of the colors of  $C(x_{n-1})$ ; suppose  $x_{n-1}, \ldots, x_{n-1}$  are colored, then choose a color from  $C(x_{n-i-1})$ , which is different from the colors of previously colored neighbors of  $x_{n-i-1}$ . This is possible, if  $i \neq n-1$  since then  $x_{n-i-1}$  has  $d_G(x_{n-i-1})$  neighbors previously colored and  $|C(x_{n-i-1})| \geq 1$  $\geq d_G(x_{n-i-1})$ . But it is also possible for i=n-1, because  $x_0$  has  $d_G(x_0)$ neighbors and  $|C(x_0)| > d_G(x_0)$ . Thus we get a desired coloration.

(b) We can find two neighboring points a, b with  $C(a) \neq C(b)$ . We may assume  $C(a) \supseteq C(b)$ . Since G - b is connected, we can arrange its points in a sequence  $(x_0 = a, \ldots, x_{n-2})$  so that  $x_j$  is adjacent to some  $x_i$  with i < j for any  $1 \le j \le n-2$ . Let  $b = x_{n-1}$ , it also has this property.

Let us give  $x_{n-1}$  a color not in C(a) (which exists, because  $C(a) \supseteq C(b)$ ). Going backwards as in the previous solution, we get  $x_{n-2}, \ldots, x_1$  colored. However, we can find a suitable color for  $x_0 = a$  as well; since it has  $d_G(a) = |C(a)|$ neighbors but one of these, b, has a color  $\alpha$ , which does not come into consideration since  $\alpha \notin C(a)$ . Therefore, we can find a color in C(a), which is different from the color of the neighbors of a.

13. (a)  $x_0, x_{n-2}, x_{n-1}$  with the property given in the hint exist, as otherwise the relation of being adjacent would be an equivalence relation and G would be complete.

Since G is 3-connected,  $G-x_{n-2}-x_{n-1}$  is connected and its points can be arranged in a sequence  $x_0, \ldots, x_{n-3}$  so that  $x_i$  is joined to an  $x_i$  with i < j

 $(j = 1, \ldots, n - 3).$ 

Now color  $x_{n-1}$ ,  $x_{n-2}$  with color 1. Working backwards again, we can give each  $x_i$  ( $i \ge 1$ ) one of the colors  $1, \ldots, k$ , since  $x_i$  has less than k neighbors colored previously. Reaching  $x_0$ , we can find a color for it too, since although  $x_0$  has k previously colored neighbors, two of these,  $x_{n-1}$  and  $x_{n-2}$ , have the same color.

(b) We may assume G is 2-connected; for otherwise, if we can k-color the blocks of it, we can put together these colorations in the obvious way (permuting the colors, if necessary, to get the same color for the cutpoints).

We may also assume that G is k-regular otherwise 9.12a applies.

Now suppose that G is not 3-connected, i.e.  $G = G_1 \cup G_2$  with  $V(G_1) \cap V(G_2) = \{x_1, x_2\}$ ,  $|V(G_i)| \geq 3$ . Since  $k \geq 3$ ,  $G_1$  and  $G_2$  have more than three points. Also, we may assume  $d_{G_1}(x_1) > 1$  otherwise we could replace  $x_1$  by its neighbor in  $G_1$ . Similarly, we may assume  $d_{G_2}(x_2) > 1$ . Then  $G_1 + (x_1, x_2)$  and  $G_2 + (x_1, x_2)$  have points of degree at most k and so, they can be k-colored (by induction or by 9.12a, whichever you prefer). In these k-colorations  $x_1, x_2$  get different colors, since they are adjacent. Therefore, by permuting the colors we can achieve that  $x_1$  gets the same color in both graphs and so does  $x_2$ . Now the two k-colorations can be put together to get a k-coloration of G [R.L. Brooks; see any textbook on graph theory].

14. (a) As noted in the hint, we have to show that "to be non-adjacent" is an equivalence-relation on V(G). This will imply that there is a partition  $\{V_1, V_2, \ldots\}$  of V(G) such that two points are adjacent iff they belong to distinct classes. The number of classes will be not more than k since G cannot contain a complete (k+1)-graph; the fact that there cannot be less than k

classes follows from the maximality of G.

The symmetry and reflexivity of the relation "to be non-adjacent" is trivial. Thus, what we have to show is that  $(a,b) \notin E(G)$ ,  $(b,c) \notin E(G)$  imply  $(a,c) \notin E(G)$ . By the maximality of G, there is a finite subgraph  $H_1$  of G such that  $H_1 + (a,b)$  is not k-colorable. Clearly  $a,b \in V(H_1)$ . Similarly, there exists a finite subgraph  $H_2$  of G with the property that  $H_2 + (b,c)$  is not k-colorable. Consider  $H_1 \cup H_2 + (a,c)$ ; we claim that this graph is not k-colorable. Suppose indirectly that it has a k-coloration  $\alpha$ . Then  $\alpha(a) = \alpha(b)$ , otherwise  $\alpha$  would be a k-coloration of  $H_1 + (a,b)$ . Similarly  $\alpha(b) = \alpha(c)$ . Now  $\alpha(a) = \alpha(c)$  and this is impossible as a and c are adjacent in  $H_1 \cup H_2 + (a,c)$ . Thus  $H_1 \cup H_2 + (a,c)$  is not k-colorable and so, it cannot be a subgraph of G. Thus a, c are non-adjacent in G.

(b) We may suppose G is simple. Consider all graphs G' such that V(G') = V(G), and all finite subgraphs of  $G' \cup G$  are k-colorable. Since this property depends on the finite subgraphs of G' only (i.e. G' has this property iff all finite subgraphs have), there is a maximal simple graph  $G_0$  with this property. Clearly  $G_0 \supseteq G$ . Thus by (a), V(G) has a partition  $V_1, \ldots, V_k$  such that two points are adjacent in G iff they belong to distinct classes. Clearly this partition is a good coloration of G [P. Erdős,-N.G. De Bruijn, Indag. Math. 13 (1961) 371–373; this proof is due to L. Pósa].

(c) Let  $e = (x, y) \in E(G)$ . Then the set

$$F_e = \{\alpha : \alpha(x) \neq \alpha(y)\}$$

of k-colorations of V(G) is closed, since its complement is the union of the open sets

$$\{\alpha\colon \alpha(x)=\alpha(y)=j\} \qquad (j=1,\ldots,k).$$

The set of legal colorations is just

$$\bigcap_{e \in E(G)} F_e$$
,

which is, therefore, closed as well.

To prove the Erdős–de Bruijn theorem, note that by Tihonov's Lemma the product space  $\underset{v \in V(G)}{\times} T_v$  is compact. The assumption that each finite subgraph of G is k-colorable translates into the assumption that the intersection of any finite number of sets  $F_e$  is non-empty. The proposition that G is k-colorable is equivalent to the assertion that the intersection of all of them is non-empty. But this follows by the Riesz Intersection Theorem.

15. I. Clearly  $\mathcal K$  contains some complete graphs. Let  $K_{k+1}$  be the least complete graph in  $\mathcal K$ . We claim that  $\mathcal K$  is precisely the class of non-k-colorable graphs. First let  $G \in \mathcal K$  and suppose indirectly that G is k-colorable. Then the k-coloration of G defines a homomorphism of it into  $K_k$ , thus by (i)  $K_k \in \mathcal K$ , a contradiction.

II. We now show that every non-k-colorable graph belongs to  $\mathcal{K}$ . Suppose G would be a counterexample. We may suppose G is such that connecting any two non-adjacent points of G by a new edge, we get a graph in  $\mathcal{K}$ . We claim that "to be non-adjacent" is an equivalence-relation on V(G). Suppose indirectly that  $(a, b) \notin E(G)$ ,  $(b, c) \notin E(G)$  but  $(a, c) \in E(G)$  (the other conditions of an equivalence-relation are trivially fulfilled). By the maximality of G,  $G + (a, b) \in \mathcal{K}$ ,  $G + (b, c) \in \mathcal{K}$ . Hence by (ii)  $G \in \mathcal{K}$ , a contradiction.

Thus "to be non-adjacent" is an equivalence-relation and hence, V(G) has a partition  $\{V_1, \ldots, V_m\}$  such that two points are adjacent iff they belong to distinct classes. Since G is not k-colorable,  $m \geq k + 1$ . But then G contains

a  $K_{k+1}$ , a contradiction as  $K_{k+1} \in \mathcal{H}$  but  $G \notin \mathcal{H}$ .

16. It is straightforward to show that operations  $(\alpha)$ ,  $(\beta)$ ,  $(\gamma)$  yield non-k-

colorable graphs if the initial graphs are not k-colorable.

Let  $\Re$  denote the class of all graphs arising by repeated application of  $(\alpha)$ ,  $(\beta)$ ,  $(\gamma)$  from the complete (k+1)-graph. We claim that  $\Re$  satisfies the conditions of 9.15. (i) is trivial. Let H be a graph, a, b,  $c \in V(H)$  such that  $H + (a, b) \in \Re$ ,  $H + (b, c) \in \Re$ . Let H' be a graph isomorphic with H and denote by x' the point of H' corresponding to  $x \in V(H)$ . Then  $H' + (b', c') \in \Re$ . Let us identify b and b' and connect a to c'; then identify x with x' ( $x \in V(H)$ ) and cancel the multiplicities of edges to get a simple graph. From H + (a, b) and H' + (b', c'), by operations  $(\gamma)$ ,  $(\beta)$  we obtain H this way. This proves that  $H \in \Re$ .

Thus  $\mathcal{H}$  consists of all non-r-colorable graphs for some r. Obviously, r = k. [G. Hajós; see B, S].

17. (a) By 5.3, a graph is 2-colorable iff it contains no odd circuit. Hence the critically 3-chromatic graphs are precisely the odd circuits.

(b) Assume for the sake of simplicity that n is odd; the even case could be treated similarly. Consider a complete bipartite graph  $K_{n,n}$  with color classes A, B. Take two disjoint odd circuits  $C_A$ ,  $C_B$  of length n and connect each point of  $C_A$  to a point of A, each point of  $C_B$  to a point of B by independent edges. Let G denote the resulting graph (Fig. 64).




Fig. 64

The graph G is not 3-colorable. In fact, every 3-coloration of  $K_{n,n}$  colors one of the sets A, B monochromatically. But then this coloration does not extend to the corresponding odd circuit.

On the other hand, G is  $\chi$ -critical. For let e = (x, y) be any edge. If  $x \in A$ ,  $y \in B$ , then color x and y red, all points of  $A - \{x\}$  blue, all points of  $B - \{y\}$  green. In this coloration neither A nor B is monochromatic, hence it extends to a 3-coloration of the whole graph G - e. We obtain by similar reasoning that G - e is 3-colorable for every edge e.

Since G has more than  $n^2$  edges, the desired graph is found [B. Toft, Studia Sci. Math. Hung. 5 (1970) 461–470].

(c) Again suppose for the sake of simplicity that n is odd. Take two disjoint n-circuits  $C_1$ ,  $C_2$  and connect each point of  $C_1$  to each point of  $C_2$ . The resulting graph is not 5-colorable, for we need 3 colors to color  $C_1$  and 3 different colors to color  $C_2$ . It is straightforward to find 5-colorations of the graphs G - e,  $e \in E(G)$ , thus G is  $\chi$ -critical. It trivially fulfils the degree requirement [G.A. Dirac, J. London Math. Soc. 27 (1952) 85–92].

18. Set  $\chi(G) = k$ . We show first that  $\chi(G') > k$ . Suppose indirectly that there is a k-coloration  $\alpha$  of G'. Let  $\alpha(y) = 1$ ; define a coloration  $\beta$  of V(G) by

$$\beta(x) = \begin{cases} \alpha(x) & \text{if } \alpha(x) \neq 1, \\ \alpha(x') & \text{if } \alpha(x) = 1. \end{cases}$$

Then  $\beta$  is a legitimate coloration of G. In fact, if  $(x, z) \in E(G)$ , then one of them, say x, has  $\alpha(x) \neq 1$ , so  $\beta(x) = \alpha(x)$ . Now no matter whether  $\beta(z) = \alpha(z)$  or  $\alpha(z')$ ,  $\beta(z)$  will be different from  $\beta(x)$  since both z, z' are neighbors of x.  $\beta$  uses only k-1 colors, which is a contradiction.

Now let  $e \in E(G')$ , we show that  $\chi(G' - e) \ge k$ . If  $e \in E(G)$ , then take a (k-1)-coloration of G - e, color the points x'  $(x \in V(G))$  with color k and y with

color 1. If e = (u, v'), take a (k-1)-coloration  $\beta$  of G - (u, v). Define

$$\alpha(z) = \begin{cases} \beta(z) & \text{if} \quad z \in V(G) - \{v\}, \\ k & \text{if} \quad z = v \text{ or } z = y, \\ \beta(x) & \text{if} \quad z = x'. \end{cases}$$

Finally, if e = (y, v'), let  $\beta$  be a (k-1)-coloration of G - v and define

$$\alpha(z) = \begin{cases} \beta(z) & \text{if} \quad z \in V(G) - \{v\}, \\ k & \text{if} \quad z = v, v' \text{ or } y, \\ \beta(x) & \text{if} \quad z = x'. \end{cases}$$

In all cases we have found a k-coloration of G' - e. Thus G' is  $\chi$ -critical and  $\chi(G') = k + 1$  as stated [J. Mycielsky; see S].

19. (a) Assume  $G_0$  is a subgraph of a 4-chromatic  $\chi$ -critical graph G. Then  $G - \{x_3, x_4\}$  is 3-colorable. Consider any 3-coloration of it. Then  $x_1, x_5$  have two different colors; therefore both  $x_2, x_4$  have the third color and so,  $x_3, x_4$  have different colors. But this implies our coloration is a good coloration of G as well, which is a contradiction.

(b) Suppose  $G_0 \subseteq G$ , where G is  $\chi$ -critical and  $\chi(G) = k + 1$ . Let  $e \in E(G_0)$ . Then G - e has a (k - 1)-coloration  $\alpha$ . Since this is not a good coloration of G, the endpoints of e have the same color. However, this means that  $\alpha$  yields a k-coloration of  $G_0/e$ .

Conversely, suppose that  $\chi(G_0/e) \leq k$  for every  $e \in E(G)$ . We show that  $G_0$  can be embedded into a critically (k+1)-chromatic graph even as an induced subgraph. The assumption means that for each  $e \in E(G)$ , there exists a k-coloration of V(G), which associates the same color with the endpoints of e, but different colors with the endpoints of any other edge of  $G_0$ . Let  $P_e$  be the partition of S induced by  $\alpha_e$ . By 9.8, we can find a graph G such that  $S \subseteq V(G)$  and the k-colorations of G induce the partitions  $P_e$  ( $e \in E(G_0)$ ) of S and no other partition. Set  $G' = G \cup G_0$ .

Now  $\chi(G') > k$ . For if there existed a k-coloration of G', this would induce a  $P_e$  on S, which is impossible as e joins two points in the same class of  $P_e$ . Therefore, G' contains a (k+1)-chromatic  $\chi$ -critical graph G''. All we have to show is that to obtain G'' from G' we do not have to remove any edge of  $G_0$ . Suppose indirectly that  $e \in E(G_0)$ ,  $e \notin E(G'')$ . Let  $\alpha_e$  be a k coloration of G' inducing  $P_e$  on S. By the definition of  $P_e$ ,  $P_e$  is a good coloration of  $P_e$  in  $P_e$ 

**20.** The first assertion is trivial: for any edge e of G, G - e is k-colorable and this yields a k-coloration of G' - e. Hence either G' itself is k-colorable or it is critically (k + 1)-chromatic.

If k=2, then the critically 3-chromatic graphs are the odd circuits. Split-

ting any point of these, we obtain a 2-chromatic graph (a path).

We show by example that for  $k \ge 3$  the second possibility can, in fact, occur. Put 3 copies of  $K_{k+1}$  together as in Hajós' construction shown in Fig. 65. Then by 9.16 (or trivially) the resulting graph G' is not k-colorable. On the other hand, even if we identify the points x and x', the resulting graph will be




Fig. 65

critically (k+1)-chromatic. For let us remove any edge e, for example from the  $K_{k+1}$  containing x. We can then color the points of this  $K_{k+1}$  in such a way that only the endpoints of e have the same color; in particular u and v are differently colored. Color the points of the other  $K_{k+1}$  (containing x') such that only the colors of t and w are the same. Color v' with the color of u. The remaining k-2 points can be colored with the k-2 colors different from the colors of v', w' and u. Similar arguments yield a k-coloration of G-e, when e is any other edge.

21. Assume indirectly that some  $m \leq k-1$  edges  $e_1, \ldots, e_m$  separate G into two pieces  $G_1, G_2$ . We may assume this is a minimal cutset, so that each  $e_i$  connects  $G_1$  to  $G_2$ . Consider a k-coloration of  $G_1$  with color-classes  $T_1, \ldots, T_k$  and a k-coloration of  $G_2$  with color-classes  $S_1, \ldots, S_k$  (these exist by the criticality of G).

We want to match each  $T_i$  with some  $S_i$  such that no edge connects  $T_i$  and  $S_i$ . Since there are only  $m \leq k-1$  edges to consider, there certainly exists an  $S_i$  not connected to  $T_1$  by any edge. We can clearly select this  $S_i$  such that either  $T_1$  or  $S_i$  should be incident with one of the edges  $e_1, \ldots, e_m$ . Let this index be 1'. Then we have to match the remaining k-1 classes T with the remaining k-1 classes S, and the number of edges  $e_i$  between these classes is now  $\leq m-1 \leq k-2$ . So we can proceed in the same way as above. (Actually what we have shown is that removing k-1 edges of  $K_{k,k}$ , the remaining bipartite graph has a 1-factor. This would of course follow from the König-Hall theorem immediately.)

Now the partition  $\{T_1 \cup S_1, \ldots, T_k \cup S_k\}$  is a k-coloration of G, a contradiction. [G. Dirac; B.]

**22.** Suppose first that G is separable, i.e.  $G = G_1 \cup G_2$  with  $V(G_1 \cap G_2) = \{x\}$ ,  $|V(G_i)| \geq 2$ . Since G is  $\chi$ -critical,  $\chi(G_i) < \chi(G) = k + 1$ . Consider a k-coloration  $\alpha_i$  of  $G_i$ . By permuting the colors we may assume that  $\alpha_1(x) = \alpha_2(x)$ . Then putting  $\alpha_1$  and  $\alpha_2$  together we obtain a k-coloration of G, a contradiction.

Now suppose that  $G = G_1 \cup G_2$  with  $|V(G_i)| \ge 3$ ,  $V(G_1 \cap G_2) = \{x, y\}$ . Let  $\alpha_i$  be a k-coloration of  $G_i$  (i = 1, 2). It is impossible that  $\alpha_1(x) \ne \alpha_1(y)$  and  $\alpha_2(x) \ne \alpha_2(y)$ ; since then, by permuting colors, we could assume that  $\alpha_1(x) = \alpha_2(x)$ ,  $\alpha_1(y) = \alpha_2(y)$  and so, we could put  $\alpha_1$ ,  $\alpha_2$  together to get a k-coloration.

tion of G. We reach a contradiction similarly, if  $\alpha_1(x) = \alpha_1(y)$  and  $\alpha_2(x) = \alpha_2(y)$ . So we have that, choosing the indices appropriately, any k-coloration of  $G_1$  paints x, y with the same color, but no k-coloration of  $G_2$  does so. Hence,  $G_1' = G_1 + (x, y)$  is not k-colorable and neither is  $G_2'$  obtained from  $G_2$  by identifying x and y.

We claim that  $G'_1$ ,  $G'_2$  are  $\chi$ -critical. Obviously,  $\chi(G'_1) = \chi(G'_2) = \chi(G)$ . Let  $e \in E(G'_1)$ . Now G - e is k-colorable; from above, this coloration associates different colors with x and y ( $G_2$  forces this) and therefore, it yields a k-colora-

tion of  $G'_1 - e$ . Thus  $G'_1$  is  $\chi$ -critical. Similarly,  $G'_2$  is  $\chi$ -critical.

Note that the k-colorability of  $G_2$  does not follow from the fact that  $G'_2$  is critically (k+1)-chromatic (cf. 9.20). Therefore, we can summarize the result as follows:

Every critically (k+1)-chromatic graph G, which is not 3-connected arises by the following construction. We take two critically (k+1)-chromatic graphs  $G'_1$ ,  $G'_2$ . We remove an edge  $(x_1, y_1)$  of  $G'_1$ . We split a point of  $G'_2$  into two points  $x_2$ ,  $y_2$  in such a way that the resulting graph is k-colorable. Finally, we identify  $x_1$  with  $x_2$  and  $y_1$  with  $y_2$ . It is easy to verify that the converse is true as well: every graph arising by this construction is critically (k+1)-chromatic [G.A. Dirac, J. Reine Angew. Math. 214/215 (1964) 43–52].

**23.** Let  $\alpha_i$  be a k-coloration of  $G - V(G_i)$ . Then  $\alpha_i$  induces a partition  $P_i$  of S into at most k classes. We claim that  $P_i \neq P_j$  for  $i \neq j$ . For if  $P_i = P_j$  then, by permuting colors, we could arrange for  $\alpha_i$  and  $\alpha_j$  to color S the same way. Now use  $\alpha_i$  on  $G - V(G_i)$  and  $\alpha_j$  on  $G_i$  (where it is defined, as  $i \neq j$ ) to get a k-coloration of G. This contradicts  $\chi(G) = k + 1$ . Thus the  $P_i$ 's are different and hence  $N \leq P_{m,k}$ .

If we want to construct a  $\chi$ -critical graph G such that an  $S \subseteq V(G)$ , |S| = m separates it into exactly  $p = P_{m,k}$  components, let  $P_1, \ldots, P_p$  be the partitions of S into at most k classes. For each i, 9.8 yields a graph  $G_i$  containing S such that the k-colorations of  $G_i$  induce all partitions of S into at most k classes except  $P_i$ . We may assume  $V(G_i) \cap V(G_j) = S$ . Let

 $G = G_1 \cup \ldots \cup G_p$ .

G is in general not  $\chi$ -critical. But certainly, it is not k-colorable, for a k-coloration of G would induce a partition  $P_i$  of S and this contradicts the definition of  $G_i$ . So we can consider a  $\chi$ -critical subgraph G' of G with  $\chi(G') = k + 1$ . We show that G' has the desired property: it suffices to verify

- (a)  $S \subseteq V(G')$ ;
- (b) G' meets every  $G_i S$ .
- (a) Suppose indirectly that  $a \in S$ ,  $a \notin V(G')$ . Let say,  $P_1 = \{\{a\}, S -\{a\}\}, P_2 = \{S\}$ . Let  $\alpha_1$  be a k-coloration of  $G_1$ , which induces the partition  $P_2$  of S and let  $\alpha_i$  be a k-coloration of  $G_i$  inducing  $P_1$  for  $i = 2, \ldots, p$ . Since  $P_1$  and  $P_2$  coincide on S a, we may assume the  $\alpha_i$ 's can be put together to get a k-coloration of G a; therefore,  $G' \subseteq G a$  is k-colorable, a contradiction.
- (b) Suppose G' does not meet  $G_i S$ . Let  $\alpha_j$  be a k-coloration of  $G_j$  inducing  $P_i$  of S  $(j \neq i)$ . Then the  $\alpha_j$ 's can be put together to get a k-coloration of  $G (V(G_i) S)$ . Since G' is a subgraph of this, G' is k-colorable, a contradiction.

- **24.** Let  $(x, y) \in E(G)$ . Since G is  $\chi$ -critical, G (x, y) has a k-coloration  $\alpha$ . Since G itself is not k-colorable,  $\alpha(x) = \alpha(y) = 1$  (say). Consider another color i and the graph  $H_i$  induced by those points of colors 1 and i. Let  $H'_i$  be the component of  $H_i$  containing x. Then we claim that  $y \in V(H'_i)$ ; for otherwise, we could switch colors 1 and i in  $V(H'_i)$ ; the arising coloration  $\alpha'$  is legitimate and has  $\alpha'(x) \neq \alpha'(y)$ , i.e. it is a good coloration of G, which is impossible. So there are (x, y)-paths in  $H_i$ ; let  $P_i$  be a shortest one. Since  $P_i$  contains alternating points with colors 1 and i, it is of even length; by minimality, it spans no chord. Since each edge of  $P_i$  connects a point of color 1 to a point of color i, the paths  $P_2, \ldots, P_k$  must be edge-disjoint. This proves the assertion. We remark that 9.21 follows easily from this.
- 25. (a) Let  $G_3$  be a 5-circuit and, if  $G_r$  is defined, let  $G_{r+1}$  be the graph obtained by the construction of problem 9.18. Then it follows by the assertion of this problem that  $G_r$  is an r-chromatic graph ( $\chi$ -critical too) and it is easy to check that it has no triangles.
- (b) Let  $G_3$  be a 7-circuit. Suppose  $G_n$   $(n \ge 3)$  is defined such that  $\chi(G_n) \ge n$ . Take a set S of  $n(|V(G_n)|-1)+1$  points. With each  $|V(G_n)|$ -element subset F of S, we associate a copy  $G_F$  of  $G_n$  (these copies should be disjoint), and join it to the points of F by a matching. Let  $G_{n+1}$  be the resulting graph, then  $\chi(G_{n+1}) > n$ . Suppose  $G_{n+1}$  could be n-colored, then some  $|V(G_n)|$  points of S would get the same color 1; let F be the set of these points. As remarked in the hint, the points of  $G_F$  cannot get the color 1, so they are, in fact, (n-1)-colored; this, however, contradicts the definition of  $G_n$ .

It is immediate that  $G_n$  contains no 3-, 4-, or 5-circuits. [W.T. Tutte (Blanche

Descartes); see S.]

26. Let  $\chi(L(G)) = k$  and consider a k-coloration  $\alpha$  of L(G). Then  $\alpha$  is a k-coloration of the edges of G such that the edges (x, y) and (y, z) cannot have the same color. Consider the red edges. For each point x, either all red edges adjacent to x have their tail at x, or all of them have their head at x. This yields a bipartition of the graph formed by the red edges. Hence G can be decomposed into k bipartite graphs  $G_1, \ldots, G_k$ . Let  $\alpha_i$  be a 2-coloration of  $G_i$ ,  $i = 1, \ldots, k$ . Then using the vector  $(\alpha_1(x), \ldots, \alpha_k(x))$  to color x, we obtain a  $2^k$ -coloration of G. Hence

$$\chi(G) \leq 2^k$$
.

27. Let  $\vec{G}_{r,n}$  be the r-times iterated line-graph of  $T_n$ , the transitive tournament. By the preceding problem,

$$\chi(\vec{G}_{r,n}) \ge \underbrace{\log \log \ldots \log n}_{r}.$$

Now let  $G_{r,n}$  be  $\vec{G}_{r,n}$  with the orientations ignored. We show that  $G_{r,n}$  has no odd circuits of length at most 2r+1. We use induction on r. Let C be a closed path in  $G_{r,n}$  of length 2s+1. Since  $\vec{G}_{r,n}$  is acyclic, C decomposes into arcs  $P_1$ ,  $Q_1, P_2, Q_2, \ldots, P_t$ ,  $Q_t$  such that  $P_1, \ldots, P_t$  are directed paths in  $\vec{G}_{r,n}$  going in the same direction around C and  $Q_1, \ldots, Q_t$  are directed paths going in the opposite direction; moreover, they follow each other on the walk as listed.

Since  $\bar{G}_{r,n}$  is acyclic, it is easy to see that the points of  $P_i$  (or  $Q_i$ ) correspond to the edges of a path  $P'_i$  (or  $Q'_i$ ) in  $\bar{G}_{r-1,n}$ . Then the last edge of  $P'_1$  coincides with the last edge of  $Q'_2$  (with respect to orientation in  $G_r$ ), etc. Thus, if we remove these last edges the rest of  $P'_1, Q'_1, \ldots, P'_t, Q'_t$  forms a walk of length 2s + 1 - 2t. Clearly, this walk contains an odd circuit of length  $l \leq 2s + 1 - 2t \leq 2s - 1$ . By the induction hypothesis  $l \geq 2r$ , whence 2s + 1 > 2r + 1 as stated.

Second solution. Consider the surface of the k-dimensional unit ball. Connect two points of it, if their distance, measured on the surface, is at least  $(1-1/l)\pi$ . The resulting graph is not k-colorable. In fact, a theorem of Borsuk asserts that in any k-coloration of the surface of the k-dimensional ball, one of the color-classes has diameter 2; this color-class then contains adjacent points.

On the other hand, let  $(x_0, \ldots, x_{2p})$  be any odd circuit in the graph. If d denotes the spherical distance, then  $d(x_i, x_{i+1}) \ge \left(1 - \frac{1}{l}\right)\pi$ , whence  $d(x_i, x_{i+2}) \le 2\pi/l$ , and hence by induction  $d(x_0, x_{2p}) \le 2p\pi/l$ . On the other hand, this distance is at least  $\left(1 - \frac{1}{l}\right)\pi$ . Hence  $2p + 1 \ge l$ . So our graph contains no short odd circuits.

This graph is infinite but by the Erdős-de Bruijn theorem 9.14, it has a finite subgraph with similar properties [P. Erdős-A. Hajnal, Ann. N.Y. Acad. Sci. 157 (1970) 115-124].

- 28. (a) Let, say,  $I_1 = [b, a]$  be the interval whose endpoint a is the first from left. The system  $I_2, \ldots, I_n$  determines the graph  $G x_1$ . By induction,  $\chi(G x_1) = \omega(G x_1) \leq \omega(G)$ . Observe, moreover, that if,  $x_i$  is adjacent to  $x_1$ , i.e.  $I_i$  meets  $I_1$ , then  $a \in I_i$  (since otherwise the right endpoint of  $I_i$  would lie to the left of a). Since at most  $\omega(G)$  intervals contain a (including  $I_1$ ), the degree of  $x_1$  is at most  $\omega(G) 1$ . Therefore, we can extend the  $\omega(G)$ -coloration of  $G x_1$  to  $x_1$ . Hence,  $\chi(G) \leq \omega(G)$ . Since, obviously,  $\chi(G) \geq \omega(G)$ , we are done.
- (b) Let  $I_1, I_2, \ldots, I_r$  denote the intervals containing a. Then  $x_1, x_2, \ldots, x_r$  form an independent set in  $\overline{G}$ . Consider  $\overline{G} = \{x_1, \ldots, x_r\}$ . If  $x_{i_1}, \ldots, x_{i_q}$  form a clique in  $\overline{G} = \{x_1, \ldots, x_r\}$ , then  $I_{i_1}, \ldots, I_{i_q}$  are disjoint intervals not containing a. By the definition of a,  $I_{i_1}, \ldots, I_{i_q}$  are to the right of a and therefore  $I_1$  is disjoint from them. Hence  $\{x_1, x_{i_1}, \ldots, x_{i_q}\}$  is a clique in  $\overline{G}$ . This proves that  $\chi(\overline{G} = \{x_1, \ldots, x_r\}) \leq \omega(\overline{G}) = 1$ . Thus,  $\overline{G} = \{x_1, \ldots, x_r\}$  can be colored by  $\omega(\overline{G}) = 1$  colors and we can use the same new color for  $x_1, \ldots, x_r$ . Hence  $\chi(\overline{G}) \leq \omega(\overline{G})$ . The equality is obvious as before.
- (c) Let  $x_2$ ,  $x_3$  be the neighbors of  $x_1$  on the circuit. Then, as we have already seen,  $I_2$ ,  $I_3$  contain a and therefore, they meet each other. Thus,  $(x_2, x_3)$  is a chord of the circuit (unless  $C = (x_1, x_2, x_3)$ ) [G. Hajós, *Intern. Math. Nachr.* 11 (1957) Sondernummer 65].
- **29.** (a) If  $G_1$  is a component of G-S, then it is adjacent to all points of S, since otherwise a set smaller than S (the set of neighbors of  $G_1$ ) would separate G.

Let  $u, v \in S$  and  $G_1, G_2$  two components of G - S. Let  $P_i$  be a minimal path, which connects u, v through  $G_i$ . Then  $P_i$  has no chords by minimality, except possibly (u, v). Set  $C = P_1 \cup P_2$ , then C is longer than 3 and therefore, C has a chord. This cannot join an inner point of  $P_1$  to an inner point of  $P_2$  as these points are in distinct components of G - S. Hence, it must join two points of the same  $P_i$ . This is only possible, if it is (u, v). Thus, u, v are adjacent. [G. Dirac]

(b) First, let G be defined through the subtrees as formulated, and consider

a circuit C of G. We may assume C is a Hamiltonian circuit.

Let us remove points of degree 1 from T as long as the remainders of  $F_1, \ldots, F_n$  have at least one point. We cannot change the graph G, i.e. if  $F_i \cap F_j \neq \emptyset$  and x is a point of degree 1, then  $(F_i - x) \cap (F_i - x) \neq 0$ ; since, if x is a common point of  $F_i$  and  $F_j$ , then certainly so is its neighbor. When we get stuck, we have an  $F_i$  with one point; let  $F_1$  be this, say. Let  $F_2$ ,  $F_3$  be the two neighbors of  $F_1$  on C, then  $F_2$ ,  $F_3$  both contain the (unique) point of  $F_1$ , and hence,

they intersect. So  $(x_2, x_3)$  is a chord of C.

Conversely, suppose G is a graph without chordless circuits of length at least 4; we show by induction on |V(G)| that it is representable in the mentioned way. Let S be a minimum cutset of G, then by the previous result, S induces a complete graph H. Let  $G = G_1 \cup G_2$ ,  $G_1 \cap G_2 = H$ ,  $G_1$ ,  $G_2 \neq H$ . By the induction hypothesis,  $G_1$ ,  $G_2$  can be represented in the given way, through certain subtrees of trees  $T_1$ , and  $T_2$ , respectively. Let  $F_1, \ldots, F_{|S|}$ ;  $F'_1, \ldots, F'_{|S|}$  be the subtrees corresponding to the points of S in the representation of  $G_1$  and  $G_2$ , respectively. By 6.18,  $F_1, \ldots, F_{|S|}$  have a point v in common and similarly,  $F'_1, \ldots, F'_{|S|}$  have a point w in common. Set

$$T = T_1 \cup T_2 + (v, w), F_i'' = F_i \cup F_i' + (v, w), \qquad (i = 1, \dots, |S|),$$

then the subtrees  $F_i''$  and those corresponding to points of  $G_1 - S$  and  $G_2 - S$  yield a representation of G.

(c) We give here proofs using the representation (b). As we have seen in the first part of that proof, we may remove points of degree 1 from T, and still get a system of trees representing the same G until finally, we have a tree  $F_1$  with one point. Let  $x_1$  be the corresponding point of G. Then  $G-x_1$  has  $\chi(G-x_1)=\omega(G-x_1)\leq \omega(G)$ . Moreover, the neighborhood of  $x_1$  forms a complete graph and so,  $x_1$  has degree at most  $\omega(G)-1$ . Thus coloring  $G-x_1$  by  $\omega(G)$  colors we can find a color for  $x_1$ .

To show  $\chi(\overline{G}) = \omega(\overline{G})$ , remove  $x_1$  and all its G-neighbors  $x_2, \ldots, x_r$ . Any set which forms a clique in  $\overline{G} - \{x_1, \ldots, x_r\}$ , i.e. which is independent in  $G - \{x_1, \ldots, x_r\}$  can be completed with  $x_1$ . So  $\omega(\overline{G} - \{x_1, \ldots, x_r\}) \leq \omega(\overline{G}) - 1$ . Color  $\overline{G} - \{x_1, \ldots, x_r\}$  by  $\omega(\overline{G}) - 1$  colors by induction; then using a further color for  $x_1, \ldots, x_r$  we get the desired coloration of  $\overline{G}$ . [A. Hajnal, J. Surányi;

C. Berge; see B.]

**30.** Let G have m isolated points  $x_1, \ldots, x_m$ . By 7.2,  $V(G - \{x_1, \ldots, x_m\})$  can be covered by

 $\alpha(G - \{x_1, \ldots, x_m\}) = \alpha(G) - m$ 

edges. So V(G) can be covered by  $\alpha(G)$  edges and points of G. We may assume that these edges are disjoint, since if two have an endpoint in common, we can replace one of them by its other endpoint. Then these  $\alpha(G)$  edges and points of

G yield  $\alpha(G) = \omega(\overline{G})$  independent subsets of  $\overline{G}$  covering the points, i.e. an  $\omega(\overline{G})$ -coloration of G. Hence,  $\chi(G) \leq \omega(G)$ . Since the converse is true as well, we have equality.

**31.** Let  $H_i$  be a maximum complete subgraph of  $G_i$  (i=1,2), then  $V(H_1)\times$  $\times V(H_2)$  spans a complete subgraph of  $G_1 \cdot G_2$  of size  $|V(H_1)| \cdot |V(H_2)| =$  $= \omega(G_1) \cdot \omega(G_2). \text{ Hence } \omega(G_1 \cdot G_2) \geq \omega(G_1) \cdot \omega(G_2).$  On the other hand, let  $\alpha_i$  be a  $\chi(G_i)$ -coloration of  $G_1$  (i = 1, 2). Define a col-

oration a by

$$\alpha(x_1, x_2) = (\alpha_1(x_1), \alpha_2(x_2)) \quad (x_i \in V(G_i)),$$

then it is easy to verify that  $\alpha$  is a good coloration of  $G_1 \cdot G_2$  and uses  $\chi(G_1) \cdot \chi(G_2)$  colors. Therefore  $\chi(G_1 \cdot G_2) \leq \chi(G_1) \cdot \chi(G_2)$ . Thus,

$$\chi(G_1 \cdot G_2) \leq \chi(G_1) \cdot \chi(G_2) = \omega(G_1) \cdot \omega(G_2) \leq \omega(G_1 \cdot G_2)$$

and, the converse inequality being trivial, we have equality as stated.

- 32. (a) The exact formulation of the hint is this. Let  $\alpha(x)$  denote the maximum length of a chain  $x = x_1 < x_2 < \ldots < x_a$ . Then  $1 \le \alpha(x) \le \omega(G)$ , because  $\omega(G)$  is the maximum size of a totally ordered subset, i.e. a chain. Moreover,  $\alpha$  is a legitimate coloration. For suppose x and y are adjacent, say x > y. Then a chain  $x = x_1 < x_2 < \ldots < x_{\alpha(x)}$ , of maximum length  $\alpha(x)$  can be extended over y, showing  $\alpha(y) \geq \alpha(x) + 1$ .
- (b) Define a directed graph G by orienting the edge (x, y) from x to y if x < y. Then, obviously,  $\omega(\overline{G}) = \alpha(\overline{G})$ . By 8.4, we can cover the points of G by  $\alpha(\vec{G})$  disjoint paths. A path of  $\vec{G}$  corresponds to a chain in P. So, it spans a complete subgraph of G; i.e. an independent set in  $\overline{G}$ . Thus,  $\overline{G}$  can be partitioned into  $\alpha(\bar{G}) = \omega(\bar{G})$  independent sets, i.e.  $\chi(\bar{G}) \leq \omega(\bar{G})$ . The converse is, as always, trivial. [This is an equivalent form of Dilworth' theorem; see B.]
- 33. In 9.28, 29, 30, 32, the graphs considered are such that all induced subgraphs of them belong to the same class, thus they satisfy  $\chi(G) = \omega(G)$  as well. However, we do not get perfect graphs from 9.31 in general. To show this, let  $G_1, G_2$  be two 3-paths, then  $G_1 \cdot G_2$  contains a chordless 7-cycle as shown in Fig. 66.




Fig. 66

**34.** Suppose first that G is perfect. Let G' be any induced subgraph of G. Let S be a color-class of a  $\chi(G')$ -coloration of G'. Then  $\chi(G'-S)=\chi(G')-1$  and so, S meets all maximum cliques of G'.

Suppose that, conversely, every induced subgraph of G has an independent set, which meets all maximum cliques. We prove that  $\chi(G') = \omega(G')$  for the induced subgraphs by induction on |V(G')|. Let S be an independent set, which meets all maximum cliques of G'. Since S is independent,

$$\chi(G') \le \chi(G' - S) + 1$$

and by its definition,

$$\omega(G') \ge \omega(G' - S) + 1.$$

The two right-hand sides are equal by the induction hypothesis, thus  $\chi(G') \le \omega(G')$ . The converse inequality is trivial.

**35.** We may substitute the graphs  $G_x$  for the points of G one by one, and it suffices to show that perfectness is preserved at each step. So we may assume that a non-trivial  $G_{x_0}$  is substituted for one point  $x_0$ , while the other points are not touched (or we may say one-point graphs are substituted for them).

Also, it suffices to show that the maximum cliques of G' can be covered by an independent set; for the induced subgraphs of G' arise in the same way as G' and so, the previous exercise yields that G' is perfect. Now let us  $\chi(G)$ -color G and take the color-class S containing  $x_0$ . Also, take an independent subset

 $S_1$  of  $G_{x_0}$ , which meets all maximum cliques of  $G_{x_0}$ .

Then clearly  $(S - x_0) \cup S_1$  is independent in G'. We show it covers all maximum cliques of G'. Let T be a maximum clique of G'. If T does not meet  $G_{x_0}$ , then it is a maximum clique of G and so  $S - x_0$  covers it. If T meets  $G_{x_0}$ , then, obviously, it contains a maximum clique of  $G_{x_0}$  and  $G_{x_0}$  covers it [L. Lovász, Discrete Math. 2 (1972) 253–267].

**36.** If P is a partition of V(G) into independent sets, then, obviously, it can arise in

$$\lambda(\lambda-1)\ldots(\lambda-|P|+1)$$

ways as a partition induced by  $\lambda$ -colorations of G; this clearly also holds, if  $|P| > \lambda$ . Hence

(1) 
$$P_G(\lambda) = \sum_{P} \lambda(\lambda - 1) \dots (\lambda - |P| + 1),$$

where the summation is over all partitions of V(G) into independent sets. Thus

 $P_{\mathcal{C}}(\lambda)$  is a polynomial indeed.

Obviously, each term in (1) has degree at most n = |V(G)| and the only term with this degree belongs to the partition into one-element sets. Thus, if one-element sets are independent, i.e. G has no loops, then  $P_G(\lambda)$  is of degree n and  $\lambda^n$  has coefficient 1.

If G has a loop, then it has no good coloration at all, so  $P_G(\lambda) = 0$ . [Concerning chromatic polynomials see O, B, Wi; R.C. Read, J. Comb. Theory 4 (1968) 52–71; W.T. Tutte, Comb. Structures and their Appl. Gordon and Breach 1970, 439–453.]

37. To get all  $\lambda$ -colorations of G, consider all mappings of V(G) into  $\{1, \ldots, \lambda\}$ . We have to exclude those, which map the two endpoints of an edge onto the same point. The inclusion-exclusion formula (2.2) yields

$$P_{\mathit{G}}(\lambda) = \sum_{T \subseteq \mathit{E}(\mathit{G})} (-1)^{|T|} p(T,\lambda),$$

where  $p(T, \lambda)$  denotes the number of those mappings which identify the endpoints of each edge of T. Note that a mapping does this if and only if it maps each component of the graph  $\langle V(G), T \rangle$  onto a single number; hence,  $p(T, \lambda) = \lambda^{c(T)}$  and the formula is proved.

38. If a  $\lambda$ -coloration of G-e associates different colors with the endpoints of e, it is a good  $\lambda$ -coloration for G, and conversely. Thus  $P_{G-e}(\lambda)-P_G(\lambda)$  is the number of those  $\lambda$ -colorations of G-e, which associate the same number with the endpoints of e. Now note that such a  $\lambda$ -coloration of G-e yields a  $\lambda$ -coloration of G/e and conversely. Hence,

$$P_{G-e}(\lambda) - P_{G}(\lambda) = P_{G/e}(\lambda)$$

for every natural number  $\lambda$ . Since we have polynomials on both sides, this must hold for every  $\lambda$ .

**39.** Since  $K_n$  has only one partition into independent sets, the formula in the solution of 9.36 implies that

$$P_{K_n}(\lambda) = \lambda(\lambda - 1) \dots (\lambda - n + 1).$$

Let F be a tree and x a point of degree 1 of F. Any  $\lambda$ -coloration of F - x can be extended to x in  $\lambda - 1$  ways; hence

$$P_F(\lambda) = (\lambda - 1) P_{F-r}(\lambda)$$
.

Since the chromatic polynomial of the one-point tree is  $\lambda$ , we have

$$P_F(\lambda) = \lambda(\lambda - 1)^{n-1}$$
.

Let  $C_n$  be a circuit of length n, and let e be an edge of it. Then

$$P_{C_n}(\lambda) = P_{C_n-e}(\lambda) - P_{C_n/e}(\lambda).$$

Now note that  $C_n - e$  is a path on n points, while  $C_{n/e}$  is an (n-1)-circuit  $C_{n-1}$ . Thus

$$P_{C_n}(\lambda) = \lambda(\lambda - 1)^{n-1} - P_{C_{n-1}}(\lambda)$$

and repeating this reduction,

$$P_{C_n}(\lambda) = \lambda(\lambda - 1)^{n-1} - \lambda(\lambda - 1)^{n-2} + \dots + (-1)^{n-2}\lambda(\lambda - 1),$$

since  $P_{C_1}(\lambda) = 0$ . Summing this series,

$$P_{C_n}(\lambda) = (\lambda - 1)^n + (-1)^n (\lambda - 1).$$

Finally, let  $W_n$  be the wheel with n points. Color the center arbitrarily, then the rim has to be colored by the remaining  $\lambda - 1$  colors. Thus

$$P_{W_n}(\lambda) = \lambda P_{C_{n-1}}(\lambda - 1) = \lambda(\lambda - 2)^{n-1} + \lambda(-1)^{n-1}(\lambda - 2).$$

**40.** Let  $G_1, \ldots, G_m$  be the components of G. Then we can combine  $\lambda$ -colorations of them arbitrarily to get a  $\lambda$ -coloration of G, thus

$$P_{G}(\lambda) = P_{G_1}(\lambda) \dots P_{G_m}(\lambda).$$

Now suppose that G is connected and let  $B_1, \ldots, B_m$  be its blocks. We may assume  $B_m$  is a block, which has only one point  $x_0$  in common with the rest and set  $G' = B_1 \cup \ldots \cup B_{m-1}$ .

Take a  $\lambda$ -coloration  $\alpha_0$  of G'. Now those and only those  $\lambda$ -colorations of  $B_m$  yield a  $\lambda$ -coloration of G, which color  $x_0$  with  $\alpha(x_0)$ . Since this only means permutation of colors, the number of such  $\lambda$ -colorations of  $B_m$  is

$$\frac{1}{\lambda}P_{B_m}(\lambda).$$

Hence

$$P_G(\lambda) = P_{G'}(\lambda) \frac{1}{\lambda} P_{B_m}(\lambda)$$

and by induction we obtain

$$P_G(\lambda) = \frac{1}{\lambda^{m-1}} P_{B_1}(\lambda) \dots P_{B_m}(\lambda).$$

**41.** Let G be the graph obtained from the system  $\{[a_1, b_1], [a_2, b_2], \ldots, [a_n, b_n]\}$  of intervals. We may assume  $b_1 < b_2 < \ldots < b_n$ . Let  $x_i$  be the point of G corresponding to  $[a_i, b_i]$ . Then, as observed and used before, the neighbors of  $x_1$  form a complete graph. Hence, if we take any  $\lambda$ -coloration of  $G - x_1$ , this excludes exactly  $d(x_1)$  colors for  $x_1$  (we assume  $\lambda$  is large). Thus,

$$P_G(\lambda) = (\lambda - d(x_1)) P_{G-x_1}(\lambda).$$

Let  $\mu_i$  denote the number of intervals containing  $b_i$ , then  $d(x_1) = \mu_1 - 1$ , so we may write

$$P_G(\lambda) = (\lambda - \mu_1 + 1)P_{G-x_1}(\lambda).$$

Continuing in a similar manner, we get

$$P_G(\lambda) = (\lambda - \mu_1 + 1)(\lambda - \mu_2 + 1)\dots(\lambda - \mu_n + 1)$$

(we have to remark that the first j intervals do not contain  $b_{j+1}$  by definition, so if we remove them,  $b_{j+1}$  will be contained in the same number  $\mu_{j+1}$  of intervals).

Note that 9.28 is an immediate consequence; if  $\lambda \ge \omega(G) = \max \mu_i$ , then  $P_G(\lambda) \ne 0$ , so there exists a  $\lambda$ -coloration.

**42.** We use induction on |E(G)|. For the edgeless graph on n points we have  $P_G(\lambda) = \lambda^n$ . We may assume that G is simple since reducing multiple edges to simple ones does not influence  $P_G(\lambda)$ . So G/e has no loops. By 9.38,

$$P_{G}(\lambda) = P_{G-e}(\lambda) - P_{G/e}(\lambda)$$

for every edge e of G. By the induction hypothesis

$$P_{G-e}(\lambda) = \lambda^n - a'_{n-1}\lambda^{n-1} + a'_{n-2}\lambda^{n-2} - \dots$$
  
 $P_{G/e}(\lambda) = \lambda^{n-1} - a''_{n-2}\lambda^{n-2} + a''_{n-3}\lambda^{n-3} - \dots$ 

where  $a'_i$ ,  $a''_i \geq 0$ . From the recurrence relation we get

$$P_G(\lambda) = \lambda^n - (a'_{n-1} + 1) \lambda^{n-1} + (a'_{n-2} + a''_{n-2}) \lambda^{n-2} - \dots,$$

which proves the assertion. It is seen that  $a_{n-1} = a'_{n-1} + 1$ , thus it follows by induction that  $a_{n-1}$  is the number of edges of G (provided G is simple).

To find an interpretation for  $a_1$ , consider the formula given in 9.37. The linear terms correspond to those subgraphs, which are connected spanning subgraphs; thus,  $(-1)^{n-1}a_1$  is the difference between the numbers of connected spanning subgraphs with an even and an odd number of edges, respectively. It is remarkable that  $a_1 \geq 0$  implies an inequality between these two numbers. The reader may find it interesting to give a direct proof.

**43.** We use induction on |E(G)|. If the graph is a tree the statement is true by 9.40 (and the binomial theorem). Suppose G is a simple connected graph, which is not a tree, then it has an edge e such that G - e is connected. Obviously so is G/e and they both have no loops. By the previous solution (and with the notation as there),

$$a_i \geq a'_i > 0$$
.

Moreover,  $a'_{n-1} < a'_{n-2} < \ldots < a'_{\lfloor \frac{n}{2} \rfloor}$ ,  $a''_{n-2} < \ldots < a''_{\lfloor \frac{n-1}{2} \rfloor}$  by the induction hypothesis, thus

$$a_{n-1} < \ldots < a_{\left[\frac{n}{2}\right]}$$

44. By 9.36,

$$P_G(\lambda) = \sum_{P} \lambda(\lambda - 1) \dots (\lambda - |P| + 1),$$

where P ranges over all partitions of V(G) into independent sets. If  $\lambda > n-1$  all terms are positive.

**45.** We may suppose that G is connected. We show  $(-1)^{n-1}P_G(\lambda) > 0$ , if  $0 < \lambda < 1$ . The statement is true for trees. Let e be an edge of G such that G - e is connected. Then

$$(-1)^{n-1}P_{G-e}(\lambda) > 0, \quad (-1)^{n-2}P_{G/e}(\lambda) > 0$$

by the induction hypothesis, so

$$(-1)^{n-1}P_G(\lambda) = (-1)^{n-1}\{P_{G-e}(\lambda) - P_{G/e}(\lambda)\} =$$

$$= (-1)^{n-1}P_{G-e}(\lambda) + (-1)^{n-2}P_{G/e}(\lambda) > 0.$$

**46.** (a) If G is connected, 9.43 implies 0 is a simple root of  $P_G$ . Thus by 9.40, the multiplicity of the root 0 is equal to the number of components. Let G be 2-connected, we show by induction on |E(G)| that 1 is a simple root; we show that, in fact,

$$(-1)^n \frac{P_G(\lambda)}{\lambda - 1} > 0.$$

To this end let us first remark if G has an edge, then 1 is a root by definition; so if G is connected but not 2-connected and  $G \neq K_2$ , then 1 is a multiple root, which follows immediately from 9.40. Thus, if e is any edge of G,

$$(-1)^n \frac{P_{G-e}(\lambda)}{\lambda - 1} \bigg|_{\lambda = 1} \ge 0,$$

$$(-1)^n \frac{P_{G/e}(\lambda)}{\lambda - 1} \bigg|_{\lambda = 1} \ge 0,$$

since, if G - e and/or G/e are 2-connected, it follows from the induction hy-

pothesis, otherwise the left-hand side is 0.

Since for  $|V(G)| \leq 3$  the assertion is trivially true, we may assume that  $|V(G)| \geq 4$ . We are going to show that one of G-e, G/e is 2-connected. Suppose G/e is separable; clearly only the image of e can be a cutpoint of it, whence the two endpoints x, y of e form a cutset. Let  $P_1$ ,  $P_2$  be (x, y)-paths through two components of G-x-y. Then  $P_1 \cup P_2$  is a circuit and e is a chord of it. But as in 6.35, this implies G-e is 2-connected.

(b) Let  $\sigma(G)$  denote the number of acyclic orientations of G. Consider an acyclic orientation of G-e. This can be extended to an acyclic orientation of G in one or two ways according as it is an acyclic orientation of G/e. Thus

$$\sigma(G) = \sigma(G - e) + \sigma(G/e).$$

Hence the result follows by induction and 9.38 [R. Stanley, *Discrete Math.* 5 (1973) 171–178].

**47**. (a) By 9.38,

$$\begin{split} P_{G_1}(\lambda) &= P_{G_1-e_1}(\lambda) - P_{G_1/e_1}(\lambda) = P_G(\lambda) - P_{G_1/e_1}(\lambda), \\ P_G(\lambda) &= P_{G_1}(\lambda) + P_{G_1/e_1}(\lambda), \end{split}$$

and similarly

$$P_G(\lambda) = P_{G_2}(\lambda) + P_{G_2/e_2}(\lambda);$$

thus,

$$P_{G_1}(\lambda) + P_{G_2/e_2}(\lambda) = P_{G_2}(\lambda) + P_{G_2/e_2}(\lambda).$$

(b) Consider the three graphs G in Fig. 67, if there is a linear relation, then it is satisfied by them. So

$$a\lambda(\lambda - 1)(\lambda^2 - 3\lambda + 3) + b\lambda(\lambda - 1)(\lambda - 2)^2 + c\lambda(\lambda - 1)(\lambda - 2)^2 = 0,$$

$$a\lambda(\lambda - 1)(\lambda - 2)^2 + b\lambda(\lambda - 1)(\lambda - 2)^2 + c\lambda(\lambda - 1)(\lambda - 2)(\lambda - 3) = 0,$$

$$a\lambda(\lambda - 1)(\lambda - 2)^2 + b\lambda(\lambda - 1)(\lambda - 2)(\lambda - 3) + c\lambda(\lambda - 1)(\lambda - 2)^2 = 0.$$

If this system of equations has a non-trivial solution, its determinant is 0. But its determinant is

$$\lambda^{3}(\lambda-1)^{3}(\lambda-2)^{2}\begin{vmatrix}\lambda^{2}-3\lambda+3&\lambda-2&\lambda-2\\(\lambda-2)^{2}&\lambda-2&\lambda-3\\(\lambda-2)^{2}&\lambda-3&\lambda-2\end{vmatrix}==\lambda^{3}(\lambda-1)^{3}(\lambda-2)^{2}(\lambda^{2}-3\lambda+1).$$

So if  $\lambda \neq 0, 1, 2, \frac{3 \pm \sqrt{5}}{2}$ , there is no linear relation (in particular, there is none, which would hold for every  $\lambda$ ). For  $\lambda = 0, 1, 2$ , we have trivial relations like  $P_{G_i}(\lambda) = 0$ .




Fig. 67

(c) We use induction on |E(G)|. If V(G) = V(F), we have only three graphs to check; those in Fig. 67 (since multiple edges can be ignored and if G has a loop, everything is 0). Fortunately, there are only four numbers to cal-

culate, namely setting  $\xi = \tau + 1 = \frac{3 + \sqrt{5}}{2}$ , we have

$$\begin{split} \xi(\xi-1)(\xi^2-3\xi+3) &= (\xi^2-\xi)\cdot 2 = 4\xi-2, \\ \xi(\xi-1)(\xi-2)^2 &= (\xi^2-\xi)(\xi^2-4\xi+4) = (2\xi-1)(3-\xi) = \xi-1, \\ \xi(\xi-1)(\xi-2)(\xi-3) &= \xi(\xi-3)\cdot (\xi-1)(\xi-2) = \\ &= (\xi^2-3\xi)(\xi^2-3\xi+2) = -1 \end{split}$$

(we have made use of the fact throughout that  $\xi^2 = 3\xi - 1$ ). With these values, and the chromatic polynomials determined in the last solution, it is easy to check that the relation holds as stated.

Now suppose that  $V(G) \neq V(F)$ . If there are only isolated points in G besides the points of F, they only mean that the equations already verified are multiplied by a power of  $\xi$ . So suppose there is an edge e with at most one endpoint on F. Set G' = G - e,  $G'_1 = G' + e_1$ ,  $G'_2 = G' + e_2$  and similarly, G'' = G/e,  $G''_1 = G'' + e_1$ ,  $G''_2 = G'' + e_2$ . Then by the induction hypothesis,

$$egin{align} P_{G'}(\xi) &= \xi ig( P_{G'_1}(\xi) + P_{G'_2}(\xi) ig), \ P_{G''}(\xi) &= \xi ig( P_{G''_1}(\xi) + P_{G''_2}(\xi) ig). \ \end{array}$$

Subtracting we get the desired equality by 9.38.

- 48. Let  $q(x) = x^2 3x + 1$ ; the roots of q(x) are  $\xi = \frac{3 + \sqrt{5}}{2}$  and  $\xi' = \frac{3 \sqrt{5}}{2}$ . Then  $q \nmid P_G$ , because by 9.45,  $\xi'$  is not the root of any chromatic polynomial. Since q is irreducible over the rationals, this implies it has no root in common with  $P_G$ , i.e.  $P_G(\xi) \neq 0$ .
- 49. (a) We use the notations of 9.47. By 9.38 and 9.47c,

$$P_{G_1}(\tau+1) + P_{G_2}(\tau+1) + P_{G_1/e_1}(\tau+1) + P_{G_2/e_2}(\tau+1) =$$
  
=  $2(\tau+1) \left( P_{G_1}(\tau+1) + P_{G_2}(\tau+1) \right),$ 

whence

(1) 
$$P_{G_1/e_1}(\tau+1) + P_{G_2/e_2}(\tau+1) = (2\tau+1) (P_{G_1}(\tau+1) + P_{G_2}(\tau+1)) =$$
  
=  $\tau^3 (P_{G_1}(\tau+1) + P_{G_2}(\tau+1)).$ 

Also by 9.47a,

(2) 
$$P_{G_1/e_1}(\tau+1) - P_{G_2/e_2}(\tau+1) = P_{G_1}(\tau+1) - P_{G_2}(\tau+1).$$

and multiplying (1) and (2) together we obtain

(3) 
$$P_{G_1/e_1}^2(\tau+1) - P_{G_2/e_2}^2(\tau+1) = \tau^3(P_{G_1}^2(\tau+1) - P_{G_2}^2(\tau+1)).$$

Now suppose that the maps  $G_2$ ,  $G_1/e_1$ ,  $G_2/e_2$  satisfy the identity given in the problem. Then (3) yields upon multiplying by  $(\tau + 2)\tau^{3n-13}$ ,

$$P_{G_1/e_1}(\tau+2) - P_{G_2/e_2}(\tau+2) = (\tau+2)\tau^{3n-10}P_{G_1}^2(\tau+1) - P_{G_2}(\tau+2).$$

By 9.47a, this implies  $G_1$  also satisfies the identity.

Now let  $G_1$  be a minimal triangulation, which does not satisfy the identity and suppose  $G_1$  has a point x of degree as large as possible. Let (x, y, z) be any triangle adjacent to x, and let (x', y, z) be the second triangle with side (y, z). Then x' = x. In fact, if  $x' \neq x$ , then letting  $G = G_1 - (y, z)$  and  $G_2 = G + (x, x')$ ,  $G_1/(y, z_j)$ ,  $G_2/(x, x')$  and  $G_2$  will satisfy the identity by the minimality assumption on  $G_1$  and the maximality assumption on  $d_{G_1}(x)$ . Hence by the above,  $G_1$  also satisfies the identity, a contradiction.

The above argument shows that, if a face is adjacent to x, then so are all the neighboring faces. Hence all faces are adjacent to x. This implies that x is adjacent to all other points and G-x has no circuits. Since a triangulation is always 2-connected (if x were a cut-point, then consider a face which meets two components of G-x; the fact that this face is a triangle implies that there is an edge connecting two components of G-x, a contradiction), G-x is a tree.

Now if we want to  $\lambda$ -color  $\overline{G}$ , we can give x any of the  $\lambda$  colors and  $(\lambda - 1)$ -color the remaining points. Hence

$$P_G(\lambda) = \lambda P_{G-x}(\lambda - 1) = \lambda(\lambda - 1)(\lambda - 2)^{n-2}$$

and

$$P_G(\tau + 2) = (\tau + 2)(\tau + 1)\tau^{n-2} = (\tau + 2)\tau^n$$

as  $\tau + 1 = \tau^2$ . Also,

$$\begin{aligned} (\tau+2)\tau^{3n-10}P_G^2(\tau+1) &= (\tau+2)\,\tau^{3n-10}(\tau+1)^2\,\tau^2(\tau-1)^{2(n-2)} = \\ &= (\tau+2)\,\tau^{3n-10}\,\tau^4\,\tau^2\,\tau^{-2(n-2)} = (\tau+2)\tau^n \end{aligned}$$

as  $\tau - 1 = \tau^{-1}$ . Thus G satisfies the identity, a contradiction.

(b) We use induction on the number of points and "backwards" induction on the number of edges; we may suppose there are no 2-gonal faces. If we have a triangulation, (a) implies the assertion. Otherwise, we have a face with more than 3 points. Put in a diagonal e into this face. Then we have, by the induction hypothesis,

$$P_{(G+e)/e}(\tau+2) > 0,$$
  
 $P_{G+e}(\tau+2) > 0.$ 

9.50

Now by 9.38,

$$P_G(\tau+2) = P_{(G+e)/e}(\tau+2) + P_{G+e}(\tau+2) > 0.$$

**50.** We use induction on |V(G)|. We may assume G is simple. By 5.25a, the number of edges of G is at most 3 |V(G)| - 6, therefore G has a point x of degree at most 5. If the degree of x is at most 4, we can remove it, 5-color the rest and extend this coloration to x. So suppose d(x) = 5. x has two neighbors, which are non-adjacent, for otherwise we would get a  $K_6$  in the graph. Let y, z be two such neighbors of x. Contract the edges (x, y) and (x, z) and consider the resulting graph G. This has no loops, thus by the induction hypothesis, it is 5-colorable. Let us consider the corresponding 5-coloration of G. This is not legitimate, because x, y and z have the same color; but only the edges (x, y) and (x, z) are monochromatic. Now note that x has only 3 other neighbors, so we can find a color, which is different from the colors of all neighbors of x, and re-color x with this color.

51. Suppose first that the faces are 4-colorable; then it is easy to check that the

3-coloration of the edges defined in the hint is a good one.

Secondly, let us consider a 3-coloration of the edges. Let  $G_1$  be the map formed by red and blue edges and  $G_2$  the map formed by red and green edges. Clearly,  $G_1$  and  $G_2$  consists of disjoint circuits and hence, the faces of  $G_1$  can be 2-colored with red and green; also the faces of  $G_2$  can be 2-colored with "light" and "dark". Now use "light red", "light green", "dark red" and "dark green" to color the faces of  $G_2$  each face F of G is contained in a face  $F_1$  of  $G_1$  and in a face  $F_2$  of  $G_2$  and color it "light red", if  $F_1$  is red and  $F_2$  is light etc. This defines a good coloration of the faces of G.

Remark: One could also prove the second half by using 5.4 [P.G. Tait; see S, OF, Wi].

52. If the faces are 4-colorable the edges are 3-colorable by 9.51; let 1, 2, 3 be their colors. Consider the assignment described in the hint. Go around the boundary circuit of a face F counterclockwise, and consider the "clock" in Fig. 68. Whenever we pass through a point with value +1, the color of the next edge will be the next mark clockwise; if we pass through a point with -1, the color of the next edge is the next mark counterclockwise. Since going around the whole circuit we get back to the same color we started with, the sum of +1's and -1's is divisible by 3.

Conversely, suppose we have an assignment of +1's and -1's as required. Consider L(G), this is also planar and has two kinds of faces: For each  $x \in V(G)$ 




Fig. 68




Fig. 69

it has a triangular face  $F_x$  surrounding x and for each face F of G, it has a face F', which lies in F and has the same number of edges (Fig. 69).

Orient L(G) by orienting the triangles  $F_x$  clockwise, and define the work v(e) on an edge  $e \in E(L(G))$  as the value assigned to the corresponding point of G.

So the three edges of a triangular face of L(G) corresponding to a point of G have the same work. Thus, the sum of work of edges around any face is  $\equiv 0 \pmod{3}$ . But, then this holds for any circuit C. For go around each face inside this circuit clockwise and write up the sum of the work. This is 0. On the other hand, the sum of these sums is exactly the work around C (clockwise), because the inside edges are counted twice, and in fact, in different direction. So the work needed to go around C is 0. By 5.4, we can find a potential p(x) of V(L(G)) such that for each edge (x, y),  $v(x, y) \equiv p(y) - p(x) \pmod{3}$ . Obviously, we may assume p takes only the values 0, 1, 2. Then p(x) is a good 3-coloration of L(G). [P. J. Heawood, see S. OF.]

53. The faces inside the Hamiltonian circuit C can be 2-colored. To show this, note that the corresponding piece of the dual graph  $G^*$  contains no circuit, for such a circuit would isolate a point of G from C, which is clearly impossible. Thus, the piece of  $G^*$  inside C is a forest, and this is 2-colorable.

Similarly, 2 other colors suffice to color the faces outside C.

**54.** (a) We use induction on |V(G)|. If this number is at most 4 the assertion is obvious.

Let  $d_G(x) \leq 4$ . By the induction hypothesis, G - x has a 4-coloration  $\alpha$ . If  $d_G(x) \leq 3$ , this 4-coloration can obviously be extended to x. So suppose  $d_G(x) = 4$ . Let y, z, u, v be the neighbors of x. We think of G as embedded in the plane and assume the edges (x, y), (x, z), (x, u), (x, v) leave x in this cyclic order.

If one of the colors does not occur among  $\alpha(y)$ ,  $\alpha(z)$ ,  $\alpha(u)$ ,  $\alpha(v)$ , we can color x with this color. So suppose  $\alpha(y) = 1$ ,  $\alpha(z) = 2$ ,  $\alpha(u) = 3$ ,  $\alpha(v) = 4$ .

Consider the subgraph spanned by colors 1 and 3 and the component  $G_1$  of it containing y. If  $u \notin V(G_1)$ , we can exchange the colors of points of  $G_1$  and get a 4-coloration of G-x, where 1 does not occur among the colors of y, z, u, v. Then, as before, x can get color 1. So we may suppose  $u \in V(G_1)$  i.e. there is a (y, u)-path  $P_1$  whose points have only colors 1 and 3.




Fig. 70

Similarly, we may assume that there is a (z, v)-path  $P_2$  having only colors 2 and 4. But since G is planar,  $P_1$  and  $P_2$  cross, and their common point cannot have any of the colors 1, 2, 3, 4, a contradiction (Fig. 70) [G. Dirac; see S, OF. This method is called  $Kempe\ chaining$ ].

(b) Suppose first that all faces are triangles. Then it is easy to see that we must have the graph of the icosahedron which, in fact, admits a 4-coloration (Fig. 71).




Fig. 71

We may assume the graph is 3-connected, this follows by an argument similar to the one in the solution of  $9.13b^{\dagger}$ . If F is a non-triangular face, then F has two points x, y which are at distance 2 on the boundary C of F; but, then x, y are not adjacent in G since otherwise, they would separate the two arcs of C connecting them.

† Note the analogy between 9.13a-b and this problem,

Identify x and y, and cancel the arising multiplicities of edges. The resulting graph G' has at least one point of degree at most 4: this is the common neighbor of x and y. It has degrees at most 5 except that the image x' of x (and y) has

higher degree.

If G' is 4-colorable clearly so is G. So suppose G' is not 4-colorable and consider a critically 5-chromatic subgraph G'' of it. The argument in part (a) shows that G'' can have no point of degree at most 4. Hence G'' is a proper subgraph of G' and, moreover, all edges of G' incident with G'' must be incident with X'; otherwise, their endpoints in G'' would have degree at most 4. But, then X' is a cutting point which is impossible, because as G is 3-connected, G' is 2-connected [J.M. Aarts and J. de Groot; see S].

55. Clearly, we may assume that G is 2-connected; otherwise, add another edge

joining two blocks of G, not parallel to the old edges.

Consider a face F of G, which is not a triangle, and let  $C = (x_1, \ldots, x_k)$  be its boundary circuit. Either  $(x_1, x_3)$  or  $(x_2, x_4)$  is not an edge of G, for otherwise they ought to cross each other outside C. Say  $(x_1, x_3) \notin E(G)$ . Then add  $(x_1, x_3)$  to G; we have broken F into two smaller faces this way. Carrying on, we can extend G into a simple triangulation. If this triangulation can be colored in the desired way, i.e. so that no face of the triangulation is monochromatic, then no face of G is monochromatic.

Thus, we may assume G is a triangulation. Then  $G^*$  is 3-regular and since

G is simple,  $G^*$  is 2-edge-connected.

By Petersen's theorem (7.29),  $G^*$  has a 1-factor F. Let  $\Phi$  be the set of corresponding edges of G. Then  $\Phi$  contains exactly one edge from each face of G. Therefore, the faces of  $G - \Phi$  are quadrilaterals, i.e. by the dual of 5.26,  $G - \Phi$  is bipartite. Consider the 2-coloration of  $G - \Phi$ ; this 2-coloration meets the requirements for G [M. Schäuble, Beiträge zur Graphentheorie, Teubner, Leipzig (1968) 137–142].

56. Suppose every point has even degree. Let us 2-color the faces (5.26) with red and blue and orient the edges, so that they have red on their right side. Let C be any circuit. Count the number of edges on and inside C in two different ways. Let  $\alpha$  edges of C have red face incident from the inside and  $\beta$  edges from the outside. Let, moreover, k red and k blue faces lie inside k. Then the number of edges on and inside k is

$$3k + \beta$$
,

and also

$$3l + \alpha$$
.

Therefore, the work needed to go around C counterclockwise is

$$\alpha-\beta=3k-3l\equiv 0\ (\mathrm{mod}\ 3).$$

This implies, by 5.4, that there is potential p(x) defined on V(G) such that if  $(x, y) \in E(G)$ , then  $p(y) - p(x) \equiv 1 \pmod{3}$ . This p(x) yields a 3-coloration of G [H. Whitney; OF, Wi].

57. Let  $(x_1, y_1), \ldots, (x_n, y_n)$  be the intersection points. We may suppose the direction of axes is chosen so that  $x_1, \ldots, x_n$  are different, say  $x_1 < x_2 < \ldots$ 

...  $< x_n$ . Now 3-color the points in the above order. When we get to  $(x_i, y_i)$  at most two neighbors of it have previously been colored and so, we can find a color for it. Thus we can 3-color the graph [H. Sachs].

## § 10. Extremal problems for graphs

1. We may assume that G is 2-connected. For if  $G = G_1 \cup G_2$  with  $|V(G_1) \cap V(G_2)| \leq 1$ ,  $|V(G_i)| \geq 2$ , then one of  $G_1$ ,  $G_2$  satisfies

$$|E(G_i)| > \frac{3}{2} (V|(G_i)| - 1),$$

and we can use induction on n.

So suppose that G is 2-connected. Clearly, it is not a single circuit, so by 6.33,  $G = P_0 \cup P_1 \cup \ldots \cup P_k$   $(k \ge 1)$ , where  $P_0$  is a circuit and  $P_{i+1}$  is a  $(P_0 \cup \ldots \cup P_i, P_0 \cup \ldots \cup P_i)$ -path for  $i = 0, \ldots, k-1$ . Then  $P_0 \cup P_1$  is a  $\Theta$ -graph.

The sharpness of the result is shown by any graph whose blocks are triangles.

2. (a) Let  $P = (x_0, x_1, \ldots, x_p)$  be a longest path in G.  $x_0$  is adjacent to two points other than  $x_1$ ; since P is maximal these two points must lie on P; let, say,  $x_0$  be adjacent to  $x_i$  and  $x_j$ ,  $2 \le i < j \le p$ . Then  $(x_0, \ldots, x_j)$  is a circuit in G which has the chord  $(x_0, x_i)$  [J. Czipszer].

(b) We use induction on n. The assertion is clear for n = 4. If G has degrees


at least 3, then it contains a circuit with a chord by (a).


Thus we may assume G has a point x with degree at most 2. Then G - x contains at least 2n - 3 - 2 = 2(n - 1) - 3 edges and thus, by the induction hypothesis, G - x contains a circuit with a chord. But, then so does G.

 $K_{2,n-2}$  shows that the result of (b) is sharp [L. Pósa].

- 3. (a) We prove by induction on |V(G)| the assertion of the hint.
- I. Assume first that G has a point x of degree at most 2. If x has degree 1, then G-x has at most one point of degree at most 2 and we are done by induction. If x has degree 2, i.e. it is adjacent to two points y, z and  $(y, z) \notin E(G)$ , then remove x and join y to z. The resulting graph has degrees at least 3 and thus, it contains a subdivision of  $K_4$ . If we subdivide (y, z) by x, we get G back; hence G contains a subdivision of  $K_4$  too. If y, z are adjacent, then we may assume  $d_G(y) = d_G(z) = 3$  (otherwise, G x satisfies the assumption). If the third neighbors of y and z are the same point w, then G x y z w satisfies the assumption; if y is adjacent to u, z is adjacent to v(x, y, z, u, v) being distinct) then contracting x, y, z to a single point x', we get a graph G' in which only x' has degree at most 2. Hence G' contains a subdivision of  $K_4$ . Since subdividing the edge (x', u) of G' we get a subgraph of G, G also contains a subdivision of  $K_4$  (see Fig. 72).

II. Now assume that every point of G has degree at least 3. We may assume G has a point x of degree 3, otherwise we could remove edges as long as all but one of the points of G have degree at least 3. Let u, v, w be the points adjacent to x. If u, v, w are adjacent by pairs, then x, u, v, w spans a  $K_4$ . Thus we may sup-





pose  $(u, v) \notin E(G)$ . Remove x and join u to v. The resulting graph G' has degrees at least 3 except possibly that w has degree 2. Thus G' contains a subdivision of  $K_4$  by induction. Since subdividing the edge (u, v) by x we get a subgraph of G, G also contains a subdivision of  $K_4$ .

(b) If every point of G has degree at least 3, we are done by (a). If there is

a point of degree 2, we can remove it and proceed by induction.

The graph shown in Fig. 73 has exactly two points of degree 2 and 2n-3 edges, it does not contain a subdivision of  $K_4$  (prove!). Thus (a) and (b) are sharp [G. Dirac, *Math. Nachr.* 22 (1960) 61–85].

4. First we prove, as suggested in the hint, that G does not contain a subdivision H of  $K_4$  and an edge (u, v) disjoint from it. Suppose it does. Consider the component  $G_1$  of G - V(H) containing (u, v). Trivially  $G_1$  is a tree. Let x, y be




Fig. 74

two endpoints of  $G_1$ . Then x (respectively y) must have two neighbors  $p_1, p_2$  (respectively  $q_1, q_2$ ) in H. No two of  $p_1, p_2, q_1, q_2$  can be contained in the "subdivided star" of one of the principal points of H as center or inner point of the "rays", for such a position would yield two disjoint circuits (Fig. 74a). But then, as is easily verified,  $p_1, p_2, q_1, q_2$  must be the principal points of H and we again get two disjoint circuits as shown in Fig. 74b.

Now we have to distinguish 5 cases.

Case 1. Assume there is in G a subdivision H of  $K_4$  and a point v of degree at least 4 not in H. A similar argument to the one above shows that v has degree exactly 4 and is adjacent to the four principal points  $x_1, x_2, x_3, x_4$  of H. Exchanging the role of  $x_i$  and v we can use the assertion of the hint and see that  $x_i$  must be adjacent to  $x_j$   $(j \neq i)$  and v, and to no other point. Hence  $G = K_5$ .

Case 2. Assume G contains a subdivision H of  $K_4$  and a point v of degree 3 not in H but connected to three principal points  $x_1$ ,  $x_2$ ,  $x_3$  of H. Interchanging the role of v and the fourth principal point  $x_4$  we get that  $\Gamma(x_4) = \{x_1, x_2, x_3\}$ . We show that every edge (r, s) of G contains one of  $x_1, x_2, x_3$ . Suppose not, then  $r, s \neq x_1, x_2, x_3, x_4, v$ . But  $x_1, x_2, x_3, x_4$  and v, together with any one of the subdivided edges  $(x_1, x_2)$ ,  $(x_2, x_3)$ ,  $(x_1, x_3)$ , form a subdivision of  $K_4$ . Hence (r, s) must meet each of these subdivided edges at inner points, which is impossible. Thus  $x_1, x_2, x_3$  in fact cover all edges. But this clearly means we have graph (iii).

Case 3. There is a subdivision H of  $K_4$  and a point v not in H of degree 3, connected to at most two principal points of H. By the assertion of the hint, the neighbors of v must be in H and it is easy to see that at least two of them,  $x_1$  and  $x_2$  say, must be principal points. The third one, being not principal, must be an inner point p of the subdivided  $(x_3, x_4)$ -edge. Interchanging the roles of  $x_3$  and v we get  $\Gamma(x_3) = \{x_1, x_2, p\}$  and similarly,  $\Gamma(x_4) = \{x_1, x_2, p\}$ . We claim that  $x_1, x_2, p$  cover all edges; then we shall be able to conclude as before. Since  $x_1, x_2, x_3, x_4, p$ , v induce a subdivision of  $K_4$ , each edge must meet one of these points. But those meeting  $x_3, x_4$  or v also meet  $x_1, x_2$  or p. This settles this case.

Case 4. So we may assume each subdivision of  $K_4$  is a spanning subgraph. By 10.3a, G contains a subdivision H of  $K_4$ . Let  $x_1, x_2, x_3, x_4$  be its principal points. Assume that this subdivided  $K_4$  can be chosen so that  $x_1$  has degree at least 4. Let  $(x_1, u)$  be an edge adjacent to  $x_1$  but not in H. Since H is a spanning subgraph, u is a point of H; trivially u must lie on the circuit C of  $H - x_1$ . Replacing the subdivided edge  $(x_1, x_i)$  (i = 2, 3, 4) by  $(x_1, u)$ , we must get spanning subdivided  $K_4$ 's. Hence  $x_1$  is adjacent to  $x_2, x_3$  and  $x_4$ . Any point of C can, trivially, be adjacent to  $x_1$  only (except for its neighbors on C). Thus G is a wheel.

Case 5. Only one more case is left, when each subdivided  $K_4$  in G is a spanning subgraph and its principal points have degree 3 in G. If  $G = K_4$ , then it is a wheel. So suppose H has a chord e, which must then connect inner points of two subdivided edges of H. Clearly these subdivided edges must be disjoint, e.g.  $(x_1, x_2)$  and  $(x_3, x_4)$ . Now G contains no further point since removing such a point the rest of H + e would still contain a subdivision of  $K_4$ , contrary to the assumption. Hence G has six points, a trivial case.

Remark: By similar arguments we could determine all (not necessarily simple) graphs, which contain no two disjoint circuits. Given any such graph, we can remove points of degree one and "smooth out" points of degree 2, so it suffices to describe these graphs with degrees at least 3. The result is as follows: the three types of graphs in the problem with the modification that the spokes of the wheel and the edges in the 3-element class of  $K_{3,\,n-3}$  may be multiple; and there is one new class: take a forest F, and a point x with possibly some loops at x, and connect x to F arbitrarily [G.A. Dirac, Canad. Math. Bull. 8 (1965) 459–463; L. Lovász, Mat. Lapok 16 (1965) 289–299].

- 5. (a) By 10.3b, G contains a subdivision of  $K_4$ . If this subdivision is proper, i.e. at least one of the subdivided edges has an inner point we easily find a subdivision of  $K_{2,3}$ . So we may suppose G contains four pairwise adjacent points x, y, z, u. Since  $n \geq 5$ , there is at least one more point v. Since G is 2-connected there are two independent paths connecting v to two distinct points of x, y, z, u. Now x, y, z, u, v are principal points of a subdivision of  $K_{2,3}$  (Fig. 75).
- (b) Since  $|E(G)| \geq 3n-5$ , 5.26 implies that G is non-planar and hence, by Kuratowski's theorem 5.39d, G contains a subdivision of  $K_{3,3}$  or  $K_5$ . In the first case we are done, so suppose G contains a subdivision of  $K_5$  with principal points  $x_1, \ldots, x_5$ . As in part (a), we distinguish whether or not this subdivision is proper.
- 1° Suppose the subdivided  $(x_1, x_2)$ -edge Q (say) contains inner points.  $G \{x_1, x_2\}$  being connected, we have a path P joining an inner point y of Q





Fig. 75







Fig. 76

to a point z outside Q. z may have three essentially different positions but in either case we easily find a subdivision of  $K_{3,3}$  easily (see Fig. 76, p. 379).

 $2^{\circ}$ . Suppose  $x_1, \ldots, x_5$  are mutually adjacent. Since  $n \geq 6$ , we must have a further point v. Since G is 3-connected, there are 3 independent paths connecting v to  $x_1, x_2, x_3$  (say). Then again a subdivision of  $K_{3,3}$  is obviously found (Fig. 77).



(c) We claim that if a graph has at least 2n-2 edges and contains no subdivision of  $K_{3,2}$ , then it is connected and its blocks are  $K_4$ 's. We use induction on n.

Suppose first that a graph G with 2n-2 edges is not connected, i.e.  $G=G_1 \cup G_2$ ,  $G_1 \cap G_2 = \emptyset$ . Let  $|V(G_i)| = n_i > 0$  (i = 1, 2). Then

$$|E(G_1)| + |E(G_2)| \ge 2n - 2 = (2n_1 - 2) + (2n_2 - 2) + 2,$$

whence, e.g.

$$|E(G_1)| \geq 2n_1 - 1.$$

Thus by induction,  $G_1$  is connected and all blocks of it are  $K'_4$ s. But, then  $|E(G_1)| = 2n_1 - 2$ , a contradiction. So G is connected.

If  $\tilde{G}$  is 2-connected the assertion follows from (a). So suppose it has a cut-point x. Let  $G = G_1 \cup G_2$ ,  $V(G_1 \cap G_2) = \{x\}$ ,  $|V(G_i)| = n_i < n$ . Then

$$|E(G_1)| + |E(G_2)| \ge 2n - 2 = (2n_1 - 2) + (2n_2 - 2).$$

As above, we get a contradiction, if

$$|E(G_i)| > 2n_i - 2$$

for any i. Thus it follows that

$$|E(G_1)| = 2n_1 - 2, |E(G_2)| = 2n_2 - 2$$

and hence by induction hypothesis,  $G_1$ ,  $G_2$  are connected graphs whose blocks are  $K_4$ 's. Thus so is G.

A similar argument shows that, if G is a graph with n points and 3n-5 edges containing no subdivision of  $K_{3,3}$ , then G is 2-connected and it is composed of complete 5-graphs by the following recursive rule. We take a graph G (already constructed) and a complete 5-graph  $K_5$ , and identify an edge of G with an edge of G (see Fig. 79a).

Thus the answer to (c) is: A graph with n points and

2n - 1 edges if  $n \equiv 1 \pmod{3}$ ,


2n - 2 edges if  $n \equiv 0.2 \pmod{3}$ 


contains a subdivision of  $K_{2,3}$ ; a graph with n points and

3n - 4 edges if  $n \equiv 2 \pmod{3}$ ,

3n - 5 edges if  $n \equiv 0.1 \pmod{3}$ 


contains a subdivision of  $K_{3,3}$ . These bounds are best possible as shown by the graphs in Figures 78 and 79 [L. Pósa].






(b) any triangulation of a circuit

Fig. 78





(b) any planar triangulation

Fig. 79

- **6.** We show: (a') If  $G_0$  has degrees at most 3, then any graph contractible onto  $G_0$  contains a subdivision of  $G_0$ ; (b') If  $G_0$  has a point with degree at least 4 this is no longer true.
- (a') It suffices to show on the one hand that, if G/e ( $e = (u, v) \in E(G)$ ) contains a subdivision H of  $G_0$ , then so does G. If the point x of G/e, which is the

image of e is not a principal point of H, then this is clear; suppose it is a principal point. By the assumption, there are at most 3 subdivided edges of H starting from x. If we pull u and v apart again, some of these will start from u and some will start from v. We may assume at most one starts from v; then adding e to this subdivided edge and considering u as a principal point we find a subdivision of  $G_0$  in G.

(b') We show that there is a graph  $G_1$ , contractible onto  $G_0$ , with degrees at most 3. Let x be a point of  $G_0$  with degree at least 4. Let us split x into d points  $x_1, \ldots, x_d$  of degree 1 and connect these by a circuit. Do so for each point x with degree at least 4, and denote the resulting graph by  $G_1$ . Then  $G_1$  has degrees at most 3 and is contractible onto  $G_0$ . If  $G_0$  has a point with degree at least 4, then, clearly,  $G_1$  contains no subdivision of  $G_0$ .

A 4-connected counterexample in the case of  $K_5$  is  $K_{4.4}$ .

- 7. Let  $a \in V(G)$ . Contract edges and cancel the arising multiplicities repeatedly so that
  - (\*) the contracted edge is adjacent to the image of a,

$$(**) \frac{|E(G')|}{|V(G')|} \ge \frac{m}{n}$$
 holds for the resulting graphs.

Suppose, we get stuck with the graph  $G_0$ . Let  $a_0$  be the image of a in  $G_0$ . It follows from (\*) above that the subgraph  $G_1$  of G mapped onto  $a_0$  is connected and  $G - V(G_1) = G_0 - a_0$ .

Let x be any point adjacent to  $a_0$  in  $G_0$  (i.e. adjacent to  $V(G_1)$  in G). Let  $\vartheta$  denote the number of points of  $G_0$  adjacent to  $a_0$  and x (this is the degree of x in the subgraph of G induced by the neighbors of  $G_1$ ). Try to contract  $(a_0, x)$  and cancel one edge from each of the arising  $\vartheta$  parallel pairs. Then (\*\*) must fail to hold for the resulting graph  $G_0$ . Since

$$\begin{split} |V(G_0')| &= |V(G_0)| - 1, \\ |E(G_0')| &= |E(G_0)| - 1 - \vartheta, \end{split}$$

we have

$$\frac{|E(G_0)|-\vartheta-1}{|V(G_0)|-1}<\frac{m}{n}$$

or, equivalently,

$$\vartheta > |E(G_0)| - 1 - \frac{m}{n} (|V(G_0)| - 1).$$

Since

$$\frac{|E(G_0)|}{|V(G_0)|} \ge \frac{m}{n},$$

we have

$$\vartheta > \frac{m}{n} |V(G_0)| - 1 - \frac{m}{n} (|V(G_0)| - 1) = \frac{m}{n} - 1,$$

which proves the assertion [W. Mader; see W].

8. We use induction on m. For m=3 the assertion is clear. Let  $m\geq 4$ . By the preceding problem, G contains a connected subgraph  $G_1$  such that the subgraph  $G_2$  induced by the neighbors of  $G_1$  has degrees  $> \frac{|E(G)|}{n} - 1 \geq 2^{m-3} - 1$ , so  $\geq 2^{m-3}$ . Hence

$$|E(G_2)| \geq 2^{m-4} V(G_2),$$

and thus, by the induction hypothesis,  $G_2$  can be contracted onto  $K_{m-1}$ . Carrying out the same contraction on G and also contracting  $G_1$ , we obtain a graph which contains  $K_m$  [ibid.].

**9.** Induction on k: for k = 1 the assertion is trivial.

Again, let  $G_1$  be a connected subgraph as in 10.7, i.e. let the subgraph  $G_2$  induced by the neighbors of  $G_1$  have degrees  $\geq 2^k$ . Then

$$|E(G_2)| \ge 2^{k-1} |V(G_2)|$$

and thus,  $G_2$  contains a subdivision H of F-e for any  $e \in E(F)$ . Let  $x_1, x_2$  be the principal points of H corresponding to the endpoints of e. Since  $x_i \in V(G_2)$ , it is adjacent to some point  $y_i \in V(G_1)$  (i = 1, 2). Since  $G_1$  is connected it contains a  $(y_1, y_2)$ -path P. Then

$$H' = H + (x_1, y_1) + (x_2, y_2) + P$$

is a subdivision of F contained in G. [ibid.]

10. I. Let  $x_1$  be a point of the 3-regular graph G with girth 4, and let  $x_2$ ,  $x_3$ ,  $x_4$  be its neighbors. Then  $x_2$ ,  $x_3$ ,  $x_4$  must be independent, because G contains no triangle. Let  $x_5$ ,  $x_6$  be two further neighbors of  $x_2$ . Thus  $|V(G)| \ge 6$ ; equality holds only if  $x_3$ ,  $x_4$  are also adjacent to  $x_5$  and  $x_6$ , i.e.  $G = K_{3,3}$ .

II. Let  $(x_1, \ldots, x_p)$  be the least circuit in  $G, p \geq 5$ . Let  $y_i$  be the third neighbor of  $x_i$ . By the minimality of the circuit considered,  $y_i$  does not lie on it and  $y_i \neq y_j$  for  $i \neq j$ . Hence  $|V(G)| \geq 2p \geq 10$ . If equality holds, then p = 5. The only points at distance at least 4 from  $y_1$  are  $y_3$  and  $y_4$ , so these must be the two further neighbors of  $y_1$  besides  $x_1$ . Similarly,  $y_2$  is adjacent to  $y_4$  and  $y_5$ , and  $y_3$  to  $y_5$ , i.e. we get the Petersen graph (Fig. 80).




Fig. 80

11. I. Assume first that g is odd. Let  $x_0 \in V(G)$ , and denote by  $S_i$  the set of points at distance i from  $x_0 \left(i=0,\ldots,\frac{g-1}{2}\right)$ . From each point of  $S_i$  there is exactly one edge to  $S_{i-1}$ , since two such edges would yield two paths from x to  $x_0$  of length i, and these would form a circuit of length less than g. Thus  $|S_{i+1}| = (r-1) \cdot |S_i| \left(i=1,\ldots,\frac{g-3}{2}\right)$ , and hence

$$|V(G)| \geq |S_0| + |S_1| + \dots + |S_{rac{g-3}{2}}| = 1 + r + r(r-1) + \dots + r(r-1)^{rac{g-3}{2}}.$$

II. Let g be even and consider two adjacent points x, y. Denote by  $S_i$  the set of points at distance i from the set  $\{x, y\}$   $(i = 1, ..., \frac{g}{2} - 1)$ . The result follows by the same counting as above [W.T. Tutte; P. Erdős-H. Sachs; see S].

12. Assume we have constructed G(r, g) and G(r', g-1), where r' = |V(G(r, g))|. Split each point of G(r', g-1) into r' points of degree one and identify these r' points with the points of a copy of G(r, g). The resulting graph G' is, obviously, (r+1)-regular. We claim that it has girth g. If we consider a minimum circuit in a copy of G(r, g), this has length g; hence G' has girth at most g. Any other circuit in copies of G(r, g) has length at least g. Let G' be a circuit of length g not in the copies of G(r, g). Contract each copy of G(r, g), then G' is mapped onto G(r', g-1). G' is mapped onto a non-empty subgraph with even degrees, hence the image of G' contains a circuit G'. Now G' contains at least one edget of a copy of G(r, g) and hence, G' has less than G' edges. Thus G' = G(r+1, g).

Now since G(r, 2) as well as G(2, g) exist trivially (r parallel edges and a g-gon, respectively) we get that G(3,3), G(4,3)...; G(3,4), G(4,4),...; ...; G(3,g),

 $G(4, g), \ldots$  also exist. [P. Erdős-H. Sachs; see S, WV].

13. (a) Suppose indirectly that  $a, b \in V(G)$  are at distance > g. Remove a and b and add r new edges, matching the r neighbors of a with the r neighbors of b. The resulting graph G' is, obviously, r-regular.

We show that G' has girth at least g. Let C be a circuit in G'. If C does not contain any of the r new edges, then it is a circuit in G and thus, the length of C

is at least g. So suppose C contains some new edges.

Now consider a path P in  $G - \{a, b\}$  joining two endpoints of the new edges. If P joins a neighbor of a to a neighbor of b, then, since a, b are at distance at least g + 1, P has length at least g - 1. If P joins two neighbors of a (or b), then with two further edges it forms a circuit of G through a (or b) and hence, it has length at least g - 2.

Now if C contains only one new edge, then it contains a path joining the two endpoints of this edge. By the above argument, this path has at least g-1

edges and thus, C has at least q edges.

<sup>†</sup> At least s/2 such edges, in fact; but we do not use this observation.

On the other hand, if C contains at least two new edges, then it also contains at least two paths connecting endpoints of these. Hence its length is at least

$$2+2(g-2)\geq g.$$

Thus G' is r-regular and has girth at least g. Since |V(G')| < |V(G)|, this is a contradiction.

(b) I. Assume r=2l. Let  $x\in V(G)$ . Remove x and join the 2l neighbors of it by new independent edges. The resulting graph G' is clearly r-regular. Since |V(G')|<|V(G)|, G' must have girth less than g, i.e. it must contain a circuit C with length less then g. Obviously, C contains some (at least one) new edges; let  $e_1,\ldots,e_s$  be these edges and let them split C into arcs  $P_1,\ldots,P_s$ . We observe that  $P_i$  forms a circuit in G when we add the two edges joining its endpoints to x, thus  $|E(P_i)| \geq g-2$ . Hence

$$g > |E(C)| \ge s(g-2) + s = s(g-1).$$

Thus s=1, i.e. C consists of a new edge  $e_1$  and a path  $P_1$  connecting its endpoints, such that  $P_1 \subseteq G - \{x\}$ . It also follows that  $P_1$  has g-2 edges, i.e. joining its endpoints to x we get a circuit of length g in G.

II. Assume r = 2l + 1. If g = 2 or 3, then G consists of r parallel edges

and  $G = K_{r+1}$  respectively. Thus we may assume  $g \geq 4$ .

Let  $(x, y) \in E(G)$ . Remove x and y from G and pair up the 2l remaining neighbors of x as well as the 2l remaining neighbors of y. The resulting graph G' is r-regular again. Since |V(G')| < |V(G)|, G' contains a circuit G of length less than g. Again, let  $e_1, \ldots, e_s$  be the new edges on G and let them divide G into arcs G, G, G, and G into arcs G, G, and G is a constant.

$$|E(P_i)| \geq \begin{cases} g-2 & \text{if } P_i \text{ joins two neighbors of } x \text{ or two neighbors of } y, \\ g-3 & \text{if } P_i \text{ connects a neighbor of } x \text{ to a neighbor of } y. \end{cases}$$

Thus,

$$g > |E(C)| \ge s(g-3) + s = s(g-2),$$

whence s = 1 as  $g \ge 4$ . Now we conclude as before that  $P_1$  is contained in a circuit of G of length g, which passes through x or y.

(c) Let  $x \in V(G)$ . By (a), any point of G can be reached from x by a path of length at most g. But there are

$$\leq r(r-1)^{j-1}$$

paths of length j starting from x (some of them may terminate at the same point; some may come back to x) and thus,

$$|V(G)| \le 1 + r + r(r-1) + \dots + r(r-1)^{g-1} =$$
  
=  $1 + \frac{r}{r-2}((r-1)^g - 1) < \frac{r}{r-2}(r-1)^g$ 

[ibid.].

14. Take a maximal graph G', which arises from G by adding independent edges and which has girth at least g. Assume indirectly that G' has a point u of degree r. Then for reasons of parity, it has another such point v.

Let S be the set of points of G' at distance at most g-1 from  $\{u, v\}$ . Each point w of G' outside S must be of degree r+1, otherwise the edge (u, w)could be added to G'. Since by a computation similar to that in the solution of 10.11 we have

$$|S| \leq 2(1 + r + \ldots + r^{g-1}) < 2r^g \leq \frac{1}{2}|V(G)|,$$

it follows that there are two points x, y outside S connected by an edge of E(G') = E(G). Now G' = (x, y) + (x, u) + (x, v) is a graph with girth at least q, arising from G by addition of independent edges, and larger than G', a contradiction.

- 15. (a) Let U be the set of points and W the set of lines of the projective plane over GF(p). Connect  $u \in U$  to  $w \in W$  iff  $u \in w$ . The resulting bipartite graph is clearly (p+1)-regular and has  $2(p^2+p+1)$  points. To show it has girth 6, note that it contains no quadrilaterals; in fact, if  $(u_1, w_1, u_2, w_2)$  were a circuit, then the lines  $w_1, w_2$  would have two points in common  $u_1, u_2$ . It contains no circuits of length 3 or 5, because it is bipartite. Any triangle in the plane vields a hexagon in the graph.
- (b) Consider the hypersurface  $\mathcal{F} = \{x: x^T x = 0\}$  in the 4-dimensional projective space over GF(p). First we prove some geometric properties of  $\mathcal{F}$ .
- (i) F contains no plane. Suppose indirectly that  $\pi$  is a plane contained in F and let  $u, v \in \pi$ . Then  $u^T u = 0$ ,  $v^T v = 0$  and  $(u + v)^T (u + v) = 0$  (since u, v and u + v are vectors representing points of  $\pi$ ). Hence it follows that  $u^T v = 0$ . Hence if V denotes the linear subspace of the 5-dimensional vector space formed by homogeneous coordinate quintuples of points of  $\pi$ , we have  $V \subset V^{\perp}$ . But this implies that dim  $V^{\perp} \geq \dim V = 3$ , contradicting 5.31.
- (ii) There are exactly p+1 lines of F through each point b of F. Let  $\Sigma=$  $= \{x: b^T x = 0\}$  be the "tangent hyperplane" at b. Each line through b is contained in  $\Sigma$ ; for if v is a point of such a line then as in (i), it follows that  $b^T v = 0$ . Conversely, if  $v \in \mathcal{F} \cap \Sigma$ , then all points of the line spanned by b and v belong to  $\mathcal{F} \cap \Sigma$ .

Let  $\pi$  be any plane in  $\Sigma$  avoiding b. Then  $\pi \cap \mathcal{F}$  is a conic section, which is non-degenerate since if it contained a line, then this, together with b, would give a plane contained in  $\mathcal{F}$ , contradicting (i). So  $\pi \cap \mathcal{F}$  has p+1 points as is well known. Now the lines in  $\mathcal{F}$  through b are exactly those lines connecting b to the points of  $\pi \cap \mathcal{F}$ , so their number is p+1.

Let us note that these considerations together with (i) also imply that no three lines in F form a triangle.

(iii)  $|\mathcal{F}| = p^3 + p^2 + p + 1$ . Observe that given any line  $\Lambda$  and point b on  $\mathscr{F} = \Lambda$ , there is a unique line on  $\mathscr{F}$  containing b and meeting  $\Lambda$ . In fact, the tangent hyperplane  $\Sigma$  at b cannot contain  $\Lambda$ , so  $\Sigma$  intersects  $\Lambda$  in a single point a, and then the line ab is this unique line. Thus, if we consider any line  $\Lambda$ . there are p+1 points on  $\Lambda$ , p further lines incident with each point of  $\Lambda$ , and any point of F is incident with exactly one of these lines. Hence |F - A| = $= p^2(p+1)$ , which proves the assertion.

Since each line contains p+1 points and each point is incident with p+1lines of F, the number of lines of F is the same. Let L denote the set of lines of F.

Now form a bipartite graph on  $\mathcal{F} \cup \mathcal{L}$  by connecting  $b \in \mathcal{F}$  to  $A \in \mathcal{L}$  iff  $b \in A$ . The resulting bipartite graph is (p+1)-regular by (ii) and has  $2(p^3+p^2+p^2)$ + p + 1) points by (iii). It contains no 4- or 6-circuit, since a 4-circuit would correspond to two lines on F meeting in two points and a 6-circuit to 3 lines on F forming a triangle, both impossible.

16. (a) If Z represents all circuits, then V(G) - Z spans a forest and hence, it spans at most n-|Z|-1 edges. Since each point has degree at least 3, there are 3(n-|Z|) edges leaving the points of V(G)-Z. The edges spanned by this set are counted here twice but still we have at least

$$3(n-|Z|) - 2(n-|Z|-1) = n-|Z|+2$$

edges connecting V(G) - Z to Z. On the other hand, a point of Z is incident with not more than d edges, hence the number of (V(G)-Z,Z)-edges is at most d. |Z|. Thus

$$d \cdot |Z| \ge n - |Z| + 2$$

or, equivalently,

$$|Z| \ge \frac{n+2}{d+1} \, .$$

(b) We may suppose  $g \ge 3$ , as g = 1, 2 are trivial. Let d denote the maximum degree in G. The same counting as in 10.11 yields

$$n \ge 1 + d + 2d + \ldots + 2^{\left[\frac{g-3}{2}\right]} \cdot d = 1 + d(2^{\left[\frac{g-1}{2}\right]} - 1).$$

Now by (a), any set Z representing all circuits satisfies

$$|Z| \geq \frac{n+2}{d+1} \geq \frac{3 + d(2^{\left\lceil \frac{g-1}{2} \right\rceil} - 1)}{d+1} \geq \frac{3 + 3(2^{\left\lceil \frac{g-1}{2} \right\rceil} - 1)}{3+1} \geq \frac{3}{8} \cdot 2^{g/2}$$

[H.-J. Voß, M. Simonovits; see WV].

17. Let  $G_1, G_2, \ldots, G_{\nu}$  be defined as in the hint;  $G_{\nu+1} = \emptyset$ . One may assume that the girth of each  $G_i$  is at most g, otherwise we could consider  $G_i$  instead of G. Hence  $|V(C_1) \cup \ldots \cup V(C_n)| < \nu q$ .

On the other hand, it is easy to verify that  $V(C_1) \cup \ldots \cup V(C_{\nu})$  represents

all circuits of G. Thus part (b) of the preceding problem yields

$$|V(C_1) \cup \ldots \cup V(C_v)| \ge \frac{3}{8} 2^{g/2}.$$

Hence

$$v \ge \frac{3}{8g} \, 2^{g/2},$$

as stated [ibid.].

- 18. (a) v = 1 means that any two circuits of G have a point in common. If G is not simple, i.e. it contains a loop or two parallel edges, then all circuits are covered by 1 or 2 points, respectively. So we may assume G is simple. Then by 10.4, G is one of the graphs displayed in Fig. 8. The circuits of  $K_5$  are covered by any 3 points of it; those of the wheel are covered by the center and any one point of the rim; those of the third example  $(K_{3,n-3}$  with some edges added in the 3-element class) are covered by any two points of the 3-element class [B. Bollobás, L. Pósa].
- (b) We use induction on V(G). We may assume that G has no points of degree 1 or 2, because these could be removed and "smoothed out", respectively. Remove the points of a shortest circuit. By 10.16b, the length of this satisfies

$$g \leq 4 \log_2 \tau$$
.

Let  $\tau'$  denote the minimum number of points of covering all circuits of the remaining graph. Clearly

$$\tau \leq \tau' + g$$
.

So using the induction hypothesis,

$$v \geq rac{ au'}{4\log au'} + 1 \geq rac{ au'}{4\log au} + 1 \geq rac{ au - g}{4\log au} + 1 \geq rac{ au}{4\log au}.$$

(c) Let G be a 3-regular graph with girth g and minimum number n of points. By 10.13c,

$$n \leq 3 \cdot 2^g$$
, or  $g \geq \log_2 \frac{n}{3}$ .

By 10.16a, we need at least n/4 points to represent all circuits, i.e.  $n \leq 4\tau$ . Furthermore,

$$v \le \frac{n}{g} \le \frac{n}{\log_2(n/3)} \le \frac{4\tau}{\log_2 \tau}$$
.

- [P. Erdős-L. Pósa; this proof is due to H.-J. Voß, in: *Theory of Graphs* (Akadémiai Kiadó, Budapest 1968) and M. Simonovits, *Acta Math. Acad. Sci. Hung.* 18 (1967) 191–206; also see WV.]
- 19. Consider  $G_1 = G x$ . Since G is 2-connected,  $G_1$  is connected.
- 1°. If  $G_1$  is 2-connected, then let  $x_1$  be any point adjacent to x. Since  $d_{G_1}(z) \ge k-1$  for every  $z \ne x_1$ , y (for every  $z \ne y$ , in fact) there is an  $(x_1, y)$ -path of length at least k-1 in  $G_1$ . Adding  $(x, x_1)$  we get an (x, y)-path of length at least k.
- 2°. Assume  $G_1$  is not 2-connected and let  $G_1 = A \cup B$ ,  $|V(A) \cap V(B)| = 1$ . Choose the notation so that  $y \in V(A)$  and suppose B is minimal. Then B is, obviously, 2-connected. Let  $\{y_1\} = V(B) \cap V(A)$ . Since G is 2-connected, x must be adjacent to a point  $x_1 \in V(B) \{y_1\}$ .

Now each point  $z \neq y_1$  of B has degree at least k-1, hence B contains an  $(x_1, y_1)$ -path P of length at least k-1. Also, let  $P_2$  be a  $(y_1, y)$ -path in A; then

$$P = P_1 + P_2 + (x, x_1)$$

is an (x, y)-path in G of length at least k.

20. Let X be defined as in the hint. Obviously, all paths arising by repeated deformation from P have the same points, thus  $X \subseteq V(P)$ . Let  $X = \{x_0,$  $x_{i_1}, \ldots, x_{i_t}, x_m$   $\{0 < i_1 < \ldots < i_t < m\}$ . We claim that

$$\Gamma(X) \subseteq \{x_1, x_{m-1}, x_{i\nu \pm 1} : \nu = 1, \ldots, t\} \cup X$$

(whence  $|\Gamma(X) - X| \leq 2t - 2$ ).

Let  $u \in \Gamma(X)$ ,  $u \notin X$ . Then u is adjacent to a point  $v \in X$ . Since, by definition, v is an endpoint of a maximum path whose points are the points of P, u must

belong to  $\bar{P}$ . Let  $u = x_i$ .

By the definition of v, there are paths  $P_0 = P, P_1, \ldots, P_s$  such that  $P_{i+1}$  arises by deformation from  $P_i$   $(i = 0, \ldots, s-1)$  and v is an endpoint of  $P_s$ . If both edges  $(x_i, x_{i-1})$  and  $(x_i, x_{i+1})$  belong to  $P_s$  then let, say,  $(x_i, x_{i+1})$  be the first edge of the (u, v)-arc of  $P_s$ . Then  $x_{i+1}$  is an endpoint of a path arising from  $P_s$  by deformation, i.e.  $x_{i+1} \in X$  and we are finished.

So suppose, e.g. that  $(x_{i+1}, x_i) \notin E(P_s)$ . Then there is an index  $j, 0 \le j \le s-1$  such that  $(x_{i+1}, x_i) \in E(P_j)$  but  $(x_{i+1}, x_i) \notin E(P_{j+1})$ . Since  $P_{j+1}$  arises from  $P_j$  by deformation, this can only happen if one of  $x_{i+1}, x_i$  is the endpoint of  $P_{j+1}$ . Since  $x_i \notin X$ , we must have  $x_{i+1} \in X$  and we are done again. Thus we know that

$$|\Gamma(X) - X| \leq 2|X| - 2,$$

whence, by the assumption,  $|X| \ge k + 1$ . Let  $X_1 \subset X$ ,  $|X_1| = k$ . Then by the assumption

$$|\Gamma(X_1) - X_1| \ge 2|X_1| - 1 = 2k - 1$$

and since

$$X_1 \cup \Gamma(X_1) \subseteq V(P),$$

we have

$$|V(P)| \ge |X_1| + |\Gamma(X_1) - X_1| \ge k + 2k - 1 = 3k - 1.$$

The assertion is sharp for any k as is shown by any graph consisting of disjoint complete (3k-1)-graphs [L. Pósa, Discrete Math. 14 (1976) 359–364].

21. (a) Suppose indirectly that G has no Hamiltonian circuit. Let us add edges to G as long as we do not form a Hamiltonian circuit. Since condition (a) remains valid (by the way, so do (b), (c), (d)) we may suppose G is already saturated, i.e. adding any new edge to it, a graph containing a Hamiltonian circuit arises. Let  $(x, y) \notin E(G)$ , and let H be a Hamiltonian circuit of G ++(x, y). Then G contains a Hamiltonian path  $P=(z_1=x, z_2, \ldots, z_n=y)$ connecting x to y.

Let  $z_{i_1}, \ldots, z_{i_k}$  be the neighbors of x on P,  $2 = i_1 < i_2 < \ldots < i_k \le n$ . Then y cannot be adjacent to  $z_{i\nu-1}$   $(1 \le \nu \le k)$ ; otherwise,  $(z_1, \ldots, z_{i\nu-1}, z_n, z_{n-1}, \ldots, z_{i\nu})$  would be a Hamiltonian circuit. Hence

$$d_G(y) \le n-1-k = n-1-d_G(x) \le n-1-\frac{n}{2} < \frac{n}{2}$$

a contradiction.

We next assume (d) holds. We again may assume G contains no Hamiltonian circuit but joining any two non-adjacent points by an edge a Hamiltonian circuit arises. Let  $x_k$ ,  $x_l$  be a non-adjacent pair with k+l maximal (k < l).

10.22

Then  $x_k$  is adjacent to  $x_{l+1}, \ldots, x_n$ , hence

$$(1) d_k \ge n - l;$$

also,  $x_l$  is adjacent to  $x_{k+1}, \ldots, x_{l-1}, x_{l+1}, \ldots x_n$ , i.e.

$$(2) d_1 \ge n - k - 1.$$

The same argument as used in (a) yields

$$(3) d_k + d_l \le n - 1.$$

Now from (2) and (3),

$$(4) d_k \le n - 1 - d_l \le (n - 1) - (n - k - 1) = k.$$

Set  $m = d_k$ . Then by (4),  $m \ge k$  and thus,  $d_m \ge d_k = m$ . Also, (3) implies that

$$m=d_k<rac{n}{2}$$
 .

Thus by the assumption,

$$d_{n-m} \ge n - m = n - d_k \ge d_l + 1.$$

It follows that

$$n - d_k = n - m > l,$$

or, equivalently,

$$d_k < n - l$$
,

which contradicts (1).

If (b) holds, then, obviously, (d) holds. Suppose (c) holds, and let  $d_k < k \le \frac{n}{2}$ ,

l=n-k. Then by (c), either  $d_l \geq l$  or  $d_k+d_l \geq l+k=n$ , whence  $d_l > l$ . Thus (d) holds again and the existence of a Hamiltonian circuit follows [see B].

**22.** Let G be a simple graph, which does not contain a Hamiltonian circuit through the edges in F. We may assume that, adding any edge to G, we already have such a Hamiltonian circuit (because in the complete graph on V(G), there is a Hamiltonian circuit through F, as F consists of disjoint paths). Let  $(x, y) \notin E(G)$ . Then there is a Hamiltonian circuit in G + (x, y) through F, i.e. there is a Hamiltonian path P in G through F connecting x to y. Let  $P = (x = z_1, z_2, \ldots, z_n = y)$ . Let x be adjacent to  $z_{i_1}, \ldots, z_{i_k}$  ( $2 = i_1 < \ldots < i_k < n$ ). As before, it follows that  $z_{i_{p-1}}$  ( $1 \le v \le k$ ) can be adjacent to y only, if  $(z_{i_{p-1}}, z_{i_p}) \in F$ . Thus, at most n-1-k+q points are adjacent to y. Hence

$$(1) d_G(y) + d_G(x) \le n + q - 1.$$

This is clearly impossible.

Remark: One could give analogous generalizations as in the preceding problem [see B].

**23.** Add a point y and connect it to all points. The resulting graph on 2n + 2 points has degrees at least n + 1 and has a Hamiltonian circuit by Dirac's

theorem 10.21a. Removing y we still have a Hamiltonian path  $(x_0, \ldots, x_{2n})$ in G. Suppose G has no Hamiltonian circuit, then we have the following rule:

if  $x_0$  is adjacent to  $x_i$ , then  $x_{2n}$  is not adjacent to  $x_{i-1}$ .

Since the degrees of  $x_0$  and  $x_{2n}$  are n, it must hold that if  $x_0$  is not adjacent to  $x_i$  then  $x_n$  is adjacent to  $x_{i-1}$ .

Suppose first that  $x_0$  is adjacent to  $x_1, \ldots, x_n$  and  $x_{2n}$  is adjacent to  $x_n, \ldots$  $\dots, x_{2n-1}$ . There is an  $i, 1 \le i \le n$  such that  $x_i$  is not adjacent to  $x_i$ ; then  $x_i$  is




Fig. 81

adjacent to  $x_i$  for some  $n < j \le 2n - 1$ , because  $d_G(x_i) = n$ . Then the circuit  $(x_i, x_{i-1}, \dots, x_0, x_{i+1}, \dots, x_{j-1}, x_{2n}, \dots, x_j)$  is a Hamiltonian circuit (Fig. 81). Now let  $1 \le i \le 2n-1$  be such that  $x_{i+1}$  is adjacent to  $x_0$  but  $x_i$  is not. By the above argument,  $x_{i-1}$  is adjacent to  $x_{2n}$ . Thus G contains a (2n)-circuit

 $(x_{i-1}, \ldots, x_0, x_{i+1}, \ldots, x_{2n})$ . Let  $C = (y_1, \ldots, y_{2n})$  be a (2n)-circuit in G, and let  $y_0$  be the last point. Since C is a maximum circuit in G,  $y_0$  cannot be adjacent to two neighboring points of C; so it must be adjacent to every second point on C, to  $y_1, y_3, \ldots$ ...,  $y_{2n+1}$ , say. Replacing  $y_{2i}$  by  $y_0$  we get another maximum circuit and so,  $y_{2i}$  must also be adjacent to  $y_1, y_3, \ldots, y_{2n-1}$ . Now we observe that  $y_1$  is adjacent to  $y_0, y_2, \ldots, y_{2n}$ , i.e.

$$d_G(y_1) \ge n+1,$$

a contradiction [C.St.J.A. Nash-Williams, Proc. Amer. Math. Soc. 17 (1966) 466-467].

**24.** Let  $x, y \in V(G)$ , we prove that they can be connected by a Hamiltonian path. We may assume  $(x, y) \in E(G)$ , since connecting x, y by a new edge influences neither the assumption nor the conclusion.

Subdivide the edge (x, y) by a new point z. Then it is easy to see that the resulting graph G' has a Hamiltonian circuit, if and only if G contains a Hamil-

tonian path connecting x to y.

Now the degrees of 
$$ar{G}'$$
 are  $2 < d_2 < \ldots < d_{n+1},$ 

where  $d_2 \leq \ldots \leq d_{n+1}$  are the degrees of G; hence  $d_2 \geq \frac{n+1}{2}$  by the assumption. Since |V(G')| = n + 1, Pósa's condition (b) in 10.21 is satisfied and hence, G' has a Hamiltonian circuit. Thus, G contains a Hamiltonian path connecting x to y [see B].

**25.** Let  $C = (x_1, \ldots, x_m)$  be a longest cycle in G. Then  $|V(C)| > \frac{n}{2}$ . In fact,

let  $Q = (z_0, \ldots, z_p)$  be a longest (directed) path and  $z_{i_1}, \ldots, z_{i_k}$  ( $i_1 < \ldots < i_k$ ) be those points of G for which  $(z_{i_{\nu}}, z_0) \in E(G)$  (they obviously lie on Q). Then  $i_k \ge k \ge n/2$  and thus, the cycle  $(z_0, \ldots, z_{i_k})$  has length > n/2. Suppose G is not Hamiltonian and let  $P = (y_0, \ldots, y_l)$  be a longest path

Suppose C is not Hamiltonian and let  $P = (y_0, \ldots, y_l)$  be a longest path in G - V(C). We have  $(u, y_0) \in E(G)$  for at least  $\frac{n}{2} - l$  points u not on P.

All these points belong to C; let us denote them by  $x_{i_1}, \ldots, x_{i_l}, t \ge \frac{n}{2} - l$ . Similarly, there are points  $x_{j_1}, \ldots, x_{j_s}, s \ge \frac{n}{2} - l$  such that  $(y_l, x_{j_r}) \in E(G)$ ,  $1 \le v \le s$ .

Now observe that if  $x_{i\nu} \neq x_{j\mu}$ , then the  $(x_{i\nu}, x_{j\mu})$ -arc of C must be at least as long as l+2; for otherwise, this arc of C could be replaced by  $(x_{i\nu}, y_0) + P + (y_l, x_{j\mu})$ . Thus if we consider an arc  $A_{\nu}$  of C, starting at  $x_{i\nu+1}$   $(x_{m+1} = x_1)$  and having length l,  $(1 \leq \nu \leq t)$ , then

$$x_{j\mu} \notin \bigcup_{v=1}^{t} V(A_{v}) \qquad (1 \leq \mu \leq s).$$

But it is easy to see that  $\left|igcup_{
u=1}^t V(A_
u)
ight| \ge t+l$  and thus

$$s \leq m - (t+l)$$

or, equivalently,

$$m \ge s + t + l \ge \left(\frac{n}{2} - l\right) + \left(\frac{n}{2} - l\right) + l = n - l$$

But P has l+1 points out of n-m, thus

$$l+1 \leq n-m, \ m \leq n-l-1,$$

a contradiction [C.St.J.A. Nash-Williams; see B].

**26.** Let C be a maximum circuit in G. Suppose indirectly that C is not a Hamiltonian circuit. Then G-V(C) is non-empty; let  $G_1$  be a component of it. Let  $x_1, \ldots, x_s$  be those points of C adjacent to  $G_1$ . Observe that no two  $x_i$  are neighboring points on C by the maximality of C. This implies that  $\{x_1, \ldots, x_s\}$  separates G and hence,  $s \geq k$ . Going around C in a given direction, let  $y_1, \ldots, y_s$  be the points following  $x_1, \ldots, x_s$ .

Now we claim that  $y_1, \ldots, y_s$  are independent. Suppose indirectly that  $y_i$  and  $y_j$  are adjacent, then remove  $(x_i, y_i)$  and  $(x_j, y_j)$  from C and add  $(y_i, y_j)$  and an  $(x_i, x_j)$ -path through  $G_1$ ; the resulting circuit is longer than C, a contra-

diction.

Also, no  $y_i$  is adjacent to  $G_1$  and thus, selecting a  $y_0 \in V(G_1)$ , the set  $S = \{y_0, y_1, \ldots, y_s\}$  will be independent. But

$$|S| = s + 1 \ge k + 1,$$

a contradiction [P. Erdős-V. Chvátal; see B].

**27.** (a) Let  $P = (x_0, x_1, \ldots, x_m)$  be a longest path. Then all neighbors of  $x_0$  are on P; since  $d_G(x_0) \geq k$ , one of these neighbors is  $x_i$  with  $k \leq i \leq m$ . Then  $C = (x_0, \ldots, x_i)$  is a circuit of length  $i + 1 \geq k + 1$ .

(b) Let P be a longest path as above. Suppose first that there are points  $x_i, x_j$  such that  $i < j, x_i$  is adjacent to  $x_m$  and  $x_j$  is adjacent to  $x_0$ . We may assume j - i is minimal among such pairs of indices. Let  $C = (x_0, \ldots, x_i, x_m, x_{m-1}, \ldots, x_j)$ . If j = i + 1, then C has length m + 1 and must be a Hamiltonian circuit, since otherwise there would be a point outside C connected to C, which would yield a path longer than P. So we may assume that  $j \ge i + 2$ . Then  $x_{i+1}, \ldots, x_{j-1}$  are not adjacent to  $x_0$  or  $x_m$ , hence C contains  $x_m$ , all neighbors of  $x_m$ , and all points  $x_v$  for which  $x_{v+1}$  is adjacent to  $x_0$ , except  $x_{j-1}$ . These points are distinct and hence C has at least 2k points.

So suppose that the last point  $x_i$  adjacent to  $x_0$  comes before the first point  $x_j$  adjacent to  $x_m$  (possibly  $x_i = x_j$ ). Since G is 2-connected, there are two disjoint paths  $P_1$ ,  $P_2$  connecting the circuits  $C_1 = (x_0, \ldots, x_i)$  and  $C_2 = (x_j, \ldots, x_m)$ . We may assume that one of them starts at  $x_i$ ; for otherwise we can walk on the path  $(x_i, \ldots, x_j)$  till we hit either  $C_2$  or  $P_v$ , and replace an appropriate piece of  $P_v$  by this path. Similarly, we may assume that one of  $P_1$ ,  $P_2$  ends at  $x_j$ . It may be that the same  $P_v$  ends at  $x_i$  and  $x_j$  or not; but in both cases we get a circuit which contains  $x_0$ ,  $x_m$  and all neighbors of them as shown in Fig. 82. This circuit is longer than 2k [G.A. Dirac, Proc. London Math. Soc. 2 (1952) 69-81].




Fig. 82

28. For  $n \le k$  the assertion is void, so suppose n > k. If G has a point x with degree at most k/2, G - x has more than  $\frac{(n-2)k}{2}$  edges and so, by the induction hypothesis, G - x contains a circuit of length greater than k. Thus we may assume that every point of G has degree at least  $\frac{k+1}{2}$ .

Also, if G is not 2-connected, say  $G = G_1 \cup G_2$  with  $|V(G_1) \cap V(G_2)| \leq 1$  then, by

$$|E(G_1)| + |E(G_2)| > \frac{k(n-1)}{2} \ge \frac{k}{2} (|V(G_1)| + |V(G_2)| - 2),$$

it follows that for i = 1 or 2,

$$|E(G_i)| > \frac{k}{2} (|V(G_i)| - 1)$$

and induction works again. Thus we may suppose that G is 2-connected. Then 10.27 implies that G contains a circuit of length at least k+1.

The result is sharp for graphs whose blocks are  $K_k$ 's. [P. Erdős—T. Gallai,

Acta Math. Acad. Sci. Hung. 10 (1959) 337-356].

**29.** Let P = (0, 1, ..., N) be any path in G. Let  $R_1$  be a (P, P)-path with endpoints  $i_1 = 0$  and  $j_1$  such that  $j_1$  is maximal. Suppose the (P, P)-paths  $R_1, ..., R_k$  have been selected,  $R_v$  has endpoints  $i_v < j_v$  and  $j_k < N$ . Let  $R_{k+1}$  be a (P, P)-path connecting  $\{0, ..., j_k - 1\}$  to  $\{j_k + 1, ..., N\}$  (such a path exists as G is 2-connected) and let  $j_{k+1}$  be as large as possible. If  $j_k = N$  we stop; let s be this value of k.

It follows from the definition that

$$0 < j_1 < j_2 < \ldots < j_s = N.$$

Also,

$$i_{k+1} \geq j_{k-1},$$

since otherwise  $R_{k+1}$  could have been chosen for  $R_k$ . Thus we have

$$0 = i_1 < i_2 < j_1 \le i_3 < j_2 \le \ldots \le i_s < j_{s-1} < j_s.$$

Similar reasoning shows that  $R_1, \ldots, R_s$  are independent paths. Thus  $P \cup$ 

 $\bigcup R_1 \bigcup \ldots \bigcup R_s$  has the structure indicated in the hint.

Suppose that s=2p+1 is odd (the even case can be treated similarly and is left to the reader). Then the arc between  $i_{p+1}$  and  $j_{p+1}$  is not longer than l-1 since it forms a circuit with  $R_{p+1}$ . Similarly, the  $(i_p, i_{p+1})$ -arc and  $(j_{p+1}, j_{p+2})$ -arc of P form a circuit with  $R_p$ ,  $R_{p+1}$ ,  $R_{p+2}$  and the  $(j_p, i_{p+2})$ -arc of P (this latter may degenerate; Fig. 83) and hence their total length is at most l-3. Similarly we get that the sum of lengths of the  $(i_q, i_{q+1})$ -arc and the  $(j_{2p+1-q}, j_{2p+2-q})$ -arc is at most l-(p+1-q). Thus

$$|E(P)| \le (l-1) + (l-3) + \ldots = \frac{l^2}{4}$$
,

which proves the assertion [H.-J. Voß, G. Dirac, see WV].




Fig. 83

**30.** Let  $x, y \in V(G)$ ,  $(x, y) \in E(G)$ . Any  $z \in V(G) - \{x, y\}$  is adjacent to at most one of x, y. Hence

$$(d(x)-1)+(d(y)-1) \le n-2, d(x)+d(y) \le n.$$

Sum this over all edges (x, y), then we get each d(x) exactly d(x) times on the left-hand side, i.e.

$$\sum_{x} d^{2}(x) \leq n \cdot |E(G)|.$$

Here

$$\sum_{x} d^{2}(x) \ge \frac{1}{n} \left( \sum_{x} d(x) \right)^{2} = \frac{4}{n} |E(G)|^{2},$$

thus

$$\frac{4}{n} |E(G)|^2 \le n|E(G)|, \quad |E(G)| \le \frac{n^2}{4}.$$

[W. Mantel, Wiskundige Opgaven 10 (1906) 60-61. Other simple proofs can be obtained by specializing the solutions of Turán's theorem 10.34 and 10.35.]

31. Let x be any point of G. Then no two neighbors of x can be adjacent, hence  $d(x) \leq \alpha(G)$ .

Now let S be a minimum point cover of G. Then each edge of G is represented by a point in S, thus

$$|E(G)| \le \sum_{x \in S} d(x) \le \alpha(G) \cdot |S| = \alpha(G) \cdot \tau(G).$$

Since

$$lpha(G)\cdot au(G) \leq \left(rac{lpha(G)\,+\, au(G)}{2}
ight)^2 = \left(rac{n}{2}
ight)^2 = rac{n^2}{4}\,,$$

the result in the preceding problem also follows from this.

32. Let us call a triple  $\{x, y, z\}$  bad if it does not span a triangle of G or  $\overline{G}$ . Then the number of bad triples such that exactly one edge spanned by them contains the point x is d(x)(n-1-d(x)), so

$$\sum_{x} d(x) (n-1-d(x))$$

counts the bad triples twice. The number of triangles in G and  $\bar{G}$  together is, therefore,

$$\binom{n}{3} - \frac{1}{2} \sum_{x \in V(G)} d(x) (n - 1 - d(x)).$$

(a) If G is k-regular this formula reduces to

$$\binom{n}{3} - \frac{1}{2} nk(n-1-k)$$

as stated.

(b) We have

$$d(x)(n-1-d(x)) \le \left(\frac{n-1}{2}\right)^2$$

and hence, the number of triangles in G and  $\overline{G}$  is at least

$$\binom{n}{3} - \frac{n}{2} \left( \frac{n-1}{2} \right)^2 = \frac{n(n-1)(n-5)}{24}.$$

Note: This is positive for n > 5, which is a very special case of Ramsey's theorem (§ 14) [A.W. Goodman, Amer. Math. Monthly 66 (1959) 778–783].

**33.** Let  $(x, y) \in E(G)$ . Then at least d(x) + d(y) - n points are adjacent to both x and y, i.e. at least this many triangles contain (x, y). Hence

$$\frac{1}{3} \sum_{(x,y) \in E(G)} \left( d(x) + d(y) - n \right)$$

estimates the total number of triangles from below. Since d(x) occurs in this sum exactly d(x) times, the number of triangles in G is at least

$$\frac{1}{3} \Big( \sum_{x \in V(G)} d(x)^2 - nm \Big).$$

By the Cauchy-Schwartz inequality, this is

$$\geq rac{1}{3} \left(rac{\left(\sum\limits_{x} d(x)
ight)^2}{n} - nm
ight) = rac{4m}{3n} \left(m - rac{n^2}{4}
ight).$$

[ibid.]

**34.** We use induction on m. For m = 1 the assertion is void.

Let H be a complete k-graph in G (such a subgraph exists by induction on k, or let us "saturate" G by adding edges as long as no complete (k+1)-graph is produced, and consider this saturated graph rather than G). If G contains no complete (k+1)-graph, each  $x \in V(G) - V(H)$  is adjacent to no more than k-1 points in H. Hence the graph  $G_1 = G - V(H)$  has more than

$$\binom{k}{2} m^2 - (k-1)(m-k) - \binom{k}{2} = \binom{k}{2} (m-1)^2$$

edges. By the induction hypothesis, this implies that  $G_1$  contains a complete (k+1)-graph and hence, so does G.

The equality is attained when  $V(G) = V_1 \cup \ldots \cup V_k$ ,  $|V_i| = m$  and two point

points are adjacent iff they belong to different sets  $V_i$ .

If n=mk+r then we claim that the most number of edges in a simple graph without triangles is attained when  $G=H_{n,k},\ V(H_{n,k})=V_1\cup\ldots\cup V_k,\ |V_1|=\ldots=|V_r|=m+1,\ |V_{r+1}|=\ldots=|V_k|=m$  and again, two points are adjacent iff they belong to different classes. This number of edges is

$$\binom{r}{2}(m+1)^2 + r(k-r)m(m+1) + \binom{k-r}{2}m^2.$$

This claim is trivially true if m = 0 and can be proved by induction on m in the same way as above [P. Turán; see B, S].

**35.** Let x be a point with maximum degree and denote by  $G_0$  the subgraph induced by the neighbors of x. Then there is a graph  $H_0$  on  $V(G_0)$ , which is (k-1)-chromatic and satisfies

$$d_{H_0}(z) \ge d_{G_0}(z)$$
  $(z \in V(G_0));$ 

this follows by induction on k since  $G_0$  contains no complete k-graph. Define H on V(G) by connecting all points of  $V(G) - V(G_0)$  to all points of  $V(G_0)$  and also all pairs of points adjacent in  $H_0$ .

Obviously, H is k-chromatic. Also,

Note that Turán's theorem is an immediate consequence. For let  $A_1, \ldots, A_k$  be the color-classes of H,  $|A_i| = a_i$ . Then  $a_1 + \ldots + a_k = n$  and

$$|E(G)| = \frac{1}{2} \sum_{x \in V(G)} d(x) \le \frac{1}{2} \sum_{x \in V(H)} d_H(x) = |E(H)| \le \sum_{1 \le i < j \le k} a_i a_j.$$

The right-hand side is maximal if the  $a_i$ 's are as nearly equal as possible, i.e.  $a_1 = \ldots = a_r = m+1, a_{r+1} = \ldots = a_k = m$ . This gives Turán's bound [P. Erdős,  $Mat.\ Lapok\ 21\ (1970)\ 249-251$ ].

**36.** (a) Suppose G is a simple graph on n points containing no 4-circuit. Let us count those pairs  $\{x, y\}$  of points both adjacent to a third point z.

For a fixed z, we count  $\binom{d(\bar{z})}{2}$  such pairs. On the other hand, each pair x, y is counted at most once since if it were counted both with  $z_1$  and with  $z_2$ , then  $(x, z_1, y, z_2)$  would form a quadrilateral. Hence

$$\sum_{z \in V(G)} \binom{d(z)}{2} \leq \binom{n}{2}.$$

Here by Jensen's inequality,

$$\sum_{z \in V(G)} \binom{d(z)}{2} \ge n \left(\frac{2 |E(G)|}{n}\right) = |E(G)| \frac{2 |E(G)| - n}{n}$$

and thus

$$|E(G)|^2 - \frac{n}{2} |E(G)| \leq \frac{n^3 - n^2}{4},$$

whence

$$|E(G)| \leq \sqrt{\frac{n^3}{4} - \frac{3n^2}{16}} + \frac{n}{4} = \frac{n}{4} \left(1 \, + \, \sqrt{4n - 3}\right) \sim \frac{1}{2} \, n^{3/2}.$$

(b) Let the points of V(G) be the points of the projective space over GF(p), and connect points [x, y, z] and [u, v, w] iff xu + yv + zw = 0 (this means, if they are conjugate with respect to the conic section  $x^2 + y^2 + z^2 = 0$ ; we use homogeneous coordinates).

The points adjacent to any given point u form a line of the geometry (the polar of u), which goes through u iff u is on the conic  $x^2 + y^2 + z^2 = 0$ . Hence G has p + 1 points of degree p and the remaining  $p^2$  points of degree p + 1.

Thus

$$|E(G)| = \frac{1}{2} (p(p+1) + (p+1)p^2) = \frac{1}{2} p(p+1)^2 =$$

$$= \frac{n-1}{4} (\sqrt{4n-3} - 1).$$

On the other hand, G contains no 4-circuit. In fact, if u, v are two points, then their polars meet in a single point (since they are trivially distinct lines), so any two points have at most one common neighbor [I. Reiman,  $Acta\ Math.\ Acad.\ Sci.\ Hung.\ 9\ (1959)\ 269-279$ ].

**37.** Let m = |E(G)|,  $V(G) = \{x_1, \ldots, x_n\}$ ,  $d_G(x_i) = d_i$ . Since G contains no  $K_{r,r}$ , any given r-tuple in V(G) is contained in the neighborhood of at most r-1 points. The neighborhood of  $x_i$  contains  $\binom{d_i}{r}$  r-tuples. Thus

$$\sum_{i=1}^n \binom{d_i}{r} \leq (r-1) \, \binom{n}{r} \, .$$

We are going to estimate the left-hand side from below using Jensen's inequality. Unfortunately,  $\binom{x}{r}$  is convex only for  $x \ge r - 1$ . But set

$$f(x) = \begin{cases} \binom{x}{r} & \text{if } x \ge r - 1, \\ 0 & \text{if } x \le r - 1, \end{cases}$$

then f(x) is convex and, since  $d_i$  is an integer, we have

$$\sum_{i=1}^{n} \binom{d_i}{r} = \sum_{i=1}^{n} f(d_i) \ge n \cdot f\left(\frac{d_1 + \ldots + d_n}{n}\right) = nf\left(\frac{2m}{n}\right).$$

Now, if  $\frac{2m}{n} < r - 1$ , we have nothing to prove. Suppose  $\frac{2m}{n} \ge r - 1$ , then we

have

$$\left((r-1)\binom{n}{r} \ge \sum_{i=1}^{n} \binom{d_i}{r} \ge nf\left(\frac{2m}{n}\right) = n\binom{\frac{2m}{n}}{r}.$$

Here

$$(r-1) {n \choose r} < (r-1) \frac{n^r}{r!}, n {2m \choose r} > n \frac{\left(\frac{2m}{n} - r + 1\right)^r}{r!}$$

and thus

$$\begin{split} \left(\frac{2m}{n} - r + 1\right)^r &< (r - 1)n^{r - 1}\,,\\ 2m &< \sqrt[r]{r - 1}\,\cdot\,n^{2 - \frac{1}{r}} + (r - 1)n &< C\cdot n^{2 - \frac{1}{r}} \end{split}$$

[K. Zarankiewicz's problem: T. Kővári—V.T. Sós—P. Turán, Collogu, Math. 3 (1954) 50-571.

**38.** (a) We use induction on k. For k=1 the assertion is true. Let  $k\geq 2$  and  $s = \left[\frac{1}{\varepsilon}t\right]^*$ . If n is large enough then, by the induction hypothesis, we find k disjoint s-sets  $A_1, \ldots, A_k$  such that any two points in distinct sets  $A_i$  are adjacent.

Now let W denote the set of those points in  $U = V(G) - A_1 - \ldots - A_k$  adjacent to at least t points in each  $A_i$ . Then there are at least

$$(|U|-|W|)(s-t) \ge (|U|-|W|)(1-\varepsilon)s = (n-ks-|W|)(1-\varepsilon)s$$
 edges missing between  $U$  and  $A_1 \cup \ldots \cup A_k$ . On the other hand, there are at most  $\left(\frac{1}{k}-\varepsilon\right)n$  edges missing from a given point and hence, there are at most

$$ks \left(\frac{1}{k} - \varepsilon\right) n = (1 - k\varepsilon) s n$$

edges missing between U and  $A_1 \cup \ldots \cup A_k$ . Thus

$$(n-ks-|W|)(1-\varepsilon)s \leq (1-k\varepsilon)sn,$$

whence

$$|W| \ge \frac{k}{1-\varepsilon}n - ks.$$

Thus W gets large if n is large.

Select t points adjacent to  $w \in W$  from each  $A_i$ , for all  $w \in W$ . If

$$|W| > {s \choose t}^k (t-1),$$

then we necessarily select the same t-tuples for t distinct points in W. But these t points together with the k t-tuples belonging to them form a desired subgraph.

(b) Remove a point with degree less than  $\left|1 - \frac{1}{L} + \frac{\varepsilon}{2}\right| \cdot |V(G)|$  (if any); do so again for the resulting graph and so on. Suppose we get stuck, i.e. we get a graph H with all degrees at least  $\left(1-\frac{1}{k}+\frac{\varepsilon}{2}\right)|V(H)|$ . Then if N=|V(H)|is large enough, the preceding problem implies the assertion. Thus it suffices to show that N cannot be too small (and that we get stuck at all, i.e.  $N \neq 0$ ). In the construction of H we removed altogether at most

$$\sum_{j=N+1}^n j \bigg(1 - \frac{1}{k} + \frac{\varepsilon}{2}\bigg) = \bigg(\binom{n+1}{2} - \binom{N+1}{2}\bigg) \bigg(1 - \frac{1}{k} + \frac{\varepsilon}{2}\bigg) \leq$$

$$\leq \left( {n \choose 2} - {N \choose 2} \right) \left( 1 - rac{1}{k} + rac{arepsilon}{2} 
ight) + (n-N)$$

edges. The remainder has at most  $\binom{N}{2}$  edges, thus

$$|E(G)| = \left(1 - \frac{1}{k} + \frac{\varepsilon}{\varepsilon}\right) \binom{n}{2} \leq \left(1 - \frac{1}{k} + \frac{\varepsilon}{2}\right) \left(\binom{n}{2} - \binom{N}{2}\right) + (n - N) + \binom{N}{2}$$

or, equivalently,

$$\left|\frac{\varepsilon}{2} {n \choose 2} \right| \le \left(\frac{1}{k} - \frac{\varepsilon}{2}\right) {N \choose 2} + (n-N).$$

Hence N gets large if n is large and the assertion follows.

(c) Set  $\chi(G_0) = k + 1$ . Consider Turán's graph  $H_{n,k}$  as in the solution of 10.34. This is k-chromatic and hence it does not contain a  $G_0$ . Hence

$$M(n,G_0) \geq |E(H_{n,\,k})| \sim \frac{n^2}{2} \left[1 \, -\frac{1}{k}\right],$$

i.e.

$$\lim\inf\frac{M(n,G_0)}{n^2}\!\geq\!\frac{1}{2}\!\left(\!1\,-\frac{1}{k}\!\right),$$

Now suppose indirectly that there were arbitrarily large graphs G containing no copy of  $G_0$  with

 $rac{|E(G)|}{n^2}$   $> rac{1}{2} \left(1 - rac{1}{k}
ight) + \, arepsilon.$ 

Then by (b), an appropriately large one would contain k+1 disjoint sets of cardinality  $|V(G_0)|$  such that any two points in two different such sets are adjacent. But then  $G_0$  is a subgraph of this subgraph already, a contradiction. [P. Erdős-A.H. Stone, Bull. Amer. Math. Soc. 52 (1946) 1089–1091. This proof is a specialization of P. Erdős-M. Simonovits, Studia Sci. Math. Hung. 1 (1966) 51–57.]

- **39.** Set  $n_0 = |V(G_0)|$ . First we prove the assertion formulated in the hint, more exactly:
- (\*) If  $G \neq H_{n,k}$  is a simple graph with  $n \geq N(k, n_0)$  points,  $|E(H_{n,k})|$  edges and G contains no  $G_0$ , then it contains an induced subgraph  $G_1$  on  $n_1 = n 2kn_0$  points such that

$$|E(G_1)| - |E(H_{n_1,k})| > |E(G)| - |E(H_{n_1,k})|.$$

In fact  $N = N(k, n_0)$  is chosen so that a graph on  $n \ge N$  points and with  $|E(H_{n,k})|$  edges should contain k disjoint classes  $A_1, \ldots, A_k$ ,  $|A_i| = 2n_0$  such that any two points in different sets  $A_i$  are adjacent. Such an N exists by 10.38b since

$$|E(H_{n, \lfloor k 
vert})| \sim inom{k}{2} inom{n}{\lfloor k 
vert} \geq iggl(1 - rac{1}{k - \lfloor 1 
vert} + rac{1}{k^2}iggr) rac{n^2}{2}.$$

Now for each  $x \notin A_1 \cup \ldots \cup A_k$ , there must be at least  $2n_0$  points of  $A_1 \cup \ldots \cup A_k$  non-adjacent to x. Otherwise, x would be adjacent to at least one point in each class and to at least  $n_0$  points in all but one of the classes. Hence we could select a set  $B_i \subseteq A_i$ ,  $|B_i| = n_0$  such that x is adjacent to all points in, say,  $B_1 \cup \ldots \cup B_{k-1}$  and to at least one point in  $B_k$ . Since  $G_0 - e$  is k-chromatic, the subgraph induced by  $B_1 \cup \ldots \cup B_k \cup \{x\}$  contains  $G_0$ , a contradiction.

Also observe that equality can hold only if x is adjacent to all points of all but one  $A_i$  and to no point of this  $A_i$ . Suppose that the equality holds for every x. Then the x's can be divided into k classes  $C_1, \ldots, C_k, x \in C_i$  being adjacent to all points of  $A_j, j \neq i$  but to no point of  $A_i$ . No two points in the same  $C_i$  can be adjacent since then we could find a  $G_0$  in G in the same way as above. Hence, G is k-chromatic. Among k-chromatic graphs, obviously,  $H_{n,k}$  has the most edges, which contradicts  $G \neq H_{n,k}$  and  $|E(G)| \geq |E(H_{n,k})|$ .

Thus some x is adjacent to less than  $(2k-2)n_0$  points of  $A_1 \cup \ldots \cup A_k$ .

Hence the removal of  $A_1 \cup \ldots \cup A_k$  destroys less than  $M = 4 \begin{bmatrix} k \\ 2 \end{bmatrix} n_0^2 + (n - 1)^2 n_0^2 + (n$ 

 $-2kn_0(2k-2)n_0$  edges. The removal of  $2n_0$  points from each class of  $H_{n,k}$  destroys exactly M edges and results in  $H_{n,k}$ . Thus

$$|E(G - A_1 - \ldots - A_k)| - |E(H_{n_1,k})| > |E(G)| - |E(H_{n_k,k})|.$$

This proves (\*).

Now suppose indirectly that there is a simple graph  $G\neq H_{n,k}$  with  $n\leq 2kn_0\binom{N}{2}+N$  points, at least  $|E(H_{n,k})|$  edges and containing no  $G_0$ . Set  $n_i=n-2kin_0$ . Select induced subgraphs  $G\supseteq G_1\supseteq G_2\ldots$  such that  $|V(G_i)|=n_i$  and  $|E(G)|-|E(H_{n,k})|<|E(G_1)|-|E(H_{n,k})|<\ldots$  Then

$$|E(G_i)| \ge |E(H_{n_i,k})| + i.$$

Let us see when the sequence  $G_1, G_2, \ldots$  stops. Suppose this happens at the  $t^{\text{th}}$  step. By (\*) we can find a  $G_{t+1}$  unless  $G_t \cong H_{n_t,k}$  or  $|E(G_t)| < |E(H_{n_t,k})|$  or  $|V(G_t)| < N$ . (1) rules out the first two possibilities immediately. But also, if

$$|V(G_t)| = n_t = n - 2kt n_0 < N$$

then

$$t > \frac{n-N}{2k n_0} \ge {N \choose 2}$$

and so,

$$|E(G_t)| \geq t > {N \choose 2} > {n_t \choose 2},$$

a contradiction. [M. Simonovits, in: Theory of Graphs (P. Erdős–G. Katona, ed.) Akadémiai Kiadó, Budapest, 1968, 270–319.]

**40.** (a) Let  $A_1, \ldots, A_{N_k}$  be the complete k-graphs and  $B_1, \ldots, B_{N_{k-1}}$  be the complete (k-1)-graphs in G. Let  $A_i$  be contained in  $a_i$  complete (k+1)-graphs; also, let  $B_i$  be contained in  $b_i$  complete k-graphs.

Let x be a point not in  $A_i$  which does not form a complete (k+1)-graph with  $A_i$ . Then x is non-adjacent to some point  $y \in A_i$ . Let  $U_1, \ldots, U_{k-1}$  be those k-subsets of  $A_i \cup \{x\}$  containing  $\{x, y\}$ . Then any pair  $(A_i, U_j)$  consists of a complete k-graph and a non-complete k-graph, and has  $|A_i \cap U_j| = k - 1$ . So

$$\sum_{i=1}^{N_k} (k-1)(n-k-a_i)$$

is a lower bound for the number of pairs (A, U) such that A is a complete k-graph, U is a non-complete k-graph and  $|A \cap U| = k - 1$ .

On the other hand, the number of such pairs (A, U) is, obviously,

$$\sum_{i=1}^{N_{k-1}} b_i (n-k+1-b_i).$$

Thus we have

(1) 
$$\sum_{i=1}^{N_k} (k-1)(n-k-a_i) \leq \sum_{i=1}^{N_{k-}} b_i(n-k+1-b_i).$$

We know that

(2) 
$$\sum_{i=1}^{N_k} a_i = (k+1)N_{k+1}, \sum_{i=1}^{N_{k-1}} b_i = kN_k.$$

The left-hand side of (1) is equal to

$$(k-1)(n-k)N_k - (k^2-1)N_{k+1}$$

while the right-hand side can be estimated by Jensen's inequality

$$egin{aligned} \sum_{i=1}^{N_{k-1}} b_i(n-k+1-b_i) & \leq N_{k-1} rac{\sum\limits_{i=1}^{N_{k-1}} b_i}{N_{k-1}} \Biggl(n-k+1 - rac{\sum\limits_{i=1}^{N_{k-1}} b_i}{N_{k-1}} \Biggr) = \\ & = k(n-k+1)N_k - rac{k^2 \, N_k^2}{N_{k-1}} \, . \end{aligned}$$

whence (a) follows [J.W. Moon—L. Moser, Mat. Kut. Int. Közl. 7 (1962) 283–286].

(b) Estimation (\*) in the hint follows from (a) by a straightforward induction. Now from (\*),

$$N_k \ge n \prod_{i=2}^k \frac{\vartheta - i + 1}{\vartheta} \frac{n}{i} = \binom{\vartheta}{k} \left(\frac{n}{\vartheta}\right)^k.$$

41. (a) Any triple of points of  $T_n$  spans either a 3-cycle or a transitive triangle. If two edges have a common tail their points form a transitive triangle and each transitive triangle contains one such pair of edges. Hence the number of transitive triangles is

$$\sum_{i=1}^n inom{d_i}{2} \geq ninom{n-1}{2} \ 2 = rac{n(n-1)(n-3)}{8}$$

by Jensen's inequality, since  $\Sigma d_i = \binom{n}{2}$ . Thus the number of 3-cycles is

$$\leq {n \choose 3} - \frac{n(n-1)(n-3)}{8} = \frac{(n+1)n(n-1)}{24} = \frac{1}{4} {n+1 \choose 3}.$$

(b) Let  $x \in V(T)$ , and denote by X and Y the sets of points y such that  $(y, x) \in E(G)$  or  $(x, y) \in E(G)$ , respectively. There must be an edge joining Y to X, otherwise T would not be strongly connected. This edge forms, with x, a 3-cycle.

We prove that the number of 3-cycles is at least n-2 by induction on n.

For n=3 this is obvious.

By 6.13 we can find a point x such that T-x is strongly connected; thus, by the induction hypothesis, T-x contains at least n-3 3-cycles. As observed above, there is at least one 3-cycle incident with x; hence T contains at least n-2 3-cycles.

Remark: Both assertions are essentially sharp: if n is odd, then  $K_n$  has an orientation such that each point has outdegree  $\frac{n-1}{2}$  by 5.13 and this tournament has exactly  $\frac{1}{4} \binom{n+1}{3}$  3-cycles. On the other hand, consider a transitive tournament and invert the edge e connecting its "first" and "last" point. Then we obtain a strongly connected tournament in which each 3-cycle contains e, thus their number is n-2 [see S, M].

**42.** First we show by induction on n that for each  $3 \le k \le n$ , there is a k-cycle through each point x. For n = k this is true since T contains a Hamiltonian circuit by 6.11. Suppose k < n.

By 6.13, there is a point  $y \neq x$  such that T - y is strongly connected. Thus

T-y contains a k-cycle through x and hence, so does T.

Now it follows by an easy induction like the one in part (b) of the preceding solution that T contains at least n-k+1 k-cycles for  $3 \le k \le n$ . Hence the total number of cycles is

$$\geq \sum_{k=3}^{n} (n-k+1) = \sum_{\nu=1}^{n-2} \nu = {n-1 \choose 2}.$$

The assertion is sharp for the tournament obtained from a transitive one by inverting the edges of the Hamiltonian path [see M].

43. We already know there is a Hamiltonian path (see 5.20). Let  $e_1, \ldots, e_{\left[\frac{n}{2}\right]}$  be independent edges. A Hamiltonian path whose  $(2i-1)^{\text{st}}$  edge is  $e_i$   $\left(i=1,\ldots,\left[\frac{n}{2}\right]\right)$  is uniquely determined. Hence, the number of Hamiltonian paths is not greater than the number of ways to select  $\left[\frac{n}{2}\right]$  independent edges (and order them).

We have  $\binom{n}{2}$  choices for  $e_1$ ;  $\binom{n-2}{2}$  choices for  $e_2$ , etc. Thus the number of

ways to select  $e_1, \ldots, e_{\lceil \frac{n}{2} \rceil}$  is

$$\binom{n}{2}\binom{n-2}{2}\ldots\binom{n-2\left[\frac{n}{2}\right]+2}{2} = \frac{n!}{2^{\left[\frac{n}{2}\right]}}. M].$$

**44.** (a) We may assume  $n = 2^{k-1}$  and want to prove T contains a transitive subtournament with k points, using induction on k. For k = 1 the assertion is trivial.

Suppose k > 1 and choose an  $x \in V(T)$ . Let X and Y denote the sets of points z such that  $(z, x) \in E(T)$  and  $(x, z) \in E(T)$ , respectively. Since  $|X| + |Y| = 2^{k-1} - 1$ , we may assume, e.g. that  $|X| \ge 2^{k-2}$ . Then, by the induction hypothesis, X spans a transitive subtournament  $T_1$  with k-1 points. Together with  $x, T_1$  yields a transitive subtournament with k points.

(b) We prove the assertion by induction on k. For k = 1 it is obvious again. Suppose k > 1. Define

$$f(x) = \begin{cases} \prod_{j=0}^{k-2} \left( \frac{x+1}{2^j} - 1 \right) & \text{if } x \ge 2^{k-2} - 1, \\ 0 & \text{if } x \le 2^{k-2} - 1. \end{cases}$$

Then f(x) is a convex function.

Let  $V(T) = \{x_1, \ldots, x_n\}$ ,  $x_i$  having outdegree  $d_i$ . By the induction hypothesis, there are at least  $f(d_i)$  transitive (k-1)-tournaments spanned by the  $d_i$  points  $x_i$  is joined to; this gives  $f(d_i)$  transitive k-tournaments with sources in  $x_i$ . Thus the number of transitive k-tournaments is

$$\geq \sum_{i=1}^n f(d_i) \geq n \cdot f\left(rac{n-1}{2}
ight) \ \left( ext{and since} rac{n-1}{2} \geq 2^{k-2} - 1
ight) \ \geq n \prod_{j=0}^{k-2} \left(rac{n-1}{2^j} + 1
ight) = n \prod_{j=1}^{k-1} \left(rac{n+1}{2^j} - 1
ight),$$

which proves the assertion.

(c) This time we use induction on n to prove the assertion of the hint. For  $n \le k$  the assertion is true. Let n > k. By 6.13, there is a point x such that T - x is strongly connected. Thus, there are at least  $\binom{n-3}{k-2}$  k-tuples in T - x containing a 3-cycle. Also, by the solution of 10.40, there is a 3-cycle through x and there are  $\binom{n-3}{k-3}$  further k-tuples containing this triangle. This gives

$$\binom{n-3}{k-3} + \binom{n-3}{k-2} = \binom{n-2}{k-2}$$

k-tuples containing a 3-cycle as stated.

Since a transitive k-tournament contains no 3-cycle, their number is

$$\leq \binom{n}{k} - \binom{n-2}{k-2}$$
.

Note that equality holds for the tournament in the solution of 10.41 [see M].

## § 11. Spectra of graphs

1. An eigenvector of  $A_G$  assigns numbers  $x_1, \ldots, x_n$  to the points  $v_1, \ldots, v_n$  in such a way that the sum of values assigned to the neighbors of  $v_i$  is  $\lambda x_i$ . Now the graphs in consideration have such a simple structure that it is easy to guess such assignments. For the complete graph we have  $(1, \ldots, 1)$ , yielding the eigenvalue (n-1); we also have vectors of form  $(1, 0, \ldots, 0, -1, 0, \ldots, 0)$ , which are linearly independent and yield the eigenvalue (-1) with multiplicity n-1. Since there are n eigenvalues altogether, this is all.

For the star, we look for eigenvectors, which associate the value 1 with the center. If  $\lambda$  is the corresponding eigenvalue we must have  $1/\lambda$  associated with

each of the other points, so

$$(n-1)\frac{1}{\lambda}=\lambda, \qquad \lambda=\pm\sqrt{n-1}.$$

This gives two eigenvectors and two eigenvalues. To get the rest, we have to consider eigenvectors associating 0 with the center. Then the eigenvalue belonging to it must be 0. It follows that the eigenvectors have arbitrary numbers at the other points, the only restriction being that the sum of entries must be 0 [as it has to give  $0 \times (\text{number in the center})$ ]. There are n-2 linearly independent such vectors, thus we have all eigenvalues again.

Finally, if  $C = (c_1, \ldots, c_n)$  is an *n*-circuit, then let

$$x_{\mu}=\,arepsilon^{\mu},\qquad \mathbf{x}=egin{pmatrix}x_{1}\ dots\ x_{n}\end{pmatrix},$$

where  $\varepsilon$  is any  $n^{\text{th}}$  root of unity. Then we have

$$A\mathbf{x} = (x_{\nu}')_{\nu=1}^n,$$

where

$$x'_{\nu} = x_{\nu-1} + x_{\nu+1} = \varepsilon^{\nu-1} + \varepsilon^{\nu+1} = \left(\varepsilon + \frac{1}{\varepsilon}\right)\varepsilon^{\nu} = (\varepsilon + \overline{\varepsilon})x_{\nu}.$$

Thus  $\varepsilon + \bar{\varepsilon}$  is an eigenvalue and considering the *n* roots of unity, we get all of them. Hence the eigenvalues of  $C_n$  are

2, 
$$2\cos\frac{2\pi}{n}$$
,  $2\cos\frac{4\pi}{n}$ , ...,  $2\cos\frac{2(n-1)\pi}{n}$ .

**2.** (a)  $\mathbf{j} = (1, \ldots, 1)$  is an eigenvector of G with eigenvalue d, the degree of G. This follows immediately, because G is regular. Similarly,  $\mathbf{j}$  is an eigenvector of  $\overline{G}$  with eigenvalue n-1-d.

Now let  $\mathbf{v}_1 = \mathbf{j}, \mathbf{v}_2, \dots, \mathbf{v}_n$  be a basis of the *n*-space consisting of orthogonal eigenvectors of G. We claim that  $\mathbf{v}_1, \dots, \mathbf{v}_n$  are eigenvectors of  $\overline{G}$ . For

$$A_{\overline{G}}\mathbf{j}=(n-d-1)\mathbf{j},$$

which, if written out, means only that  $\bar{G}$  is (n-d-1)-regular. For the other eigenvectors we have

$$A_{\overline{G}}\mathbf{v}_i = (J - I - A_G)\mathbf{v}_i = J\mathbf{v}_i - \mathbf{v}_i - A_G\mathbf{v}_i = -\mathbf{v}_i - A_G\mathbf{v}_i = (-1 - \lambda_i)\mathbf{v}_i$$

since, by  $\mathbf{j}\mathbf{v}_i = 0$ ,  $J\mathbf{v}_i = 0$ . Thus  $\mathbf{v}_i$  is an eigenvector indeed and thus, if  $\lambda_1 \geq \ldots \geq \lambda_n$  are the eigenvalues of G, then  $n-1-\lambda_1, -1-\lambda_2, \ldots, -1-\lambda_n$  are the eigenvalues of G.

(b) The two equalities in the hint are easily verified. We use now the well-known fact that if C is an  $(m \times n)$  matrix and D is an  $(n \times m)$  matrix, then

$$\det (xI - CD) = x^{m-n} \det (xI - DC).$$

Let G have m edges and n points. Then


$$\det (xI - A_{L(G)}) = \det ((x+2)I - B_G^T B_G) =$$

 $=(x+2)^{m-n}\det\left((x+2)I-B_GB_G^T\right)=(x+2)^{m-n}\det\left((x+2-d)I-A_G\right).$  Thus if  $m\geq n$ , the eigenvalues of L(G) are

 $d-2+\lambda$  for each eigenvalue  $\lambda$  of G, with the same multiplicity, and with multiplicity m-n.

(If -d is an eigenvalue of G, then further -2's occur among the eigenvalues of the first type.) If m < n, then we have to remove n - m numbers -2 from the numbers listed first to get the spectrum of L(G) [H. Sachs; see Biggs].

(c) Figure 84 shows how to assign the edges of the complete 5-graph to the points of the Petersen graph in such a way that adjacent edges correspond to



non-adjacent points, and vice versa. Hence the Petersen graph is  $\overline{L(K_5)}$ , and by 11.1 and (a), (b), its eigenvalues are -2, -2, -2, -2, 1, 1, 1, 1, 1, 3.

3. The expansion of det  $(\lambda I - A_T)$  by its first k rows contains only two nonzero terms: One, where the  $1^{\rm st}$ ,  $2^{\rm nd}$ , ...,  $k^{\rm th}$  columns go with the first k rows, the other, where the  $1^{\rm st}$ , ...,  $(k-1)^{\rm st}$ ,  $(k+1)^{\rm st}$  columns do so (any other combination yields a column of zeros in one subdeterminant or the other.) Now the first term yields  $p_{T_1}(\lambda)$   $p_{T_2}(\lambda) = p_{T-e}(\lambda)$ ; the second term is the product of two determinants  $p_1$ ,  $p_2$ , which do not correspond to characteristic polynomials directly. But if we expand  $D_1$  (the subdeterminant composed of the  $1^{\rm st}$ , ...,  $k^{\rm th}$  rows and  $1^{\rm st}$ , ...,  $(k-1)^{\rm st}$ ,  $(k+1)^{\rm st}$  columns) by its last column we get  $p_{T_1-x_k}(\lambda)$ ; similarly,  $D_2=p_{T_2-x_{k+1}}(\lambda)$  and so, taking the sign into consideration, the second term is

$$-p_{T_1-x_k}(\lambda) p_{T_2-x_{k+1}}(\lambda) = p_{T-x_k-x_{k+1}}(\lambda)$$

[A. Mowshowitz, J. Comb. Th. B 12 (1972) 177–193; L. Lovász—J. Pelikán, Periodica Math. Hung. 3 (1973) 175–182].

**4.** We use induction on |E(T)|.

The statement is true for forests with no edges. Let  $e = (x, y) \in E(T)$ . By the previous exercise,

$$p_T(\lambda) = p_{T-e}(\lambda) - p_{T-x-y}(\lambda)$$

and by the induction hypothesis,

$$p_{T-e}(\lambda) = \lambda^{n} - a'_{1} \lambda^{n-2} + a'_{2} \lambda^{n-4} - \dots,$$
  

$$p_{T-x-y}(\lambda) = \lambda^{n-2} - a''_{1} \lambda^{n-4} + a''_{2} \lambda^{n-6} - \dots,$$

where  $a'_k$   $(a''_k)$  is the number of k-element matchings in T-e (T-x-y, resp.). Thus

$$p_T(\lambda) = \lambda^n - (a_1' + 1)\lambda^{n-2} + (a_2' + a_1'')\lambda^{n-4} - (a_3' + a_2'')\lambda^{n-6} + \dots$$

To prove the formula it suffices to show that  $a_k' + a_{k-1}'' = a_k$  (and  $a_1' + 1 = a_1$ ). Since  $a_1$  is the number of edges of T,  $a_1'$  is the number of edges of T - e,  $a_1' + 1 = a_1$  obviously. Moreover,  $a_k'$  is the number of k-element matchings in T - e, i.e. the number of those k-element matchings of T not containing e. On the other hand, the (k-1)-element matchings of T - x - y are in an obvious one-to-one correspondence with those k-element matchings of T containing e. Thus,  $a_k' + a_{k-1}''$  gives the total number of k-element matchings in T as stated [ibid.].

$$p_{P_n}(\lambda) = \lambda^n - a_1 \lambda^{n-2} + a_2 \lambda^{n-4} - \dots,$$

where  $a_k$  is the number of k-element matchings in  $P_n$ ; this means the number of ways to select the k non-neighboring edges from n-1, which is  $\binom{n-k}{k}$  by 1.31.

(b) In 1.29 the recurrence relation

$$p_{P_n}(\lambda) = p_{P_{n-1}}(\lambda) - p_{P_{n-2}}(\lambda)$$

was deduced and formula (b) was proved by it (note that this recurrence relation is a consequence of 11.3 as well). This yields a simple inductive proof of (a) whose details are left to the reader. Also in 1.29 the eigenvalues of the path were found to be

$$2\cos\frac{k\pi}{n+1}, \qquad k=1,\ldots,n.$$

[ibid.]




Fig. 85

6. Trees with no 3 independent edges have one of the forms shown in Fig. 85. Let  $T_1$  be one from the first class,  $T_2$  be one from the second class. Let  $a_k, a'_k$  denote the number of k-element matchings in  $T_1, T_2$  respectively. Then

$$a_1 = a + b + 1,$$
  $a'_1 = c + d + 2,$   $a_2 = ab,$   $a'_2 = cd + c + d.$ 

If we find values a, b > 0, c, d > 0 such that  $a_1 = a_1'$ ,  $a_2 = a_2'$ , then we are done as clearly this also implies that  $T_1, T_2$  have the same number of points  $(a_1 + 1 = a_1' + 1)$ . A little experimentation yields, e.g. the solution  $a = b = u^2 + u + 1$ ,  $c = u^2$ ,  $d = (u + 1)^2$ . (In fact, almost every tree has a "pair" with the same spectrum; see A. Schwenk, in: New Directions in the Theory of Graphs, Acad. Press, 1973, 275–307.)

7. (a) The adjacency matrix of G arises from the adjacency matrix of  $G_1$  by replacing the 1's by I, the  $|V(G_2)|$  dimensional identity matrix, and putting  $A_{G_2}$  into the diagonal. Hence  $\lambda I - A_G$  arises by substituting -I for the -1's in  $\lambda I - A_{G_1}$  and  $\lambda I - A_{G_2}$  for the  $\lambda$ 's:

$$\begin{pmatrix} \lambda & 0 & -1 & -1 \\ 0 & \lambda & 0 & 0 \\ -1 & 0 & \lambda & -1 \\ -1 & 0 & -1 & \lambda \end{pmatrix} \begin{pmatrix} \lambda I - A_{G_2} & 0 & -I & -I \\ 0 & \lambda I - A_{G_2} & 0 & 0 \\ -I & 0 & \lambda I - A_{G_2} & -I \\ -I & 0 & -I & \lambda I - A_{G_2} \end{pmatrix}$$

$$\lambda I - A_{G_1}$$

Since

$$\lambda I - A_{G_2}$$
 and I commute, this yields

$$p_G(\lambda) = \det (\lambda I - A_G) = \det p_{G_1}(\lambda I - A_{G_2}).$$

Let  $\lambda_1, \ldots, \lambda_n$  be the eigenvalues of  $G_1, \mu_1, \ldots, \mu_m$  be those of  $G_2$ . Then

$$p_G(\lambda) = \det \prod_{i=1}^n \left(\lambda I - A_{G_2} - \lambda_i I\right) = \prod_{i=1}^n \det \left((\lambda - \lambda_i)I - A_{G_2}\right) =$$

$$= \prod_{i=1}^n p_{G_i}(\lambda - \lambda_i) = \prod_{i=1}^n \prod_{j=1}^m (\lambda - \lambda_i - \mu_j).$$

Hence the eigenvalues of G are  $\lambda_i + \mu_i$   $(1 \le i \le n, 1 \le j \le m)$ .

(b) Let  $A'_G = A_G + I$ . With this notation, if  $x_i, y_i \in V(G_i)$ , then

$$\{((x_1, x_2), (y_1, y_2)) \in E(G_1 \cdot G_2) \text{ or } (x_1, x_2) = (y_1, y_2) \Leftrightarrow$$

$$((x_1, y_1) \in E(G_1) \text{ or } x_1 = y_1) \text{ and } ((x_2, y_2) \in E(G_2) \text{ or } x_2 = y_2),$$

which means that  $A'_{G_1 \cdot G_2}$  is the Kronecker product of  $A'_{G_1}$  and  $A'_{G_2}$ . The eigenvalues of  $A'_{G_1}$  and  $A'_{G_2}$  are  $\lambda_1 + 1, \ldots, \lambda_{n_1} + 1, \ldots, \mu_{1} + 1, \ldots, \mu_{n_2} + 1$ , where  $\lambda_1, \ldots, \lambda_{n_1}$ ;  $\mu_1, \ldots, \mu_{n_2}$  are the eigenvalues of  $G_1$  and  $G_2$ , respectively. Thus the eigenvalues of  $A'_{G_1 \cdot G_2}$  are

$$(\lambda_i+1)(\mu_j+1)$$
  $(1 \leq i \leq n_1, 1 \leq j \leq n_2)$ 

and the eigenvalues of  $G_1 \cdot G_2$  are

$$\lambda_i \mu_j + \lambda_i + \mu_j$$
  $(1 \le i \le n_1, 1 \le j \le n_2).$ 

A similar argument shows that the eigenvalues of the weak direct product are  $\lambda_i \mu_i$  [D. Cvetković, Univ. Beograd, Publ. Elektrotehn. Fak. Ser. Mat. Fiz. 354–356 (1971) 1–50].

8. The  $i^{th}$  coordinate of Av is

$$\sum_{j=1}^{n} a_{ij} v_{j} = \sum_{(x_{i}, x_{j}) \in E(G)} \chi(\gamma_{x_{1}, x_{j}}) = \sum_{(x_{i}, x_{j}) \in E(G)} \chi(\gamma_{x_{i}, x_{j}}) \chi(\gamma_{x_{1}, x_{i}}).$$

Take here a term and set  $\gamma_{x_ix_j}(x_1) = x_k$  (i.e.  $\gamma_{x_ix_j} = \gamma_{x_1x_k}$ ). We claim that  $(x_i, x_j) \in E(G)$  if and only if  $(x_1, x_k) \in E(G)$ . For consider  $\gamma_{x_1x_k}$ . Then  $\gamma_{x_1x_k}(x_1) = x_i$ by definition. On the other hand,

$$\gamma_{x_1x_i}(x_k) = \gamma_{x_1x_i}(\gamma_{x_ix_i}(x_1)) = \gamma_{x_ix_i}(\gamma_{x_1x_i}(x_1)) = x_i$$

(by the commutativity of  $\Gamma$ ) and so,  $(x_1, x_k)$  is mapped onto  $(x_i, x_i)$  by  $\gamma_{x_i, x_i}$ .

Thus, they are both edges or non-edges.

Thus if  $x_j$  ranges over all neighbors of  $x_i$ , the point  $x_k = \gamma_{x_i x_j}(x_1)$  ranges over all neighbors of  $x_1$  (it is different for different points  $x_j$ , because  $\Gamma$  is regular). Hence, the formula for the  $i^{\text{th}}$  coordinate of  $A\mathbf{v}$  can be rewritten as

$$\left\{\sum_{(x_1, x_k) \in E(G)} \chi(\gamma_{x_1, x_k})\right\} \chi(\gamma_{x_1, x_i})$$

showing that v is an eigenvector with eigenvalue

$$\sum_{(x_1, x_k) \in E(G)} \chi(\gamma_{x_1, x_k}).$$

We remark that, since a commutative group  $\Gamma$  has exactly  $|\Gamma|$  distinct characters (and they all have  $\chi(1) = 1$ , so that they do not give parallel vectors), the eigenvalues found above are all the eigenvalues of G. (For the case of non-commutative groups see L. Lovász, Periodica Math. Hung. 6 (1975) 191–195.)

**9.** First solution. Let us represent the points of  $Q_n$  by 01-vectors of length n, two being adjacent iff they differ at exactly one place. For each 01-vector  $(a_1, \ldots, a_n)$ , the mapping

$$\alpha_{a_1,\ldots,a_n}$$
:  $(x_1,\ldots,x_n)\mapsto (x_1+a_1,\ldots,x_n+a_n)$   
 $((x_1,\ldots,x_n)\in V(Q_n); \text{ addition mod 2})$ 

is an automorphism of  $Q_n$ . Geometrically, these are those automorphisms of  $Q_n$  preserving the directions of edges; they are reflexions in certain affine subspaces. They form a commutative, regular group  $\Gamma$ , which is (abstractly) isomorphic with  $(Z_2)^n$ .

Now the characters can be obtained this way: let  $1 \le i_1 < i_2 < \ldots < i_k \le n$ ,

and

$$\chi_{i_1,\ldots,i_k}(\alpha_{a_1,\ldots,a_n})=(-1)^{a_{i_1}+\cdots+a_{i_k}}.$$

It is easy to verify that these are characters and since their number is  $2^n = |\Gamma|$ , they are all the characters. Thus, we get an eigenvalue as follows:

$$\lambda_{i_1,\ldots,i_k} = \sum_{((a_1,\ldots,a_n),(0,\ldots,0))\in E(Q_n)} \chi_{i_1,\ldots,i_k}(\alpha_{a_1,\ldots,a_n}) =$$

$$= \sum_{((a_1,\ldots,a_n),(0,\ldots,0))\in E(Q_n)} (-1)^{a_{i_1}+\cdots+a_{i_k}}.$$

By the definition of  $Q_n$ , the sequences  $(a_1, \ldots, a_n)$  considered here are those having exactly one non-zero entry. Hence

$$\lambda_{i_1,\ldots,i_k}=n-2k.$$

Thus, the eigenvalues of  $Q_n$  are

$$n, n-2, \ldots, -n$$
; the multiplicity of  $n-2k$  is  $\binom{n}{k}$ .

Second solution.  $Q_n$  is the Cartesian product of n copies of  $K_2$ . Hence by 11.7a, its eigenvalues arise in the form  $\lambda_1 + \ldots + \lambda_n$ , where  $\lambda_i = \pm 1$  are eigenvalues of  $K_2$ . We conclude as before.

10. The mappings

$$\varphi_{v}: \varepsilon^{k} \to \varepsilon^{k+v}$$

are automorphisms of G, forming a cyclic group  $\Gamma$  of order p, which acts transitively on V(G). The characters of  $\Gamma$  are given by

$$\chi_t(\varphi_v) = \varepsilon^{tv}$$
  $(t=0,\ldots,p-1; \quad v=0,\ldots,p-1),$ 

where  $\varepsilon = e^{\frac{2\pi i}{p}}$ . Thus by 11.8, the eigenvalues of G are given by

$$\lambda_t = \sum_{i=1}^k \varepsilon^{ta_i}$$
  $(t = 0, \dots, p-1),$ 

where 1 is adjacent to  $\varepsilon^{a_1}, \ldots, \varepsilon^{a_k}$ . Similarly, the eigenvalues of H are

$$\mu_t = \sum_{i=1}^k \varepsilon^{tb_i} \qquad (t=0,\ldots,p-1),$$

where  $\varepsilon^{b_1}, \ldots, \varepsilon^{b_k}$  are the neighbors of 1. Since G and H must have the same spectra, we have

or, equivalently,

$$\lambda_1 = \mu_t \quad (0 \le t \le p - 1)$$

$$\sum_{i=1}^k (\varepsilon^{a_i} - \varepsilon^{tb_i}) = 0.$$

Let  $\overline{a}$  denote the residue of  $a \mod p$ ,  $0 \le \overline{a} \le p-1$ . Then  $\varepsilon$  is a root of the polynomial

$$f(x) = \sum_{i=1}^{k} (\bar{x^{a_i}} - \bar{x^{tb_i}}).$$

Since the minimal polynomial of  $\varepsilon$  is  $1+x+\ldots+x^{p-1}$ , it follows that

$$1+x+\ldots+x^{p-1}|f(x).$$

Since  $\deg f \leq p-1$ , we get;

$$f(x) = c(1 + x + \ldots + x^{p-1}).$$

From f(0) = 0, we deduce that c = 0, i.e. f(x) = 0. But this means that in f(x) all terms cancel out, i.e.

$$\{\overline{a}_j: 1 \leq j \leq k\} = \{t\overline{b}_j: 1 \leq j \leq k\}$$

proving the assertion [D.Z. Djoković, Acta Math. Acad. Sci. Hung. 21 (1970) 267–277; B. Elspas–J. Turner, J. Comb. Th. 9 (1970) 297–307].

11. Let  $V(G) = \{x_1 \dots x_n\}$ .

If det  $A \neq 0$ , we have a non-zero expansion term of det A; say  $a_{1,i_1} \dots a_{n,i_n} \neq 0$ , where  $(i_1, \dots, i_n)$  is a permutation of  $(1, \dots, n)$ . Thus  $(x_v, x_{iv}) \in E(G)$   $(i = 1, \dots, n)$ . If say,  $x_1, \dots, x_k$   $(k \geq n/2)$  form one of the two color-classes of G, then the edges

$$(x_1, x_{i_1}), \ldots, (x_k, x_{i_k})$$

are independent and their number is  $k \ge \frac{n}{2}$ , thus they have to form a 1-factor of G.

12. Let  $G_1, \ldots, G_m$  be vertex-disjoint subgraphs as shown in Fig. 10 (see p. 72), which cover all points. Let S be the set of endpoints of  $G_1 \cup \ldots \cup G_m$  and T = V(G) - S; also set  $V(G) = \{v_1, \ldots, v_n\}$  (n = 6m). Then each point is adjacent to two points of S and to one point of T. Hence if we set

$$x_i = \left\{ egin{array}{cccc} 2 & ext{if} & v_i \in T, \ & & & \mathbf{x} = \left(egin{array}{c} x_1 \ dots \ x_n \end{array}
ight), \end{array} 
ight.$$

we have

$$A\mathbf{x} = 0$$

showing that  $\lambda = 0$  is an eigenvalue [H. Sachs. Wiss. Z.TH Ilmenau, 19 (1973) 83–99].

13. We may assume that G' is a spanning subgraph of G; otherwise, we can add the remaining points of G to G' as isolated points and this only adds 0's to the spectrum of G', thus does not change the maximum eigenvalue of it.

Let  $\mathbf{v}$  be an eigenvector of  $A_{G'}$ , with eigenvalue  $\Lambda'$  and with non-negative entries. We may assume  $|\mathbf{v}| = 1$ , then

$$A' = \mathbf{v}^T A_{G'} \mathbf{v} \leq \mathbf{v}^T A_G \mathbf{v}$$

as v has non-negative entries and  $A_G \geq A_{G'}$ . On the other hand,

$$\Lambda = \max_{|\mathbf{x}|=1} \mathbf{x}^T A_G \mathbf{x} \ge \mathbf{v}^T A_G \mathbf{v} \ge \Lambda'.$$

14. (a) I. Let  $\mathbf{j}$  be the vector of length n all of whose entries are 1. Then  $A_G \mathbf{j}$  is a vector whose entries are the degrees and so,

 $A_G \mathbf{j} \ge d\mathbf{j}, \quad \mathbf{j}^T A_G \mathbf{j} \ge d\mathbf{j}^T \mathbf{j} = dn.$ 

Thus

$$\Lambda = \max_{\mathbf{x} \neq 0} \frac{\mathbf{x}^T A_G \mathbf{x}}{\mathbf{x}^T \mathbf{x}} \ge \frac{\mathbf{j}^T A_G \mathbf{j}}{\mathbf{j}^T \mathbf{j}} \ge d.$$

II. Let G' denote the star of a point of maximum degree. Then, by 11.1, the maximum eigenvalue of G' is  $\sqrt{D}$ . By 11.13,

$$\Lambda \geq \sqrt{D}$$
.

III. We have  $A_G \mathbf{j} \leq D \mathbf{j}$  by the definition of D.

Let  $\mathbf{v}$  be an eigenvector belonging to D; we may assume the maximum coordinate of it, say the first one, is equal to 1. Then

$$\Lambda \mathbf{v} = A_G \mathbf{v} \le A_G \mathbf{j} \le D \mathbf{j},$$

so considering the first coordinates

$$\Lambda \leq D$$
.

IV. Suppose we have equality here. Then  $A_G \mathbf{v}$  and  $A_G \mathbf{j}$  have the same first coordinate D, i.e.

$$\sum_{(x_1,x_i)\in E(G)} v_i = D.$$

Since  $v_i = 1$  this can only happen if each  $x_i$  adjacent to  $x_1$  has the same value  $v_i = 1$ . Then the above argument applies to these points and yields that their neighbors  $x_j$  also have  $v_j = 1$  and so on; since G is connected, we conclude that  $v_i = 1$  for every point. Thus  $\mathbf{v} = \mathbf{j}$ , and then

$$A_G \mathbf{j} = A \mathbf{j} = D \mathbf{j}$$

yields that G is D-regular.

(b) The two identities in the hint are easily seen: If  $\lambda_1 \leq \ldots \leq \lambda_n = \Lambda$  are the eigenvalues of G, then  $\sum_{i=1}^n \lambda_i = \operatorname{Tr} A_G = 0$ ,  $\sum_{i=1}^n \lambda_i^2 = \operatorname{Tr} A_G^2 = 2m$ . Hence

$$\Lambda^{2} = (\lambda_{1} + \ldots + \lambda_{n-1})^{2} \leq (n-1)(\lambda_{1}^{2} + \ldots + \lambda_{n-1}^{2}) = (n-1)(2m - \Lambda^{2}),$$

whence the upper bound follows. The lower bound follows by the argument of I.

15. The formula

$$A_{L(G)} = B^T B - 2I$$

follows easily.

Since  $B^TB$  is positive semidefinite, all of its eigenvalues are non-negative. Hence, the eigenvalues of  $A_{L(G)}$  are  $\geq -2$ . Moreover, if |V(G)| < |E(G)|, then

$$r(B^TB) = r(B) \le |V(G)| < |E(G)|$$

(r(X) is the rank of the matrix X). So,  $B^TB$  has at least one 0 eigenvalue, i.e.  $A_{L(G)}$  has at least one -2 eigenvalue. [A.J. Hoffman, in: Beiträge zur Graphentheorie, Teubner (1968) 75–80.]

**16.** Let  $\alpha$  be an automorphism of G; define a matrix  $P = (p_{ij})$  by

$$p_{ij} = egin{cases} 1 & ext{if} & lpha(v_i) = v_j, \ 0 & ext{otherwise.} \end{cases}$$

Then P is a permutation matrix, i.e. each row and column of it contains exactly one 1. Hence  $P^{-1} = P^{T}$ .

The fact that  $\alpha$  is an automorphism is equivalent to the relation

$$P^{-1}A_GP = A_G$$

For if  $A_G = (a_{ij})$ , then the (i, j)<sup>th</sup> entry of  $P^{-1}A_GP$  is

$$\sum_{\nu} \sum_{\mu} p_{\nu i} a_{\nu \mu} p_{\mu j} = a_{i',j'},$$

where i', j' are determined by  $\alpha(x_{i'}) = x_i$ ,  $\alpha(x_{j'}) = x_j$ . Thus  $\alpha$  maps the pair  $(x_{i'}, x_{j'})$  onto the pair  $(x_i, x_j)$ , and it is an automorphism iff

$$a_{i'j'} = a_{ij} \qquad (i, j = 1, \ldots, n)$$

as stated.

Now  $A_G$  is symmetric, hence  $A_G = T^{-1}LT$ , where

$$L = egin{bmatrix} \lambda_1 & & & 0 \ & \lambda_2 & & & \ & & & \ddots & \ & & & & \lambda_n \ \end{pmatrix}$$

and T is an orthogonal matrix. Thus

$$P^{-1}T^{-1}LTP = T^{-1}LT,$$

or

$$L(TPT^{-1}) = (TPT^{-1})L.$$

Multiplying a matrix by L from the left (from the right) means to multiply its  $i^{\text{th}}$  row ( $i^{\text{th}}$  column) by  $\lambda_i$  for each  $1 \leq i \leq n$ . Since  $\lambda_i \neq \lambda_j$  for  $i \neq j$  by the assumption, this yields that the matrix  $TPT^{-1}$  is a diagonal matrix, say

$$TPT^{-1} = egin{bmatrix} t_1 & & \mathbf{0} \ & \cdot & & \ & \cdot & & \ & \mathbf{0} \ & & \cdot & \ & & \cdot & \ & & \cdot & \ & & & t_n \ \end{pmatrix},$$

Here, however, P and T are orthogonal matrices and thus, so is  $TPT^{-1}$ . Hence  $t_i = \pm 1$ . But then

$$TP^2T^{-1} = (TPT^{-1})^2 = I, \qquad P^2 = I.$$

Now it is easy to see that  $P^2$  is the permutation matrix associated with  $\alpha^2$ , whence  $\alpha^2 = 1$ . [A. Mowshowitz, in: *Proof Techn. in Graph Th.* Academic Press (1969) 109–110; M. Petersdorf—H. Sachs, in: *Combin. Th. Appl.* Coll. Math. Soc. J. Bolyai 4, Bolyai–North-Holland, (1970) 891–907. For directed graphs cf. C.Y. Chao, *J. Comb. Th.* 3 (1971) 301–302.]

17. Since by 11.16, all elements of  $\Gamma$  are of order 2,  $\Gamma$  is commutative  $((xy)^2 = 1 = x^2y^2)$  and hence, xy = yx. Suppose that  $\Gamma$  is transitive on the points, then it follows that it is regular. Thus 11.8 applies and we get that any eigenvalue of  $\Gamma$  is of the form

$$\lambda = \sum_{(\nu_1, \nu_i) \in E(G)} \chi(\gamma_{\nu_1, \nu_i}),$$

where  $\chi$  is a character of  $\Gamma$ . Moreover, since  $\chi(\gamma)^2 = \chi(\gamma^2) = \chi(1) = 1$  for each  $\gamma$ , the summands here are  $\pm 1$ 's. Thus  $\lambda$  is an integer; if d is the degree of points of G, then  $\lambda$  is the sum of d + 1's and hence,

$$-d \le \lambda \le d$$
,  $\lambda \equiv d \pmod{2}$ .

This leaves exactly d+1 values for  $\lambda$ . Since there are |V(G)| eigenvalues, all different, we have  $|V(G)| \leq d+1$ . Since G is simple, it must be a complete graph, but this is impossible again except for  $G \cong K_2$ , since  $K_n$  has n-1 equal eigenvalues (see 11.1) [ibid.].

18. Let  $G_1$ ,  $G_2$  be the subgraphs induced by a maximum clique and a maximum independent set, respectively. Then  $A_{G_1}$ ,  $A_{G_2}$  are symmetric submatrices of  $A_G$ . Since  $G_1$  has eigenvalues

$$\underbrace{-1,\ldots,-1}_{\omega(G)-1},\,\omega(G)-1,$$

we have, by the theorem mentioned in the hint,

$$\lambda_{\omega(G)-1} \leq -1, \qquad \lambda_{n-\omega(G)+1} \geq -1.$$

Similarly,  $G_2$  has eigenvalues  $0, \ldots, 0$  and therefore

$$\lambda_{\alpha(G)} \leq 0, \qquad \lambda_{\tau(G)+1} \geq 0.$$

19. (a) Suppose first that  $-\Lambda$  is an eigenvalue.

Let  $\mathbf{w} = (w_1, \dots, w_n)^T$  be the eigenvector belonging to  $-\Lambda$  and set  $\mathbf{w}' = (|w_1|, \dots, |w_n|)^T$ ,  $A_G = (a_{ij})$ . Then

$$|\mathbf{w}_{-}^{T}A_{G}\mathbf{w}| = |-\Lambda\mathbf{w}^{T}\mathbf{w}| = \Lambda\mathbf{w}^{T}\mathbf{w};$$

<sup>†</sup> Suppose  $\varphi \in \Gamma$  fixes a point x. Then by  $\varphi(\alpha(x)) = \alpha(\varphi(x)) = \alpha(x)$ , it follows that it fixes every point  $\alpha(x)$ ,  $\alpha \in \Gamma$ . Since  $\Gamma$  is transitive,  $\varphi = 1$ .

on the other hand,

(2) 
$$\mathbf{w}^T A_G \mathbf{w} = \left| \sum_i \sum_j a_{ij} w_i w_j \right| \le \sum_i \sum_j a_{ij} |w_i| |w_j| = \mathbf{w}^T A_G \mathbf{w}^T \le \Lambda \mathbf{w}^T \mathbf{w}^T = \Lambda \mathbf{w}^T \mathbf{w}.$$

(since  $\Lambda$  is the maximum of  $\mathbf{u}^T A_G \mathbf{u} / \mathbf{u}^T \mathbf{u}$ ).

Comparing (1) and (2), we find that we must have equality everywhere in (2). Thus,  $\mathbf{w}'$  is an eigenvector belonging to  $\Lambda$  and since G is connected, no entry of  $\mathbf{w}'$  is 0. Moreover, all non-zero terms in

$$\sum_{i} \sum_{j} a_{ij} w_i w_j = -\Lambda \mathbf{w}^T \mathbf{w}$$

must have the same sign and since the sum is negative, all summands are negative. Thus, if  $a_{ij} \neq 0$ ,  $w_i$  and  $w_j$  have different signs. So, if

$$S_1 = \{x_i : w_i > 0\} \quad S_2 = \{x_i : w_i < 0\},$$

then  $\{S_1, S_2\}$  is a 2-coloration of G.

Conversely, let G be bipartite and let  $\mathbf{w} = (w_1, \dots, w_n)$  be the eigenvector belonging to the eigenvalue  $\Lambda$ . Let  $S_1, S_2$  be a 2-coloration of G and set

$$w'_i = \begin{cases} w_i & \text{if } x_i \in S_1, \\ -w_i & \text{if } x_i \in S_2, \end{cases}$$
$$\mathbf{w}' = (w'_1, \dots, w'_n)^T.$$

Then the  $i^{th}$  coordinate of  $A_G \mathbf{w}'$  is

$$\sum_{i} a_{ij} w'_{j} = -\sum_{i} a_{ij} w_{j} = -\Lambda w_{i} = -\Lambda w'_{i},$$

if  $x_i \in S_1$  (since all j's here for which  $a_{ij} \neq 0$  have  $x_j \in S_2$ ), and similarly,

$$\sum_{j} a_{ij} w_i' = \sum_{j} a_{ij} w_j = \Lambda w_i = -\Lambda w_j',$$

if  $x_i \in S_2$ . Thus w' is an eigenvector and the corresponding eigenvalue is  $-\Lambda$ . The same idea proves the "only if" part of (b) too.

(b) Suppose the spectrum is symmetric with respect to the origin. If G is connected (a) proves the theorem. We are going to reduce the general case

to this, using induction on the number of components.

Note that the spectrum of G is the union of the spectra of its components (in the sense that, if  $\lambda$  occurs in the spectra of components with multiplicity  $m_1, \ldots, m_r$ , then it has multiplicity  $m_1 + \ldots + m_r$  in the spectrum of G). Let A be the largest eigenvalue and let  $G_1$  be a component such that -A is an eigenvalue of  $G_1$ . Then the largest eigenvalue of  $G_1$  is  $\geq |-A| = A$  and, since it cannot be greater than this, it is A. Hence, applying (a) to  $G_1$ , we get that  $G_1$  is bipartite. Hence, its spectrum is symmetric with respect to the origin. We obtain the spectrum of  $G - G_1$  by removing the spectrum of  $G_1$  from the spectrum of G. Both these spectra being symmetric, so is the spectrum of  $G - G_1$ . By induction,  $G - G_1$  is bipartite. So is  $G_1$  as we have seen and therefore, so is G.

20. We use induction on the number of points.

Set  $k = [\Lambda] + 1$ . Let x be a point of minimum degree d. By 11.14,  $d \leq \Lambda$ , so  $d \leq k - 1$ . Let  $\Lambda'$  be the maximum eigenvalue of G - x. By 11.13,  $\Lambda' \leq \Lambda$ , so by the induction hypothesis

$$\chi(G-x) \le \Lambda' + 1 \le \Lambda + 1$$
, so  $\chi(G-x) \le k$ .

Let us k-color G - x. Since d = k - 1, this coloration can be extended to x, and so,  $\chi(G) \le k \le A + 1$  [H.S. Wilf; see Biggs].

**21.** (a) It follows from  $\chi(G) = k$  that for a suitable choice of indices,  $A_G$  can be written in the form

$$egin{pmatrix} 0 & A_{12} & \dots & A_{1k} \ A_{21} & 0 & & A_{2k} \ \vdots & \vdots & \ddots & & & \ \vdots & \vdots & \ddots & & & \ A_{k1} & A_{k2} & & 0 \ \end{pmatrix},$$

where  $A_{ij}$  is an  $m_i \times m_j$  matrix and, of course,  $A_{ji} = A_{ij}^T$ ; here  $m_i$  is the number of points with color i.

Let **v** be an eigenvector belonging to  $\Lambda$ . Let us break **v** into pieces  $\mathbf{v}_1, \ldots, \mathbf{v}_k$  of lengths  $m_1, \ldots, m_k$ , respectively. Set

$$\mathbf{w}_i = egin{bmatrix} |\mathbf{v}_i| \ 0 \ \vdots \ \vdots \ 0 \end{bmatrix} \quad m_i \; \; ext{entries}, \quad \mathbf{w} = egin{bmatrix} \mathbf{w}_1 \ \vdots \ \mathbf{w}_k \end{bmatrix}.$$

Let  $B_i$  be any orthogonal matrix such that

$$\mathbf{w}_i B_i = \mathbf{v}_i \qquad (i = 1, \dots, k),$$

and

$$B=egin{bmatrix} B_1 & & & \mathbf{0} \ & B_2 & & & \ & & \ddots & & \ & & & \ddots & & \ \mathbf{0} & & & B_k \end{pmatrix}.$$

Then  $B\mathbf{w} = \mathbf{v}$  and

$$B^{-1}AB\mathbf{w} = B^{-1}A\mathbf{v} = AB^{-1}\mathbf{v} = A\mathbf{w},$$

so w is an eigenvector of  $B^{-1}AB$ .

Moreover,  $B^{-1}AB$  has the form

$$\begin{bmatrix} 0 & B_1^{-1}A_{12}B_2 & \dots & B_1^{-1}A_{1k}B_k \\ B_2^{-1}A_{21}B_1 & 0 & B_2^{-1}A_{2k}B_k \\ \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ B_k^{-1}A_{k1}B_1 & B_k^{-1}A_{k2}B_2 & \dots & 0 \end{bmatrix}.$$

Pick the entry in the upper left corner of each of the  $k^2$  submatrices  $B_i^{-1}A_{ij}B_j$   $(A_{ii}=0)$ , these form a  $k \times k$  submatrix D. Now observe that

$$\mathbf{u} = egin{pmatrix} |\mathbf{v}_1| \ dots \ dots \ |\mathbf{v}_k| \end{pmatrix}$$

is an eigenvector of D; for **w** is an eigenvector of  $B^{-1}AB$  and has 0 entries on places corresponding to those rows and columns of  $B^{-1}AB$ , which are to be deleted to get D. Moreover, the eigenvalue belonging to **u** is  $\Lambda$ .

Now let  $\mu_1 \leq \ldots \leq \mu_k$  be the eigenvalues of D. Since D has 0's in its main

diagonal,

$$\mu_1+\ldots+\mu_k=0.$$

On the other hand,  $\Lambda$  is an eigenvalue of D and so

$$\Lambda \leq \mu_k$$
,

while by a theorem quoted in the proof of 11.18,

$$\lambda_1 \leq \mu_1, \ldots, \lambda_{k-1} \leq \mu_{k-1}.$$

Thus

$$\lambda_1 + \ldots + \lambda_{k-1} \leq \mu_1 + \ldots + \mu_{k-1} = -\mu_k \leq -\Lambda$$

[A.J. Hoffman, in: Graph Th. Appl. Academic Press (1970) 79-91].

(b) Let the k-coloration of G have  $a_i$  points of the  $i^{\text{th}}$  color  $(i=1,\ldots,k)$ . Then the  $a_i$  rows of the matrix  $A_G$  corresponding to the  $i^{\text{th}}$  color are equal, hence the rank of  $A_G$  is at most k and n-k of the eigenvalues are 0; moreover if we denote by  $\mathbf{w}_i$  the vector with

$$\mathbf{w}_i = (w_{ij}),$$
  $w_{ij} = \begin{cases} 1 & ext{if the } j^{ ext{th point has color }} i \\ 0 & ext{otherwise,} \end{cases}$ 

then the eigenvectors belonging to the n-k 0's are orthogonal to every  $\mathbf{w}_{i}$ . We look now for eigenvectors of the form

$$\mathbf{w} = \sum_{i=1}^k x_i \mathbf{w}_i;$$

these will be different from the above mentioned ones and if we find k of them (i.e. k eigenvalues belonging to them) we have all. (1) is an eigenvector with eigenvalue  $\lambda$  if and only if

$$\sum_{\nu \neq i} a_{\nu} x_{\nu} = \lambda x_{i},$$

i.e. if  $\lambda$  is an eigenvalue and  $\begin{pmatrix} x_1 \\ \vdots \\ x_k \end{pmatrix}$  is a corresponding eigenvector of

$$A' = egin{bmatrix} 0 & a_2 & a_3 & \dots & a_k \ a_1 & 0 & a_3 & \dots & a_k \ & & & & & \ & & & & \ & & & & \ & & & & \ & & & & \ & & & & \ & & & & \ & & & & \ & & & & \ & & & & \ & & & & \ & & & & \ & & & & \ & & & \ & & & & \ & & & & \ & & & \ & & & & \ & & & \ & & & & \ & & & \ & & & & \ & & & \ & & & & \ & & & \ & & & \ & & & \ & & & \ & & & \ & & & \ & & & \ & & & \ & & & \ & & & \ & & \ & & & \ & & \ & & & \ & & \ & & & \ & & \ & & \ & & \ & & \ & & \ & & \ & & \ & & \ & \ & & \ & & \ & \ & \ & \ & \ & & \$$

We claim that every eigenvalue of A' except the largest one is non-positive. In order to show this write (2) as

$$\sum_{\nu=1}^k a_{\nu} x_{\nu} = (\lambda + a_i) x_i.$$

If  $\lambda = -a_i$  for some i, then we have nothing to show. Otherwise,

$$\sum_{\nu=1}^k a_{\nu} x_{\nu} \neq 0$$

and hence, we may assume

$$\sum_{\nu=1}^k a_{\nu} x_{\nu} = 1.$$

Then (3) yields

$$x_i = \frac{1}{\lambda + a_i}$$
  $(i = 1, \ldots, k),$ 

whence by (4)

$$\sum_{\nu=1}^{k} \frac{a_{\nu}}{\lambda + a_{\nu}} = 1.$$

Now it is easy to see that (5) has exactly one positive root. Thus all but one of the eigenvalues of A' are non-positive.

Thus if  $\lambda_1, \ldots, \lambda_{k-1}$  are the k least eigenvalues of  $A_G$  and  $\Lambda$  is the largest one, then  $\lambda_1, \ldots, \lambda_{k-1}$ ,  $\Lambda$  are exactly the eigenvalues of A'. Since A' has 0's in its diagonal,

$$\lambda_1+\ldots+\lambda_{k-1}+\Lambda=0,$$

which proves the assertion.

**22.** Let  $G = G_1 \cup \ldots \cup G_m$ , where  $G_i$  is a complete bipartite graph. We add all points of  $V(G) - V(G_i)$  as isolated points to  $G_i$ . Then

$$A_G = A_{G_1} + \ldots + A_{G_m}, \quad \mathbf{x}^T A_G \mathbf{x} = \sum_{i=1}^m \mathbf{x}^T A_{G_i} \mathbf{x}.$$

Let  $V(G) = \{1, \ldots, n\}$  and denote by  $\{A_i, B_i\}$  the 2-coloration of  $G_i$ . Then

$$\mathbf{x}^T A_{G_i} \mathbf{x} = \sum_{v \in A_i} \sum_{\mu \in B_i} x_v x_\mu = \frac{1}{4} (\sum_{v \in A_i} x_v + \sum_{\mu \in B_i} x_\mu)^2 - (\sum_{v \in A_i} x_v - \sum_{\mu \in B_i} x_\mu)^2$$

and thus

$$\mathbf{x}^T A_G \mathbf{x} = \sum_{i=1}^m \frac{1}{4} \left( \sum_{\nu \in A_i} x_{\nu} + \sum_{\mu \in B_i} x_{\mu} \right)^2 - \sum_{i=1}^m \frac{1}{4} \left( \sum_{\nu \in A_i} x_{\nu} - \sum_{\mu \in B_i} x_{\mu} \right)^2.$$

Thus  $\mathbf{x}^T A_G \mathbf{x}$  is the sum of m positive and m negative squares. Hence, as is well known,  $m \ge k$  and  $m \ge l$  both follow.

(Let U denote the subspace of vectors x such that

$$\sum_{\nu \in A_i} x_{\nu} = \sum_{\mu \in B_i} x_{\mu} \qquad (i = 1, \dots, m)$$

and V the subspace spanned by the eigenvectors with negative eigenvalue. Then clearly  $U \cap V = 0$  and so

$$l = \dim V \le n - \dim U \le n - (n - m) = m.$$

R.L. Graham, H.O. Pollak, in: Graph Th. Appl. (Lecture Notes in Math. 303, Springer 1972) 99-110.1

23. Let k be the number of distinct eigenvalues of G and assume indirectly that the diameter of G is at least k. Let  $f(x) = (x - \lambda_1) \dots (x - \lambda_k) = x^n + x^$  $+a_{n-1}x^{n-1}+\ldots+a_0$ , where  $\lambda_1,\ldots,\lambda_k$  are all the distinct eigenvalues of G. As is well known

(1) 
$$f(A_G) = A_G^k + a_{n-1} A_G^{k-1} + \dots + a_0 I = 0.$$

Now let  $v_i$ ,  $v_i$  be two points at distance k (these exist as the diameter of G is at least k). Observe that the  $(i, j)^{th}$  entry of  $A_G^m$  is the number of  $(v_i, v_j)$ -walks of length m; because it is

$$\sum_{i_1=1}^n \sum_{i_2=1}^n \dots \sum_{i_{m-1}=1}^n a_{i,i_1} a_{i_1i_2} \dots a_{i_{m-1}j}$$

and this term is 1 if  $(v_i, v_{i_1}, \ldots, v_{i_{n-1}}, v_j)$  is a walk and 0 otherwise. Thus, the  $(i, j)^{th}$  entry in  $A_G^k$  is positive, but it is 0 in all the other terms in (1), which is a contradiction [see Biggs].

- 24. We consider the following three statements:
  - (i) G is connected and d-regular.
  - (ii) There exists a polynomial f such that  $f(A_G) = J$ ;
- (iii) Every eigenvector of  $A_G$  is an eigenvector of J, i.e. either parallel or orthogonal to j.

(ii)  $\Rightarrow$  (iii). Let **u** be any eigenvector of  $A_G$ ; it is an eigenvector of any polynomial of  $A_G$ , in particular of J.

(iii)  $\Rightarrow$  (i). Since the eigenvectors of  $A_G$  span the whole space, one of them must be **j**. Let d denote the eigenvalue belonging to **j**, then

$$A_G \mathbf{j} = d\mathbf{j}$$

expresses exactly the fact that G is d-regular.

Suppose G is disconnected, say  $G = G_1 \cup G_2$ ,  $G_1 \cap G_2 = \emptyset$ . Define  $\mathbf{u} = (u_i)$  by

$$u_i = \begin{cases} 1 & \text{if} \quad v_i \in V(G_1), \\ 0 & \text{otherwise.} \end{cases}$$

Then u is an eigenvector of  $A_G$  but not of J, a contradiction.

(i)  $\Rightarrow$  (ii). Since G is d-regular,  $\mathbf{j}$  is an eigenvector with eigenvalue d. Let  $\mathbf{u}_1 = \mathbf{j}, \mathbf{u}_2, \ldots, \mathbf{u}_n$  be a basis of eigenvectors of  $A_G$ , with corresponding eigenvalues  $d = \lambda_1, \ldots, \lambda_n$ . As is well-known,  $\mathbf{j}^T \mathbf{u}_i = 0$  ( $2 \leq i \leq n$ ). Consider a polynomial f(x) such that

$$f(d) = n, \quad f(\lambda_i) = 0 \qquad (2 \le i \le n).$$

Since G is connected, d is a single eigenvalue and this definition makes sense. Then

$$f(A_G)\mathbf{j} = n\mathbf{j}, \quad f(A_G)\mathbf{u}_i = 0 \quad (2 \le i \le n),$$

and therefore, the action of  $f(A_G)$  coincides with the action of J on a basis; hence,  $f(A_G) = J$  [A.J. Hoffman, Amer. Math. Monthly 70 (1963) 30–36].

25. From the basic properties of projective planes, we get

$$A_L^2 = egin{pmatrix} J+q\,I & 0 \ 0 & J+q\,I \end{pmatrix}$$
 ,

where q is the order of the plane. Hence the eigenvalues of  $A_L^2$  are:  $(q+1)^2$  (twice) and  $q(2(q^2+q))$  times). Now the spectrum of L is symmetric with respect to the origin (see 11.19), hence L has eigenvalues

$$q+1$$
 with multiplicity 1,  $\sqrt[q]{q}$  "  $q^2+q$ ,  $-\sqrt[q]{q}$  "  $q^2+q$ ,  $-q-1$  " 1

[A.J. Hoffman, Proc. Amer. Math. Soc. 16 (1965) 297-302].

**26.** (a) (i) expresses the fact that each diagonal entry of  $A_G^2$  is d; (ii) expresse the fact that the off-diagonal entries are b. Hence

$$A_G^{2} = bJ + (d-b)I.$$

Hence the eigenvalues of  $A_G^2$  are nb+d-b, d-b, ..., d-b. Note that G is connected by (ii) and is not bipartite, since again by (ii), it contains triangles. Hence 11.19 implies that  $-\sqrt{nb+d-b}$  is not an eigenvalue of G. Thus

 $\sqrt{nb+d-b}$ ,  $\pm\sqrt{d-b}$  are the different eigenvalues of G. Since by (i), d is the largest eigenvalue of G, we must have

(1) 
$$\sqrt{nb+d-b} = d, \quad n = \frac{d^2-d}{b} + 1.$$

Suppose G has k eigenvalues equal to  $\sqrt{d-b}$  and n-1-k eigenvalues equal to  $-\sqrt{d-b}$ . Since the sum of eigenvalues of G is 0, we have

$$-(n-1-2k) \sqrt{d-b} + d = 0.$$

Hence  $\sqrt{d-b}$  is rational, consequently integer, say  $\sqrt{d-b}=r$ . Then

$$(2) d = r^2 + b$$

and

$$r(n-1-2k) = r^2 + b$$
.

Hence

Also, by (1),

$$b | d^2 - d = r^4 + 2r^2b + b^2 - r^2 - b$$

thus

(4) 
$$b | r^2(r^2 - 1)$$
.

(b) Now if b = 1 (3) implies that r = 1 and hence, by (2), d = 2. Thus

from (1) we get n = 3, i.e.  $G = K_3$ .

Note that (3), (2), and (1) imply that  $n \le b^2(b+2)$  and hence there is a finite number of such graphs for fixed b; (4) shows that one need not even consider all divisors of b in (3) [S. S. Shrikhande, see Biggs].

(c) Let the points of G be the non-vertical lines of the affine plane over  $GF(2^k)$ , and let us connect two of them if they intersect and the intersection point is on one of  $2^{k-1}$  prescribed vertical lines  $L_1, \ldots, L_{2^{k-1}}$ . Set  $L = L_1 \cup \ldots \cup L_{2^{k-1}}$ . If two non-vertical lines E, F do not intersect in L, then there are  $2^{k-1}(2^{k-1}-1)$  other non-vertical lines meeting both E, F in L. If E, F intersect in a point  $p \in L$ , then there are  $2^k-2$  other non-vertical lines through p and  $(2^{k-1}-1)(2^{k-1}-2)$  non-vertical lines avoiding p, which meet both E, F in L. This gives again

$$2^{k} - 2 + (2^{k-1} - 1)(2^{k-1} - 2) = 2^{k-1}(2^{k-1} - 1)$$

common neighbors of E, F. Thus G has the desired property.

## § 12. Automorphisms of graphs

1. Figure 63 (see p. 344) shows how to associate lines of  $K_5$  with the points of the Petersen graph P so that adjacent points are associated with non-adjacent lines. So  $P \cong \overline{L(K_5)}$  and the automorphism group of P is the same as that of  $L(K_5)$ . The automorphism group of  $L(K_5)$  is isomorphic with the

automorphism group of  $K_5$ . For obviously, each automorphism of  $K_5$  induces an automorphism of its line-graph. Conversely, let  $\alpha$  be an automorphism of  $L(K_5)$ . Observe that any 4-element clique in  $L(K_5)$  corresponds to a star in  $K_5$  and vice versa, so  $L(K_5)$  has exactly five 4-element cliques, and any point in  $L(K_5)$  is uniquely characterized as the intersection of some two of them. Thus,  $\alpha$  induces a permutation of the five 4-element cliques of  $L(K_5)$  and this induces an automorphism of  $K_5$ .

Thus, the automorphism group of P is isomorphic with the automorphism group of  $K_5$ , which is  $S_5$  [R. Frucht, Comment. Math. Helv. 9 (1936-37)

217-223].

2. (a) Each automorphism of this graph is a congruence of the dodecahedron. This follows from the observation that for any mapping of the star of a point onto the star of any other point there is a unique automorphism extending it; also, there is a unique congruence of the dodecahedron and so, the automorphism must be the same as the one induced by this congruence.

This observation also shows that the dodecahedron has 120 congruences.

Consider the cubes shown in Fig. 16, p. 127. Observe that each diagonal of any face is an edge of a unique cube and each cube contains exactly one diagonal of each face. Thus, the number of such cubes is exactly 5. Let  $Q_1, \ldots, Q_5$  be these cubes. Each congruence  $\alpha$  of the dodecahedron induces a permutation  $\overline{\alpha}$  of  $Q_1, \ldots, Q_5$ . Suppose  $\overline{\alpha} = 1$  for some  $\alpha$ . Consider any point x of the dodecahedron. This is contained in exactly two cubes  $Q_i, Q_j$ . These two cubes have only one more point in common, namely the point x' diametrically opposite to x. Thus  $\alpha(x) = x$  or x'. Moreover, if  $\alpha(x) = x$  for some x, then  $\alpha(y)$  must be a neighbor of x for every neighbor y of x, so  $\alpha(y) = y$ . Thus it follows that  $\alpha$  is either the identity or the reflection  $\rho$  about the centre of the dodecahedron.

The number of congruences of the dodecahedron is 120. By the above,  $\alpha \to \overline{\alpha}$  is a homomorphism of the congruence group  $\Gamma$  into  $S_5$ , and the image has 60 elements. The only 60-element subgroup of  $S_5$  is  $A_5$ , so this is the image of  $\Gamma$ . Also the above considerations show that the mapping is monomorphic on the subgroup  $\Gamma_0$  of direct (orientation preserving) congruences, hence this subgroup (which clearly has index 2) is isomorphic with  $A_5$ . Also it follows that  $\{1, \varrho\}$  is a normal subgroup, whence  $\varrho$  commutes with every element of  $\Gamma$ . Hence

$$\Gamma = \Gamma_0 \times \{1, \varrho\} \cong A_5 \times C_2.$$

(b) It follows as for the dodecahedron that each automorphism of the cube is, in fact, induced by a congruence and that the whole group is the direct product of  $\mathbb{Z}_2$  and the group of direct congruences of the cube. Each direct congruence of the cube induces a permutation of the 4 main diagonals. It is also easy to see that, if a direct congruence maps every main diagonal onto itself, then it is the identity. Therefore, the group of direct congruences of the cube is a subgroup of  $S_4$ . A straightforward counting shows that the cube has 24 direct congruences (each edge is mapped onto any other edge by exactly two direct congruences) and hence, the group of direct congruences is  $S_4$ .

The tetrahedron clearly has automorphism group  $S_4$ . The octahedron is the dual of the cube; since its automorphisms must be induced by congruences in the same way as in the previous cases, its automorphism group is the same as that of the cube, i.e.  $Z_2 \times S_4$ . Similarly, the automorphism group of the icosa-

hedron is isomorphic with  $Z_2 \times A_5$ .




Fig. 86

- **3.** A graph like the one in Fig. 86 has only the rotations as automorphisms [R. Frucht, Compositio Math. 6 (1938) 239-250].
- 4. The right multiplication by any fixed group element g, i.e. the mapping  $g_i \mapsto g_i g$  is an automorphism: if  $g_i$  is joined to  $g_i$  by an edge of color k (i.e.  $g_ig_i^{-1}=g_k$ , then  $(g_ig)(g_ig)^{-1}=g_k$  and hence,  $g_ig$  is joined to  $g_ig$  by an edge of color k.

Conversely, every automorphism of G arises in this way. For let  $\alpha$  be any automorphism of G and set  $g = \alpha(1)$ . We claim that  $g_i g = \alpha(g_i)$ . Since  $(g_i g) g^{-1} = g_i$ ,  $g_i g$  is joined to g by an edge of color i and  $g_i$  is the only point with this property. On the other hand,  $g_i$  is joined to 1 by an edge of color i by definition, and since  $\alpha$  is an automorphism this implies that  $\alpha(g_i)$  is joined to  $\alpha(1) = g$  by an edge of color i. Hence,  $\alpha(g_i) = g_i g_i$ 

Since it is easy to see that the multiplication of elements of  $\Gamma$  is the same as multiplication between the corresponding automorphisms of G, we conclude that the automorphism group of G is isomorphic with  $\Gamma$  [R. Frucht, ibid.].

5. We may assume  $|\Gamma| \geq 2$ . The previous problem yields a colored digraph  $G_0$  with automorphism group  $\cong \Gamma$ .

If  $g_i, g_i \in V(G_0) = \Gamma$  are joined by an edge of color k, then connect them by a path of length k+2, with paths of length 1 attached to each inner point of it except for the inner point next to  $g_i$ , where we attach a path of length 2




Fig. 87

(Fig. 87, p. 423). We do this for each pair  $g_i$ ,  $g_j$  and then remove the directed

edges. We denote the resulting graph by G.

Observe first that each automorphism of  $\vec{G}_0$  induces a unique automorphism of G. It suffices to show that, conversely, if  $\alpha$  is an automorphism of G, then  $\alpha$  is induced by some automorphism of  $\vec{G}_0$ . The points of  $V(\vec{G}_0)$  are exactly those points of G, which are neither cutpoints nor endpoints. Hence,  $V(\vec{G}_0)$  is invariant under A(G).

The new paths with the attached shorter paths are the components of  $G - V(\vec{G}_0)$ , hence  $\alpha$  maps them onto each other. A path of length k must be mapped onto a path of length k and the end of the path with the longer attached path must be mapped onto the corresponding end. Hence, if  $g_i$  is connected to  $g_j$  by an edge of color k, then so is  $\alpha(g_i)$  to  $\alpha(g_j)$ . Thus,  $\alpha$  yields an automorphism of  $\vec{G}_0$  as stated [R. Frucht, ibid.].

**6.** Let  $V_i = \{(\gamma, i) \colon \gamma \in \Gamma\}$  (i = 1, 2). We may consider  $\Gamma$  to act on  $V_1 \cup V_2$  according to

$$\delta(\gamma, i) = (\gamma \delta, i)$$
  $(\gamma, \delta \in \Gamma, i = 1, 2).$ 

Let  $h_1, \ldots, h_m$  be a minimal set of generators of  $\Gamma$ . Assume first that  $m \geq 2$ . Connect (1, 1) to  $(1, 2), (h_1, 2), \ldots, (h_m, 2), (h_1, 1)$ . Also connect  $(h_1, 2)$  to  $(h_2, 2)$  to  $\ldots$  to  $(h_m, 2)$ . Take all images of these edges under  $\Gamma$  and denote the resulting graph by G.

Clearly G admits all elements of  $\Gamma$  as automorphisms. Let us see, if it has any

other.

Let  $\alpha$  be an automorphism of G and assume first that it keeps  $V_1$ ,  $V_2$  invariant. By multiplying by a suitable element of  $\Gamma$ , we may achieve  $\alpha(1,1)=(1,1)$ . The point (1,1) is adjacent to  $(1,2),(h_2,2),\ldots,(h_m,2)$  in  $V_2$ .  $\alpha$  must keep this set invariant.

We claim that these points induce the following subgraph: (1, 2) is isolated,  $(h_1, 2), \ldots, (h_m, 2)$  form a path P. Clearly the edges of P are in G but we have to show no other edge runs between these points. More exactly we show:  $\Gamma$  does not map any pair  $((h_i, 2), (h_{i+1}, 2))$  onto a pair  $((h_r, 2), (h_s, 2))$  unless  $\{r, s\} = \{i, i+1\}$  and it does not map it onto  $((1, 2), (h_r, 2))$  at all. For assume

$$\gamma((h_i, 2), (h_{i+1}, 2)) = ((h_r, 2), (h_s, 2)), \ \gamma \in \Gamma.$$

Then  $\gamma = h_i^{-1}h_r = h_{i+1}^{-1}h_s$ . If one of r, s is different from both  $h_i$  and  $h_{i+1}$ , then it can be expressed from this relation in contradiction with the minimality of generating set  $\{h_1, \ldots, h_m\}$ . So  $\{r, s\} = \{i, i+1\}$ . The other assertion

above follows similarly.

Thus it follows that  $\alpha$  must keep (1,2) fixed, and that there are only two possibilities for  $\alpha$  to act on  $(h_1,2),\ldots,(h_m,2)$ : either  $\alpha$  is the identity on this set or  $\alpha(h_i,2)=(h_{m-i+1},2)$ . We can rule out the second possibility as follows.  $(h_1,2)$  is adjacent to  $(h_1,1)$ , which is a neighbor of (1,2). We show  $(h_m,2)$  is not adjacent to any neighbor of (1,1) in  $V_1$ . In fact, (1,1) has two (not necessarily different) neighbors  $(h_1,1)$  and  $(h_1^{-1},1)$  in  $V_1$ . If  $(h_1^{\pm 1},1)$  were adjacent to  $(h_m,2)$ , then we would have  $(h_1^{\pm 1},1)=\gamma(1,1)$ , and  $(h_m,2)=\gamma(h_r,2)$  or  $(h_m,2)=\gamma(1,2)$  for some  $\gamma \in \Gamma$  and  $1 \le r \le m$ . But this would imply  $\gamma = h_1^{\pm 1}$  and either  $\gamma = h_r^{-1}h_m$  or  $\gamma = h_m$ , which are both impossible by the minimality of the generating set  $\{h_1,\ldots,h_m\}$ .

So we have shown that if  $\alpha$  fixes (1,1), it also fixes its neighbors in  $V_2$ . We show now that  $\alpha$  acts on  $V_1$  and  $V_2$  in the same way, i.e. if  $\alpha(g,1)=(g',1)$ , then  $\alpha(g,2)=(g',2)$ . In fact,  $g \alpha g'^{-1}$  keeps (1,1) fixed which, as above, implies that it fixes (1,2). But this means that  $\alpha(g,2)=(g',2)$ . Thus  $\alpha$  fixes  $(h_1,1),\ldots,\ldots,(h_m,1)$ . It follows then that if  $\alpha$  fixes (g,1), is also fixes  $(gh_i,1)$  for each  $i=1,\ldots,m$ . Since  $\{h_1,\ldots,h_m\}$  generates  $\Gamma$  it follows that  $\alpha$  fixes every element of  $V_1$ . As we have seen, this implies that it fixes every element of  $V_2$ , i.e.  $\alpha=1$ . This proves that those automorphisms of G keeping  $V_1$  and  $V_2$  invariant are only the elements of  $\Gamma$ .

To rule out those automorphisms of G, which would not keep  $V_1$  and  $V_2$  invariant we want to arrange for the degrees in these sets to be different. If G itself fulfils this we are done. Otherwise, take the complement in  $V_1$ . Since the degrees of the subgraph induced by  $V_1$  in G are at most 2, in the complement they will become at least n-1-2>2. Thus they increase, so after complementation they will not be equal to the degrees in  $V_2$ . Since the complementation in  $V_1$  does not influence those automorphisms keeping  $V_1$  invariant, we get

the desired graph.

In the case when m=1, i.e.  $\Gamma$  is cyclic, take the graph shown in Fig. 88. The points on the rim have degree 5, the points inside have degree 3, so each automorphism maps the boundary circuit onto itself. The inner points prevent reflections [L. Babai,  $Can.\ Math.\ Bull.\ 17\ (1974)\ 467-470$ ].




Fig. 88

- 7. (a) Suppose indirectly that a tournament T has an even number of automorphisms. By Cauchy's theorem T has an automorphism  $\alpha$  of order 2. Let (xy) be a 2-cycle of  $\alpha$ . Since  $\gamma$  is a tournament, either (x, y) or (y, x) is an edge of T, but not both. Say  $(x, y) \in E(T)$ , then  $\alpha$  maps (x, y) onto  $(y, x) \notin E(T)$ , a contradiction.
- (b) Let us first note the following fact. Let V be a set such that the group  $\Gamma$  of odd order acts regularly on V. Let us consider the orbits of  $\Gamma$  on the set E of ordered pairs (x, y),  $x \neq y$ ,  $x, y \in V$ . No element of  $\Gamma$  can map (x, y) onto (y, x); for such a permutation would contain an even cycle and hence, it would be of even order.

Thus, the orbits of  $\Gamma$  on E decompose into pairs S, S' such that  $S' = \{(x, y): (y, x) \in S\}$ . Selecting one orbit from each such pair S, S', the union of these

orbits forms a tournament T, which admits all elements of  $\Gamma$  as automorphisms. Now let us take two sets  $V_i = \{(\alpha, i) : \alpha \in \Gamma\}$  (i = 1, 2), and define the action of  $\alpha \in \Gamma$  on  $V_1 \cup V_2$  by

$$\alpha(\gamma, i) = (\gamma \alpha, i) \ (\gamma \in \Gamma, i = 1, 2).$$

Let  $h_1, \ldots, h_m$  be a minimal generating set of  $\Gamma$ .

Let us connect  $(\gamma, 1)$  to  $(\delta, 2)$  if  $\delta = \gamma$  or  $\delta = \gamma \cdot h_i (1 \le i \le m)$ , and  $(\delta, 2)$  to  $(\gamma, 1)$  otherwise. Take any tournament on  $V_1$  invariant under  $\Gamma$ . To define the tournament on  $V_2$  let us observe that no pair  $((h_i, 2), (h_j, 2))$  or  $((h_i, 2), (h_j, 2))$  is mapped onto such a pair by any element  $\gamma \in \Gamma$ ,  $\gamma \ne 1$ ; for, e.g.

$$\gamma((h_i, 2), (h_j, 2)) = ((h_{\mu}, 2), (h_{\nu}, 2))$$

would imply

$$h_i = h_\mu h_\nu^{-1} h_j,$$

which implies, by the minimality of  $\{h_1,\ldots,h_m\}$ , that either  $i=v,\ j=\mu$  and thus  $(h_ih_j)^2=1$  (which has been excluded, because  $\Gamma$  has odd order) or  $i=\mu,\ j=v$  and thus  $\gamma=1$ . So we may take the edges  $(1,2),\ (h_i,2)$   $(i=1,\ldots,m)$  and  $(h_i,2),\ (h_j,2)$   $(1\leq i< j\leq m)$  and all images of them under  $\Gamma$  and, as noted at the beginning, one of each remaining pair of orbits of  $\Gamma$  on  $\Gamma$ , and obtain a tournament on  $\Gamma$  invariant under  $\Gamma$ . Thus we have defined a tournament  $\Gamma$  on  $\Gamma$  on  $\Gamma$  invariant under  $\Gamma$ .

We claim that  $A(T) = \Gamma$ . Let  $\alpha \in A(T)$ , we want to show  $\alpha \in \Gamma$ . First observe that the points in  $V_1$  have outdegree

$$\frac{n-1}{2}+m+1,$$

while the points in V<sub>2</sub> have outdegree

$$\frac{n-1}{2}+n-m-1,$$

which are different, because n is odd. Therefore,  $\alpha$  must keep  $V_1$  and  $V_2$  invariant.

We may assume  $\alpha(1,1)=(1,1)$ . Then the set of points in  $V_2$  accessible from (1,1) on an edge is also invariant under  $\alpha$ . This set is  $(1,2), (h_1,2), \ldots, h_m, 2)$ , which is a transitive tournament. Therefore  $\alpha$  must fix all these points. Hence it follows as in the preceding solution that  $\alpha$  must fix every point i.e.  $\alpha=1$  [J.W. Moon, Canad. J. Math. 16 (1964) 485–489; this construction is due to L. Babai].

8. Instead of the paths shown in Fig. 87, p. 423, use the paths on Fig. 89 to connect  $g_i$  to  $g_j$  if they are joined by a directed edge of color k; where the 6-point configuration is repeated k times. In this way, we get a graph  $G_1$  which has automorphism group  $\Gamma$  (this follows exactly as before). However,  $G_1$  is not 3-regular as the points of  $V(G_0)$  have degree 2  $|\Gamma|$ .

Take any  $g_i \in V(G_0)$ . Observe that there are exactly two of the strings of Fig. 86 of length k attached to it, one with each kind of ending  $(k = 1, ..., |\Gamma|)$ . Split  $g_i$  into  $2 |\Gamma|$  points of degree 1; they can be denoted by  $a_{1,6}, ..., a_{|\Gamma|,6}$ ,  $a_{1,8}, ..., a_{|\Gamma|,8}$ , where  $a_{k,j}$  is the starting point of a string in Fig. 86 of length

Fig. 89

k and with a j-point configuration next to  $a_{k,j}$ . Now join the  $a_{k,j}$ 's by a circuit  $C_i$  in the above order. Do the same for each  $i=1,\ldots,|\Gamma|$ . Clearly, each automorphism of  $G_1$  induces an automorphism of the resulting graph G. Conversely, let  $\alpha$  be an automorphism of G. The only triangles in G are those in the strings of Fig. 86, so a maps them onto each other; also the triangles in the 6-point configurations have different neighborhoods than the triangles in the 8-point configurations, therefore α maps 6-point-configurations onto 6-point configurations and 8-point-configurations onto 8-point configurations. Thus,  $\alpha$  maps the rest, i.e. the points on circuits  $C_i$  onto each other. Since the  $C_i$  s are the components of the subgraph spanned by  $V(C_1) \cup \ldots \cup V(C_{|T|})$ ,  $\alpha$ maps each  $C_i$  onto some  $C_i$ . Hence,  $\alpha$  induces a (unique) automorphism of the graph obtained by contracting each  $C_i$ . But this graph is  $G_1$ .

Thus the automorphisms of G and  $G_1$  are in a 1-1 correspondence, which is obviously an isomorphism. Hence, the automorphism group of G is isomorphic to  $\Gamma$ . Since G is 3-regular, we are finished. [R. Frucht, Canad. J. Math. 1 (1949) 365-378. This proof is due to L. Babai. For regular graphs of any degree see

G. Sabidussi, Canad. J. Math 9 (1957) 515-525.

**9.** Let  $x_1, \ldots, x_n$  be as in the hint, and denote by  $\Gamma_i$  the group of those automorphisms fixing  $x_1, \ldots, x_i$ . Then  $\Gamma_1 = A(G)$ , because  $x_1$  is the only point of degree 2 and so each automorphism must fix it;  $\Gamma_n = \{1\}$ . We show that the

index  $|\Gamma_{i-1}:\Gamma_i|$  is 1 or 2; this will imply the assertion.

Look at the images of  $x_i$  under  $\Gamma_{i-1}$   $(i \ge 2)$ . Let j < i be such that  $x_j$  is adjacent to  $x_i$ . Then  $x_i$  has at most one other neighbor in  $\{x_i, \ldots, x_n\}$ . Since each  $\gamma \in \Gamma_{i-1}$  must map  $x_i$  onto such a neighbor of  $x_i$ , it follows that either  $x_i$  is fixed by all automorphisms in  $\Gamma_{i-1}$  or it has one other image. The index  $|\Gamma_{i-1}|$ : :  $\Gamma_i$  is, accordingly, 1 or 2.

Remark: Every group of order  $2^k$  arises as the automorphism group of such a graph [cf. L. Babai—L. Lovász, Studia Sci. Math. Hung. 8 (1973) 141-150].

10. Select an  $e \in E(G)$  and orient it arbitrarily. Let f be any other element in the orbit of e under A(G) and let  $\alpha$  be the unique automorphism mapping e onto f. Now orient f in such a way that a should preserve the orientation of e,

if  $\alpha \in \Gamma$  and conversely, otherwise. The remaining edges of G we orient in such a way that every  $\gamma \in A(G)$  preserves their orientations (this can be done as A(G) acts semiregularly on E(G)).

The resulting digraph  $\tilde{G}$  admits the elements of  $\Gamma$  as automorphisms by the construction. On the other hand, let  $\alpha$  be any automorphism of  $\tilde{G}$ . Then, clearly,  $\alpha$  is an automorphism of G, i.e.  $\alpha \in A(G)$ . Since  $\alpha(e)$  must be oriented like e, it follows that  $\alpha \in \Gamma$  [Cf. J. Nešetřil, *Monatsh. f. Math.* 76 (1972) 323–327].

11. It is easy to check that in 12.5, if we remove an edge e, which connects a point of degree 1 to a point of degree 2, we obtain a graph with no automorphisms. Taking the complement we have a connected graph  $G_2$  and an edge  $e = (u, v) \in E(G_2)$  such that  $A(G_2) = \{1\}$ ,  $A(G_2 - e) \cong \Gamma_2$ . Also the construction in 12.5 has the property that its automorphism group acts semiregularly on the points, i.e. no automorphism  $\alpha \neq 1$  fixes any point. Thus we have a graph  $G_1$  such that  $A(G_1) \cong \Gamma_1$  and only the identity fixes a

certain  $x \in V(G_1)$ .

Now construct G as follows. Let its points be the pairs  $(x_1, x_2)$ ,  $x_i \in V(G_i)$ . Moreover, connect  $(x_1, x_2)$  to  $(y_1, y_2)$ , if either  $(x_1, y_1) \in E(G_1)$  or  $x_1 = y_1$  and  $(x_2, y_2) \in E(G_2)$ . (G is called the lexicographic product of  $G_1$  and  $G_2$ .) Also let us color those edges  $((x_1, x_2), (y_1, y_2))$  of G with  $x_1 = y_1$  red and the rest black. Now  $A(G) \cong \Gamma_1$ . For obviously, each automorphism of  $G_1$  induces an automorphism of G. On the other hand, let  $\alpha \in A(G)$ . Since  $G_2$  is connected, the red coloration of edges forces that  $\alpha$  must map each class  $V_{x_1} = \{(x_1, x_2) : x_2 \in V(G_2)\}$  onto such a class  $V_{x_1}$ . From the fact that  $G_2$  has no automorphism except the identity, it follows that  $\alpha(x_1, x_2) = (x_1', x_2)$ , i.e.  $\alpha$  is induced by an automorphism of  $G_1$ .

On the other hand, let us remove the edge f = ((x, u), (x, v)). Then for each

 $\gamma \in A(G_2 - e)$  the mapping  $\gamma$  defined by

$$\overline{\gamma}(x_1, x_2) = \begin{cases} (x, \gamma(x_2)) & \text{if } x_1 = x, \\ (x_1, x_2) & \text{if } x_1 \neq x. \end{cases}$$

is an automorphism of G-f. Conversely, if  $\alpha$  is any automorphism of G-f, then it follows as above that  $\alpha$  maps  $V_{x_1}$  onto  $V_{x_1'}$ . But then  $\alpha$  must map  $V_x$  onto itself (since  $V_x$  induces fewer edges than any other  $V_{x_1}$ ). Therefore,  $\alpha$  induces an automorphism  $\tilde{\alpha}$  of  $G_1$ , which fixes x, whence  $\tilde{\alpha}=1$ , i.e.  $\alpha(V_{x_1})=V_{x_1}$  for every  $x_1 \in V(G_1)$ . Since  $V_{x_1}$  induces a graph with no proper automorphism for  $x_1 \neq x$ , we conclude that  $\alpha = \overline{\gamma}$  for some  $\gamma \in A(G_2 - e)$ . Thus  $A(G - f) \cong \Gamma_2$ .

We still have to get rid of the colorings of edges. This can easily be done by replacing each black edge by a path of length N, N large [L. Babai].

12. (a) Let us remark first that there is a tree T such that each orbit of  $\Gamma$  contains exactly one point of T. For choose a maximal tree  $T_0$  such that each orbit contains at most one point of  $T_0$ . If  $V_0 = \bigcup_{\gamma \in \Gamma} \gamma(V(T_0)) = V(G)$  we are done.

Suppose  $V_0 \neq V(G)$ , then, since G is connected, there is a point  $x \notin V_0$  adjacent to some  $y \in V_0$ . Let  $y \in \gamma(T_0)$ , then  $\gamma(T_0) + (x, y)$  is a larger tree meeting each orbit in at most one point.

So let the tree T meet each orbit in exactly one point. Let us contract each  $\gamma(T)$ ,  $\gamma \in \Gamma$ . The resulting graph G' is clearly connected, and  $\Gamma$  acts regularly

on it [L. Babai, Acta Math. Acad. Sci. Hung. 24 (1973) 215-221; cf. G. Sabidussi, Proc. AMS 9 (1958) 800-804].

(b) We use induction on |V(G)|. We may suppose  $\Gamma \neq \{1\}$ . Let  $e \in E(G)$ 

such that not both endpoints of e are fixed points of  $\Gamma$ .

Let G' be the subgraph of G formed by those edges of G of form  $\gamma(e)$ ,  $\gamma \in \Gamma$ . Then G' is a subgraph of G on which  $\Gamma$  acts edge-transitively. Moreover, different elements of  $\Gamma$  have different effects on V(G'); for the set N of those permutations  $\gamma \in \Gamma$  which have  $\gamma|_{V(G)} = 1$  is a normal subgroup of  $\Gamma$ ; by the simplicity of  $\Gamma$ ,  $N = \Gamma$  or  $N = \{1\}$ . However,  $N = \Gamma$  would imply that all points of V(G'), in particular both endpoints of e, are fixed points of  $\Gamma$ , a contradic-

tion. Thus if G' is connected we are done.

Suppose G' is disconnected; let  $G_1, \ldots, G_k$  be its components. Then  $\{V(G_1), \ldots, G_k\}$ ...,  $V(G_k)$   $\cup$   $\{x\}: x \in V(G) - V(G')\}$  is a partition of V(G) invariant under  $\Gamma$ . Let us contract each  $G_i$  onto one point. The resulting graph G'' is clearly connected. Each automorphism  $\gamma \in \Gamma$  induces an automorphism  $\tilde{\gamma} \in A(G'')$ . As above, the simplicity of  $\Gamma$  implies that either  $\tilde{\gamma} = 1$  for each  $\gamma \in \Gamma$  or  $\gamma \mapsto \tilde{\gamma}$  is one-to-one. But the former possibility cannot occur since some  $\gamma \in \Gamma$ maps  $G_1$  onto  $G_2$  and for this  $\tilde{\gamma} \neq 1$ . Hence A(G'') contains a subgroup isomorphic with  $\Gamma$ . We are finished by induction. [L. Babai, Discrete Math. 8 (1974) 13-20.7

13. If a permutation group is commutative and transitive, then it is regular

(see the footnote to the solution of 11.17).

If we fix any  $x_0$  then, by the regularity of A(G), each  $x \in V(G)$  can be written uniquely as  $\alpha(x_0)$ ,  $\alpha \in A(G)$ . So the mapping  $\varphi$  with  $\varphi(\alpha(x_0)) = \alpha^{-1}(x_0)$  is well defined and is a permutation of V(G). Moreover,  $\varphi$  is an automorphism of G, since if  $(x, y) \in E(G)$ ,  $x = \alpha(x_0)$ ,  $y = \beta(x_0)$ ,  $(\alpha, \beta \in A(G))$  then, since  $\alpha$  is an automorphism,  $(x_0, \alpha^{-1}(\beta(x_0)) = (x_0, \beta(\alpha^{-1}(x_0))) \in E(G)$ , whence  $(\beta^{-1}(x_0), \alpha^{-1}(x_0)) \in E(G)$ . Since  $\varphi$  fixes  $x_0$ , we get, by the regularity of A(G), that  $\varphi = 1$ . Thus,  $\alpha^{-1} = \alpha$  for each  $\alpha \in A(G)$ , i.e. every element of A(G) is of order 2. It is well known from group theory that this implies that A(G) is the direct product of cyclic groups of order 2. [C.Y. Chao, Proc. AMS 15 (1964) 291-292; G. Sabidussi, Monatsh. f. Math. 68 (1964) 426-438.]

(b) Let  $Q_n$  be the *n*-cube. We may assume the vertices of  $Q_n$  are 01-vectors of length n and two such vectors are adjacent iff they differ at exactly one place. Denote by  $e_I$  the characteristic vector of  $I \subseteq \{1, ..., n\}$ . If we switch the coordinates belonging to a subset  $I \subseteq \{1, \ldots, n\}$ , we get an automorphism  $\alpha_I$  of  $Q_n$ . The group  $\Gamma_0$  of all automorphisms  $\alpha_I$  is isomorphic with  $Z_2 \times \ldots \times Z_2$ .

However,  $Q_n$  has other automorphisms, e.g. exchanging the first two coordinates. To exclude these, let us modify  $Q_n$  by joining two vectors by k edges if they differ at the  $k^{th}$  place only. It is obvious that  $\alpha_I$  as defined above is still an automorphism of the resulting graph  $Q'_n$ . We show that  $Q'_n$  has no other automorphisms. If it had we could assume it had one, which fixes  $(0, \ldots, 0)$ . However, if  $\alpha$  is an automorphism, which fixes  $(0, \ldots, 0)$ , then it also fixes  $(1, 0, \ldots, 0)$  $\ldots$ , 0) since this is the only point joined to  $(0, \ldots, 0)$  by exactly one edge. Similarly, it fixes  $(0, \ldots, 0, 1, 0, \ldots, 0)$  for any position of the 1. Thus if  $\alpha$ fixes a point it fixes all neighbors of it.  $Q'_n$  being connected,  $\alpha$  must fix every point, i.e.  $\alpha = 1$ , a contradiction.

(c) We want to introduce new edges to  $Q_n$  such that the arising simple graph G still admits every  $\alpha_I$  as an automorphism, but has no other automorphism. Let H be a graph on the set of vectors  $\{\mathbf{e}_1, \ldots, \mathbf{e}_n\}$ ,  $\mathbf{e}_i = (0, \ldots, 0, 1, 0, \ldots, 0)$ .

If  $(\mathbf{e}_i, \mathbf{e}_j) \in E(H)$ , then  $\alpha_I(\mathbf{e}_i) = (\mathbf{e}_{I \triangle \{i\}})$ ,  $\alpha_I(\mathbf{e}_j) = \mathbf{e}_{I \triangle \{j\}}$ . Thus if we add all edges  $(\mathbf{e}_I, \mathbf{e}_J)$  to  $Q_n$  such that  $I \triangle J = \{i, j\}$  with  $(\mathbf{e}_i, \mathbf{e}_j) \in E(H)$ , we obtain a graph G invariant under  $\Gamma_0$ .

Let us see what properties H must have to ensure that G has no other automor-

phism.

Suppose  $A(G) \neq \Gamma_0$ . Since  $\Gamma_0$  is transitive, we have an  $\alpha \in A(G)$ ,  $\alpha \neq 1$  with  $\alpha(0, \ldots, 0) = (0, \ldots, 0)$ . Let  $H_1$  be the graph induced by the neighbors of  $(0, \ldots, 0)$  in G (this is, of course, not H since  $(0, \ldots, 0)$  has neighbors other than  $\mathbf{e}_i$ ). The points of  $H_1$  are  $\mathbf{e}_1, \ldots, \mathbf{e}_n$  and all points  $\mathbf{e}_{\{i,j\}}$  with  $(\mathbf{e}_i, \mathbf{e}_j) \in E(H)$ . Two points of form  $\mathbf{e}_i$  are adjacent iff they are adjacent in H.  $\mathbf{e}_{\{i,j\}}$  is adjacent to  $\mathbf{e}_i$ ,  $\mathbf{e}_j$  and no other  $\mathbf{e}_v$ . Finally,  $\mathbf{e}_{\{i,j\}}$  is adjacent to  $\mathbf{e}_{\{\mu,\nu\}}$  iff  $(\mathbf{e}_i, \mathbf{e}_j)$ ,  $(\mathbf{e}_\mu, \mathbf{e}_\nu)$  are adjacent edges, e.g.  $j = \nu$  and  $(\mathbf{e}_i, \mathbf{e}_\mu) \in E(G)$ .

Thus  $H_1$  arises from H by taking a new point  $p_e$  for each edge e of H, connecting it to the endpoints of e and connecting  $p_e$  to  $p_{e'}$  iff e, e' are edges of a tri-

angle in H.

Note that if H contains no triangles, then each point of  $V(H_1) - V(H)$  has degree 2. If H has no point of degree 1, then these are the only points of degree 2. Thus if in addition H is asymmetric, so is  $H_1$ .

If this can be achieved we are done, since it means that  $\alpha$  fixes the neighbors of  $(0, \ldots, 0)$  in G, and G being connected, it follows that it fixes every point, a contradiction. So all we need is an asymmetric triangle-free graph with no point of degree 1.

Figure 90 shows such a graph H for  $n \ge 8$ . [W. Imrich, Monatsh. f. Math. 73 (1969) 341–347. For graphs with non-abelian regular automorphism group

see L.A. Nowitz-M.E. Watkins, Canad. J. Math. 24 (1972) 993-1018.]

14. Let k be the edge-connectivity of G, and  $X \subseteq V(G)$  a minimal set with  $\delta_G(X) = k$ .

By the minimality of X,  $|X| \leq |V(G)|/2$ . Let  $x_1, x_2 \in X$  and  $\alpha \in A(G)$  be such that  $\alpha$  moves  $x_1$  to  $x_2$ . Let  $\alpha(X) = X'$ . Then  $\delta_G(X') = k$  (as  $\alpha$  is an auto-




Fig. 90

morphism) and  $X \cap X' \neq \emptyset$  and  $X \cup X' \neq V(G)$  (as  $|X| = |X'| \leq |V(G)|/2$ ). Hence by 6.48a,  $\delta_G(X \cap X') = k$ . Since X has been minimal, this is only

possible, if  $X \cap X' = X$ , i.e. X = X'.

Thus, those automorphisms which keep X invariant act transitively on X. This implies that each point of X has the same number a of neighbors in X. Obviously,

 $a \leq |X| - 1$ ,

and since each point of X is adjacent to r-a points outside X,

$$(r-a)|X|=k.$$

Thus

$$k = (r - a) |X| \ge (r - a)(a + 1) \ge r.$$

Since obviously  $r \geq k$ , we have r = k as stated.


15. (a) Let  $T \subseteq V(G)$ , |T| = 3,  $G_1$  a component of G - T,  $X = V(G_1)$  and

suppose T,  $G_1$  are chosen so that |X| is minimal.

Consider all images of X under automorphisms of G. By the transitivity of A(G), they will cover all points. Moreover, they are disjoint (if different). For if  $\alpha(X) \cap \beta(X) \neq \emptyset$ ,  $\alpha(X) \neq \beta(X)$  then  $\alpha(T)$  meets  $\beta(X)$  but does not contain it, which contradicts 6.60a. This also shows that T is the union of certain images of X. Since |T| = 3, this is only possible if |X| = 1 (in which case we are finished) or |X| = 3.

To exclude this latter case, observe that X is joined by edges to T only. So if  $T = \alpha(X)$  for some  $\alpha$ , then T is joined to  $\alpha(T) = \alpha^2(X)$  only; however, T is in fact joined to X and to some other points too (being a cutset).

The statement is false for 4 instead of 3 as is shown by Fig. 91.



(b) Again let T be a minimum cutset of G and  $G_1$  a component of G-T, chosen so that |X| is minimal, where  $X=V(G_1)$ . It follows as before that T is the union of certain images of X which are disjoint, and T cannot be an image of X (under automorphisms of G). Hence,

$$|T| \geq 2|X|$$
.

Since any  $x \in X$  can be joined to the points of  $T \cup (X - \{x\})$  only, we have

$$r \le |T| + |X| - 1 \le |T| + \frac{1}{2}|T| - 1 = \frac{3}{2}|T| - 1$$

or, equivalently,

$$|T| \ge \frac{2}{3}(r+1).$$

This estimate is sharp if r+1 is divisible by 3, as is shown by the strong direct product of a circuit and a complete (r+1)/3-graph (Fig. 88 for r=5) [M.E. Watkins, J. Comb. Th. 8 (1970) 223-226; W. Mader, Arch. d. Math. 21 (1970) 331-336].

- (c) Again let T be a minimum cutset of G and  $G_1$  a component of G-T. Assume T and  $G_1$  are chosen so that  $|V(G_1)|$  is minimal. We want to show that  $G_1$  consists of a single point. Suppose indirectly that  $G_1$  has an edge e. Let f be an edge connecting  $G_1$  to T, and  $\alpha$  an automorphism mapping f onto e. Then  $\alpha(T)$  meets  $G_1$  and does not contain it, a contradiction with 6.60a.
- 16. Suppose indirectly that the edge e=(x,y) of G does not occur in any 1-factor (G itself may or may not have 1-factors). Then G-x-y has no 1-factor and therefore, by Tutte's theorem, there exists an  $X_1 \subseteq V(G) \{x,y\}$  such that  $G-x-y-X_1$  has more than  $|X_1|$  odd components. For reasons of parity, the number of odd components of  $G-x-y-X_1$  is  $\geq |X_1|+2$  (since |V(G)| is even). So if we set  $X=X_1\cup\{x,y\}$ , then G-X has  $\geq |X|$  odd components.

Let  $T_1, \ldots, T_k$  be the components of G-X ( $k \ge |X|$ ) and let G be r-regular. By 12.14, G is r-edge-connected and hence, there must be at least r edges joining  $T_i$  to X ( $i=1,\ldots,k$ ). This means that at least kr edges enter X. But X has only  $|X| \le k$  points to receive them and a point of X can receive at most r. Moreover x and y can receive at most r-1 as they are incident with e too. Thus, the points of X cannot receive all the edges entering X, a contradiction.

17.  $\Gamma$  is regular (see the footnote to the solution of 11.17). Let  $x \in V(G)$  and  $\gamma \in \Gamma$  be such that  $(x, \gamma(x)) \in E(G)$ . Then  $(x, \gamma(x), \ldots, \gamma^{r-1}(x))$  is a circuit in G (where r is the order of  $\gamma$ ).

Now let  $\Gamma_1$  be a maximal subgroup of  $\Gamma$  such that the elements of  $\Gamma_1(x)$  form a circuit  $(x, x_1, \ldots, x_m)$   $(|\Gamma_1| = m + 1)$ . We claim that  $\Gamma_1(x) = V(G)$ , which will prove the assertion. Suppose  $\Gamma - \Gamma_1 \neq \emptyset$ , then since G is connected, there is an edge (a, b) with  $a \in \Gamma_1(x)$ ,  $b \notin \Gamma_1(x)$ . Then  $a = \gamma(x)$   $(\gamma \in \Gamma_1)$  and  $(x, \gamma^{-1}(b))$  is an edge as well. Clearly  $\gamma^{-1}(b) \notin \Gamma_1(x)$ ; let  $\gamma^{-1}(b) = \delta(x)$ ,  $\delta \in \Gamma - \Gamma_1$ . Let p be the least integer with  $\delta^p \in \Gamma_1$ .

By the commutativity of  $\Gamma$ ,  $(y, \delta(y)) \in E(G)$  for any  $y \in V(G)$ ; in fact,  $y = \gamma(x)$  for some  $\gamma \in \Gamma$  and then  $(y, \delta(y)) = (\gamma(x), \delta(\gamma(x))) = (\gamma(x), \gamma(\delta(x))) = \gamma(x, \delta(x)) \in E(G)$ . Also the commutativity implies that  $\Gamma_2 = \Gamma_1 \cup \Gamma_1 \delta \cup \ldots \cup \Gamma_1 \delta^{p-1}$  is a subgroup.

If p is even, then

$$(x, x_1, \ldots, x_m, \delta(x_m), \ldots, \delta(x_1), \delta^2(x_1), \ldots, \delta^2(x_m), \ldots, \delta^{p-1}(x_m), \ldots, \delta^{p-1}(x_1), \delta^{p-1}(x), \ldots, \delta(x))$$

is a circuit formed by the points of  $\Gamma_2(x)$ , a contradiction (Fig. 92a). If p is odd we have the circuit

$$(x, x_1, \ldots, x_m, \delta(x_m), \ldots, \delta(x_1), \delta^2(x_1), \ldots, \delta^2(x_m), \ldots, \delta^{p-1}(x_m), \delta^{p-1}(x), \delta^{p-2}(x), \ldots, \delta(x)),$$

a contradiction again (Fig. 92b).




Fig. 92

*Remark*: The condition that  $\Gamma$  is abelian cannot be dropped, as is shown by the Petersen graph [J. Pelikán].

18. It suffices to describe *connected* graphs with the desired properties; the disconnected ones are disjoint unions of arbitrary numbers of the same connected examples.

We distinguish two cases. If  $\Gamma = A(G)$  acts point-transitively, then by 12.15b, G is at least 2-connected unless  $G = K_2$ . Moreover if G is not 3-connected, then

it is 2-regular, i.e. a single circuit.

Assume that G is r-regular,  $r \geq 3$  and 3-connected. By 6.69 it follows that G has an essentially unique embedding into the sphere and every automorphism maps faces onto faces. By edge-transitivity, either all faces have the same size or there are only two types of faces, k-gons and l-gons, say. In the first case G can only be one of the five platonic bodies, the proof of which is found in any geometry book.

In the second case, the edge-transitivity implies that each edge must be adjacent to just one k-gon and one l-gon. Hence each point is adjacent to some a k-gons and a l-gons. Clearly  $a \ge 2$  and by 5.26,  $|E(G)| = a \cdot n \le 3n - 6$ ,

whence a=2.

The total number of k-gons is, clearly,  $\frac{2n}{k}$ , the number of l-gons is  $\frac{2n}{l}$ , thus by Euler's formula

$$n + \frac{2n}{k} + \frac{2n}{l} = 2n + 2,$$

$$\frac{1}{k} + \frac{1}{l} = \frac{1}{2} + \frac{1}{n}.$$

We may suppose  $3 \le k < l$ , whence either k = 3, l = 4, n = 12 or k = 3, l = 5, n = 30. It is easy to see that the corresponding graphs are the line-graphs of the cube and dodecahedron, respectively.

Now suppose that  $\Gamma$  is not transitive on the points. Then clearly A(G) has two transitivity classes on V(G), U and W, say. Each edge connects two points in different classes.

If the minimum degree of G is 1, it is a star. If the minimum degree of G is 2, it arises by subdividing each edge of a (not necessarily simple) planar graph  $G_1$  by one point. This graph  $G_1$  has an automorphism group transitive on both the points and the edges. Hence  $G_1$  arises from one of the graphs determined above by multiplying each edge by the same number. Thus G is one of the graphs described in the hint.

So suppose the minimum degree of G is at least 3. Then G is 3-connected by 12.15c, and thus by 6.69 it has an essentially unique embedding into the sphere. From the edge-transitivity it again follows that A(G) is either transitive on the faces or there are two types of faces. In the first case consider the dual graph  $G^*$ . This has an automorphism group transitive on both the points and the edges, and is therefore one of the graphs determined above.

So we may suppose A(G) is not transitive on the faces either. Let each edge be adjacent to a k-gonal and an l-gonal face. Let the degrees in U and W be a and b, respectively. Since the faces containing a given point are alternatingly in the two transitivity classes, a, b are even and therefore at least 4. Similarly k,  $l \geq 4$ . But then the number of points is at most |E(G)|/2, the number of faces is at most |E(G)|/2, contradicting Euler's theorem. So the last case cannot occur. Thus the list given in the hint is verified.

19. Let  $\Gamma$  be a non-cyclic simple group and assume that  $A(G) \cong \Gamma$ , where G is planar. We show that this implies  $\Gamma \cong A_5$ , even if coloring the points is allowed. Let G be a minimum graph with  $A(G) \cong \Gamma$ . Then G is connected. For, suppose that G has components  $G_1, \ldots, G_k$ . Then each automorphism of G induces a permutation of the components. Those automorphisms inducing the identity form a normal subgroup of A(G). Since A(G) is simple, either no automorphism induces the identity or all of them do. In the first case the components of G are asymmetric and hence A(G) is a direct product of symmetric groups (permuting the isomorphic components), which is never a non-cyclic simple group. So each automorphism leaves the components invariant. Let  $G_1$  be a component, which is not asymmetric, then those automorphisms keeping the points of  $G_1$  fixed form a normal, proper subgroup of A(G). Since A(G) is simple it follows that




Fig. 93

only the identity does so, i.e.  $A(G) \simeq A(G_1)$ , a contradiction with the minimality of G.


Furthermore A(G) has no fixed points. For removing a fixed point and coloring its neighbors with a new color we obtain a smaller colored graph with the same

automorphism group.

Now let us contract an appropriate connected subgraph of G onto a graph  $G_0$  as in 12.12b such that  $A(G_0)$  contains a subgroup  $\Gamma' \cong A(G)$ , which acts edge-transitively. Observe that the construction in 12.12b is such that if A(G) has no fixed points, neither does  $\Gamma'$ .  $G_0$  is clearly planar and so it is one of the graphs described in the preceding hint. The stars are ruled out since  $\Gamma'$  is fixed point free. The remaining examples are the platonic bodies or graphs, which are easily seen to have the same automorphism groups as the platonic bodies. Using 12.2 it is easy to see that the automorphism groups of the platonic bodies have no non-cyclic simple subgroups other than  $A_5$ .

Remark: Fig. 93 shows a planar graph with automorphism group  $A_5$  [L. Babai, in: Infinite and Finite Sets, Coll. Math. Soc. J. Bolyai 10 Bolyai–North-Holland (1975) 29–84].

**20.** (a) Using 12.12a, we may assume that  $\Gamma$  acts regularly. Then it is easy to see that G' contains the Cartesian product of two (p-1)-paths and an edge. (In fact, it contains the Cartesian product of three p-circuits.) Let  $G \supset P \oplus P' \oplus Q$ ,  $P = (x_1, \ldots, x_p)$ ,  $P' = (y_1, \ldots, y_p)$ ,  $Q = (z_1, z_2)$ . It suffices to show that  $P \oplus P' \oplus Q$  can be contracted onto  $K_p$ . To this end contract the subgraphs spanned by  $P \oplus \{y_i\} \oplus \{z_1\}$  and  $\{x_i\} \oplus P' \oplus \{z_2\}$  (which are clearly connected). This results in a graph isomorphic to  $K_{p,p}$  which, in turn, is trivially contractible onto  $K_p$  (Fig. 94).



(b) For reasons similar to those in the solution of the previous problem we may assume that the action of  $\Gamma'$  is edge-transitive. By (a) we may suppose that the action of  $\Gamma$  is not semiregular. Let  $\gamma_0 \in \Gamma$  have fixed points. Since G is connected it has two adjacent points  $x_0$ ,  $y_0$  such that  $\gamma_0(x_0) = x_0$ , but  $\gamma_0(y_0) \neq y_0$ . Then  $y_0, \gamma_0(y_0), \ldots, \gamma_0^{p-1}(y_0)$  are different neighbors of  $x_0$ . Hence  $d_G(x_0) \geq p$ .

If A(G) is point-transitive, then each point has degree at least p and we are done by Mader's theorem 10.8. So suppose A(G) is not point-transitive. By the edge-transitivity, it follows then that  $\Gamma'$  has two transitivity classes, namely  $U = \Gamma'(x_0)$  and  $W = \Gamma'(y_0)$ . Each edge connects U to W. It is trivial

that the degrees in U are all at least p and those in W are all at least 2. If the degrees in W are at least  $\sqrt[3]{p}$ , then we can still conclude as before by 10.8. So

suppose the degrees in W are less than  $\sqrt{p}$ .

Form a graph  $G_1$  on  $V(G_1) = U$  by connecting two points, if they are at distance 2 in G. Each  $\gamma \in \Gamma'$  induces a  $\overline{\gamma} \in A(G_1)$ . Clearly  $\gamma \mapsto \gamma'$  is a homomorphism and since  $\Gamma'$  is simple, it is either a monomorphism or maps onto 1. However, the latter is impossible since  $\Gamma'$  acts transitively on U. Moreover,  $\gamma_0$  has fixed points, but is not the identity, thus as before, it follows that the degrees of  $G_1$  are at least p.

For each edge  $e = (u, v) \in E(G_1)$ , let f(e) be one of the common neighbors of

u, v in G. Clearly  $f(e) \in V.$  A point  $y \in V$  occurs at most

$$inom{d_G(y)}{2} \leq inom{\sqrt[3]{p}}{2} < rac{1}{2} \, p^{2/3}$$

times as f(e) ( $e \in E(G_1)$ ) and so, the number of points of form f(e) is at least

$$\frac{2|E(G)|}{p^{2/3}} \ge \frac{p \cdot |U|}{p^{2/3}} = |U| \sqrt[3]{p}.$$

For each point x of the form f(e) select an edge  $(u_x, v_x) = e_x \in E(G_1)$  such that

 $f(e_x) = x$  and contract the edge  $(u_x, x)$  of G.

Identify those points of W not of form f(e) with an arbitrary neighbor of them. In this way G is mapped onto a subgraph G' of  $G_1$ , which contains all edges of the form  $e_x$  (and possibly more). So

$$|E(G')| \ge |U| \sqrt[3]{p}.$$

Thus by 10.8 G' can be contracted onto  $K_m$ .

(c) Let N be a large enough number; we show that the alternating group  $A_N$  is not isomorphic to the automorphism group of any graph from  $\mathcal{K}$ , even, if coloring of the points is considered. More exactly we show that if a graph G (with possibly colored points) has automorphism group  $A_N$ , then a suitable subgraph of it can be contracted onto  $K_m$  provided  $N \geq 3p$ , where p is a prime greater than  $8^m$ . Since for large enough m,  $\mathcal{K}$  certainly does not contain any graph contractible onto  $K_m$ , this will prove the assertion.

For reasons similar to those in the solution of 12.19, we may suppose that G is connected and A(G) has no fixed points. Then disregarding the coloration of the points, we obtain a graph, which satisfies the conditions of (b). This implies that G can be contracted onto  $K_m$  [L. Babai, Discrete Math. 8 (1974) 13–20].

**21.** Let G be a graph with  $A(G) \cong \Gamma$ ,  $V(G) \cap \Omega = \emptyset$ . On the basis of the construction in 12.5 we may assume A(G) acts on G semiregularly, i.e. no element

 $\alpha \neq 1$  has a fixed point.

Also by the construction in 12.5 we may assume  $|V(G)| \ge |\Gamma| \cdot |\Omega|$ ; then A(G) has at least  $|\Omega|$  orbits on V(G). Finally, by taking the complement of the construction in 12.5 we can arrange for each degree in G to be greater than  $|\Gamma|$ . Let  $\gamma \to \overline{\gamma}$  be an isomorphism of  $\Gamma$  onto A(G). Now let  $\Omega_1, \ldots, \Omega_k$  be the orbits of  $\Gamma$  on  $\Omega$  and select a  $y_i \in \Omega_i$   $(i = 1, \ldots, k)$ . Let  $x_1, \ldots, x_k \in V(G)$ 

be points in different orbits. For every  $\gamma \in \Gamma$  and  $1 \le i \le k$ , connect  $\gamma(y_i)$  to  $\overline{\gamma}(x_i)$ . The resulting graph G' admits all permutations  $\gamma'$  as automorphisms, where

$$\gamma'(x) = \begin{cases} \gamma(x) & \text{if} \quad x \in \Omega, \\ \overline{\gamma}(x) & \text{if} \quad x \in V(G). \end{cases}$$

For, clearly  $\gamma'$  preserves adjacency in V(G) and if  $(\gamma_0(y_i), \overline{\gamma}_0(x_i))$  is any new

edge, then  $\gamma'(\gamma_0(y_i), \overline{\gamma_0(x_i)}) = ((\gamma_0\gamma)(y_i), (\overline{\gamma_0\gamma})(x_i))$  is also a new edge.

On the other hand, G' has no other automorphisms. For let  $\alpha \in A(G')$ . Then  $\alpha$  preserves V(G), as the set of points with degree greater than  $|\Gamma|$ . Therefore  $\alpha|_{V(G)} = \overline{\gamma}|_{V(G)}$  for some  $\gamma \in \Gamma$ . Let  $y \in \Omega$ , then  $y = \delta(y_i)$  for some  $\delta \in \Gamma$  and  $1 \le i \le k$ . Since  $\alpha$  is an isomorphism,  $\alpha(y)$  is adjacent to  $\alpha(\overline{\delta}(x_i)) = (\overline{\delta \gamma})(x_i)$ . But the only point of  $\Omega$  adjacent to  $(\overline{\delta \gamma})(x_i)$  is  $(\delta \gamma)(y_i)$ . So  $\alpha(y) = (\delta \gamma)(y_i) = \gamma(y)$ . Thus we conclude that  $\alpha = \gamma'$ .

This proves that G' has the desired properties [I.Z. Bouwer, J. Comb. Th. 6 (1969) 378–386; Z. Hedrlin-A. Pultr, Illinois J. Math. 10 (1966) 392–405].

- 22. The assertion of the hint easily follows from these two observations:
- (a) If  $\eta$  is any endomorphism of the Petersen graph and C is a pentagon in it, then  $\eta$  is one-to-one on C. For the image of C contains an odd circuit (it cannot be 2-colorable as this would yield a 2-coloration of C), and the shortest odd circuit in the Petersen graph is a pentagon. Thus  $\eta(C)$  contains a pentagon and so  $\eta$  is one-to-one on C.
- (b) Any two points of the Petersen graph are contained in a pentagon. Therefore, no endomorphism can identify two distinct points. Thus, the endomorphism semigroup is  $\cong S_5$  by 12.1.
- 23. Consider the graph shown in Fig. 95, where s>a+b+c is odd. This graph G has three circuits of length s and one circuit of length 3s-2(a+b+c)>s. Therefore, any endomorphism  $\eta$  of G must be one-to-one on the s-circuits. In particular, the three neighbors of x lie pairwise on circuits of length s, thus  $\eta(x)$  must have three distinct neighbors, which also lie pairwise on such circuits. Hence  $\eta(x)=x$ . It follows trivially that if a,b,c are distinct, then  $\eta$  must keep the three "spokes" fixed. Also, the arc of length s-a-c on the "rim" cannot be mapped onto any other arc with the same endpoints,




Fig. 95

for there are only two other paths with the same parity, namely the paths of length c+b+(s-a-b) and a+b+(s-b-c), but these are longer. Thus  $\eta=1$  [Z. Hedrlin-A. Pultr, *Monatsh. f. Math.* 69 (1965) 318-322].

**24.** (a) Let  $\eta_1, \ldots, \eta_n$  be the elements of  $\Sigma$ . Choose these as points of a colored digraph G. Connect  $\eta_i$  to  $\eta_j$  by a directed edge of color k if  $\eta_i = \eta_k \eta_j$ . Since we only have a semigroup it may happen that certain pairs are connected by several edges of different colors or by no edge at all.

For any  $\eta_m$ , defining  $\overline{\eta}_m(\eta_j) = \eta_j \eta_m$ , we obtain an endomorphism of G; for if an edge of color k connects  $\eta_i$  to  $\eta_j$ , i.e.  $\eta_i = \eta_k \eta_j$ , then  $\eta_i \eta_m = \eta_k \eta_j \eta_m$ , i.e. an edge of color k connects  $\overline{\eta}_m(\eta_i)$  to  $\overline{\eta}_m(\eta_j)$ . Moreover, the  $\overline{\eta}_m$ 's are distinct endomorphisms of G; for if  $\overline{\eta}_m(\eta_i) = \overline{\eta}_l(\eta_i)$  for each i, then in particular,

 $\bar{\eta}_m(1) = \bar{\eta}_l(1)$ , i.e. m = l.

We claim G has no other endomorphisms. Let  $\varepsilon \in \operatorname{End}(G)$  and set  $\eta_m = \varepsilon(1)$ . We claim that  $\varepsilon = \overline{\eta}_m$ . Let  $\eta_i \in V(G)$ . Then  $\eta_i$  is connected to 1 by an edge of color i, as  $\eta_i = \eta_i \cdot 1$ . Therefore,  $\varepsilon(\eta_i)$  is connected to  $\varepsilon(1) = \eta_m$  by an edge of color i, since  $\varepsilon$  is an endomorphism. Hence

$$\varepsilon(\eta_i) = \eta_i \eta_m$$
, i.e.  $\varepsilon = \overline{\eta}_m$ .

(b) First we observe that there is an arbitrarily large number of simple graphs, which are rigid and have no homomorphisms into each other. Just take a large enough s and different choices of a, b, c with a+b+c=s-1 in the construction in the solution of 12.20. Let  $G_1, \ldots, G_N$  be these rigid graphs, where  $N=2 \mid \Sigma \mid$ . Each point of each  $G_i$  is contained in an odd circuit of length s, which is shortest in each of them.

Now we replace each edge (x, y) of color k in G (constructed in part (a)) by a chain as shown in Fig. 96, where the two edges incident with a  $G_i$  are supposed to be attached to it at the same two points in each case. Loops are replaced

similarly, except that x and y coincide.

The resulting graph G' clearly has endomorphisms  $\tilde{\eta}_i$  corresponding to each endomorphism  $\bar{\eta}_i$  of G. We claim it has no other endomorphisms. Since G' has no other s-circuits or shorter circuits than those in the  $G_i$ 's it follows that any  $\varepsilon \in \operatorname{End}(G)$  maps the points of  $G_i$ 's onto points of  $G_i$ 's. Since these are rigid and cannot be mapped into each other it follows that  $\varepsilon$  maps each  $G_i$  into another (or the same) copy of the same graph. The points of G are adjacent to points in different  $G_i$ 's. Therefore,  $\varepsilon(V(G)) \subseteq V(G)$ . It is also easy to see that if x and y are connected in G by an edge of color k, i.e. they are connected in G' by a chain of  $G_{2k-1}$ 's and  $G_{2k}$ 's (see Fig. 92), then so are  $\varepsilon(x)$  and  $\varepsilon(y)$ , and there is a unique way to map the (x, y)-chain onto the  $(\varepsilon(x), \varepsilon(y))$ -chain. Therefore,  $\varepsilon = \tilde{\eta}_i$  for some i. [Z. Hedrlin–A. Pultr, Monatsh. f. Math. 68 (1964) 213–217; see also Z. Hedrlin–J. Lambek, J. of Alg. 11 (1969) 195–212].




Fig. 96

**25.** (a) Suppose End  $(G) = \{1, \alpha, \omega\}$ , where  $\omega$  is a 0-element:  $\alpha\omega = \omega\alpha = \omega$ , 1 is the identity and  $\alpha^2 = 1$ . Let us remove the edge e = (x, y). If  $\omega$  does not remain an endomorphism, there exists an edge  $(x_1, y_1) \in E(G - e)$  with  $\omega(x_1) = x$ ,  $\omega(y_1) = y$ . Similarly, if  $\alpha$  does not remain an endomorphism, then there is an edge  $(x_2, y_2) \in E(G - e)$  with  $\alpha(x_2) = x$ ,  $\alpha(y_2) = y$ . But then

$$x_2 = \alpha(x) = \alpha(\omega(x_1)) = \omega(x_1) = x, \quad y_2 = y,$$

which means that  $(x_2, y_2)$  is not an edge of G - (x, y), a contradiction.

(b) Consider the construction in the solution of 12.21b, but choose s such that we have one more rigid graph  $G_{N+1}$  of girth s having no endomorphism into  $G_1, \ldots, G_N$  and conversely. If we add  $G_{N+1}$  as a new component to G' we obtain a graph G'' with End  $(G'') \cong \operatorname{End}(G') \cong \Sigma$ . However, if we connect the point 1 of V(G) to  $G_{N+1}$  by an edge, the resulting graph is rigid [L. Babai, J. Nešetřil].

## § 13. Hypergraphs

1. Consider  $G_H$  as defined in the hint. Obviously,  $G_H$  is connected. Moreover, observe that a circuit of H corresponds to a circuit of  $G_H$  and conversely. Thus H has no circuits, iff  $G_H$  is a tree, which is, in turn, equivalent to

$$|E(G_H)| = |V(G_H)| - 1.$$

Obviously,  $|E(G_H)|$  is the sum of degrees in W, i.e.

$$|E(G_H)| = \sum_{E \in E(H)} |E|.$$

On the other hand,

$$|V(G_H)| = |V(H)| + |E(H)|.$$

Thus (\*\*) is the same as

$$\sum_{E \in E(H)} |E| = |V(H)| + |E(H)| - 1,$$

which is the same as (\*) [see B].

2. (a) We first verify the assertion of the hint. Suppose  $H' = (H - \{E\}) \setminus E$  is connected. Let F be an edge such that  $|E \cap F|$  is maximal. We claim that each point of E - F has degree 1 in H. Let  $x \in E \cap F$ ,  $y \in E - F$  (if such x or y does not exist we have nothing to prove), and assume indirectly that  $y \in E'$  for some  $E' \in E(H)$ ,  $E' \neq E$ . By the assumption F - E, E' - E are non-empty, thus by the connectivity of H' there exists a sequence  $a_0, \ldots, a_k \in V(H) - E$  of points and  $E_0 = E', E_1, \ldots, E_{k+1} = F$  of edges such that  $a_i \in E_i \cap E_{i+1}$  ( $i = 0, \ldots, k$ ). If k is as small as possible, we clearly have that  $a_i \notin E_j$  unless j = i or i + 1. Then the circuit  $(y, E_0, a_0, \ldots, E_k, a_k, E_{k+1}, x, E)$  is not balanced, a contradiction.

Now to prove the problem we use induction on |E(H)|. We may assume H is connected, otherwise we may apply the induction hypothesis with one of its

components.

Let  $x \in V(H)$  and suppose that x is not contained in every edge (if no such point exists the assertion is trivial). Choose an  $E \in E(H)$  such that  $x \notin E$  and

the connected component  $H_1$  of  $(H-\{E\}) \setminus E$  containing x is maximal. If  $(H-\{E\}) \setminus E$  is connected, we already know the assertion is true; so suppose there are edges of H, which do not meet  $V(H_1)$  other than E and let  $H_2$  be the hypergraph formed by all edges F not meeting  $V(H_1)$ . Clearly  $E \in E(H_2)$  and  $|E(H_2)| \geq 2$ . Let T be the set of those points in E which belong to edges not in  $H_2$ . We claim that every  $F \in E(H_2)$  contains T. Suppose there are  $F \in E(H_2)$  and  $t \in T - F$ . Then consider  $(H - \{F\}) \setminus F$ . In this hypergraph the component containing x clearly contains  $V(H_1)$  and also t, a contradiction.

Thus  $\cap$   $F \supseteq T$ . Clearly  $\check{H}_2$  is totally balanced (each partial hypergraph

of a totally balanced one is totally balanced) and so, there are  $E_1$ ,  $F_1 \in E(H_2)$  such that all points of  $E_1 - F_1$  have degree 1 in  $H_2$ . Since the edges of  $H_1'$  meet in T only, these points have degree 1 in H.

(b) We use induction on |E(H)|. Let  $E_0$ ,  $F_0$  be two edges such that each point of  $E_0 - F_0$  has degree 1. Consider the hypergraph H' arising from H by the removal of  $E_0$  and the points of  $E_0 - F_0$ .

Obviously, this is totally balanced and satisfies  $|E_1 \cap E_2| \leq p$  for any two

 $E_1, E_2 \in E(H')$ . So we have, by induction,

$$\sum_{E \in E(H')} (|E| - p) \leq |V(H')| - p.$$

Now observe that  $|E_0 \cap F_0| \le p$  by the assumption. So  $|E_0 - F_0| \ge |E_0| - p$  and so,

$$\sum_{E \in E(H)} (|E| - p) = |V(H')| - p + |E_0| - p \le |E| + |V(H')| + |E| + |E|$$

$$\leq |V(H')| - p + |E_0 - F_0| = |V(H)| - p.$$

3. Consider the points of P to be linearly ordered. Let  $C = (x_1, P_1, \ldots, x_k, P_k)$  be a circuit in  $H, k \geq 3$ . We may assume  $x_1$  is neither the least nor the largest point in C, i.e.  $x_i < x_1, x_j > x_1$  for some i, j. We may suppose i < j; then there is an  $i \leq \mu < j$  such that

$$x_{\mu} < x_1 < x_{\mu+1}$$

Now  $P_{\mu}$  contains  $x_{\mu}$  and  $x_{\mu+1}$  and since it is a path, it contains  $x_1$ . Thus C is balanced.

**4.** If H has no circuit of length at least 3, then it is totally balanced and thus' it satisfies (\*) in 13.2b with p = 2. But (\*) simplifies to

$$|E(H)| \leq |V(H)| - 2,$$

a contradiction [L. Lovász, Beiträge zur Graphentheorie, Teubner (1968) 99-106].

5. (a) The given condition is obviously necessary. Suppose it is satisfied, we show H has a system of distinct representatives.

Define a bipartite graph G on the set of points  $V(H) \cup E(H)$ , by joining  $v \in V(H)$  to  $E \in E(H)$  iff  $v \in E$ . Then, for any  $X \subseteq E(H)$ ,

$$|\Gamma(X)| = |\bigcup_{E \in X} E| \ge |X|$$

by the assumption, so by 7.4, G has a matching, which covers all points in E(H).

The endpoints of edges of this matching in V(H) form a system of distinct representatives of H [P. Hall; see B, Mi].

(b) Again it is trivial that the given condition is necessary. We show it is sufficient. Form the bipartite graph G as above. Then the condition means that

$$(1) |\Gamma(X)| \ge |X| + 1$$

holds for  $\emptyset \neq X \subseteq E(H)$ . By 7.6, there is a subgraph G' of G on the same set of points such that each point of E(H) has degree 2 and G' also satisfies (1). For each  $E \in E(H)$ , let f(E) be the set of points in V(H) adjacent to E in G'. Then |f(E)| = 2, and (1) implies

$$|\bigcup_{E \in X} f(E)| \ge |X| + 1$$

for every  $\emptyset \neq X \subseteq E(H)$ . Hence the pairs f(E) form the edges of a forest on V(H). [L. Lovász, Acta Math. Acad. Sci. Hung. 21 (1970) 443–446.]

- **6.** First solution. Suppose there exists a function  $f: E(H) \to V(H)$  such that  $f(E) \in E$ , f is one-to-one and  $f(E(H)) \supseteq T$ . Then, setting  $h = f^{-1}|_T$ , we can observe that
  - (a) H has a system of distinct representatives,
- (b) h defines a system of distinct representatives of the hypergraph  $H_T^*$ , defined by  $V(H_T^*) = E(H)$ ,  $E(H_T^*) = \{U_x : x \in t\}$ , where  $U_x = \{E \in E(H) : x \in E\}$ .

Conversely, suppose g and h are mappings defining systems of distinct representatives of H and  $H_T^*$ , respectively. We claim H has a system of distinct representatives containing T.

For each  $x \in T$ ,  $x \notin R(g)$  consider the sequence x, h(x), g(h(x)), h(g(h(x))), ... until we get stuck; this clearly happens when  $g(h \ldots g(h(x))) \in T$ . Let  $H_1$  be the set of edges in these sequences. Define

$$f(E) = \begin{cases} h^{-1}(E) & \text{for } E \in H_1, \\ g(E) & \text{for } E \in E(H) - H_1. \end{cases}$$

Then f defines a system of distinct representatives. Clearly,  $f(E) \in E$ . If f(E) = f(E'), then we must have  $E \in H_1$ ,  $E' \notin H_1$  (or conversely) and  $g(E') = h^{-1}(E)$ ; but this means that g(E') and E' are the elements preceding E in the chain containing it, i.e.  $E' \in H_1$ , a contradiction. Finally, every  $x \in T$  belongs to R(f). For if x belongs to one of the chains, then x = f(h(x)); if x does not belong to the chains, then  $x \in R(g)$  and the edge  $g^{-1}(x) \in H_1$ , so  $x = f(g^{-1}(x))$ .

Thus (a) and (b) together are a necessary and sufficient condition for the existence of a system of distinct representatives containing T. By the preceding

problem, they can be rewritten in the form

- (a')  $|V(H')| \ge |E(H')|$  for every partial hypergraph H';
- (b') every  $T' \subseteq T$  meets at least |T'| edges [This is a version of Bernstein's theorem on equicardinality of sets; see B, Mi].

Second solution. Set F = V(H) - T, and add F as a new edge |V(H)| - |E(H)| times to H. Then it is easy to verify that the resulting hypergraph H' has a system of distinct representatives iff H has one containing T. Applying the Hall condition (preceding problem) to H' we obtain a condition, which is equivalent to (a'), (b') above. Details are left to the reader.

7. Take two new points a, b. Define a digraph G on the set  $\{a\} \cup \{b\} \cup E(H_2) \cup U(H_1) \cup V(H_1)$  as follows: join a to all elements of  $E(H_1)$ ; join each  $E \in E(H_1)$  to each  $x \in E$ ; join each  $x \in V(H_1) = V(H_2)$  to all  $F \in E(H_2)$  containing x; finally, join each element of  $E(H_2)$  to b.

Observe that  $H_1$  and  $H_2$  have a common system of distinct representatives iff G has m independent (a, b)-paths. By Menger's theorem, this is equivalent to the property that every set of points of G separating a and b has at least

m points.

Now let  $X \cup Y \cup Z$  separate a and b,  $X \subseteq E(H_1)$ ,  $Y \subseteq V(H_1)$ ,  $Z \subseteq E(H_2)$ . Let  $H'_1, H'_2$  be the hypergraphs formed by the edges in  $E(H_1) - X$  and  $E(H_2) - Z$ , respectively. Obviously, Y must contain  $V(H'_1) \cap V(H'_2)$  and does not have to contain any other point. So the fact that every set of points separating a and b has cardinality at least k is equivalent to saying that

$$|E(H_1) - E(H_1')| + |E(H_2) - E(H_2')| + |V(H_1') \cap V(H_2')| \ge m$$

for every two partial hypergraphs  $H'_1$ ,  $H'_2$  of  $H_1$  and  $H_2$ , respectively. This is the same as the formula given in the problem [R. Rado, B, Mi].

8. (a) For a fixed  $E_0 \in E(H)$ , the sets

$$E_0 \triangle F(F \in E(H))$$

are distinct, as  $E_0 \triangle F_1 = E_0 \triangle F_2$  implies that

$$F_1 = (E_0 \triangle F_1) \triangle E_0 = (E_0 \triangle F_2) \triangle E_0 = F_2.$$

(b) Let  $H_1$  be the hypergraph obtained from  $H \setminus x$  by removing one of each double edge. Obviously,  $H \setminus x$  has |E(H')| double edges, where H' is defined as in the hint. So setting

 $m_1 = |E(H_1)|, m_2 = |E(H')|,$ 

we have

$$m_1 + m_2 = m.$$

By induction on |V(H)|, we find  $m_1$  pairs  $\widetilde{E}_1, \widetilde{F}_1; \ldots; \widetilde{E}_{m_i}, \widetilde{F}_{m_1}$  such that  $\widetilde{E}_i, \widetilde{F}_i \in H_1$  and  $\widetilde{E}_1 - \widetilde{F}_1, \ldots, \widetilde{E}_{m_1} - \widetilde{F}_{m_1}$  are distinct. Similarly, we find  $m_2$  pairs  $E_1^*, F_1^*; \ldots; E_{m_2}^*, F_{m_3}^*$  such that  $E_i^*, F_i^* \in H'$  and  $E_1^* - F_1^*, \ldots$  ...,  $E_{m_2}^* - F_{m_3}^*$  are distinct.

There are sets  $E_i$ ,  $F_i \in H$   $(1 \le i \le m_1)$  such that  $\widetilde{E}_i = E_i$  or  $\widetilde{E}_i = E_i - \{x\}$ ,  $\widetilde{E}'_i = F_i$  or  $\widetilde{F}_i = F_i - \{x\}$ . Then, obviously,  $E_i - F_i$   $(i = 1, \ldots, m_1)$  are distinct

For each  $E_j^*$   $(1 \le j \le m_2)$ , we will denote by  $E_j'$  one of  $E_j^*$  and  $E_j^* \cup \{x\}$  (both are edges of H), by the following rule. If  $E_j^* - F_j^*$  does not occur among  $E_i - F_i$   $(i = 1, \ldots, m_1)$ , then let  $E_j' = E_j^*$ . If  $E_j^* - F_j^* = E_{i_0} - F_{i_0}$ , then let  $E_j' = E_j^* \cup \{x\}$ . Then  $E_j' - F_j^*$  will be different from the sets  $E_i - F_i$ ;

for if  $E'_j - F^*_j = E_i - F_i$ , then  $E_i - F_i = E_{i_0} - F_{i_0}$  which is impossible. Thus, if we consider the differences  $E_1 - F_1, \ldots, E_{m_1} - F_{m_1}, E'_1 - F^*_1, \ldots, E'_{m_2} - F^*_{m_2}$ , they are distinct and their number is  $m_1 + m_2 = m$  [J. Marica, J. Schönheim, Canad. Math. Bull. 12 (1969) 635–637.]

(c) If m=2 the assertion is obviously true. Suppose  $m\geq 3$  and use induction on m.

Let  $x \in V(H)$ ; we may assume x belongs to at least one edge  $E_0$ , but not to all. Suppose first that x has degree 1. Then  $H_1 = H - x$  has m-1 edges, so there are at least m-1 edges representable in one of the forms  $E \cap F$ ,  $E \cup F$  ( $E, F \in E(H_1), E \neq F$ ). None of these sets contains x. Thus, the set  $E_0 \cup F$  yields an  $m^{\text{th}}$  set or the desired form for an arbitrary F.

Secondly, let the degree of x be between 2 and m-2 (inclusively). Then

consider

$$H_1 = H - x$$
,  $H_2 = H - E(\dot{H}_1)$ 

They both have at least two edges, so, by the induction hypothesis, there are at least  $|E(H_i)|$  sets representable in the form  $E \cup F$  or  $E \cap F$ ,  $E \neq F \in E(H_i)$ , (i = 1, 2). Moreover,  $x \notin E \cap F$ ,  $x \notin E \cup F$ , if  $E, F \in E(H_1)$  but  $x \in E \cap F$ ,  $E \cup F$ , if  $E, F \in E(H_2)$ . Thus we have at least  $|E(H_1)| + |E(H_2)| = m$  sets of the required form altogether.

Finally, if x belongs to all edges but  $E_1$ , then set  $H_2 = H - \{E_1\}$ .  $H_2$  has at least two edges, hence there are at least  $|E(H_2)| = m-1$  sets of the form  $E \cap F$  or  $E \cup F$ ,  $E \neq F \in E(H_2)$ . All these sets contain x, thus  $E_1 \cap F$  yields an  $m^{\text{th}}$  for any  $F \in E(H_2)$  [D.E. Daykin, L. Lovász, J. London Math. Soc. 12

(1976) 225-230].

**9.** If  $H_1$ ,  $H_2$  are defined as in the hint, then (by induction on |V(H)|) we may assume there are permutations  $\sigma_i$  of  $E(H_i)$  (i=1,2) such that  $E \cap \sigma_i(E) = \emptyset$  for each  $E \in E(H_i)$ .

Now define  $\sigma$  as follows: Let  $E \in E(H)$ . If  $x \notin E$ , then let

$$\sigma(E) = \begin{cases} \sigma_1(E) & \text{if } \sigma_1(E) \cup \{x\} \notin E(H), \\ \sigma_1(E) \cup \{x\} & \text{if } \sigma_1(E) \cup \{x\} \in E(H). \end{cases}$$

If  $x \in E$ , then let

$$\sigma(E) = \sigma_2(E - \{x\}).$$

It is easy to check that  $E \cap \sigma(E) = \emptyset$  for each E. We show that  $\sigma$  is a permutation. Let  $E \in H$ . If  $x \in E$ , then  $E = \sigma(\sigma_1^{-1}(E - \{x\}))$ . If  $x \notin E$  but  $E \cup \{x\} \in E(H)$ , then  $E = \sigma(\sigma_2^{-1}(E))$ . Finally, if  $E \cup \{x\} \notin E(H)$ , then  $E = \sigma(\sigma_1^{-1}(E))$ . So each edge of H is the image of some edge under  $\sigma$ , i.e.  $\sigma$  is a permutation. [P. Erdős, J. Herczog, J. Schönheim, Israel J. Math. 8 (1970) 408-412.]

10. (a) First solution. We may assume  $n \ge 2$ . Then there is a point v contained in some edge but not in all.  $X = \{v\}$  satisfies the requirement of the assertion in the hint for k = 1.

Suppose we have a set X' such that |X'| = k - 1 and  $H_{X'}$  has at least k distinct edges. If  $H_{X'}$  has more than k distinct edges, then so does  $H_{X' \cup \{y\}}$  for any  $y \in V(H) - X'$ , and we can set  $X = X' \cup \{y\}$ . Thus we may assume that  $H_{X'}$  has exactly k distinct edges. Since k < n,  $H_{X'}$  has two distinct edges

 $E, \ F \ \text{with} \ E \cap X' = F \cap X'.$  Let  $y \in E \triangle F$ , and  $X = X' \cup \{y\}$ . Obviously,  $H_X$  has more distinct edges than  $H_{X'}$ , as  $E \cap X \neq F \cap X$ . Thus, we have (for k = n - 1) a set  $X = V(H) - \{x\}$  such that  $H_X = H \setminus x$  has n distinct edges.

Second solution. Let x be any point. If  $H \setminus x$  has fewer distinct edges than H, then there must be an  $E_x \in E(H)$  such that  $x \in E_x$  and  $F_x = E_x - \{x\} \in E(H)$ . Let us fix one such pair  $(E_x, F_x)$  of edges for each point x (if, for

some x, no such pair exists we are done).

Define a graph G on V(G) = E(H) by joining E to F iff  $E = E_x$ ,  $F = F_x$  for some x. This simple graph G has n points and n edges, so it contains a circuit. Let  $(E_1, \ldots, E_k)$  form a circuit in G ( $E_i \in E(H)$ ,  $k \geq 3$ ) and let  $E_1 = E_x$ ,  $E_k = F_x$  (say). Then  $x \in E_1$ ,  $x \notin E_k$  and thus, there is an index j such that  $x \in E_j$ ,  $x \notin E_{j+1}$ . But  $E_j$  and  $E_{j+1}$  are adjacent, so  $\{E_j, E_{j+1}\} = \{E_y, F_y\}$  for some y. Since  $E_y$  and  $F_y$  differ at the element y only, we must have y = x and  $E_j = E_y - E_x$ ,  $E_{j+1} = F_y = F_x$ . This is, however, a contradiction as  $(E_j, E_{j+1}) \neq (E_1, E_k)$  [J.A. Bondy, J. Comb. Theory B 12 (1972) 201–202].

(b) Suppose indirectly that  $H \setminus x$  contains at most m-2 distinct edges. Construct a graph G on E(H) as follows. For every  $x \in V(H)$ , select two pairs  $E_x \supseteq F_x$  and  $E_x' \supseteq F_x'$  of edges such that  $E_x - F_x = E_x' - F_x' = \{x\}$ . Such pairs exist otherwise  $H \setminus x$  has at least m-1 distinct edges. Now connect the pairs  $(E_x, F_x)$  and also the pairs  $(E_x, F_x)$  in G. Thus

$$|V(G)| = m, |E(G)| = 2n.$$

1° G is bipartite. In fact

$$V_1 = \{ E \in E(H) : |E| \text{ is odd} \},\$$
  
 $V_2 = \{ E \in E(H) : |E| \text{ is even} \}$ 

defines a 2-coloration of it.

2° No two points of G are connected by three independent paths. Suppose indirectly that E, E' are two points of G connected by three independent paths  $P_1$ ,  $P_2$ ,  $P_3$ . Let  $x \in E - E'$  (say). Then, on each path  $P_i$  we find a last point  $E_i$  containing x. So if  $F_i$  is the point following  $E_i$  on  $P_i$ , then  $x \in E_i - F_i$ . Since  $E_i$ ,  $F_i$  are adjacent they differ at one point only, i.e.  $(E_i, F_i) = (E_x, F_x)$  or  $(E'_x, F'_x)$ . This cannot hold for all three values of i. This proves 2°.

Now we estimate the number of edges in G. It follows from  $2^{\circ}$  that each block in G is a single edge or a single circuit. Let G have c components, a cutting

edges and b other blocks, of sizes  $k_1, \ldots, k_b$ . By 1°,  $k_i \geq 4$ . Also

$$m = |V(G)| = c + a + \sum_{i=1}^{b} (k_i - 1),$$
  $2n = |E(G)| = a + \sum_{i=1}^{b} k_i.$ 

Thus

$$m \ge 1 + a + \sum_{i=1}^{b} (k_i - 1) \ge 1 + \frac{3}{4}a + \sum_{i=1}^{b} \left(\frac{3}{4}k_i\right) = 1 + \frac{3}{4}(2n) = 1 + \frac{3}{2}n,$$

a contradiction [B. Bollobás].

(c) We use induction on n. Let  $H_1$ ,  $H_2$  be as in the hint. If

$$|E(H_1)| > \sum_{i=0}^{k-1} {n-1 \choose i}$$

then, by the induction hypothesis, it has a set X of n points such that each subset of X can be written as  $X \cap E$ ,  $E \in E(H_1)$ . Then, obviously, the same set X will be good for H as well. Thus we may assume

$$|E(H_1)| \leq \sum_{i=1}^{k-1} \binom{n-1}{i}$$

Then

$$|E(H_2)| = |E(H) - E(H_1)| > \sum_{i=0}^{k-1} \binom{n}{i} - \sum_{i=0}^{k-1} \binom{n-1}{i} = \sum_{i=0}^{k-2} \binom{n-1}{i}$$

and so, by the induction hypothesis,  $H_2$  contains a set X' of k-1 elements such that each subset of X' can be written in the form  $X' \cap E$ ,  $E \in E(H_2)$ . Now set  $X = X' \cup \{x\}$ . If  $Y \subseteq X'$ , then  $Y = X' \cap E = X \cap E$  for some  $E \in E(H_2) \subseteq E(H)$ . If  $Y \subseteq X$ ,  $x \in Y$ , then we know  $Y - \{x\} = X' \cap E$  for some  $E \in E(H_2)$ . Now by definition of  $H_2$ ,  $E = F - \{x\}$  for some  $F \in E(H)$ ,  $x \in F$ . Then  $X \cap F = Y$ . Thus we have shown that every subset Y of X can be represented in the form  $X \cap E$ ,  $E \in E(H)$  [N. Sauer, J. Comb. Theory 13 (1972) 145–147].

11. We use induction on n. Let  $x \in V$  and

$$\mathcal{R}' = \{A \in E(H) : x \in A\}, \ \mathcal{R}'' = \{A \in E(H) : x \notin A\},$$
 
$$\mathcal{B}' = \{B \in E(K) : x \in B\}, \ \mathcal{B}'' = \{B \in E(K) : x \notin B\}.$$

Then clearly

$$E(H) = \mathcal{R}' \cup \mathcal{R}'', \ E(K) = \mathcal{B}' \cup \mathcal{B}'', \ E(H) \cap E(K) = (\mathcal{R}' \cap \mathcal{B}') \cup (\mathcal{R}'' \cup \mathcal{B}'').$$
 By the induction hypothesis,

$$|\mathcal{A}' \cap \mathcal{B}'| \leq \frac{1}{2^{n-1}} |\mathcal{A}'| \cdot |\mathcal{B}'|$$

and omitting x from all members of  $\mathcal{A}''$  and  $\mathcal{B}''$ , we similarly get

$$|\mathscr{R}''\cap \mathscr{B}''| \leq rac{1}{2^{n-1}}|\mathscr{R}''|\cdot |\mathscr{B}''|.$$

So we know that

$$|\mathit{E}(\mathit{H}) \cap \mathit{E}(\mathit{K})| \leq \frac{1}{2^{n-1}} (|\mathcal{R}'| \cdot |\mathcal{B}'| + |\mathcal{R}''| \cdot |\mathcal{B}''|)$$

and it suffices to prove that

$$|\mathcal{A}'|\cdot|\mathcal{B}'|+|\mathcal{A}''|\cdot|\mathcal{B}''|\leq \frac{1}{2}(|\mathcal{A}'|+|\mathcal{A}''|)(|\mathcal{B}'|+|\mathcal{B}''|).$$

This can be re-written as

$$(|\mathcal{A}'| - |\mathcal{A}''|)(|\mathcal{B}'| - |\mathcal{B}''|) \le 0.$$

Observe that, if  $A \in \mathcal{A}''$ , then  $A - \{x\} \in \mathcal{A}'$ , thus  $|\mathcal{A}'| \geq |\mathcal{A}''|$ . Analogous reasoning gives  $|\mathcal{B}'| \leq |\mathcal{B}''|$ . This proves (\*) [D. Kleitman, J. Comb. Theory 1 (1966) 153-155].

12. (a) Let f(X) denote the number of edges contained in X or disjoint from X, for  $X \subseteq V(H)$ . Set |V(H)| = n. Then

$$\sum_{X\subseteq V(H)} f(X) = |E(H)| 2^{n-r+1},$$

because each given edge E is counted  $2^{n-r+1}$  times (there are  $2^{n-r}$  sets X disjoint from E and  $2^{n-r}$  sets X containing E). Thus,

$$\frac{1}{2^n} \sum_{X \subseteq V(H)} f(X) = \frac{|E(H)|}{2^{r-1}}.$$

The left-hand side being the average of f(X), there must be a set X below average, i.e. such that

 $f(X) \leq \frac{|E(H)|}{2r-1}$ 

as stated.

(b) Let  $\varphi \colon V(H) \to \{1, \ldots, r\}$  be an r-coloration of V(H) and denote by  $||\varphi||$  the number of those edges E all of whose points are differently colored by  $\varphi$ . Then

 $\sum_{m} ||\varphi|| = r! r^{n-r} |E(H)|,$ 

because there are r! ways to color a given edge E and  $r^{n-r}$  ways to color the rest (n is again |V(H)|). Thus the average of  $|\varphi|$  is

$$\frac{1}{r^n} \sum_{\varphi} ||\varphi|| = \frac{r!}{r^r} |E(H)|.$$

Since there is a  $\varphi$  such that  $||\varphi||$  is at least equal to the average, this proves the assertion.

13. Let d(x) denote the degree of x in H. Then, for any  $E \in E(H)$ ,

$$\sum_{x \in E} d(x) = \sum_{F \in E(H)} |E \cap F| = r + \sum_{F \in E(H) - \{E\}} |E \cap F| \le r + (m-1)k.$$

Sum this over all edges E, then

ever all edges 
$$E$$
, then
$$\sum_{E \in E(H)} \sum_{x \in E} d(x) = \sum_{x \in V(H)} d(x)^2 = |V(H)| \frac{\sum_{x \in V(H)} d(x)^2}{|V(H)|} \ge$$

$$\geq |V(H)| \left( \frac{\sum_{x \in V(H)} d(x)}{|V(H)|} \right)^2 = \frac{\left( \sum_{E \in E(H)} |E| \right)^2}{|V(H)|} = \frac{m^2 r^2}{|V(H)|}.$$

Hence

$$\frac{m^2r^2}{\mid V(H)\mid} \leq m \big(r + (m-1)\,k\big)$$

or, equivalently,

$$|V(H)| \ge \frac{mr^2}{r + (m-1)k}.$$

[K. Corrádi, Problem at the Schweitzer Competition, Mat. Lapok 20 (1969) 159–162.]

14. Supposing m > n, we have  $m - d(x) > n - d(x) \ge n - |E|$  for any pair x, E such that  $x \notin E$ , and hence

$$\frac{d(x)}{m - d(x)} < \frac{\mid E \mid}{n - \mid E \mid}$$

(because  $|E| \neq 0$ , n). Summing for every pair  $x \notin E$ , we obtain

$$\sum_{\mathbf{x}} \left( m - d(\mathbf{x}) \right) \frac{d(\mathbf{x})}{m - d(\mathbf{x})} < \sum_{E} \left( n - |E| \right) \frac{|E|}{n - |E|}$$

or, equivalently,

$$\sum_{x}d(x)<\sum_{E}|E|,$$

which is not the case, because here we have equality. The contradiction proves that  $m \leq n$  [cf. P. Crawley-R.P. Dilworth, Algebraic Theory of Lattices, Prentice-Hall, 1973, proof of 14.2].

15. (a) If  $\emptyset \in E(H)$ , then clearly |E(H)| = 1 and we are done. So suppose  $\emptyset \notin E(H)$ . Clearly  $V(H) \notin E(H)$ . Let  $x \in V(H)$ ,  $E \in E(H)$ ,  $x \notin E$ . By the preceding problem, it suffices to prove that

$$d(x) \leq |E|$$
.

Let  $E_1, \ldots, E_d$  be those edges containing x. Then  $E_1, \ldots, E_d$  meet E by the assumption and their intersections with E are disjoint since they cannot have any common element other than x. Thus,  $|E| \geq d = d(x)$  as stated.

(b) Let 
$$V(H) = \{x_1, \ldots, x_n\}$$
,  $E(H) = \{E_1, \ldots, E_m\}$ ,  $a_{ij} = \begin{cases} 1 & \text{if } x_i \in E_j, \\ 0 & \text{otherwise,} \end{cases}$ 

$$\mathbf{a}_j = \begin{pmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{nj} \end{pmatrix}.$$

We claim that  $a_1, \ldots, a_m$  are linearly independent. This clearly also proves that  $m \leq n$ . Suppose

(1) 
$$\sum_{j=1}^{m} \eta_{j} \mathbf{a}_{j} = \mathbf{0};$$

we show every  $\eta_i$  is 0. Multiplying (1) by  $\mathbf{a}_k$ , we get

$$\sum_{j=1}^m \eta_j \mathbf{a}_j \mathbf{a}_k = 0.$$

13.16

Since by assumption  $\mathbf{a}_i \mathbf{a}_k = \lambda$  for  $i \neq k$ , this yields

$$\lambda \sum_{j=1}^m \eta_j + \eta_k(\mathbf{a}_k^2 - \lambda) = 0.$$

If  $\mathbf{a}_k^2 = |E_k| = \lambda$ , then  $E_k$  is contained in every other edge, which is excluded except when  $E_k = \emptyset$ , in which case  $m = 1 \le n$ . So suppose  $\mathbf{a}_k^2 > \lambda$  for every k. Then

(2) 
$$\eta_k = \frac{-\lambda}{\mathbf{a}_k^2 - \lambda} \sum_{j=1}^m \eta_j$$

and so

$$\sum_{k=1}^{m} \eta_k = \left(\sum_{j=1}^{m} \eta_j\right) (-\lambda) \sum_{k=1}^{m} \frac{1}{\mathbf{a}_k^2 - \lambda}$$

or, equivalently,

$$\left(\sum_{j=1}^m \eta_j\right) \left(1 + \lambda \sum_{k=1}^m \frac{1}{\mathbf{a}_k^2 - \lambda}\right) = 0.$$

Since the second factor is strictly positive we derive

$$\sum_{j=1}^{m} \eta_j = 0$$

and hence by (2),  $\eta_k = 0$  as stated [This is a generalized version of Fisher's inequality; see Hall].

16. Let  $x \in V(H)$ , and let a and b denote the degrees of x in H and K, respectively. Then x is counted in

$$a(m'-b) + b(m-a)$$

terms of the left-hand side and

$$a(m-a) + b(m'-b)$$

terms of the right-hand side. Their difference is

$$a(m'-b) + b(m-a) - a(m-a) - b(m'-b) =$$

$$= (a-b)^2 - (m'-m)(a-b) \ge 0,$$

since a-b is an integer and m'-m=0 or 1 [J.B. Kelly, Combin. Structures and Appl. Proc. Calgary Conference, Gordon and Breach, 1970, 201–207].

17. (a) As in the hint, assume that  $x \in F_1, \ldots, F_d, x \notin F_{d+1}, \ldots, F_m$ . Define two hypergraphs H, K on  $V = \bigcup_{i=1}^m F_i$ , by letting E(H) and E(K) consist of m-d copies of  $F_1-\{x\},\ldots,F_d-\{x\}$  and d copies of  $F_{d+1},\ldots,F_m$ , respectively.

Then |E(H)| = |E(K)| = d(m-d). Also we have

$$egin{align} \sum_{A \in E(H)} \sum_{B \in E(K)} |A igtriangle B| &= (2k-1)\, d^2(m-d)^2, \ \sum_{\{A,A'\} \subseteq E(H)} |A igtriangle A'| &= 2k(m-d)^2iggl( rac{d}{2} iggr), \ \sum_{\{B,B'\} \subseteq E(K)} |B igtriangle B'| &= 2kd^2iggl( rac{m-d}{2} iggr). \end{aligned}$$

Thus by the inequality in 13.16,

$$(2k-1)\,d^2(m-d)^2\geq 2kigg[(m-d)^2iggl(d\ 2iggr)+d^2iggl(m-d\ 2iggr)iggr],$$

whence the inequality of the problem follows after cancellations.

(b) Replacing  $F_i$  by  $F_i \triangle F_m$  the conditions remain unchanged. Thus we may suppose  $|F_1| = \ldots = |F_{m-1}| = 2k$ ,  $F_m = \emptyset$ . Then, if d is the degree of any point, then by (a)

 $d(m-d) \leq km.$ 

Assume first that  $k+2 \le d \le m-k-2$ . Then

$$km \ge d(m-d) \ge (k+2)(m-k-2),$$

i.e.

$$m \leq \left[\frac{(k+2)^2}{2}\right] \leq k^2 + k + 2,$$

and we are finished. Hence we may suppose that  $d \leq k+1$  or  $d \geq m-k-1$ 

holds for each degree d.

Let  $x_1, \ldots, x_l$  be those points with degree at least m-k-1. Suppose first that  $l \geq k+1$ . Clearly there is at most one  $F_j$  containing all elements of  $\{x_1, \ldots, x_{k+1}\}$ . Thus, m-2  $F_j$ 's miss at least one point of  $\{x_1, \ldots, x_{k+1}\}$ ; one misses k+1 of them (the empty set  $F_m$ ) and one may contain all of them. Thus, there are at most k(m-2)+k+1 incidences between these points and the sets  $F_j$ . On the other hand, the number of such incidences is at least (m-k-1)(k+1). Thus

$$k(m-2) + k + 1 \ge (m-k-1)(k+1),$$

whence

$$m \le k^2 + k + 2$$

and we are finished again. So we may assume that there are  $l \leq k$  points  $x_1, \ldots, x_l$  only with degree at least m - k - 1. By our previous assumption, the remaining points have degree at most k + 1.

There is a set  $F_j$ ,  $1 \le j \le m-1$  such that  $F_j$  contains less than k elements of  $\{x_1, \ldots, x_l\}$ . Otherwise, we would have l=k and  $x_1, \ldots, x_k$  would be common elements of  $F_1, \ldots, F_{m-1}$ . But, then the sets  $F_1 - \{x_1, \ldots, x_k\}$   $\ldots, F_{m-1} - \{x_1, \ldots, x_k\}$  are disjoint (by  $F_i \triangle F_j = 2k$ ) and so we would have a situation which we have excluded.

Thus we may assume that, e.g.  $F_1$  contains k+1 (or more) points whose degrees are at most k+1. Hence the number of incidences between points in  $F_1$  and the sets  $F_1, F_2, \ldots, F_m$  is at most  $(k-1)(m-1)+(k+1)^2$ . On the other hand, this number is exactly 2k+(m-2)k, since each of  $F_2, \ldots, F_{m-1}$  intersects  $F_1$  in exactly k elements. Thus

$$2k + (m-2)k \le (k-1)(m-1) + (k+1)^2, \ m \le k^2 + k + 2$$

as stated.

Equality holds if k is such that there exists a finite projective plane of order k. Let us add k-1 further points and let  $F_1, \ldots, F_{k^2+k+1}$  be the lines of the plane together with these k-1 points; let  $F_{k^2+k+2} = \emptyset$ . Then these sets satisfy our conditions [M. Deza, J. Comb. Theory 16 (1974) 166-167].

- 18. (a) Arrange all subsets of V(H) into pairs (X, V(H) X). Then we will have  $2^{n-1}$  pairs and E(H) can contain at most one member of each. Hence the assertion follows.
- (b) First solution. Let H' be the hypergraph with V(H') = V(H) and  $E(H') = \{X : X \subseteq E \in E(H)\}$ . Consider a permutation  $\sigma$  of the edges of H' for which  $E \cap \sigma(E) = \emptyset$  ( $E \in E(H')$ ); such a  $\sigma$  exists by 13.9. The edges  $\sigma(E)$ ,  $E \in E(H)$  cannot be edges of H since  $E \cap \sigma(E) = \emptyset$ , while any two edges of H intersect. Thus

 $(1) \qquad |E(H')| \ge 2 |E(H)|.$ 

Observe that no two edges of H' cover V(H); for if  $E, F \in E(H')$  have  $E \cup F = V(H)$ , then choose  $E_1, F_1 \in E(H)$  so that  $E \subseteq E_1, F \subseteq F_1$  and then  $E_1 \cup F_1 = V(H)$ , a contradiction. Hence E(H') contains at most one of X, V(H) - X for any  $X \subseteq V(H)$  and, therefore,

$$|E(H')| \le 2^{n-1}.$$

(1) and (2) prove the assertion [J. Schönheim, Combinatorics, London Math.

Soc. Lecture Notes Series 13 (1974) 139-140].

Second solution. Consider the hypergraph  $H_1$  with  $V(H_1) = V(H)$ ,  $E(H_1) = \{X : X = E - F, E, F \in E(H)\}$ , and set  $H_2 = H \cup H_1$ . No edge of  $H_1$  is an edge of H, since  $E - F = E_0$  ( $E, F, E_0 \in E(H)$ ) would imply  $F \cap E_0 = E(H)$ , which is not allowed. Moreover,  $|E(H_1)| \ge |E(H)|$  by 13.8b. Thus

(3) 
$$|E(H_2)| \ge 2 |E(H)|$$
.

Now no two edges of  $H_2$  cover V(H), because each edge of  $H_2$  is contained in an edge of H. Hence, as before,

$$(4) |E(H_2)| \le 2^{n-1},$$

and (3), (4) imply the assertion of the problem [D.E. Daykin, L. Lovász, J. London Math. Soc. 12 (1976) 225-230].

Third solution. Let  $\mathcal{A} = \{X \subseteq V(H) : X \subseteq E \in E(H)\}$ ,  $\mathcal{B} = \{X \subseteq V(H) : X \supseteq E \in E(H)\}$ . Then by (a),  $|\mathcal{A}| \leq 2^{n-1}$  and  $|\mathcal{B}| \leq 2^{n-1}$ . By 13.11,

$$|E(H)| \le |\mathcal{A} \cap \mathcal{B}| \le \frac{1}{2^n} |\mathcal{A}| |\mathcal{B}| \le 2^{n-2}$$

[D. Kleitman].




Fig. 97

19. We may assume each one-element set belongs to E(H), as well as  $\emptyset$  and V(H). If  $E[\neq V(H) \text{ or } \emptyset]$  is any edge of H, then there is a unique minimal edge  $E_1$  properly containing E; for if  $E_1$ ,  $E_2$  were two such edges, then we would have  $E_1 \subset E_2$ ,  $E_2 \subset E_1$  or  $E_1 \cap E_2 = \emptyset$  by the assumption; however, any of these is impossible. If  $E = \emptyset$ , then it is contained in n minimal members of H; E = V(H) is contained in none. This way we count m + n - 2 edges.

Consider any  $E \in E(H)$ ,  $|E| \ge 2$  and consider the maximal edges properly contained in E. They cover E and are disjoint, so their number is at least 2. So E is counted at least twice. If |E| = 1 E is counted once;  $E = \emptyset$  is not counted. Hence

$$m+n-2 \ge 2(m-n-1)+n$$
,

or, equivalently,

$$m \leq 2n$$
.

This number can be attained; in fact, if G is any arborescence with outdegrees 2 at all points except n endpoints and, if for each point x of G, we consider the set  $E_x$  of endpoints accessible from x, then the hypergraph formed by edges  $E_x$  and the empty set has n points and 2n edges (Fig. 97).

**20.** For n=1 the problem is obvious. We use induction on n. Suppose we have the chains  $K_1, \ldots, K_t$  of  $2^S$ , where |S|=n and add an  $(n+1)^{st}$  point x. For each chain  $K_i=\{E_r,\ldots,E_{n-r}\}$  we define two new chains

$$K'_i = \{E_r, E_r \cup \{x\}, E_{r+1} \cup \{x\}, \ldots, E_{n-r} \cup \{x\}\}\$$

and

$$K_i'' = \{E_{r+1}, \ldots, E_{n-r}\}$$
 (provided  $|K_i| \ge 2$ ).

Obviously, both  $K_i'$  and  $K_i''$  are symmetric chains of subsets of  $S \cup \{x\}$ . Moreover, any subset X of  $S \cup \{x\}$  belongs to exactly one of them. For if  $X \subseteq S \cup \{x\}$  and  $x \in X$ , then let  $X - \{x\} \in K_i$ ; we will have  $X \in K_i'$ . If  $X \subseteq S$ , then let  $X \in K_i$ ; if X is the first member of  $K_i$ , then  $X \in K_i'$ ; if X is a later member, then  $X \in K_i'$ . Similarly one sees that no X belongs to more than one  $K_i'$  or  $K_i''$ .

Thus the chains  $K'_i$ ,  $K''_i$  yield a decomposition of all subsets of  $S \cup \{x\}$  into symmetric chains and the induction is complete [N.G. de Bruijn, C. van E. Tengbergen, D. Kruijswijk, Nieuw Arch. Wiskunde 23 (1949-51) 191-193].

**21.** (a) First proof: Let  $K_1, \ldots, K_t$  be the chains of subsets of V(H) defined

in 13.20. Since each subset X with  $|X| = \left[\frac{n}{2}\right]$  belongs to exactly one  $K_i$  and conversely, we have  $t = \left(\left[\frac{n}{2}\right]\right)$ . Any  $K_i$  contains at most one edge of H, as

any two edges of  $K_i$  are comparable (one contains the other). Hence  $|E(H)| \leq t$ . [E. Sperner, Math. Z. 27 (1928) 544-548. This proof is due to de Bruijn,

C. A. van E. Tengbergen and D. Kruijswijk, ibid.]

Second proof: Let us consider those permutations  $(x_1, \ldots, x_n)$  of V(H) for which  $\{x_1, \ldots, x_k\} \in E(H)$  for some k. For any edge E with |E| = k, we get k!(n-k)! such permutations by putting the elements of E in the first k places and the other elements in the last (n-k) places. No permutation is counted more than once; for if  $(x_1, \ldots, x_n)$  is a permutation, then it cannot happen that  $E_1 = \{x_1, \ldots, x_{k_1}\}$  and  $E_2 = \{x_1, \ldots, x_{k_2}\}$  are both edges since then one of  $E_1$ ,  $E_2$  would contain the other.

Thus we count k!(n-k)! permutations with any edge E(k=|E|), and,

altogether, we count at most n! permutations. Since

$$k!(n-k)! = \frac{n!}{\binom{n}{k}} \ge \frac{n!}{\binom{n}{\frac{n}{2}}} = \left[\frac{n}{2}\right]! \left[\frac{n+1}{2}\right]!,$$

this implies that

$$|E(H)| \cdot \left[\frac{n}{2}\right]! \left[\frac{n+1}{2}\right]! \le n!,$$
 $|E(H)| \le \frac{n!}{\left[\frac{n}{2}\right]! \left[\frac{n+1}{2}\right]!} = \binom{n}{\left[\frac{n}{2}\right]}.$ 

[D. Lubell, J. Comb. Th. 1 (1966) 299].

(b) Let H be a clutter on n points with exactly  $\binom{n}{\left\lceil \frac{n}{2} \right\rceil}$  edges. Then we must

have equality in all estimations used in the second proof of part (a). In particular, if  $E \in E(H)$ , |E| = k, then

$$k!(n-k)! = \left[\frac{n}{2}\right]! \left[\frac{n+1}{2}\right]!$$
, i.e.  $\frac{n}{\binom{n}{k}} = \frac{n!}{\binom{n}{\frac{n}{2}}}$ ,

whence 
$$k = \left[\frac{n}{2}\right]$$
 or  $\left[\frac{n+1}{2}\right]$ .

Now there are two cases:

 $1^{\circ}$  n=2p. Then |E|=p for any  $E\in E(H)$  and so, E(H) consists of all ptuples of V(H).

 $2^{\circ}$  n=2p+1. Then |E|=p or p+1 for any  $E\in E(H)$ . We claim that every edge has the same cardinality, i.e. E(H) consists of all p-tuples or of all (p+1)-tuples.

If  $(x_1, \ldots, x_n)$  is any permutation of the points, then this permutation must be counted in the second solution; i.e. either  $x_1, \ldots, x_p$  or  $x_1, \ldots, x_{p+1}$  is an edge. Thus, if  $X \subset Y \subset V(H)$ , |X| = p, |Y| = p + 1, then one of X, Y

is an edge.

Now suppose there is an edge  $E \in E(H)$  with |E| = p + 1 and a set  $X \subset V(H)$ , |X| = p + 1, which is not an edge (if all (p + 1)-tuples are edges then, obviously, no p-tuple is an edge and we have nothing to prove). We can find such a pair E, X with  $|E \cap X| = p$ ; for let  $E = \{x_1, \ldots, x_{p+1}\}$ ,  $E \cap X = \{x_i, \ldots, x_{p+1}\}, X = \{x_i, \ldots, x_{i+p}\}.$  Then consider the sets  $\{x_{\nu}, \ldots, x_{\nu+p}\}$   $(\nu = 1, \ldots, i)$ . There is a last one among these, which is an edge, say  $\{x_{\nu}, \ldots, x_{\nu+p}\}$ . Then we can consider  $\{x_{\nu}, \ldots, x_{\nu+p}\}$  instead of E and  $\{x_{\nu+1}, \ldots, x_{\nu+1+p}\}$  instead of X. Now since  $E \cap X \subset E$ ,  $E \cap X \notin E(H)$ ; on the other hand,  $E \cap X \subset X$ ,

 $|E \cap X| = p$ , |X| = p + 1 and so by our remark above,  $E \cap X \in E(H)$ ,

a contradiction.

Thus E(H) consists of all p-tuples or of all (p+1)-tuples of V(H).

**22.** Suppose A, B are maximum antichains with m elements and define

$$A \vee B = \{x \in A \cup B : x \lessdot y \text{ for every } y \in A \cup B\},$$
  
$$A \wedge B = \{x \in A \cup B : x \gtrdot y \text{ for every } y \in A \cup B\}.$$

Then  $A \vee B$ ,  $A \wedge B$  are antichains by their definition. Moreover

$$(1) (A \vee B) \cap (A \wedge B) \supseteq A \cap B.$$

In fact, if  $x \in A \cap B$ , then there is no  $y \in A \cup B$  such that x < y for if  $y \in A$  (say), then  $x \in A$  also and A is an antichain. Thus,  $x \in A \vee B$  and similarly,  $x \in A \wedge B$ .

$$(A \lor B) \cup (A \land B) = A \cup B.$$

The inclusion  $\subset$  is obvious. Let  $x \in A \cup B$ , say  $x \in A$ . If  $x \notin A \vee B$ , then there is a  $y_1 \in A \cup B$  such that  $x < y_1$ . Obviously,  $y_1 \notin A$ ; so  $y_1 \in B$ . Similarly, if  $x \notin A \land B$ , then there exists a  $y_2 \in B$  with  $y_2 < x$ . Now  $y_2 < y_1$ , a contradiction. Thus  $x \in (A \vee B) \cup (A \wedge B)$ .

From (1) and (2) we deduce that

(3) 
$$|A \lor B| + |A \land B| = |(A \lor B) \cup (A \land B)| + |(A \lor B) \cap (A \land B)| \ge |A \cup B| + |A \cap B| = |A| + |B| = 2m.$$

Since A, B were maximum antichains we have

$$(4) |A \vee B| \leq m, |A \wedge B| \leq m.$$

Now from (3) and (4) we see that equality must hold throughout, in particular in (4). This proves that  $A \vee B$ ,  $A \wedge B$  are maximum antichains.

mum antichain which is, obviously, invariant under the automorphisms of

It is easy to verify that the set  $\mathcal{L}$  of maximum antichains forms a lattice with respect to the operations  $\wedge$  and  $\vee$ . Let E be the unity of  $\mathcal{L}$ , then E is a maximum

(S, <).

Now consider  $S = 2^{V(H)}$  and the partially ordered set  $(S, \subseteq)$ . E(H) is an antichain in  $(S, \subseteq)$ , so we want to determine the maximum size of an antichain. By the above, there is a maximum antichain  $\mathcal{E}$ , which is invariant under the automorphisms of  $(S, \subseteq)$ . Let  $E \in \mathcal{E}$ , |E| = k. Any permutation of V(H) induces an automorphism of S, and these permutations map E onto all k-element subsets. Thus, all k-element subsets belong to S. Obviously, S cannot have any other member, thus

$$|\mathcal{E}| = \binom{n}{k} \le \left(\left[\frac{n}{2}\right]\right)$$

[R.P. Dilworth, Combin. Analysis, Proc. Symp. Appl. Math. AMS (1960) 85]. 23. (a) The hypergraph H with  $V(H) = \{1, 2, 3, 4\}$ ,  $E(H) = \{\{1, 2, 3\}, \{3, 4\}, \{1, 4\}, \{2\}\}$  is cross-cutting but contains no other cross-cutting hypergraph, in particular, no cross-cutting clutter (Fig. 98).




Fig. 98

(b) Let H be a minimal cross-cutting hypergraph. Then for any  $E \in E(H)$  there exists a set  $S_E \subseteq V(H)$ , which is not comparable with any edge of  $H - \{E\}$ . Clearly,  $E \subseteq S_E$  or  $S_E \subseteq E$ . Denote by  $E^*$  the larger of E,  $S_E$ , and let

$$V(H^*) = V(H), E(H^*) = \{E^* : E \in E(H)\}.$$

We claim that  $H^*$  is a clutter. Assume indirectly that  $E^* \subseteq F^*$ . If  $E^* = S_E$ ,  $F^* = F$ , then  $S_E \subseteq F$ , which is a contradiction since  $S_E$  is comparable only with E by definition. We get a similar contradiction, if  $E^* = E$ ,  $F^* = S_F$ . If  $E^* = E$ ,  $F^* = F$ , then  $S_E \subseteq E$  and  $E \subseteq F$ , so  $S_E \subseteq F$ . If  $E^* = S_E$ ,  $F^* = S_F$ , then  $E \subseteq S_E \subseteq S_F$ . We get a contradiction in both cases again.

Thus  $H^*$  is a clutter and so by Sperner's theorem

$$|E(H)| = |E(H^*)| \le \left(\left[\frac{n}{2}\right]\right)$$

[P. Erdős]

24. Following the hint, let

$$p = \left\lceil \frac{n-k+1}{2} \right\rceil.$$

Then

$$p+k-1 = \left\lceil \frac{n+k-1}{2} \right\rceil$$

and hence, p + (p + k - 1) = n or n - 1. Therefore  $\binom{n}{p}, \ldots, \binom{n}{p + k - 1}$ 

are the k largest binomial coefficients (if n-k is even there is another such set:

we could replace  $\binom{n}{p}$  by  $\binom{n}{p-k+2}$  but this is unimportant. Form the hypergraph M on V(H) consisting of all  $p,\ldots,(p+k-1)$ -tuples as in the hint. Let  $K_1,\ldots,K_t$  be a decomposition of  $2^{V(H)}$  into disjoint symmetric chains as in 13.20. Obviously,

$$|E(M) \cap K_i| = \min(k, |K_i|);$$

on the other hand, E(H) contains no chain of length k, whence

$$|E(H) \cap K_i| \leq \min(k, |K_i|).$$

Since this holds for every  $i = 1, \ldots, t$ , we conclude that

$$|E(H)| \leq |E(M)| = \binom{n}{p} + \cdots + \binom{n}{p+k-1}$$

[P. Erdős, Bull. Amer. Math. Soc. 51 (1945) 898-902].

**25.** (a) Suppose indirectly that the edges of H cannot be covered by one point. Let  $E = \{x_1, \ldots, x_r\} \in E(H)$ . Then  $x_i$  does not cover all edges, so we find an edge  $F_i$  such that  $x_i \notin F_i$   $(i = 1, \ldots, r)$ . Now  $E, F_1, \ldots, F_r$  have no point in common; for such a point ought to be in E but  $x_i$  does not cover  $F_i$ .

(b) We construct j edges  $E_1, \ldots, E_i$  such that

$$|E_1 \cap \ldots \cap E_j| \leq r - (j-1)(\tau(H)-1)$$

for  $j=1,\ldots,k$ . For  $j=1,E_1$  can be chosen arbitrarily. Assume  $E_1,\ldots,E_j$  have been selected  $(j\leq k-1)$ . Since every other edge intersects  $E_1\cap\ldots\cap E_j$  this set must be of cardinality at least  $\tau(H)$ . Let  $X\subseteq E_1\cap\ldots\cap E_j$ ,  $|X|=\tau(H)-1$ . Since X cannot cover all edges, there must be an edge  $E_{j+1}$  such that  $E_{j+1}\cap X=\emptyset$ . Thus

$$|E_1 \cap \ldots \cap E_{j+1}| \leq |(E_1 \cap \ldots \cap E_j) - X| =$$
  
=  $|E_1 \cap \ldots \cap E_j| - (\tau(H) - 1) \leq r - j(\tau(H) - 1),$ 

i.e.  $E_{j+1}$  has the desired property. Now  $E_1 \cap \ldots \cap E_k \neq \emptyset$ , i.e.

$$1 \leq |E_1 \cap \ldots \cap E_k| \leq r - (k-1)(\tau(H) - 1),$$

whence the inequality of the problem follows immediately.

**26.** Take any edge F. If  $|E \cap F| \ge 2$  for every edge E we can take S = F. Suppose  $|F' \cap F| = 1$  for some F', say  $F' \cap F = \{x\}$ . Let F'' be an edge of H - x (such an edge exists since  $\tau(H) \ge 2$ ), and set  $S = F \cup F' \cup F''$ . Clearly

 $|S| \leq 3r - 3$ .

On the other hand, let  $E \in E(G)$ . If  $x \notin E$ , then E must intersect F and F' in distinct points so  $|E \cap S| \ge 2$ . If  $x \in E$ , then E must meet F'' in a point different from x so  $|E \cap S| \ge 2$  again.

27. Let  $(x_1, \ldots, x_p)$  be any ordering of W, where W is a minimal set such that any two edges of  $H_W$  meet (|W| = p). Then there is at most one point  $x_i$  such that both  $\{x_1, \ldots, x_i\}$  and  $\{x_i, \ldots, x_p\}$  contain edges E, E' of  $H_W$  (clearly  $E \cap E' = \{x_i\}$ ). For if  $j \neq i$  and  $\{x_1, \ldots, x_j\}$  contained  $F, \{x_j, \ldots, x_p\}$  contained F' ( $F, F' \in E(H_W)$ ), then assuming, e.g. i < j we would have  $E \cap F' = \emptyset$ , a contradiction. So if we carry out the counting for every ordering of W, we count at most p! points.

Now every element of W is counted many times. In fact, if  $x \in W$ , then there are edges E,  $F \in E(H_W)$  such that  $E \cap F = \{x\}$ , by the minimality of W. Set |E| = s, |F| = t. Let us arrange the points of E, then put x, then put the points of F; finally, "stick in" the remaining points arbitrarily. This can

be done in

$$(t-1)!(s-1)!\binom{p}{s+t-1}(p-s-t+1)! = p!\frac{(s-1)!(t-1)!}{(s+t-1)!}$$

ways. Now  $s, t \leq r$ , hence

$$\frac{(s+t-1)!}{(s-1)!(t-1)!} = s \begin{pmatrix} s+t-1 \\ t-1 \end{pmatrix} \le r \begin{pmatrix} 2r-1 \\ r-1 \end{pmatrix}.$$

Also we can exchange the role of E and F. Thus every point in W is counted at least  $\frac{2p!}{r\binom{2r-1}{r-1}}$  times. Hence the number of points in W is at most

$$p! \left| \frac{2p!}{r \binom{2r-1}{r-1}} = \frac{r}{2} \binom{2r-1}{r-1} = (2r-1) \binom{2r-3}{r-1}.$$

The second assertion follows by considering the following hypergraph. Let |U| = 2r - 2. With each partition  $\{X, Y\}$  of U such that |X| = |Y| = r - 1 we associate a new point  $v_{XY}$ ; let V be the set of these points  $v_{XY}$  and  $V(H) = U \cup V$ . Moreover, let E(H) consist of all r-tuples in U and also of all sets  $X \cup \{v_{XY}\}$ ,  $Y \cup \{v_{XY}\}$ . It is easy to check that any two edges of H intersect but that no subhypergraph has this property. Thus the minimal W is V(H), and

$$|V(H)| > |V| = \frac{1}{2} {2r-2 \choose r-1} = {2r-3 \choose r-1}.$$

[A sharpened version of a theorem of M. Calczynska-Karlowicz, Bull. Acad. Polon. Sci. ser. math. astr. phys. 12 (1964) 87–89.]

28. (a) Let x be any point of C. Then x is the endpoint of at most one  $A_i$ ; for if  $A_i$ ,  $A_i$  had a common endpoint then, since they are distinct, they ought to start in different directions and, as  $|V(C)| \geq 2k$ , they would be edgedisjoint.

Now observe that, since each  $A_i$  (j = 2, ..., t) has an edge in common with  $A_1$ , one of the endpoints of  $A_j$  is an inner point of  $A_1$ . Hence there are at most k-1 arcs  $A_j$  with  $2 \le j \le t$  and this implies that  $t \le k$ .

(b) Let  $\pi = (x_1, \ldots, x_n)$  be a cyclic permutation of V(H). We consider a circuit C of length n and associate  $x_1, \ldots, x_n$  with the edges of C (in the same cyclic order). Then if  $E \in E(H)$  consists of consecutive points in  $\pi$ , then it corresponds to an arc A of C of length r. Any two of these arcs have an edge in common. Thus, by the preceding "lemma", there are at most r such arcs, i.e., for any cyclic permutation  $\pi$ , there are at most r edges, which consist of consecutive points. Since there are (n-1)! cyclic permutations, this way we count not more than

$$r(n-1)!$$

edges. Let us see, how often a given edge E is counted here. To specify a cyclic permutation for which E consists of consecutive points, we have to order E and V(H) - E. Thus E is counted r!(n-r)! times. So the number of edges is

$$m \leq \frac{r(n-1)!}{r!(n-r)!} = \binom{n-1}{r-1}.$$

- [P. Erdős, Chao Ko, R. Rado, Quart. J. of Math. Oxford, II. 12 (1961) 313-320, this proof is due to G.O.H. Katona, J. Comb. Theory 13 (1972) 183-184.]
- **29.** (a) Let  $F_1, \ldots, F_t$  be all the sets which are unions of  $\nu$  disjoint edges of H. The fact that H is  $\nu$ -critical means that

$$F_1 \cap \ldots \cap F_t = \emptyset.$$

By 13.25a, we find rv+1 sets among  $F_1, \ldots, F_t$ , say  $F_1, \ldots, F_{rv+1}$ , whose intersection is void. Let  $S=F_1\cup\ldots\cup F_{rv+1}$ . We claim that  $|S\cap E|\geq 2$  for every edge E of H. In fact,  $S\cap E$  meets every one of  $F_1, \ldots, F_{rv+1}$ (otherwise E would be disjoint from  $F_i$  for some i and then it could be added to the v disjoint edges forming  $F_i$  as a  $(v+1)^{st}$  one). Since by their construction  $F_1, \ldots, F_{r\nu+1}$  cannot be covered by one point it follows that  $|S \cap E| \geq 2$ . Now  $|S| \leq (rv+1)rv$ . Thus the number of edges E is at most

$$\binom{|S|}{2}\binom{n}{r-2} \leq \binom{(rv+1)\,rv}{2}\binom{n}{r-2} = O(n^{r-2}).$$

(b) We use induction on  $\nu$ . If H is  $\nu$ -critical, (a) settles the question. If His not v-critical, let  $x \in V(H)$  be such that v(H - x) < v(H). Then

$$|E(H-x)| \le {n-2 \choose r-1} + \cdots + {n-\nu \choose r-1}$$

by the induction hypothesis and since x cannot be contained in more than  $\begin{pmatrix} n-1\\r-1 \end{pmatrix}$  edges, the assertion follows.

Equality holds for the hypergraph on n points consisting of those r-tuples, which meet a given v-element set [A. Hajnal, B. Rothschild, J. Comb. Theory 15 (1973) 359–362].

30. Let  $\{t_x\}$   $(x \in V(H))$  be an optimum fractional cover of H. Then

$$\sum_{x \in E} t_x \ge 1$$

holds for every edge E. Summing this for every edge of  $H_i = H - x_1 - \ldots - x_i$ , we obtain

$$\sum_{E \in E(H_i)} \sum_{x \in E} t_x \ge |E(H_i)|.$$

The left-hand side can be written as

$$\sum_{\mathbf{x}} t_{\mathbf{x}} \sum_{\substack{E \ni \mathbf{x} \\ E \in E(H_i)}} 1 \leq \left(\sum_{\mathbf{x}} t_{\mathbf{x}}\right) d_{i+1} = \tau^* d_{i+1}.$$

So

$$|E(H_i)| \leq \tau^* d_{i+1}.$$

Now suppose the procedure stops after t steps, i.e.  $x_1, \ldots, x_t$  cover all edges,  $E(H_{t+1}) = \emptyset$ . Then clearly

$$\tau(H) \leq t$$
.

We know that

$$|E(H_i)| - |E(H_{i+1})| = d_i$$

SO

$$t = \sum_{i=1}^{t} \frac{|E(H_i) - |E(H_{i+1})|}{d_i} = \sum_{i=2}^{t} |E(H_i)| \left(\frac{1}{d_i} - \frac{1}{d_{i-1}}\right) + \frac{|E(H_1)|}{d_1}.$$

Since  $d_i \leq d_{i+1}$  we can estimate  $|E(H_i)|$  by (1) and obtain

$$t \leq \tau^* \left\{ \sum_{i=2}^t d_i \left( \frac{1}{d_i} - \frac{1}{d_{i-1}} \right) + \frac{1}{d_1} \right\} = \tau^* \sum_{i=1}^t \frac{d_i - d_{i+1}}{d_i} \leq \tau^* \sum_{k=1}^d \frac{1}{k} < \tau^* (1 + \log d)$$

[L. Lovász, Discrete Math. 13 (1975) 383-390].

**31.** (a) Clearly  $u \ge w + 1$  since

$$\binom{u}{r} = \binom{v}{r} + \binom{w}{r-1} \ge \binom{w}{r} + \binom{w}{r-1} = \binom{w+1}{r}.$$

Let w = t + r - 1,  $u = t + u_1$ ,  $v = t + v_1$ . Then by 1.42i and b,

Hence

$$(1) 0 = \begin{pmatrix} v \\ r \end{pmatrix} + \begin{pmatrix} w \\ r-1 \end{pmatrix} - \begin{pmatrix} u \\ r \end{pmatrix} = \sum_{j=0}^{r-1} \begin{pmatrix} t+j-1 \\ j \end{pmatrix} \varphi_{r-j},$$

where

$$arphi_i = inom{v_1-r+i}{i} + 1 - inom{u_1-r+i}{i}.$$

Similarly

in case t = 0, we define  $\frac{0}{t} = 0$ .

We claim that if  $\varphi_i$  is positive then so are  $\varphi_{i-1}, \ldots, \varphi_1$ . In fact,

$$egin{aligned} arphi_{i-1} &= rac{i}{v_1-r+i}inom{v_1-r+i}{i}+1-rac{i}{u_1-r+i}inom{u_1-r+i}{i} \geq \ &\geq rac{i}{u_1-r+i}inom{v_1-r+i}{i}+rac{i}{u_1-r+i}-rac{i}{u_1-r+i}inom{u_1-r+i}{i}= \ &= rac{i}{u_1-r+i}\,arphi_i>0. \end{aligned}$$

(since  $u_1 \ge v_1$  and  $u_1 = u - t \ge w + 1 - t = r$ ). Thus we have a k (0  $\le k \le r$ ) such that

$$\varphi_1,\ldots,\varphi_k>0,\,\varphi_{k+1},\ldots,\varphi_r\leq 0.$$

Then replacing  $\frac{j}{t}$  by  $\frac{r-k}{t}$  on the right-hand side of (2) we decrease it. Therefore by (1),

$$egin{pmatrix} w \ r-1 \end{pmatrix} + egin{pmatrix} w \ r-2 \end{pmatrix} - egin{pmatrix} u \ r-1 \end{pmatrix} \geq rac{r-k}{t} \sum_{j=0}^{r-1} egin{pmatrix} t+j-1 \ j \end{pmatrix} arphi_{r-j} = 0.$$

(b) We use induction on |E(H)|. It suffices to treat the case k=r-1. Let x be a point with minimum degree d. We may assume that there are no isolated points, i.e.  $d \ge 1$ ; also we may suppose there are at least two edges, so that  $d < \binom{u}{r}$ .

Let  $H_1 = H - x$ . Then  $|E(H_1)| = \binom{u}{r} - d$ . Let us write this number in the form  $\binom{v}{r}(v \ge r)$ ; then, by the induction hypothesis, there are at least  $\binom{v}{r-1}$ 

(r-1)-tuples covered by the edges of  $H_1$ . Note that none of these contains x. Let us define  $H_2$  by  $V(H_2) = V(H) - \{x\}$ ,  $E(H_2) = \{E - \{x\} : x \in E \in E(H)\}$ . Then  $|E(H_2)| = d$ , and  $H_2$  is (r-1)-uniform. Let us write d in the form  $\binom{w}{r-1}$ ,  $w \ge r-1$ , then by the induction hypothesis, there are at least  $\binom{w}{r-2}$  (r-2)-tuples covered by the edges of  $H_2$ . Adding x to these (r-2)-

tuples, we obtain at least  $\binom{w}{r-2}(r-1)$ -tuples contained in edges of H. Since each of these (r-1)-tuples contains x, they are different from the ones

found previously. Thus the number of (r-1)-tuples contained in edges of H is at least

$$\begin{pmatrix} v \\ r-1 \end{pmatrix} + \begin{pmatrix} w \\ r-2 \end{pmatrix}$$
.

Using (a) we will be done if we can show that  $v \geq w$ . To this end observe that the average degree is

$$r\frac{|E(H)|}{|V(H)|} = r\frac{\binom{u}{r}}{|V(H)|}$$

and here, obviously,  $|V(H)| \geq u$ . Thus

$$\binom{w}{r-1} = d \le r \frac{\binom{u}{r}}{u} = \binom{u-1}{r-1},$$

whence

$$w \leq u - 1$$
.

Thus

$$\binom{v}{r} = \binom{u}{r} - \binom{w}{r-1} \ge \binom{u}{r} - \binom{u-1}{r-1} = \binom{u-1}{r}, \quad v \ge u-1 \ge w,$$

as stated. [J.B. Kruskal, in: *Math. Opt. Techniques*, Univ. of Calif. Press 1963, 251–278; G. Katona, in: *Theory of Graphs* (Akadémiai Kiadó, 1966, 187–207. See these references for the exact optimum when *u* is not an integer.]

(c) Let H have m distinct edges, which are r-tuples and mutually intersecting. Let H' be defined by

$$V(H') = V(H)$$
  
 $E(H') = \{V(H) - E : E \in E(H)\}.$ 

Then by assumption no edge of H can occur as a subset of any edge of H'. Let us write

$$m = \begin{pmatrix} u \\ n-r \end{pmatrix}$$
  $(u \ge n-r, \text{ real}),$ 

then by (b), there are at least  $\binom{u}{r}$  r-tuples among the subsets of edges of H'.

Hence

$$\binom{u}{n-r} + \binom{u}{r} \le \binom{n}{r}.$$

Since

$$\binom{n-1}{n-r} + \binom{n-1}{r} = \binom{n}{r},$$

it follows that  $u \leq n-1$  and thus

$$m = \binom{u}{n-r} \le \binom{n-1}{n-r} = \binom{n-1}{r-1}$$

[D.E. Daykin, J. Comb. Th. A 17 (1974) 254-255].

**32.** (a) Observe that if  $(x_1, \ldots, x_n)$  is any permutation of V(H), then there is at most one index i for which every element of  $A_i$  has smaller index than every element of  $B_i$ . So we count at most n! pairs  $(A_i, B_i)$ .

Moreover, if i is any given index, then the number of permutations of V(H)

for which every element of  $A_i$  anticipates every element of  $B_i$  is

$$\binom{n}{p+q} p! q! (n-p-q)! = \frac{n!}{\binom{p+q}{p}}.$$

Hence, a given pair  $(A_i, B_i)$  is counted this number of times. Hence

$$m \le n! \left| \frac{n!}{\binom{p+q}{p}} = \binom{p+q}{p} \right|.$$

(b) Let  $E_1, \ldots, E_m$  be the edges of H. Since H is  $\tau$ -critical,  $H - \{E_i\}$  has a  $(\tau(H) - 1)$ -element point-cover  $T_i$ . Then, obviously,  $E_i \cap T_i = \emptyset$  while  $T_i \cap E_i \neq \emptyset$ , if  $i \neq j$ . Thus (a) implies that

$$m \leq \binom{r + (\tau(H) - 1)}{r}$$

[B. Bollobás, G.O.H. Katona, F. Jaeger, C. Payan; see B].

**33.** Let  $V(H) = \{v_1, \ldots, v_n\}$ . We color  $v_1, v_2, \ldots, v_n$  by one of the colors red and blue, so that we do not color all points of any edge with the same color. Suppose  $v_1, \ldots, v_i$  are colored (1 < i < n). If we cannot color  $v_{i+1}$  red, then there is an edge  $E \subset \{v_1, \ldots, v_{i+1}\}$ ,  $v_{i+1} \in E$  such that all the other points of E are red. Similarly, if  $v_{i+1}$  cannot be given color blue, there must be an edge  $F \subset \{v_1, \ldots, v_{i+1}\}$ ,  $v_{i+1} \in F$  all of whose points except  $v_{i+1}$  are blue. Now  $E \cap F = \{v_{i+1}\}$ , a contradiction. Thus we can color  $v_{i+1}$  either red or blue.

**34.** Suppose first that H has one point  $x_1$  of degree 1 and all other points have degree 2. Let  $x_2$  be a point adjacent to  $x_1$ ,  $x_3$  a point adjacent to  $x_2$  or  $x_3$ , ..., let  $x_{i+1}$  be a point, different from  $x_1, \ldots, x_i$  and adjacent to one of them. Such

an  $x_{i+1}$  exists, because H is connected. Now going "back" we can color  $x_n$ ,  $x_{n-1}, \ldots, x_1$  successively with red and blue in the same way as in the previous solution.

Now suppose that every point of H has degree 2. We may assume H is not a graph. Then it contains an edge E with  $|E| \geq 3$ . Let  $x \in E$ ,  $E' = E - \{x\}$  and  $H' = H - \{E\} + \{E'\}$ . Now if H' is connected, then, since x has degree 1 in H', by the first part of the solution H' is 2-chromatic. A 2-coloration of H' is a 2-coloration of H. If H' is not connected, then  $H = H_1 \cup H_2$ ,  $V(H_1) \cap V(H_2) = \{x\}$ . 2-coloring  $H_1$  and  $H_2$  (which have points with degree 1) we can put these 2-colorations together after exchanging colors in  $H_1$ , if necessary.

- 35. Suppose we have no 1-element edge (which is a trivial case). Also remove isolated points. We verify the assertions in the hint.
- (a) Let  $x \in V(H)$ ,  $E \in E(H)$  be such that  $x \in E$ . Color  $E \{x\}$  red,  $V(H) E \cup \{x\}$  blue. Clearly there will be no red monochromatic edge. Since H is not 2-colorable, there must be a blue monochromatic edge F. F meets E in a single point and this point must be blue; hence  $x \in F$ . Thus x has degree at least 2.
- (b) Let  $x \in E_1, \ldots, E_d \in E(H)$ ,  $d \ge 4$ , and suppose indirectly that  $|E_d| \ge 3$ . Let  $y_i \in E_i \{x\}$   $(i = 2, \ldots, d 1)$ . Since  $d \ge 4$ , there is at most one edge containing  $y_2, \ldots, y_{d-1}$ . So we can select a point  $y_d \in E_d \{x\}$  such that no edge contains  $y_2, \ldots, y_d$ . Now let us color  $E_1 \{x\}, y_2, \ldots, y_d$  red, all other points blue. It is easy to check that no monochromatic edge can occur.

This settles the case when there is a point x of degree at least 4; more generally, if there is a point x which is adjacent to 2-element edges only, then consider an edge E not containing x (such an edge clearly exists). E must contain the (single) point of  $F - \{x\}$  for every edge  $F \ni x$ . This shows that E is unique and so  $V(H) = E \cup \{x\}$ ,  $E(H) = \{E\} \cup \{\{x,y\} : y \in E\}$ .

So suppose each point belongs to at least one edge of cardinality at least 3. Then there is no 2-element edge. For let  $\{x, y\} \in E(H)$  and let E, F be  $\geq$  3-element edges containing x and y, respectively. Let u be the (unique) common point of E and F, and let  $v \in E - \{u, x\}, w \in F - \{u, y\}$ . By (a), there are other edges A, B containing v and w. Since A, B must meet  $\{x, y\}$ , we have

 $y \in A, x \in B$ .


Now color every point in  $E \cup F - \{u\} - \{y\}$  red, the remaining points blue. Then no red monochromatic edge occurs; for such an edge X must meet  $\{x, y\}$  in a red point, i.e. in x; it must meet A in a red point, i.e. in v; but then X = E, which is not monochromatic. Also no blue monochromatic edge arises; for such an edge X must meet  $\{x, y\}$  in a blue point, i.e. in y; it must meet E in a blue point, i.e. in u; but then X = F, which is not monochromatic. Thus E is 2-colorable, a contradiction.

Now we are prepared to prove (c). Let  $x, y \in V(H)$ , and select edges E, F such that  $x \in E$ ,  $y \in F$ . We may assume  $x \notin F$  and  $y \notin E$ , otherwise we have nothing to prove. Let us color  $E \cup F - \{x, y\}$  red, everything else blue. Then no monochromatic red edge can arise; for such an edge could only consist of one point of  $E - \{x\}$  and one point of  $F - \{y\}$ , but 2-element edges are excluded. But then a blue monochromatic edge A arises, which must meet

E in x and F in y, which proves (c).

It follows that every edge is a 3-tuple. For if  $E \in E(H)$ ,  $|E| \ge 4$ , then let  $x \in V(H) - E$ . For each point y in E, there exists an edge containing x, y and these edges are, obviously, distinct. But then x has degree at least 4, a contradiction.

Now let E, F, G be three edges such that  $E \cap F \subset G$ . Let  $E \cap F = \{x\}$ ,  $F \cap G = \{y\}$ ,  $E \cap G = \{z\}$ . Then: there are unique further points u, v, w



y t u

Fig. 99

of E, F, G. There is a unique edge  $\{x, w, t\}$  containing x, w. There is a unique edge containing y, u, which must also contain t since this is the only point in which it can meet  $\{x, w, t\}$ . Similarly,  $\{z, v, t\}$  and  $\{u, v, w\}$  are edges and there are no other edges. This hypergraph (usually called the *Fano configuration*, or 7-point-plane) is 3-chromatic. Thus we have found the hypergraphs shown in Fig. 99 as the only possibilities.

**36.** Suppose indirectly that H has a 2-coloration with red and blue. Let  $n_1$  and  $n_2$  denote the numbers of red and blue points, respectively. Then the number of red-red-blue edges is

 $a \begin{pmatrix} n_1 \\ 2 \end{pmatrix}$ ,

where a is the number of edges containing a given pair. Similarly, there are

 $a \binom{n_2}{2}$ 

blue-blue-red edges. Thus

$$|E(H)| = a \left[ \binom{n_1}{2} + \binom{n_2}{2} \right].$$

On the other hand, each edge contains exactly two pairs (x, y), where x is blue, y is red. Thus,

(2) 
$$2|E(H)| = an_1n_2.$$

From (1) and (2) we get

$$n_1 n_2 = n_1^2 + n_2^2 - n_1 - n_2 = n^2 - n - 2n_1 n_2$$

or, equivalently,

$$3n_1n_2 = n^2 - n.$$

But we have

$$n_1 n_2 \le \left(\frac{n_1 + n_2}{2}\right)^2 = \frac{n^2}{4},$$

so (3) implies that

$$\frac{n^2}{4} \ge n^2 - n \quad \text{or} \quad n \le 4,$$

a contradiction [L. Lovász, Proc. 4th SE Conf. on Comb. Graph Theory and Computing, Utilitas Math. 1973, 1-12].

**37.** Consider the polyhedron P in n-dimensional space described by

$$\left[\frac{|E_j|}{2}\right] \leq \sum_{i=1}^n a_{ij} x_i \leq \left[\frac{|E_j|}{2}\right]^* \qquad (j = 1, \dots, m),$$

$$0 \leq x_i \leq 1 \qquad (i = 1, \dots, n).$$

P is non-empty since the vector  $\left(\frac{1}{2}, \dots, \frac{1}{2}\right)^T$  belongs to it. Also the matrix  $\binom{A^T}{I}$  is totally unimodular and the constants are integers, so by the Hoffman-Kruskal theorem, P contains a lattice point. Let  $(\xi_1, \ldots, \xi_n)^T$  be this lattice point. Color  $v_i$  red, if  $\xi_i = 0$  and blue, otherwise. This coloration is a legitimate one, if H contains no 1-element edges. More exactly, there are at least  $\left| \frac{|E_j|}{2} \right|$ and at most  $\left|\frac{|E_j|}{2}\right|^*$  blue points in every edge  $E_j$ . This proves the "only if" part of the assertion.

Now suppose that each subhypergraph has a bicoloration as formulated. Then we show that every subdeterminant of the incidence matrix A is 0 or  $\pm 1$ . Since a square submatrix of A is the incidence matrix of some partial subhypergraph which, obviously, has the same property it suffices to show that

if n = m, then det A = 0 or  $\pm 1$ .

We claim that there exists a  $W \subseteq V(H)$ ,  $W \neq \emptyset$  such that  $|W \cap E_1|, \ldots$  $|W \cap E_{n-1}|$  are even; in fact, consider the vectors

$$\mathbf{a}_j = egin{pmatrix} a_{1j} \ \vdots \ a_{nj} \end{pmatrix}, \quad ext{where } a_{ij} = egin{bmatrix} 1 & ext{if } v_i \in E_j, \ 0 & ext{otherwise}, \end{bmatrix}$$

 $(i=1,\ldots,n-1)$  over GF(2). They do not generate the whole space, so there is a 01-vector

$$\mathbf{b} = \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix} \neq 0$$

å

orthogonal to all of them by 5.32f. Now set

$$W = \{v_i : b_i = 1\}.$$

Then  $|W \cap E_j| \equiv \mathbf{b} \cdot \mathbf{a}_j \equiv 0 \pmod{2}$   $(j = 1, \dots, n-1)$ .  $|W \cap E_n|$  may be odd or even.

We may assume  $W = \{v_1, \ldots, v_t\}$ . By the assumption  $W = A_0 \cup A_1, A_0 \cap A_1 = \emptyset$  so that

$$\left\lceil \frac{|E_j \cap W|}{2} \right\rceil \leq |E_j \cap A_0| \leq \left\lceil \frac{|E_j \cap W|}{2} \right\rceil^* \ (j = 1, \ldots, n).$$

Note that, by the choice of W,

$$|E_j \cap A_0| = \frac{|E_j \cap W|}{2}$$

for  $j=1,\ldots,n-1$ . Let, say,  $A_0=\{v_1,\ldots,v_s\}$ . Then add the  $2^{\rm nd},\ldots,s^{\rm th}$  columns to the first one but subtract the  $(s+1)^{\rm st},\ldots,t^{\rm th}$  columns from it. This way we get 0's in all but the last entry of the first column, and get 0 or  $\pm 1$  in the last entry. If this last entry is 0, then det  $A_1=0$ . Otherwise, we can expand by the first column and proceed by induction [A. Ghouila-Houir; see B].

**38.** (a) First we prove the statement of the hint. It suffices to consider the case k=2. Let  $E\in E(H),\ E_1,\ E_2\subset E,\ E_1\cap E_2=\emptyset$ . Consider

$$H' = H - \{E\} + \{E_1, E_2\}.$$

We claim that H' has no odd circuits. For let C be any circuit of H'. If C contains none or one of  $E_1$ ,  $E_2$ , then it corresponds to a circuit of H in the natural way and hence it is even. Suppose it contains both of  $E_1$ ,  $E_2$ . Then it is of the form

$$(E_1, x_1, F_1, \ldots, x_k, E_2, y_1, G_1, \ldots, y_l),$$

where  $k \geq 2$ ,  $l \geq 2$  as  $E_1 \cap E_2 = \emptyset$ . Now

$$(E, x_1, F_1, \ldots, x_k)$$
 and  $(E, y_1, G_1, \ldots, y_l)$ 

are circuits of H, hence k and l are even. Then k+l is even, i.e. C is even.

Now replace each edge E by  $\left\lceil \frac{|E|}{2} \right\rceil$  disjoint pairs. The resulting graph is bipartite by the above result and thus it possesses a 2-coloration with red and blue. Now each edge E contains at least  $\left\lceil \frac{|E|}{2} \right\rceil$  red and at least  $\left\lceil \frac{|E|}{2} \right\rceil$  blue points, which proves the assertion.

- (b) This follows trivially by (a) and 13.37 [C. Berge; see B].
- **39.** Suppose indirectly that H is not balanced. Then it contains an odd circuit  $(x_1, E_1, \ldots, x_{2k+1}, E_{k+1})$  which is not balanced. This means that there is no other incidence between  $x_1, \ldots, x_{2k+1}$  and  $E_1, \ldots, E_{2k+1}$ , than is indicated by

the definition of circuits. Hence A contains a  $(2k+1)\times(2k+1)$  submatrix of the form

which has determinant 2, a contradiction [see B].

40. Suppose indirectly that there are balanced hypergraphs, which are not 2-chromatic and consider a minimum counterexample H. Define a graph G by V(G) = V(H),  $E(G) = \{E \in E(H) : |E| = 2\}$ .

We claim that G is a connected graph. Suppose indirectly that  $V(G) = V_1 \cup V_2$ ,  $V_1 \cap V_2 = \emptyset$ ,  $V_1$ ,  $V_2 \neq \emptyset$  such that each  $E \in E(G)$  is contained in  $V_1$  or  $V_2$ . Consider  $H_{V_1}$  and  $H_{V_2}$  and remove the one-element edges of them. This way

we obtain two balanced hypergraphs  $H_1$ ,  $H_2$ . By the minimality of H,  $H_1$  and  $H_2$  are 2-chromatic. Let us 2-color each of them by red and blue, we claim this is a good 2-coloration of H. Let  $E \in E(H)$ . If |E|=2, then E is spanned by one of  $V_1$  or  $V_2$  and therefore it gets both colors. If  $|E|\geq 3$ , then  $|E\cap V_i|\geq 2$  for one of  $i=1,\ 2$  and so,  $E\cap V_i$ gets both colors in the 2-coloration of  $H_i$ . Thus we have obtained a 2-coloration of H, a contradiction.

Now G has no odd circuits, because H is balanced and since it is connected, it has an (essentially unique) 2-coloration with red and blue. We claim that this is a legitimate 2-coloration of H. Assume indirectly that there were a monochromatic edge E, |E| > 1. Since G is connected, it contains a path  $P = (x_0, F_1, x_1, \ldots, F_k, x_k)$  with  $x_0 \in E$ ,  $x_k \in E$ . We may assume  $x_i \notin E$  for  $i = 1, \ldots, k-1$ . The points  $x_0, \ldots, x_k$  are alternately red and blue, hence k is even. Thus

$$(x_0, F_1, x_1, \ldots, x_k, E)$$

is an unbalanced odd circuit, a contradiction. Thus the 2-coloration we considered is a good 2-coloration of H, a contradiction again [C. Berge; see B].

**41.** Let  $E(H) = \{E_1, \ldots, E_m\}$   $(m \le 2^{r-1})$  and color the points of H with red and blue at random, independently of each other and with probability 1/2. Let  $A_i$  denote the event that  $E_i$  is monochromatic. Then

$$P(A_i) = \frac{1}{2^{r-1}}$$

as there are  $2^r$  ways to color  $E_i$  and 2 of these come into consideration. So the probability that a random coloration contains a monochromatic edge is

$$P(A_1 + \ldots + A_m) < \sum_{i=1}^m P(A_i) = \frac{m}{2^{r-1}} \le 1,$$

(the first strict inequality follows from the fact that  $A_1, \ldots, A_m$  are not mutually exclusive: all occur, when all points are red) [P. Erdős; see ES].

**42.** Let H be defined as in the hint. Then

$$|E_P| = inom{|X|}{r} + inom{|Y|}{r} \ge 2inom{r^2}{r}$$

for any edge  $E_P$  ( $P = \{X, Y\}$ ). Therefore,

$$t_{\mathrm{x}} = \frac{1}{2 \binom{r^2}{r}}$$

defines a fractional cover. Thus

$$au^*(H) \leq rac{inom{2r^2}{r}}{2inom{r^2}{r}}.$$

(It would not be difficult to show that one has equality here.)

Now let T be any set of points of H covering every edge  $E_P$ . Then T is a collection of r-tuples, so  $V(H_0) = S$ ,  $E(H_0) = T$  defines an r-uniform hypergraph. The fact that T covers  $E_P$  means that P does not give a legitimate 2-coloration of  $H_0$ ; so  $H_0$  is not 2-chromatic. Thus if T is a minimum cover we have by 13.30,

$$|E(H_0)| = \tau(H) \le (1 + \log d(H))\tau^*(H).$$

Here, however,

$$au^*(H) = rac{inom{2r^2}{r}}{2inom{r^2}{r}},$$

$$d(H)=2^{r^2-r},$$

and so

$$|E(H_0)| \le \left(1 + (r^2 - r) \log 2\right) \frac{{2r^2 \choose r}}{2{r^2 \choose r}} < r^2 \frac{{2r^2 \choose r}}{{r^2 \choose r}} =$$
 $= r^2 \cdot \frac{2r^2}{r^2} \cdot \frac{2r^2 - 1}{r^2 - 1} \cdot \dots \cdot \frac{2r^2 - r + 1}{r^2 - r + 1} < c \cdot r^2 \cdot 2^r$ 

[P. Erdős; see ES].

43. Let us color the points of H at random again, as in the solution of 13.41. Observe that if  $E_1, \ldots, E_p$  are disjoint from  $E_q$ , then  $A_1, \ldots, A_p$  and any polynomial in them are independent of  $A_q$ . So if we form L(H) and associate the event  $A_j$  with  $E_i$ , then this graph and the associated events satisfy the first

condition of 2.18. Moreover, by the assumption, each point of L(H) has degree at most  $2^{r-3}$ , hence the second condition is satisfied as well. Therefore

13.44

$$P(\bar{A}_1 \ldots \bar{A}_m) > 0,$$

i.e. there is a coloration in which no edge is monochromatic [P. Erdős–L. Lovász, in: *Infinite and Finite Sets*, Coll. Math. Soc. J. Bolyai 10, Bolyai–North-Holland (1974) 609–627].

**44.** (a) Suppose H is not 2-chromatic. Then by 13.43, there is an edge E which is intersected by more than  $2^{r-3}$  other edges. Then one of the points of E belongs to more than  $2^{r-3}/r$  of these. Thus, the maximum degree d of H is at least  $2^{r-3}/r$ .

Now let x be a point with maximum degree d. Then the edges containing x have no other point in common by the assumption and so, they cover at

$$1 + \frac{(r-1)2^{r-3}}{r} > 2^{r-4}$$

least points, a contradiction.

(b) For  $E \in E(H)$ , let  $\varphi(E)$  denote the point of E which has largest degree in H (or one of these) and set  $E' = E - \{\varphi(E)\}$ ,

$$H' = (V(H); \{E' : E \in E(H)\}).$$

Then H' must be at least 3-chromatic since any 2-coloration of H' would yield a 2-coloration of H. Thus by the solution of (a), there is a point x with degree  $t > 2^{r-4}/(r-1)$  in H'. Let  $E'_1, \ldots, E'_t$  be the edges of H' containing x. Then, by definition, the degree of  $\varphi(E_i)$  in H is at least the degree of x in H, which is at least the degree of x in H', which is equal to t. Moreover, the points  $\varphi(E_1)$ , ...,  $\varphi(E_t)$  are distinct, as  $E_1, \ldots, E_t$  have no point in common other than x. Thus we have t points with degree at least t, which proves the assertion.

(c) Suppose indirectly that H has chromatic number at least 3. Then, by the preceding problem, there are  $2^{r-4}/r$  points with degree  $> 2^{r-4}/r$ . If we count the edges adjacent to these points we get  $4^{r-4}/r^2$ , and each edge is counted not more than r times. Hence

$$|E(H)| > 4^{r-4}/r^3$$
.

[ibid.]

- **45.** (a) By 13.5b, we can select a pair f(E) from each edge E of H so that the hypergraph (in fact, graph)  $G = (V(H), \{f(E) : E \in E(H)\})$  is a forest. Hence G is 2-colorable and a 2-coloration of G gives a 2-coloration of H [M. Las Vergnas, L. Lovász; see B].
- (b) Let  $H_2$  be the triangle and suppose we already have an r-uniform hypergraph  $H_r$  with the desired property. Define  $H_{r+1}$  as follows; for every edge  $E \in E(H_r)$ , take an (r+1)-element set E'; let the sets E' be disjoint from each other and from  $V(H_r)$ . Define

$$V(H_{r+1}) = \bigcup_{E \in E(H_r)} E' \cup V(H_r),$$

$$E(H_{r+1}) = \{E' : E \in E(H_r)\} \cup \{E \cup \{x\} : E \in E(H_r), x \in E'\}.$$

Then, obviously,  $H_{r+1}$  is (r+1)-uniform. Moreover,  $H_{r+1}$  is not 2-colorable. For let  $\alpha$  be a 2-coloration of  $H_{r+1}$ . Then there is an edge  $E \in E(H_r)$ , which is monochromatic since  $H_r$  is not 2-colorable. Let, e.g. all points of E be red. Then for any  $x \in E'$ ,  $E \cup \{x\}$  is an edge of  $H_{r+1}$ , so x must be blue. But then E' is a monochromatic edge of  $H_{r+1}$ , a contradiction.

Let H' be any partial hypergraph of  $H_{r+1}$ , we claim that  $|V(H')| \ge |E(H')|$ . Suppose not, and let H' be a minimal counterexample. Then every point in H' has degree at least 2, since otherwise it could be removed. Therefore, if  $x \in E'$  ( $E \in E(H_r)$ ) is a point of H', then  $E' \in E(H')$ , and also  $E \cup \{y\} \in E(H')$ 

for each  $y \in E'$ .

Let A denote the set of edges of H' of form E',  $E \in E(H_r)$ . For every  $E \in E(H') - A$ , let  $E^* = E \cap V(H_r)$ . Clearly  $E^* \in E(H_r)$ . Thus the hypergraph H'' defined by

$$V(H'') = V(H') \cap V(H_r),$$
  
 $E(H'') = \{E^* : E \in E(H') - A\}$ 

(each set  $E^*$  is taken only once) is a partial hypergraph of  $H_r$ , and thus

$$|V(H'')| \ge |E(H'')|.$$

Also we have

$$|E(H')| = (r+1) |E(H'')|,$$
  $|V(H')| = |V(H'')| + r|E(H'')|$ 

and so

$$|V(H')| \ge (r+1) |E(H'')| = |E(H')|.$$

This proves that  $H_{r+1}$  has the desired properties. We remark that every partial hypergraph H' of  $H_r$  even satisfies

$$|V(H')| \ge |E(H')| + 1$$

except for  $H' = H_r$  [D.R. Woodall, in: *Combinatorics*, Proc. Conf. Southendon-Sea, 1972, 322–340].

**46.** (a) Let  $H_r$  be an r-uniform non-2-colorable hypergraph, without multiple edges, in which any two edges meet. We construct an (r+1)-uniform hypergraph  $H_{r+1}$  with similar properties. Let |U|=r+1,  $U\cap V(H_r)=\emptyset$ . Define  $H_{r+1}$  by

$$\begin{split} V(H_{r+1}) &= V(H_1) \, \cup \, U, \\ E(H_{r+1}) &= \{U\} \, \cup \, \{E \, \cup \, \{x\} : E \in E(H_r), \ x \in U\}. \end{split}$$

It is easy to verify that  $H_{r+1}$ , has the desired properties. Also,

$$|E(H_{r+1})| = (r+1) |E(H_r)| + 1,$$

whence

$$|E(H_r)| = [(e-1)r!] > r!.$$

(b) Let H, H' be two hypergraphs. Define H[H'] as follows. Let

$$V(H[H']) = V(H) \times V(H'),$$

$$E(H[H']) = \{\{x_1\} \times F_1 \cup \{x_2\} \times F_2 \cup \ldots \cup \{x_r\} \times F_r : \{x_1, \ldots, x_r\} \in E(H), F_1, \ldots, F_r \in E(H')\}.$$

Then, if any two edges of H and of H' meet, the same holds for H[H']. In fact, if  $\{x_1\} \times F_1 \cup \ldots \cup \{x_r\} \times F_r$  and  $\{x_1'\} \times F_1' \cup \ldots \cup \{x_s'\} \times F_s'$  are two edges of H[H'], then  $\{x_1, \ldots, x_r\}$  and  $\{x_1', \ldots, x_s'\}$  meet; let  $x_i = x_j'$ , then also  $F_i$  and  $F_j'$  meet; let  $y \in F_i \cap F_j'$ , then  $(x_i, y)$  is a common point of

the two edges of H[H'].

If H, H' are not 2-colorable, than neither is H[H']. For let us 2-color the points of H[H']. Then, for each  $x \in V(H)$ , one of the sets  $\{x\} \times F$  ( $F \in E(H')$ ) is monochromatic; choose such an edge  $F_x$  of H'. Now color x with the color of  $\{x\} \times F_x$ , then there will be an edge  $\{x_1, \ldots, x_r\} \in E(H_r)$  which is monochromatic. Then the edge

$$\{\{x_1\} \times F_{x_1}, \ldots, \{x_r\}\} \times F_{x_r}\}$$

is monochromatic.

Let K denote the triangle and set

$$K_m = \underbrace{K[K[K \dots [K] \dots]]}_{m \text{ times}}.$$

Then, by the above, any two edges of  $K_m$  meet, but it is not 2-chromatic. It is  $r=2^m$ -uniform and has  $3^{2^m-1}$  edges, as is easily seen by induction. Hence it has all the desired properties.

- (c) The hypergraph constructed in the solution of 13.27 trivially has all the desired properties [P. Erdős-L. Lovász, ibid.].
- 47. Consider such an r-uniform H without multiple edges, which is not 2-chromatic. Let  $x_1$  be a point with maximum degree, then its degree is greater than
- $\frac{m}{r}$ . For let  $E \in E(H)$ . Every other edge meets E, it meets itself in r points, hence

$$(m-1)+r \leq \sum_{F \in E(H)} |E \cap F| = \sum_{x \in E} d_H(x).$$

Thus

$$\max_{x \in R} d_H(x) \ge \frac{m}{r} - \frac{1}{r} + 1 > \frac{m}{r}$$

Now suppose that  $x_1, \ldots, x_i$  are chosen so that  $\{x_1, \ldots, x_i\}$  is contained in more than  $\frac{m}{r^i}$  edges (i < r). Try to color  $x_1, \ldots, x_i$  red, the rest blue. This cannot work, i.e. there must be an edge  $E_i$  such that  $x_1, \ldots, x_i \notin E_i$ . Any edge containing  $\{x_1, \ldots, x_i\}$  meets  $E_i$  and thus, there is a point  $x_{i+1}$  which is contained in more than  $\frac{m}{r^i}/r$  of them. Thus  $\{x_1, \ldots, x_{i+1}\}$  is contained in more than  $\frac{m}{r^{i+1}}$  edges.

We conclude that  $\{x_1, \ldots, x_r\}$  is contained in more than  $\frac{m}{r^r}$  edges. Since

 $\{x_1, \ldots, x_r\}$  is, obviously, contained in at most one edge (it is either an edge or not), we have

$$\frac{m}{r^r} < 1, \qquad m < r^r$$

which proves the assertion. [P. Erdős-L. Lovász, ibid.]

48. Since each cover defines a fractional cover, the inequality

$$\tau^*(H) \leq \tau(H)$$

is trivial. Similarly

$$\nu(H) \leq \nu^*(H)$$

is obvious.

With the notation introduced in the hint, any real vector x such that

$$A\mathbf{x} \leq \mathbf{1}, \, \mathbf{x} \geq 0$$

defines a fractional matching; any real vector y such that

$$A^T y \ge 1$$
,  $y \ge 0$ 

defines a fractional cover. Thus

$$v^*(H) = \max \{ \mathbf{1}^T \mathbf{x} \mid A\mathbf{x} \le \mathbf{1}, \, \mathbf{x} \ge 0 \},$$
  
 $\tau^*(H) = \min \{ \mathbf{1}^T \mathbf{y} \mid A^T \mathbf{y} \ge \mathbf{1}, \, \mathbf{y} \ge 0 \},$ 

and the relation

$$\nu^*(H) = \tau^*(H)$$

follows by the Duality theorem in linear programming.

**49.** (a) Let t be any k-cover of a graph G. Set

$$A = \{x: t(x) \ge k\}$$

$$B = \{x: t(x) = 0\}$$

and define

$$t'(x) = \begin{cases} 2 \text{ if } x \in A, \\ 0 \text{ if } x \in B, \\ 1 \text{ otherwise;} \end{cases}$$

$$t''(x) = t(x) - t'(x).$$

We claim that t' is a 2-cover and t'' is a (k-2)-cover. Let  $(x,y) \in E(G)$ . If  $x, y \notin B$ , then  $t'(x) + t'(y) \ge 2$ . If, say,  $y \in B$ , then  $t(x) + t(y) \ge k$  implies that  $t(x) \ge k$ ,  $x \in A$ , and  $t'(x) + t'(y) \ge 2$  again. So t' is a 2-cover. On the other hand,  $t''(x) + t''(y) \ge (t(x) - 1) + t(y) - 1 \ge k - 2$  unless, e.g.  $x \in A$ . But then  $t''(x) + t''(y) \ge t''(x) \ge k - 2$  follows again.

(b) Let w be a k-matching, and suppose k is even. Replace each edge E by w(E) parallel edges. The resulting graph G' has degrees at most k. By adding new edges (loops are allowed) we can obtain a k-regular graph G''. By 7.40 G''

has a 2-factor; the rest of its edges form a (k-2)-factor. Removing the edges of E(G'') - E(G'), we obtain a decomposition of E(G') into a 2-matching and a (k-2)-matching. These yield a decomposition of w into a 2-matching and a (k-2)-matching.

(c) Let t be an optimal fractional cover with rational weights (it is known from linear programming that such an optimal fractional cover exists). Let 2k be a common denominator of the weights in t. Then 2kt is a (2k)-cover. By (a),  $2kt = t_1 + \ldots + t_k$ , where  $t_1, \ldots, t_k$  are 2-covers. Thus

$$t = \frac{1}{k} \left( \frac{t_1}{2} + \ldots + \frac{t_k}{2} \right),$$

where  $\frac{t_1}{2}, \ldots, \frac{t_k}{2}$  are fractional covers. Since t is optimal, it follows that

 $\frac{t_1}{2}$ ,...,  $\frac{t_k}{2}$  must be optimal fractional covers too. Now, e.g.  $\frac{t_1}{2}$  consists of halves of integers.

For matchings the assertion follows similarly. Also

$$2\tau^*(H) = \sum_{x \in V(H)} t_1(x)$$

is an integer.

(d) Let t be an optimal 2-cover. Set

$$A = \{x : t(x) = 0\}.$$

Obviously, A is independent. Moreover for any point y in  $\Gamma(A)$ ,  $t(y) \ge 2$ , since there is an x in A adjacent to y and  $t(x) + t(y) \ge 2$ . Obviously,  $t(y) \le 2$  for every point, thus t(y) = 2.

Now  $t(z) \ge 1$  for every  $z \notin A \cup \Gamma(A)$ ; moreover, taking t(z) = 1 is sufficient to cover every edge of G twice. Thus t(z) = 1 by the optimality of t. So t is determined by A and

$$\sum_{x \in V(H)} t(x) = 2|\Gamma(A)| + |V(G) - \Gamma(A) - A| = |V(G)| + |\Gamma(A)| - |A|.$$

So, using (c),

$$2\tau^*(G) = \min_{t} \sum_{x} t(x) = |V(G)| + \min_{A \text{ indep.}} \{|\Gamma(A)| - |A|\}.$$

Thus (c) implies 7.37.

**50.** Let  $\{U, V\}$  be a 2-coloration of G, and let t be a k-cover. For  $1 \leq i \leq k$ , define

$$t_i(x) = egin{cases} 1 & ext{if} & x \in U ext{ and } t(x) \geq ext{i} \ & ext{or} & x \in V ext{ and } t(x) \geq k-i+1, \ 0 & ext{otherwise}. \end{cases}$$

We claim that  $t_1 + \ldots + t_k = t$  and that every edge is covered by every  $t_i$ . The first assertion is trivial. Let  $(x, y) \in E(G)$ ,  $1 \le i \le k$ . Since  $t(x) + t(y) \ge k$ ,

we have either  $t(x) \ge i$  or  $t(y) \ge k - i + 1$ . Accordingly,  $t_i(x) = 1$  or  $t_i(y) = 1$ . The assertion concerning matchings follows exactly as in the previous solution only now we can use 7.10 instead of 7.40.

**51.** (a) Let t, t' be minimum fractional covers of G, H, respectively. Define

$$\bar{t}(x, y) = t(x)t'(y)$$
  $(x \in V(G), y \in V(H)),$ 

then it is easy to check that t is a fractional cover of  $G \otimes H$ . Moreover,

$$\sum_{(x, y) \in V(G \otimes H)} \bar{t}(x, y) = \left(\sum_{x} t(x)\right) \left(\sum_{y} t'(y)\right) = \tau^*(G)\tau^*(H),$$

whence

$$\tau^*(G \otimes H) \leq \tau^*(G)\tau^*(H).$$

An analogous argument with fractional matchings yields

$$\nu^*(G \otimes H) \ge \nu^*(G)\nu^*(H),$$

and hence 13.48 implies the first relation.

Let S be a minimum point cover of  $G \otimes H$ . Let  $F \in E(H)$  and

$$S_F = \{x \in V(G) : (x, y) \in S \text{ for some } y \in F\}.$$

Then, clearly,  $S_F$  covers all edges of G, whence  $|S_F| \ge \tau(G)$ . Set  $\tau(G) = k$ . Define, for  $y \in V(H)$ ,

 $t(y) = |(V(G) \times \{y\}) \cap S|.$ 

Then

$$\sum_{y \in F} t(y) \ge |S_F| \ge k,$$

.e. t is a k-cover of H. Thus

$$\tau_k(H) \leq \sum_{y \in V(H)} t(y) = |S| = \tau(G \otimes H),$$

and thus

$$\tau(G \otimes H) \ge \tau_k(H) \ge k\tau^*(H).$$

This proves the upper bound in the second statement. The lower bound follows by a similar method as in the first step of the solution. The assertion concerning matching number follows analogously and therefore we omit details.

(b) Assume  $\tau(H) = \tau^*(H)$ . Then by (a),

$$\tau(G)\tau(H) \ge \tau(G \otimes H) \ge \tau(G)\tau^*(H),$$

which proves that

$$\tau(G\otimes H)=\tau(G)\tau(H).$$

Conversely, assume  $\tau(H) > \tau^*(H)$ . Then  $\tau_n(H) < n \cdot \tau(H)$  for some n. Let  $N = \tau_n(H)$  and define a hypergraph G on  $1, \ldots, N$  to consist of all (N - n + 1)-element subsets. Let  $\{x_1, \ldots, x_N\}$  be a minimum n-cover of H (the same point may occur several times on this list) and

$$S = \{(i, x_i) : 1 \leq i \leq N\}.$$

Then S represents all edges of  $G \otimes H$ ; for if  $E \times F$  is an edge of  $G \otimes H$   $(E \in E(G), F \in E(H))$ , then there are at least n points of S of the form  $(i, x_i)$ ,  $x_i \in F$ , and at most n-1 of these have  $i \notin E$  by the definition of G. Thus

$$\tau(G \otimes H) \leq |S| = N.$$

Since clearly  $\tau(G) = n$ , we have

$$\tau(G \otimes H) = N = \tau_n(H) < n\tau(H) = \tau(G)\tau(H)$$

[C. Berge, M. Simonovits, in: *Hypergraph Seminar*, Lecture Notes in Math. 411, Springer, 1974, 21–33].

(c) By (a),

$$\tau^*(H^p) = \tau^*(H)^p,$$

thus

$$\tau(H^p) \geq \tau^*(H)^p.$$

To estimate  $\tau(H^p)$  from above we use 13.30. We have

$$d(H^p) = d(H)^p$$

by easy computation and thus,

(2) 
$$\tau(H^p) \leq (1 + \log d(H^p))\tau^*(H^p) =$$
$$= (1 + p \log d(H))\tau^*(H)^p.$$

By (1) and (2),

$$\tau^*(H) \leq \sqrt[p]{\tau(H^p)} \leq \sqrt[p]{1+p\log d(H)} \ \tau^*(H).$$

Thus

$$\lim_{p\to\infty}\sqrt[p]{\tau(H^p)}=\tau^*(H).$$

[R.J. McEliece-E.C. Posner, Ann. Math. Stat. 42 (1971) 1706-1716].

52. Suppose H contains two edges  $E_1$ ,  $E_2$  with a common point x. Since it is  $\tau$ -critical there exists a 1-cover  $t_i$  of  $H - \{E_i\}$  such that

$$\sum_{x} t_i(x) = \tau(H) - 1.$$

Then define

$$t(y) = egin{cases} 1 & ext{if} & y = x, \ & & & \ t_1\!\left(y\right) + t_2\!\left(y\right) & ext{otherwise.} \end{cases}$$

Then, clearly, t is a fractional cover. Moreover, its size is

$$\sum_{y} t(y) = 1 + \frac{1}{2} \left( \sum_{y} t_1(y) + \sum_{y} t_2(y) \right) = \tau(H) - \frac{1}{2}.$$

Hence

$$au^*(H) \le au(H) - rac{1}{2} < au(H)$$

[L. Lovász, in: Hypergraph Seminar, Lecture Notes in Math. 411, Springer, 1974, 111–126].

53. Let  $E_0 = \{v_1, \ldots, v_k\} \in E(H)$ . Obviously,  $k \geq 2$ . Let  $w_i$  be a maximum matching of  $H - v_i$   $(i = 1, \ldots, k)$ ; then, since H is  $\nu$ -critical,

$$\sum_{E} w_i(E) = \nu(H) \qquad (i = 1, \dots, k).$$

Set

$$w(E) = egin{cases} 1 & ext{if } E = E_0, \ rac{1}{k} \left( w_1(E) + \ldots + w_k(E) 
ight) & ext{if } E 
eq E_0. \end{cases}$$

Then it is easy to check that w is a fractional matching; in fact, if  $x \notin E_0$ , then

$$\sum_{E\ni x} w(E) = \frac{1}{k} \left( \sum_{E\ni x} w_1(E) + \ldots + \sum_{E\ni x} w_k(E) \right) \le 1$$

and if  $x = v_1$  (say), then

$$\sum_{E\ni\nu_1}w(E)=\frac{1}{k}\left(1+\sum_{E\ni\nu_1}w_2(E)+\ldots+\sum_{E\ni\nu_1}w_k(E)\right)\leq 1.$$

Thus  $\nu^*(H)$  can be estimated as follows:

$$v^*(H) \ge \sum_{E} w(E) = \frac{1}{k} \left( 1 + \sum_{E \ne E_{\bullet}} \left( w_1(E) + \ldots + w_k(E) \right) \right) =$$

$$= \frac{1}{k} \left( 1 + kv(H) \right) = v(H) + \frac{1}{k} > v(H).$$

[L. Lovász, ibid.]

54. Suppose indirectly that there are balanced hypergraphs with  $\tau(H) > \nu(H)$  and consider a minimum example. Then

$$\tau(H - \{E\}) = \nu(H - \{E\})$$

for every  $E \in E(H)$  and so,

$$\tau(H - \{E\}) = \nu(H - \{E\}) \le \nu(H) < \tau(H),$$

i.e. H is  $\tau$ -critical. Since H cannot consist of disjoint edges, there are  $E_1$ ,  $E_2 \in E(H)$ ,  $x \in V(H)$  with  $x \in E_1 \cap E_2$ .

Since  $\tau(H - \{E_i\}) < \tau(H)$ , we have a  $(\tau(H) - 1)$ -element point-cover  $T_i$  in  $H - \{E_i\}$ . Obviously,  $E_i \cap T_i = \emptyset$ . Set

$$W = (T_1 \triangle T_2) \cup \{x\}.$$

By 13.38, the subhypergraph  $H_W$  has chromatic number 2 (if 1-element edges are ignored). Let  $\{A, B\}$  be a bicoloration of it,  $A \cap B = \emptyset$ ,  $A \cup B = W$ ,  $|A| \leq |B|$ . Since

$$|W| = |T_1 - T_2| + |T_2 - T_1| + 1 = 2\tau(H) - 2|T_1 \cap T_2| - 1,$$

we have  $|A| \le \tau(H) - |T_1 \cap T_2| - 1$ .

Now  $A \cup (\overline{T}_1 \cap \overline{T}_2)$  covers every edge of H. For if an edge  $E \in E(H)$ ,  $E \neq E_1$ ,  $E_2$  does not meet  $T_1 \cap T_2$ , then it meets both  $T_1 - T_2$  and  $T_2 - T_1$ , i.e. it has at least two points in W and since  $\{A, B\}$  is a bicoloration of  $H_W$ ,  $E \cap A \neq \emptyset$ . Similarly, if  $E = E_i$ , then it has two points in W and we conclude as before. So  $A \cup (T_1 \cap T_2)$  is a point-cover of H. But

$$|A \cup (T_1 \cap T_2)| = \tau(H) - 1,$$

a contradiction [C. Berge; B].

**55.** Obviously, (iii)  $\Rightarrow$  (i) and (iii)  $\Rightarrow$  (ii). Assume (i) holds but (iii) fails and consider a minimal such counterexample. Then

$$\tau(H) > \nu(H), \qquad \tau(H-x) = \nu(H-x)$$

and since

$$\tau(H-x) \ge \tau(H)-1,$$

we have

$$\nu(H-x)=\nu(H),$$

i.e. H is v-critical. By 13.53, however, it follows that

$$v^*(H) > v(H),$$

a contradiction. So (iii)  $\Rightarrow$  (i).

Similarly, it follows that, if H is a minimal hypergraph, which satisfies (i) but not (iii), then H is  $\tau$ -critical (cf. 13.54), and so, by 13.52,  $\tau(H) > \tau^*(H)$ , a contradiction.

Thus (i) ⇔ (iii) ⇔ (ii) [L. Lovász, ibid.].

56. Let us first verify the assertion of the hint.

It suffices to show that, if we add a new edge, parallel to an old one to H, then the resulting hypergraph  $H_0$  has property (iv). Let E be this new edge, parallel to  $E_0 \in E(H)$ .

It suffices to show that  $d(H_0) = q(H_0)$  since this follows similarly for the partial hypergraphs.  $d(H_0) \leq q(H_0)$  is trivial.

If  $d(H_0) = d(H) + 1$ , then

$$q(H_0) \le q(H) + 1 = d(H) + 1 = d(H_0).$$

So suppose  $d(H_0) = d(H)$ , i.e. no point of  $E_0$  has maximum degree in H. Since H has property (iv), its edges can be d(H)-colored. Let  $E_0$  have color red, say, and let  $F_1, \ldots, F_t$  be the other red edges. Let  $H' = H - \{F_1, \ldots, F_t\}$ . Let us determine d(H'). If x has degree less than d(H) in H, then it has degree less than d(H) in H'; if x has maximal degree d(H) in H', then it must be adjacent to a red edge, and since the points of  $E_0$  have degree less than d(H),  $x \notin E_0$ . Hence x has degree less than d(H) in H'. Thus d(H') < d(H), and, con-

sequently, the edges of H' can be (d(H)-1)-colored. Color the edges E,  $F_1, \ldots, F_t$  by a further color; we obtain a d(H)-coloration of the edges of  $H_0$ . This proves the assertion of the hint.

Now we turn to the solution of the problem.

(iv)  $\Rightarrow$  (i). Let w be an optimal fractional matching of H, and let k be a common denominator of the weights w(E) (we may suppose these are rational). Replace each edge of H by kw(E) parallel edges. The resulting hypergraph  $H_1$  has  $d(H_1) \leq k$  by the definition of w. Since by the lemma, it also satisfies (iv) we have

$$q(H_1) \leq k$$
,

i.e. the edges of  $H_1$  can be k-colored.

Since each color occurs at most  $v(H_1) \leq v(H)$  times, it follows that

$$|E(H_1)| \leq k\nu(H).$$

Since by the construction of  $H_1$ 

$$|E(H_1)| = k v^*(H),$$

if follows that

$$v^*(H) \leq v(H)$$
.

Since the converse inequality is trivial, (i) follows.

The relation (iii)  $\Rightarrow$  (iv) can be reduced to the relation (iv)  $\Rightarrow$  (iii) by constructing a hypergraph  $H^*$  as follows. Let  $V(H^*)$  consist of all 1-matchings of H; and, for every  $E \in E(H)$ , let the set  $E^*$  of all 1-matchings of H containing E be an edge of  $H^*$ . We omit the details.

(iii)  $\Rightarrow$  (v). Suppose H is normal, i.e. it satisfies (iii).

Let  $E_1, \ldots, E_k \in E(H)$ ,  $E_i \cap E_j \neq \emptyset$   $(1 \leq i, j \leq k)$ . Then set  $H' = (V(H), \{E_1, \ldots, E_k\})$ . We have

$$v(H') = 1$$
,

so we must have

$$\tau(H')=1$$
,

i.e.  $E_1, \ldots, E_k$  must have a point in common. Thus H has the Helly-property. Let  $G_1$  be an induced subgraph of  $\overline{L(H)}$ . Then  $G_1 \cong \overline{L(H')}$  for some partial hypergraph H' of H. We have

$$\chi(G_1) = \tau(H');$$

for a k-coloration of  $G_1$  corresponds to a partition of L(H') into k cliques and so, to a k-element point-cover of H' (since H' has the Helly-property). Similarly,

$$\omega(G_1) = \nu(H').$$

Since H is normal, we have

$$\chi(G_1) = \tau(H') = \nu(H') = \omega(G_1),$$

i.e. G is perfect.

The converse follows similarly [L. Lovász, B].

57. Suppose G is perfect. Let  $C_1, \ldots, C_N$  be the cliques of G. Take N points  $p_1, \ldots, p_N$  and, with each  $x \in V(G)$ , associate an edge  $E_x$  such that

$$p_i \in E_x \Leftrightarrow x \in C_i$$
.

Then

$$H = (\{p_1, \ldots, p_N\}; \{E_x: x \in V(G)\})$$

is a hypergraph such that  $L(H) \cong G$ . We claim that H satisfies property (iv) of normal hypergraphs. Let H' be any partial hypergraph of H. Then L(H') is an induced subgraph of G, hence L(H') is perfect, i.e.

$$\chi(L(H')) = \omega(L(H')).$$

Now observe that  $\chi(L(H')) = q(H')$ . Also,  $\omega(L(H')) = d(H')$ ; for if  $E_{x_1}, \ldots, E_{x_p}$  are edges of H' adjacent to the same point, then  $x_1, \ldots, x_p$  are points of L(H') forming a complete subgraph; hence  $\omega(L(H')) \geq d(H')$ . On the other hand, if  $x_1, \ldots, x_p$  form a complete subgraph of L(H'), then there is a  $C_j$  such that  $x_1, \ldots, x_p \in C$ , and then  $p_j$  is adjacent to  $E_{x_1}, \ldots, E_{x_p}$ , hence  $d(H') \geq \omega(L(H'))$ . So

$$\varrho(H') = \chi(L(H')) = \omega(L(H')) = d(H').$$

Thus H satisfies (iv) in 13.56 and so, it is normal. This implies that  $\overline{L(H)} = \overline{G}$  is perfect.

If  $\bar{G}$  is perfect, then the fact that G is perfect follows by interchanging the

roles of G and G. [ibid.]

## § 14. Ramsey Theory

1. (a) We use induction on k+l. For k=1 or l=1 the assertion is clear. Suppose k, l>1. Let  $x\in V(G)$ . Since

$$d_G(x)+d_{\overline{G}}\left(x
ight)=\left|V(G)
ight|-1=inom{k+l}{k}-1=inom{k+l-1}{k-1}+inom{k+l-1}{k}-1,$$

we have either

$$d_G(x) \geq {k+l-1 \choose k-1} \quad ext{or} \quad d_{\overline{G}}(x) \geq {k+l-1 \choose k}.$$

Assume, e.g. that the first inequality holds, and let  $G_1$  be the subgraph induced by the neighborhood of x. Then, by the induction hypothesis,  $G_1$  contains either a complete k-graph or l+1 independent points. In the second case we are done. If  $G_1$  contains a complete k-graph H, then  $V(H) \cup \{x\}$  spans a complete (k+1)-graph.

(b) By (a),

$$R_{\mathbf{2}}(a_{\mathbf{1}},\,a_{\mathbf{2}}) \leq \binom{a_1+a_2-2}{a_1-1}\,.$$

Now suppose that  $k \geq 3$  and that  $R_{k-1}(b_1, \ldots, b_{k-1})$  exists for each  $b_1, \ldots, b_{k-1}$ . Then we claim that

$$R_k(a_1,\ldots,a_k) \leq R_{k-1}(a_1,\ldots,a_{k-2},R_2(a_{k-1},a_k)).$$

In fact, consider a complete graph with  $R_{k-1}(a_1, \ldots, a_{k-2}, R_2(a_{k-1}, a_k))$  points and k-color its edges. Identify the  $(k-1)^{\rm st}$  and  $k^{\rm th}$  colors for the moment; then we know that either the  $i^{\rm th}$  color contains a complete  $a_i$ -graph for some  $1 \leq i \leq k-2$  (in which case we are done) or the last color contains a complete  $R_2(a_{k-1}, a_k)$ -graph K. The edges of K are colored with k-1 and k and so, by definition, either color k-1 contains a complete  $a_{k-1}$ -graph or color k contains a complete  $a_k$ -graph. This completes the proof [see H, B, ES].

2. (a) I. We use induction on k. For k = 2,  $R_2(3,3) \le 6$  follows from 14.1a (here equality holds as is shown by the pentagon).

Let x be any point of a k-colored complete graph on [ek!] + 1 points. There are [ek!] edges adjacent to this point, which are split into k classes. Since

$$[ek!] = \left[\sum_{j=0}^{\infty} \frac{k!}{j!}\right] = \sum_{j=0}^{k} \frac{k!}{j!} = 1 + k \sum_{j=0}^{[k-1]} \frac{(k-1)!}{j!} = 1 + k[e(k-1)!],$$

one of the k colors, let us call it red, contains at least [e(k-1)!] + 1 edges adjacent to x. Let S be the set of those points joined to x by a red edge.

If S spans a red edge this forms, together with x, a red triangle and we are finished. If S spans no red edge it spans a complete graph with [e(k-1)!]+1 points, whose edges are (k-1)-colored; thus by the induction hypothesis, one of these k-1 colors contains a triangle and we are done again [see B].

II. To prove  $R_3(3,3,3) = [e\cdot 3!] = 17$ , we have to 3-color the edges of a complete 16-graph avoiding monochromatic triangles. Let x be any point, it easily follows that x must be adjacent to 5 edges of each color and those points joined to x by a red edge must span a green and a blue pentagon (Fig. 100, p. 480). It is natural to assume that rotation through 120° maps red onto green and green onto blue, hence it suffices to specify the red edges. It is also quite natural to assume that rotation of each of the three pentagons by 72° keeps each color invariant. Some experimenting results in Fig. 101, p. 480 (where the point x is not shown for the sake of clearness and the three pentagons are drawn in a geometrically different way to show this latter symmetry).

Remark: This coloration can be described as follows: Let P be a pentagon and  $V(K_{16})$  the set of subsets of V(P) with even cardinality. For A,  $B \in V(K_{16})$ , let (A, B) be colored

red if  $A \triangle B$  is an edge of P, blue if  $A \triangle B$  is an edge of  $\overline{P}$ , green if  $|A \triangle B| = 4$ .




Fig. 101

It is easy to see that each edge gets one of the three colors. If  $A, B, C \in V(K_{16})$ , then

$$(A \triangle B) \triangle (B \triangle C) = A \triangle C;$$

hence it is seen that none of the three colors contains a triangle.

(b) The upper bound is trivial from 14.1a:

$$R_2(k, k) \le {2k-2 \choose k-1} + 1 < 2^{2k}.$$

To prove the lower bound we may suppose  $k \geq 4$ . Let  $n = 2^{k/2}$  and consider the hypergraph defined in the hint. Clearly this hypergraph is 2-colorable iff the edges of  $K_n$  can be 2-colored in such a way that no monochromatic complete k-graph arises, i.e. if  $n < R_2(k, k)$ .

The number of edges of H is, clearly,  $\binom{n}{k}$  and H is  $\binom{k}{2}$ -uniform. So by 13.41, H is 2-colorable provided

$$\binom{n}{k} < 2^{\binom{k}{2}-1}.$$

But this is so, because

$$\binom{n}{k} < \frac{n^k}{2^{k-1}} < \frac{2^{\frac{k^2}{2}}}{2^{k-1}} = 2^{\binom{k}{2} - \frac{k}{2} + 1} \le 2^{\binom{k}{2} - 1}$$

[P. Erdős; see ES].

**3.** We use induction on r and  $a_1 + \ldots + a_k$ . For r = 2 we already know that  $R_k^{\alpha}(a_1,\ldots,a_k)=R_k(a_1,\ldots,a_k)$  exists. Suppose r>2 (as a matter of fact,  $R_k^1(a_1,\ldots,a_k)$  trivially exists and we could start the induction with r=2). Also, if  $a_i = 1$  for any i, the assertion is trivial; so we may suppose  $a_i \geq 2$ . Now let  $x \in V(K_n^r)$ . Color every (r-1)-tuple S in  $V(K_n^r) - \{x\}$  with the color of  $S \cup \{x\}$ . Thus the edges of  $K_{n-1}^{r-1}$  are k-colored. So if

$$n-1 \ge R_k^{r-1}(b_1,\ldots,b_k),$$

then, for some  $1 \le i \le k$ , there is a  $V_i \subseteq V(K_n^r) - x$ ,  $|V_i| = b_i$  such that all r-tuples of  $V_i \cup \{x\}$  containing x have color i. Now if we choose here

$$b_i = R_k^r(a_1, \ldots, a_{i-1}, a_i - 1, a_{i+1}, \ldots, a_k)$$

(this number exists by the induction hypothesis on  $a_1 + \ldots + a_k$ ) we find that either  $V_i$  has an  $a_i$  subset all of whose k-tuples have color j for some  $j \neq i$  (in which case we are done), or else,  $V_i$  contains an  $(a_i - 1)$ -subset V'all of whose r-tuples are colored i. Thus all k-subsets of  $V' \cup \{x\}$  are colored i, and we are done again [see H, B].

(b) Let  $m = R_k^r(a)$ ,  $N = r + k^{\binom{m}{r}}$  and  $S = \{1, \ldots, N\}$ . Trivially  $N < k^{mr}$  except for trivial cases. Assume that the set of all (r+1)-element subsets of Sis k-colored. We prove that there is an a-element subset of S all of whose (r+1)-subsets have the same color. We follow the hint.

We define elements  $x_i$  and a "reservoir"  $U_i$  of points recursively such that (i)  $x_1, \ldots, x_i \notin U_i, x_{i+1}, \ldots, x_m \in U_i$ ,

- (ii) for each subset  $1 \le v_1 < \ldots < v_r \le i$ , all (r+1)-tuples  $(x_{\nu_1}, \ldots, x_{\nu_r}, y)$  with  $y \in U_i$  have the same color, and moreover

(iii) 
$$|U_i| \ge k^{\binom{m}{r} - \binom{i}{r}}$$
  $(i = 1, ..., m)$ .

All these conditions clearly hold for  $i=1,\ldots,r-1$ , if  $x_1=1,\ldots,x_{r-1}=r-1$  and  $U_i=\{i+1,\ldots,N\}$ . Now assume that  $x_1,\ldots,x_i,U_1,\ldots,U_i$   $(r-1\leq i\leq m-1)$  have been defined in accordance with (i), (ii) and (iii). Choose an arbitrary element of  $U_i$  for  $x_{i+1}$ . To define  $U_{i+1}$ , call two elements  $y,y'\in U_i-\{x_{i+1}\}$  equivalent, if the colors of

$$(x_{\nu_1}, [...], x_{\nu_{r-1}}, [x_{i+1}, [y)]$$
 and  $(x_{\nu_1}, .[...], x_{\nu_{r-1}}, x_{i+1}, [y'])$ 

are the same for every  $1 \leq \nu_1 < \ldots < \nu_{r-1} \leq i$ . There are at most  $k^{\binom{i}{r-1}}$  equivalence classes. Define  $U_{i+1}$  as the largest one of these. (i) and (ii) are trivially fulfilled. Also

$$|U_{i+1}| \ge \left[\frac{|U_i|| - |1|}{k^{\binom{i}{r-1}}}\right]^* \ge k^{\binom{m}{r} - \binom{i+1}{r}}.$$

Thus (iii) holds as well.

We now define a k-coloration of the r-subsets of  $\{x_1, \ldots, x_m\}$  by coloring  $(x_{r_1}, \ldots, x_{r_r})$   $(1 \le r_1 < \ldots < r_r \le m)$  with the common color of (r+1)-sets  $(x_{r_1}, \ldots, x_{r_r}, y)$   $(y \in U_{r_r})$ . By the choice of m, there is an a-subset A of  $\{x_1, \ldots, x_m\}$  such that all r-tuples of A have the same color, say red. But then all (r+1)-subsets of A are red in the original coloration. In fact, let  $(x_{r_1}, \ldots, x_{r_{r+1}})$  be an (r+1)-tuple of elements of A. Then by the choice of A,  $(x_{r_1}, \ldots, x_{r_r})$  is red. But then, since  $x_{r_{r+1}} \in U_{r_r}$ , the above (r+1)-tuple is red by the definition of the coloration of r-tuples. This proves that  $R_k^{r+1}$  (a)  $\le N$ . [P. Erdős–R. Rado, Proc. London Math. Soc. Series 3, 2 (1952) 417–439. Concerning the sharpness of this estimation, see P. Erdős–A. Hajnal–R. Rado, Acta[Math. Hung. 16 (1965) 93–196.]

4. For n=3 the assertion is trivial; suppose n>3. Let us remove a point x; we find, by induction, an (n-1)-circuit C in the rest, which is either monochromatic or has two points a, b such that one (a, b)-arc P is red, while the other one, Q is blue. The first case can be regarded as the special case of the latter when a=b.

We may assume edge (a, x) is red. Let u be the point on Q next to a (if Q consists of a single point, u can be any neighbor of a). Then

$$P + (a, x) + (x, u) + (Q - a)$$

is a Hamiltonian circuit in the complete n-graph such that either the two (b, x)-arcs are red and blue, respectively or the two (b, u) arcs are so [H. Raynaud; see B].

5. Let P,  $Q_1$ ,  $Q_2$  be as in the hint. By induction on k, we may assume that P has length k-1. Let a, b;  $c_1$ ,  $d_1$ ;  $c_2$ ,  $d_2$  denote the endpoints of P,  $Q_1$ ,  $Q_2$ , respectively. Suppose indirectly that there is a point  $x \notin V(P) \cup V(Q_1) \cup V(Q_2)$ . Since, from the structure of  $Q_1$ ,  $Q_2$ ,

$$|V(Q_i) \cap V(P)| = \frac{1}{2}(|V(Q_i)| - 1) = |V(Q_i) - V(P)| - 1,$$

we have

$$\begin{split} | \, V(P) - V(Q_1) - V(Q_2) | &= k + 2 - | \, V(Q_1) \cup V(Q_2) - V(P) | > \\ > k + 2 - | \, V(G) - V(P) | &= 2k + 2 - \left | \frac{3k+1}{2} \right | = \left \lceil \frac{k}{2} \right \rceil + 2. \end{split}$$

Thus  $Q_1 \cup Q_2$  misses at least  $\left\lceil \frac{k}{2} \right\rceil + 3$  points of P. This implies that there is

an edge  $(u, v) \in E(P)$  such that  $u, v \neq a, b$  and  $u, v \notin V(Q_1 \cup Q_2)$ .

One of the edges (x, u), (x, v) must be blue; otherwise, replacing (u, v) by (u, x) and (x, v) we could increase the length of P. Let, say, (x, u) be blue. Then  $(c_1, u)$  must be red otherwise  $(c_1, u)$  and (x, u) could be added to  $Q_1$ . But then  $(c_1, v)$  must be blue by the same argument as above. Similarly,  $(c_2, v)$  is blue. Now

$$\begin{aligned} Q_1' &= Q_1 + (c_1, v) + (c_2, v) + Q_2, \\ Q_2' &= \{x\} \end{aligned}$$

are two paths with total length greater than  $|E(Q_1)| + |E(Q_2)|$  satisfying the same conditions as  $Q_1$  and  $Q_2$ , a contradiction.

Thus  $P \cup Q_1 \cup Q_2$  covers all points. Let us consider the edges  $(c_1, a)$ ,  $(c_2, a)$ ,  $(d_1, b)$ ,  $(d_2, b)$ . By the maximality of P, these are blue. Hence, the circuit

$$C = Q_1 + (c_1, a) + (c_2, a) + Q_2 + (d_1, b) + (d_2, b)$$

is blue. The length of C is

$$2\left|V(G)-V(P)\right|=2\left\lceil\frac{k+1}{2}\right\rceil.$$

If k is odd this is equal to k+1 and so, removing any edge of C we have a blue path of length k. Suppose k is even. If there is a blue edge joining C to V(G)-V(C), we find such a blue path again. So suppose all edges connecting C to V(G)-V(C) are red. Then a red path with length  $2\left\lceil \frac{k+1}{2}\right\rceil \geq k$  can trivially be found (Fig. 102) [L. Gerencsér-A. Gyárfás, Annales Univ. Sci. R. Eötvös Sectio Math. X. (1967) 167-170].




Fig. 102

**6.** (a) Let  $(x_0, \ldots, x_{2k})$  be a red circuit. If  $(x_i, x_{i+2})$  is red for some  $0 \le i \le 2k$  (where  $x_{2k+1} = x_0, x_{2k+2} = x_1$  etc.), then we trivially have a red (2k)-circuit. Thus we may assume  $(x_i, x_{i+2})$  is blue for each  $0 \le i \le 2k$ .

The circuit  $(x_0, x_2, \ldots, x_{2k}, x_1, \ldots, x_{2k-1})$  is a blue (2k+1)-circuit. Therefore as above, we may assume  $(x_i, x_{i+4})$  is red for each  $1 \le i \le 2k$ . Now  $(x_0, x_4, \ldots, x_{2k}, x_2, x_1)$  is a 2k-circuit, which is red except possibly for its edge  $(x_{2k}, x_2)$ . Thus we may assume that  $(x_{2k}, x_2)$  is blue and similarly, so is  $(x_i, x_{i+3})$  for each  $0 \le i \le 2k$ . Now the circuit  $(x_0, x_3, x_6, x_8, \ldots, x_{2k}, x_1, x_{2k-1})$   $x_{2k-3}, \ldots, x_5, x_2)$  is a blue (2k)-circuit.

14.6

(b) Let  $(x_0, \ldots, x_{2k-1})$  be a red circuit, and suppose there is no monochromatic (2k-1)-circuit. As before,  $(x_i, x_{i+2})$  is blue for  $1 \le i \le 2r$  (now setting

 $x_{2k+i} = x_i).$ 

We claim that  $(x_i, x_{i+2\nu})$  is blue for each  $1 \le i \le 2k$ ,  $1 \le \nu \le k-1$ . Suppose indirectly that  $(x_0, x_{2\nu})$  is red (say). Then all edges of  $(x_0, x_{2\nu}, x_{2\nu-1}, \ldots, x_1, x_{2\nu+2}, x_{2\nu+3}, \ldots, x_{2k-1})$  are red except possibly  $(x_1, x_{2\nu+2})$ . Thus  $(x_1, x_{2\nu+2})$  must be blue. Similarly, the circuit

$$(x_0, x_1, \ldots, x_{2\nu-2}, x_{2k-1}, x_{2k-2}, \ldots, x_2)$$

implies that  $(x_{2\nu-2}, x_{2k-1})$  is blue. Now the circuit

$$(x_{2k-1}, x_{2\nu-2}, x_{2\nu-4}, \dots, x_0, x_{2k-2}, \dots, x_{2\nu+2}, x_1, x_3, \dots, x_{2k-3})$$

has length 2k-1 and is blue, a contradiction. So  $(x_i, x_{i+2\nu})$  is blue for every  $i, \nu$  and thus,  $\{x_1, \ldots, x_{2k-1}\}, \{x_0, \ldots, x_{2k-2}\}$  induce blue complete subgraphs.

(c) The case m=4 can be verified directly; suppose  $m\geq 5$ . For the two graphs  $G_1$ ,  $G_2$  formed by the red and blue edges, respect

For the two graphs  $G_1$ ,  $G_2$  formed by the red and blue edges, respectively, we have

$$\mid E(G_1) \mid + \mid E(G_2) \mid = \binom{2m-1}{2} \, ,$$

whence, e.g.

$$\mid E(G_1) \mid \geq \frac{1}{2} \binom{2m-1}{2} = \frac{m-1}{2} \left(2m-1\right) > \frac{(m-1) \left(\mid V(G_1) \mid -1\right)}{2} \, .$$

Thus by 8.26,  $G_1$  contains an M-circuit for some  $M \geq m$ .

Consider a monochromatic M-circuit C with  $M \ge m$ , M minimal; let, e.g. C be red. If M = m, we are done. Suppose indirectly that M > m. By (a), (b), we also have a monochromatic (M-1)-circuit (contradicting the choice of M) or M = 2k is even and we have two disjoint complete graphs K and H, both in blue, with  $V(K) \cup V(H) \supseteq V(C)$ ,  $|V(K)| \ge k$ ,  $|V(H)| \ge k$ . Choose here

|V(K)| + |V(H)| to be maximal.

If there are two independent blue edges connecting K to H, then obviously we will have a blue (M-1)-circuit. Thus we may suppose all blue edges between K and H have a point in common  $x \in V(K)$ . If m is even we trivially form a red m-circuit from the edges connecting H to K so suppose m is odd. Let  $v \in V(G) - V(C)$ . If v is joined by red edges to both K - x and H, we can form a red m-circuit from these two red edges and the red (K, H)-edges. So suppose that for each  $v \in V(G) - V(C)$  either all (v, K - x)-edges or all (v, H)-edges are blue. The latter cannot happen by the maximality of |V(H)| + |V(K)| and the same reasoning implies that (v, x) is red.

If  $|V(H)| \ge m$ , we are trivially done so suppose  $|V(H)| \le m-1$ . Consider the sets S = V(K-x), T = V(G) - S - V(H). Then  $|S| + |T| \ge (2m-1) - (m-1) = m$ ,  $|S| \ge k \ge \frac{m-1}{2}$ , and all edges spanned by S or joining S to T are blue. If  $|S| > \frac{m-1}{2}$  here we trivially find a blue m-circuit. Similarly,




Fig. 103

we are finished, if  $|S| = \frac{m-1}{2}$  but T induces at least one blue edge. Thus |V(H)| = m-1,  $|S| = \frac{m-1}{2}$ ,  $|T| = \frac{m+1}{2}$  and T induces a red complete graph. If a point of T is joined to at least two points of H by blue edges, we are finished again.

Thus we may suppose at most one blue edge connects each  $v \in T$  to H. Since |V(H)| = m-1, H contains  $\frac{m-3}{2}$  points, which are connected to T by red edges only and a further point, which is connected to T by at most one blue edge. If M is the set of these  $\frac{m-1}{2}$  points we easily find a red m-gon in  $V(T) \cup M$  (Fig. 103). [A. Bondy-P. Erdős, J. Comb. Theory 14 (1973) 46–54. For other Ramseyan results concerning circuits, see V. Rosta, J. Comb. Theory (B) 15 (1973) 94–104 and 105–120.]

- 7. Suppose indirectly that the points of the plane can be 3-colored with no two points at distance 1 having the same color. Let a in Fig. 17 (p. 134) have, say, color red. Then b, c must be blue and green (in some order). Therefore, f must be red. Similarly, g must be red. But then f, g form a pair at distance 1 with the same color, a contradiction [H. Hadwiger, *Elemente d. Math.* 16 (1961) 103-104].
- 8. Suppose the triangle ABC in Fig. 18 (p. 135) has red vertices. Then, if any of D, E, F is red we are done; so suppose D, E, F are blue. Now, if G or H is blue, then DEG or DFH is a monochromatic triangle; if G and H are red, then BGH is one [P. Erdős–R.L. Graham–P. Montgomery–B.L. Rothschild–J. Spencer–E.G. Straus, J. Comb. Th. 14 (1973) 341–363].

9. (a) Let us split the plane into strips of width  $\frac{\sqrt{3}}{2}$ , counting their lower edge with the strips but not the upper one. Color these strips alternately red and blue. It is easy to see that any regular triangle with sides 1 has to contain vertices from two adjacent strips and so, it must not be monochromatic.

14.9

(b) We may as well consider 1,  $\sqrt{3}$ ,  $\frac{\pi}{\sqrt{2}}$  as this only means a transformation of the unit by a scalar factor. We show that, if we 2-color the points of the plane, there will be a monochromatic regular triangle with side 1 or  $\sqrt{3}$ ; in either case, the preceding problem will imply the existence of a monochromatic regular triangle with side 1 or  $\sqrt{3}$ ; in either case, the preceding problem will imply the existence of a monochromatic regular triangle with side 1 or  $\sqrt{3}$ .

matic triangle with sides 1,  $\sqrt{3}$ ,  $\frac{\pi}{\sqrt{2}}$ .

If not all points have the same color there must be a pair a, b of points at distance 2 colored differently; for any pair of points with different colors can be connected by a polygon with sides 2 and one of these sides must have differently colored endpoints.

So we may assume a is red and b is blue in Fig. 19 (p. 135). Also we may assume without loss of generality that c is red. If d or e is red we have a monochromatic regular triangle with side 1; if both d, e are blue we have a monochromatic regular triangle with side  $\sqrt{3}$  [P. Erdős et al. in: *Infinite and Finite Sets*, Coll. Math. Soc. J. Bolyai 10 (1974) 529–583].

10. (a) Let a, b the lengths of edges of the rectangle R. Let  $v_0, \ldots, v_N$  be the vertices of a regular simplex in the N-dimensional space with side a, where  $N = k^{k+1}$ ; let  $w_0, \ldots, w_k$  be the vertices of a regular simplex in the k-dimensional space with side b. Consider the points  $(v_i, w_j)$  of the (N + k)-dimensional space  $(0 \le i \le M, \ 0 \le j \le k)$ . For each fixed i, the k+1 points  $(v_i, w_j)$  can be colored in  $k^{k+1} = N$  ways; therefore, there are two indices  $0 \le i_1 < i_2 \le N$  such that  $(v_{i_1}, w_{j_1})$  and  $(v_{i_1}, w_{j_1})$  have the same color for each  $0 \le j \le k$ . Also, there are two indices  $0 \le j_1 < j_2 \le k$  such that  $(v_{i_1}, w_{j_1})$  and  $(v_{i_1}, w_{j_2})$  have the same color. Hence,

$$(v_{i_1}, w_{j_1}), (v_{i_1}, w_{j_2}), (v_{i_2}, w_{j_1}), (v_{i_2}, w_{j_2})$$

have the same color. These four points obviously form a rectangle congruent to R.

(b) Let a, b be the two side vectors of R, then, as R is not rectangular,  $ab \neq 0$ . We may suppose ab = 1, since this only means transformation of the unit by a scalar factor. We claim n(4, R) does not exist. For any n, construct a 4-coloration of the n-dimensional Euclidean space by giving  $\mathbf{w}$  the color i  $(0 \leq i \leq 3)$ , if  $[\mathbf{w}^2] \equiv i \pmod{4}$ . We prove that there exists no monochromatic set congruent to R.

Suppose R' is such a set. Then

$$R' = \{ w, w + a', w + b', w + a' + b' \}.$$

for some w, a  $b \in E^n$ . Since this is monochromatic we have

$$\lceil \mathbf{w}^2 \rceil \equiv \lceil (\mathbf{w} + \mathbf{a}')^2 \rceil \equiv \lceil (\mathbf{w} + \mathbf{b}')^2 \rceil \equiv \lceil (\mathbf{w} + \mathbf{a}' + \mathbf{b}')^2 \rceil \pmod{4}$$

But

and so,

$$[\mathbf{w}^2] - [(\mathbf{w} + \mathbf{a}')^2] - [(\mathbf{w} + \mathbf{b}')^2] + [(\mathbf{w} + \mathbf{a}' + \mathbf{b}')^2] \equiv 0 \pmod{4}$$

$$[\mathbf{w}^2] - [(\mathbf{w} + \mathbf{a}')^2] - [(\mathbf{w} + \mathbf{b}')^2] + [(\mathbf{w} + \mathbf{a}' + \mathbf{b}')^2] =$$

$$= \mathbf{w}^2 - (\mathbf{w} + \mathbf{a}')^2 - (\mathbf{w} + \mathbf{b}')^2 + (\mathbf{w} + \mathbf{a}' + \mathbf{b}')^2 + \Theta =$$

$$= 2\mathbf{a}'\mathbf{b}' + \Theta = 2 + \Theta,$$

where  $-2 < \Theta < 2$ , and so, this number cannot be divisible by 4. [Erdős et al. ibid.]

11. If the edges of a complete (n+1)-graph are k-colored, then, by 14.2a, we find a monochromatic triangle. Thus in the complete graph on  $\{1, \ldots, n+1\}$  where, following the hint, we color the edge (i,j) with color v, if  $|i-j| \in A_v$ , we find three points i, j, k such that x = |i-j|, y = |j-k|, z = |i-k| belong to the same class. Let, say, i < j < k, then

$$x+y=z$$

which proves the assertion. [I. Schur, Jahresb. Deutschen Math.-Ver. 25 (1916) 114.]

12. First solution. Suppose  $n_0(k, r-1)$  exists (certainly so does  $n_0(k, 1)$ ), then we prove that we can take

(1) 
$$n_0(k,r) = k^{n_0(k,r-1)} + n_0(k,r-1).$$

For suppose  $n \ge n_0(k, r)$  and  $\{1, \ldots, n\}$  is k-colored. Consider the colors of i,  $i+1, \ldots, i+n_0(k, r-1)-1$ , for each  $1 \le i \le k^{n_0(k, r-1)}+1$ . There are  $k^{n_0(k, r-1)}$  sequences of colors, which can occur here, thus we have two numbers i < j such that, for each  $0 \le v \le n_0(k, r-1)-1$ , the colors of i+v and j+v are the same.

By induction, we have a number  $a, i \le a \le i + n_0(k, r-1) - 1$ , and integers  $d_1, \ldots, d_{r-1}$  such that all numbers  $a + d_{i_1} + \ldots + d_{i_p}$ ,  $1 \le i_1 < \ldots < i_p \le r-1$  have the same color. Set  $d_r = j-i$ , then, by the choice of i and j, all numbers

$$a + d_{i_1} + \ldots + d_{i_{\nu}}, \ 1 \le i_1 < \ldots < i_{\nu} \le r$$

have the same color.

Second solution. First we prove: If  $A \subseteq \{1, \ldots, n\}$ ,  $|A| \ge 2$ , then there is a number  $d \ge 1$  such that the set

has 
$$|A_d| = \{a \in A\colon a+d \in A\}$$
 
$$|A_d| > \frac{|A|^2}{4n}.$$
 In fact 
$$\sum_{d=1}^{n-1} |A_d| = {|A| \choose 2} \ge \frac{|A|^2}{4},$$
 whence 
$$\max_{1 \le d \le n-1} |A_d| \ge \frac{|A|^2}{4(n-1)} > \frac{|A|^2}{4n}.$$

Now let A be the largest color-class in a k-coloration of  $\{1, \ldots, n\}$ . Then we have a  $d_1 \ge 1$  such that

$$|A_{d_1}|\!>\!rac{|A|^2}{4n}\!\ge\!rac{n}{4k^2}\!=\!rac{4n}{(4k)^2}\quad \left(\! ext{as }|A|\!\ge\!rac{n}{k}\!
ight).$$

Similarly, we have a  $d_2 \geq 1$  such that

$$egin{align} A_{d_1,d_2} &= |\{a \in A_{d_1} : a + d_2 \in A_{d_1}\}| = |\{a \in A : \ a + d_1 \in A, \ a + d_2 \in A, \ a + d_1 + d_2 \in A\}|> \ &> rac{n}{64k^4} = rac{4n}{(4k)^4} \end{aligned}$$

and repeating this r times we get numbers  $d_1, \ldots, d_r$  such that

$$|A_{d_1,\ldots,d_r}| = |\{a \in A : a + d_{i_1} + \ldots + d_{i_{\nu}} \in A,\}|$$

for each 
$$1 \le i_1 < \ldots < i_v \le r\} | > \frac{4n}{(4k)^{2^r}}$$
.

Any element of  $A_{d_1,\dots,d_r}$  satisfies our requirements. We need, however,  $|A_{d_1,\dots,d_i}| \geq 2$  for each  $1 \leq i \leq r-1$ . This is satisfied provided

$$n > \frac{1}{4} (4k)^{2^{r-1}}.$$

13. Let n = [ek!]. Assume the non-empty subsets of  $\{1, \ldots, n\}$  are k-colored with colors  $1, \ldots, k$ . Now color the pair  $\{i, j\}$ ,  $1 \le i < j \le n+1$  by the color of the interval  $\{i, \ldots, j-1\}$ . By 14.2a, there is a monochromatic triangle. If  $1 \le p < q < r \le n+1$  are the vertices of this triangle, then  $X = \{p, \ldots, q-1\}$ ,  $Y = \{q, \ldots, r-1\}$  and  $X \cup Y = \{p, \ldots, r-1\}$  have the same color.

14. (a) Suppose that N(k, t) exists for every k ( $t \ge 1$ ); we prove N(k, t + 1) exists for every k. Since N(k, 1) exists by the preceding problem, this will prove the assertion.

Let  $a = N(k^2, t)$  and  $N(k, t+1) = N(k^{2^a}, 1) + a$ . Suppose a coloration  $\alpha$  of the set of all subsets of an N(k, t+1)-element set S is given. Let  $T \subseteq S$ , |T| = a. First we define a coloration  $\alpha^*$  of the subsets of S - T, for which

$$\alpha^*(X) = (\alpha(X \cup Y) : Y \subseteq T).$$

This coloration uses  $k^{2^s}$  colors and so by the choice of N(k, t+1), there exist two disjoint sets  $A_1$ ,  $B_1 \subseteq S - T$  such that  $\alpha^*(A_1) = \alpha^*(B_1) = \alpha^*(A_1 \cup B_1)$ , i.e.

$$\alpha(A_1 \cup X) = \alpha(B_1 \cup X) = \alpha(A_1 \cup B_1 \cup X)$$

for each  $X \subseteq T$ . Now define a coloration  $\alpha^{**}$  of the subsets of T by

$$\alpha^{**}(X) = (\alpha(X), \ \alpha(X \cup A_1)).$$

This coloration uses  $k^2$  colors and hence by the definition of a, there are 2t disjoint sets  $A_2, B_2, \ldots, A_{t+1}, B_{t+1}$  such that for each sequence  $2 \le i_1 < \ldots < i_{\nu} \le t+1$ ,

$$\alpha^{**}(C_{i_1} \cup \ldots \cup C_{i_{\nu}})$$

is the same for any choice of  $C_j = A_j$  or  $B_j$  or  $A_j \cup B_j$ . Then  $A_1, B_1, \ldots, A_{t+1}, B_{t+1}$  satisfy the requirements of our problem.

(b) We use induction on r; for r = 2 the preceding problem contains the solution.

Assume b = n(k, r) exists; we define n(k, r + 1) = N(k, b) and prove it has the desired property. Let  $\alpha$  be any k-coloration of all subsets of an n(k, r + 1)-element set S. By (a), there exist 2b disjoint sets  $A_1, B_1, \ldots, A_b, B_b$  such that for each  $1 \leq i_1 < \ldots < i_v \leq b$ , all sets of the form

$$C_{i_1} \cup \ldots \cup C_{i_{\nu}} \quad (C_j = A_j \text{ or } B_j \text{ or } A_j \cup B_j)$$

have the same color. Define a k-coloration  $\alpha^*$  of the set  $\{1, \ldots, b\}$  by

$$\alpha^*(\{i_1,\ldots,i_v\}) = \alpha(A_{i_1}\cup\ldots\cup A_{i_v}).$$

By the choice of b, there are disjoint subsets  $I_1, \ldots, I_r \subseteq \{1, \ldots, b\}$  such that the color of any non-empty union of them is the same, say red. Now set

$$X_j = \bigcup_{i \in I_j} A_i, \ Y_j = \bigcup_{i \in I_j} B_i.$$

Then  $X_1, \ldots, X_r, Y_1, \ldots, Y_r$  are disjoint and any non-empty union of them is red. In fact, let  $V, W \subseteq \{1, \ldots, r\}$  and set  $I = \bigcup\limits_{j \in V \cap W} I_j, I' = \bigcup\limits_{j \in V - W} I_j,$   $I'' = \bigcup\limits_{j \in V - W} I_j$ . Then we have

$$lphaig((igcup_{j\in V}X_j)\cup(igcup_{i\in W}Y_j)ig)=lphaig(\{igcup_{i\in I}(A_i\cup B_i)\}\cup\{igcup_{i\in I'}A_i\}\cup\{igcup_{i\in I''}B_i\}ig).$$

By the choice of  $A_i$  and  $B_i$  this color is the same as

$$\alpha(\bigcup_{i\in I\cup I'\cup I''}A_i)=\alpha^*(I\cup I'\cup I'').$$

which is red.

Since the number of sets  $X_1, \ldots, X_r, Y_1, \ldots, Y_r$  is  $2r \ge r+1$ , this proves the assertion. [R.L. Graham-B.L. Rothschild, Bull. Amer. Math. Soc. 75 (1969) 418-422; also see R.L. Graham, K. Leeb, B.L. Rothschild, Adv. in Math. 8 (1972) 417-433.]

15. Let n = n(k, r), where n(k, r) is the function in the preceding problem. Suppose  $\alpha$  is a k-coloration of  $\{1, \ldots, n\}$ . Then define a k-coloration  $\alpha^*$  of the subsets of  $S = \{1, \ldots, n\}$  by

$$\alpha^*(X) = \alpha(|X|) \ (X \subseteq S).$$

By 14.14, there are disjoint sets  $X_1, \ldots, X_r \subseteq S$  such that all non-empty unions of any of them have the same color. Then the choice  $d_i = |X_i|$  is an appropriate one [J. Folkman, J. Sanders; see ibid.].

16. (a) Take  $n_1(k, m) = k$ ; set  $S = \{x_1, \ldots, x_k\}$ . Consider the *m*-collections  $m \cdot \emptyset + (m-1) \cdot \{x_1, \ldots, x_k\}$ ;  $m\{x_1\} + (m-1) \cdot \{x_2, \ldots, x_k\}$ ,  $\ldots, m \cdot \{x_1, \ldots, x_k\} + (m-1) \emptyset$ .

Some two of these, say

$$m \cdot \{x_1, \ldots, x_i\} + (m-1) \cdot \{x_{i+1}, \ldots, x_k\}$$

and

$$m \cdot \{x_1, \ldots, x_i\} + (m-1) \cdot \{x_{i+1}, \ldots, x_k\}$$

have the same color  $(0 \le i < j \le k)$ . Let  $X = \{x_{i+1}, \ldots, x_j\}$ ,  $\mathcal{B} = m \cdot \{x_1, \ldots, x_i\} + (m-1) \cdot | \{x_{j+1}, \ldots, x_k\}$ , then the requirement is satisfied.

(b) We prove the existence of  $n_r(k, m)$  by induction on r. Set  $a = k^{m^{r-1}}$ ,  $k' = k^{m^a}$ ,  $n_{r+1}(k, m) = n_r(k', m) + a$ . We prove that this choice is correct. Let  $T \subset S$ , |T'| = a. Define a coloration of  $m^{S-T}$  by

$$\alpha^*(\mathcal{A}) = (\alpha(\mathcal{A} + \mathfrak{D}); \mathfrak{D} \in m^T).$$

Here we use  $m^a$ -tuples of the original set of colors, so we use not more than  $k^{m^a} = k'$  colors. Hence by the choice of  $n_{r+1}(k, m)$ , there exists r disjoint subsets  $X_1, \ldots, X_r$  of S - T and an m-collection  $\mathfrak{B}_1 \in m^{S - X_1 - \cdots - X_r}$  such that

$$\alpha^* \left( \sum_{i=1}^r a_i X_i + \mathcal{B}_1 \right) = \alpha^* \left( \sum_{i=1}^r \min \left( a_i, m-1 \right) X_i + \mathcal{B}_1 \right)$$

holds for every  $0 \le a_i \le m$ ; i.e. we have

(1) 
$$\alpha \left( \mathfrak{D} + \sum_{i=1}^{r} a_i X_i + \mathfrak{B}_1 \right) = \alpha \left( \mathfrak{D} + \sum_{i=1}^{r} \min \left( a_i, m - 1 \right) X_i + \mathfrak{B}_1 \right)$$

for every  $0 \le a_i \le m$  and  $\mathfrak{D} \in m^T$ . Now define a coloration  $\alpha^{**}$  of  $m^T$  by

$$\alpha^{**}(\mathfrak{D}) = \left(\alpha \left(\mathfrak{D} + \sum_{i=1}^r a_i X_i + \mathfrak{B}_1\right) : 0 \leq a_i \leq m-1\right).$$

In this way an m'-tuple of colors is a new color, so  $\alpha^{**}$  uses at most  $k^{m'} = a$  colors. Since |T| = a, we find by (a) a set  $X_{r+1} \subseteq T$  and an m-collection  $\mathcal{B}_2 \in m^{T-X_{r+1}}$  such that

$$\alpha^{**}(mX_{r+1} + \mathcal{B}_2) = \alpha^{**}((m-1)X_{r+1} + \mathcal{B}_2),$$

i.e.

(2) 
$$\alpha \left( mX_{r+1} + \sum_{i=1}^{r} a_i X_i + \mathcal{B}_1 + \mathcal{B}_2 \right) = \alpha \left( (m-1) X_{r+1} + \sum_{i=1}^{r} a_i X_i + \mathcal{B}_1 + \mathcal{B}_2 \right)$$

for any integers  $0 \le a_i \le m-1$ . Now let  $\mathfrak{B} = \mathfrak{B}_1 + \mathfrak{B}_2$  (this is an *m*-collection, as  $\mathfrak{B}_1$ ,  $\mathfrak{B}_2$  are disjoint). Then  $X_1, \ldots, X_{r+1}$ ,  $\mathfrak{B}$  satisfy the requirement. Clearly  $X_1, \ldots, X_{r+1}$  are disjoint from each other and  $\mathfrak{B}$ . Let

$$\mathcal{A} = \sum_{i=1}^{r+1} a_i X_i + \mathcal{B},$$

then

$$\alpha(\mathcal{R}) = \alpha \left( \sum_{i=1}^{r} \min \left( a_{i}, \, m-1 \right) X_{i} + a_{r+1} X_{r+1} + \mathfrak{B} \right)$$

by (1) and

$$\begin{split} &\alpha\left[a_{r+1}X_{r+1} + \sum_{i=1}^{r}\min\left(a_{i},\,m-1\right)X_{i} + \mathcal{B}\right] = \\ &= \alpha\left(\min\left(a_{r+1},\,m-1\right)X_{r+1} + \sum_{i=1}^{r}\min\left(a_{i},\,m-1\right)X_{i} + \mathcal{B}\right) \end{split}$$

by (2). This completes the proof. [ibid.]

17. We use induction on m. For m = 1 the answer follows by 14.14b (we may even take  $\mathfrak{B} = \emptyset$ ). Suppose N(k, r, m) exists and define a = N(k, r, m),

$$N(k, r, m + 1) = n_a(k, m + 1),$$

where  $n_a(k, m + 1)$  is defined as in 14.16b. Then if  $|S| \ge N(k, r, m + 1)$  we find disjoint sets  $Y_1, \ldots, Y_a \subseteq S$  and an (m + 1)-collection  $\mathcal{B}_1$  of points of  $S - Y_1 - \ldots - Y_a$  such that

(1) 
$$\alpha \left( \sum_{i=1}^{a} a_i Y_i + \mathfrak{B}_1 \right) = \alpha \left( \sum_{i=1}^{a} \min (a_i, m) Y_i + \mathfrak{B}_1 \right)$$

holds for every  $0 \le a_i \le m+1$ . For each  $\mathfrak{J} \in m^{\{1,\dots,a\}}$ , define

$$\alpha^*(\mathfrak{F}) = \alpha \left( \sum_{i=1}^a a_i Y_i + \mathfrak{B}_1 \right),$$

where  $a_i$  is the number of times i occurs in  $\mathfrak{J}$ . Now by the choice of a, there are disjoint subsets  $V_1, \ldots, V_r \subseteq \{1, \ldots, a\}$  and an m-collection  $\mathfrak{K}$  of elements of  $\{1, \ldots, a\} - V_1 - \ldots - V_r$  such that

$$\alpha^* \left( \sum_{j=1}^r b_j V_j + \Re \right)$$

is the same color, say red, for every choice of  $0 \le b_i \le m$ . Define

$$X_j = \bigcup_{i \in V_i} Y_i, \ \mathcal{B} = \mathcal{B}_1 + \sum_{i=1}^a a_i Y_i,$$

where  $a_i$  is the number of times i occurs in  $\mathcal{K}$ . Clearly  $\sum_{i=1}^{a} a_i Y_i$  is disjoint from  $X_j$  and  $\mathcal{B}_1$  so  $X_1, \ldots, X_r$ ,  $\mathcal{B}$  are mutually disjoint. Moreover, if  $0 \leq a_i \leq m+1$ , then

$$\alpha\left(\sum_{i=1}^{r}a_{i}X_{i}+\mathfrak{B}\right)=\alpha\left(\sum_{i=1}^{r}\min\left(a_{i},\,m\right)X_{i}+\mathfrak{B}\right)$$

by (1); moreover, this is equal to

$$\alpha^* \left( \sum_{i=1}^r \min\left(a_i, m\right) V_i + \mathfrak{K} \right),$$

which is red. This proves that N(k, r, m + 1) has the desired property [A. Hales-R.I. Jewett, *Trans. Amer. Math. Soc.* 106 (1963) 222-229].

18. Let w = (m-1) N(k, 1, m-1), |S| = N(k, 1, m-1). Let  $\alpha$  be any k-coloration of  $\{0, \ldots, w\}$ . We define a k-coloration of  $(m-1)^S$  by setting

$$\alpha^*(\mathcal{A}) = \alpha(|\mathcal{A}|).$$

Since  $0 \le |\mathcal{R}| \le (m-1) |S| = w$ , this is well defined. By the choice of |S| there exists a set  $X \subseteq S$  and an (m-1)-collection  $\mathcal{B}$  of points of S-X such that

$$\alpha^*(\mathfrak{B}) = \alpha^*(\mathfrak{B} + X) = \alpha^*(\mathfrak{B} + 2X) = \ldots = \alpha^*(\mathfrak{B} + (m-1)X),$$

i.e.

$$\alpha(|\mathfrak{B}|) = \alpha(|\mathfrak{B}| + |X|) = \ldots = \alpha(|\mathfrak{B}| + (m-1)|X|).$$

Thus,  $(|\mathfrak{B}|, |\mathfrak{B}| + |X|, \ldots, |\mathfrak{B}| + (m-1)|X|)$  is a monochromatic arithmetic progression [cf. R.L. Graham-B.L. Rothschild, *Proc. Amer. Math. Soc.* 42 (1974) 385-386].

19. Suppose  $\{1, \ldots, N\}$  has a k-coloration  $\alpha_N$  for which no subset has property P. Consider the colors  $\alpha_N(1)$ . Since we use only k colors, there will be one occurring infinitely many times, say

$$\alpha_{N_1^1}(1) = \alpha_{N_2^1}(1) = \dots$$

Denote this common color by  $\alpha(1)$ . Now consider the colors  $\alpha_{N_1^1}(2)$ . Again we will find an infinite subsequence  $N_1^2, N_2^2, \ldots$  of  $N_1^1, N_2^1, \ldots$  such that

$$\alpha_{N_{\epsilon}^{2}}(2) = \alpha_{N_{\epsilon}^{2}}(2) = \ldots = \alpha(2).$$

Now suppose we have defined colors  $\alpha(1), \ldots, \alpha(n)$ , so that still infinitely many indices  $N_1^n, N_2^n, \ldots$  exist with

$$\alpha_{N_i^n}(m) = \alpha(m)$$
 for  $1 \leq m \leq n$  and  $i = 1, 2, \ldots$ 

Then among the colors  $\alpha_{N^n}(n+1)$ , one color occurs infinitely often; denote this by  $\alpha(n+1)$ .

We claim the coloration  $\alpha$  of all natural numbers is such that no finite subset has property P. Assume indirectly that a subset  $S \subseteq \{1, \ldots, n\}$  has property P. Then

$$\alpha_{N^n}(m) = \alpha(m)$$

for each  $m \in S$  and thus, S has property P in the coloration  $\alpha_{N_1^n}$  too, a contradiction.

20. (a) Write each natural number m in the form  $m = 5^{\alpha_m} (5\beta_m + \gamma_m)$ ,  $1 \le \gamma_m \le 4$  and color m with color  $\gamma_m$ . Thus the set of all natural numbers is

4-colored. We claim that x+y=3z has no monochromatic solution. Suppose it has one in the  $\gamma^{\rm th}$  class, then

$$5^{\alpha_z}(5\beta_x+\gamma)+5^{\alpha_y}(5\beta_y+\gamma)=3\cdot 5^{\alpha_z}(5\beta_z+\gamma).$$

Divide here by  $5^{\alpha}$ , where  $\alpha = \min(\alpha_x, \alpha_y, \alpha_z)$ , and consider the equation mod 5. Then we get

$$\varepsilon_{x}\gamma + \varepsilon_{y}\gamma \equiv 3\varepsilon_{z}\gamma \pmod{5},$$

where  $\varepsilon_x = 1$ , if  $\alpha = \alpha_x$  and = 0, if  $\alpha < \alpha_x$  (similarly for  $\varepsilon_y$  and  $\varepsilon_z$ ). Then  $\varepsilon_x + \varepsilon_y \equiv 3\varepsilon_z \pmod{5}$ ,

which obviously, has a unique solution in  $\varepsilon_x$ ,  $\varepsilon_y$ ,  $\varepsilon_z = 0$ , 1:  $\varepsilon_x = \varepsilon_y = \varepsilon_z = 0$ . But this is a contradiction as  $\alpha$  must be equal to one of  $\alpha_x$ ,  $\alpha_y$ ,  $\alpha_z$ .

(b) We prove by induction on k that x + 2y = z has a monochromatic solution. For k = 1, this is obvious. Suppose it holds for k - 1 colors.

It follows as in 14.19 that there is a natural number N with the property that if we (k-1)-color the numbers  $\{1, \ldots, N\}$ , then one of the color classes will contain a solution of x+2y=z.

Now suppose that the natural numbers are k-colored. By van der Waerden's theorem 14.18, one of the classes, the "red" class say, contains an arithmetic progression

$$\{a, a+d, a+2d, \ldots, a+2Nd\}.$$

Let us seek a monochromatic solution of (b) in the form  $x=a, y=\varkappa d, z=a+2\varkappa d$ . Now x and z are red for each  $1\leq \varkappa \leq N$ , so if  $\varkappa d$  is red for any  $1\leq \varkappa \leq N$  we are finished. Suppose this is not the case, then we have the numbers  $d, 2d, \ldots, Nd$  colored with k-1 colors. Define a new coloration of  $\{1,\ldots,N\}$  by giving  $\varkappa$  the color of  $\varkappa d$ . Here we use k-1 colors, hence by the definition of N, there are  $x_1,y_1,z_1$  having the same color and satisfying  $x_1+2y_1=z$ . Now  $(dx_1)+2(dy_1)=(dz_1)$  and  $dx_1, dy_1, dz_1$  have the same color in the original k-coloration, which proves the assertion.

**21.** I. (i)  $\Rightarrow$  (ii). Consider a (0, 1)-solution of (\*); we may assume it looks like

$$\sum_{i=1}^{r} a_i \cdot 1 + \sum_{i=r+1}^{m} a_i \cdot 0 = 0.$$

Now since the equation

$$(**) \hspace{3.1em} a_1 \cdot x - a_1 \cdot y + \left(\sum_{i=r+1}^m a_i\right) \cdot z = 0$$

also satisfies (i) and any monochromatic solution of (\*\*) yields a monochromatic solution of (\*) with  $x_1 = x$ ,  $x_2 = \ldots = x_r = y$ ,  $x_{r+1} = \ldots = x_m = z$ , we may assume that (\*) looks like

$$(***) Ax - Ay + Bz = 0.$$

We use induction on k. The case k=1 is trivial, so suppose k>1 and (ii) holds for k-1 colors. As in 14.19, we can find a natural number N such that if  $\{1,\ldots,N\}$  is (k-1)-colored, (\*\*\*) has a solution in one of the color classes.

Now suppose that all natural numbers are k-colored. By van der Waerden's theorem, we find a monochromatic (say, red) arithmetic progression

$$a, a + d, \ldots, a + BNd$$
.

Then from

$$A(a + B \times d) - [Aa + B(\times Ad) = 0]$$

it follows that if  $\varkappa Ad$  is red for some  $1 \le \varkappa \le N$ , we are done. So suppose  $\varkappa Ad$  is never red; then by the choice of N, we find numbers  $x_1, y_1, z_1$  such that  $x_1Ad, y_1Ad, z_1Ad$  have the same color and then

$$A(x_1Ad) - A(y_1Ad)[ + B(z_1Ad) = 0.$$

II. (ii)  $\Rightarrow$  (i). Let p be a very large prime and write  $t = p^{\alpha_t}(\beta_t p + \gamma_t)$  ( $1 \le \gamma_t \le p-1$ ). Let us color t with  $\gamma_t$ . Then by (ii), (\*) has a monochromatic solution; i.e. there are non-negative integers  $\alpha_{\mathbb{R}}^{(1)}, \ldots, \alpha_{\mathbb{R}}^{(m)}, \beta^{(1)}, \ldots, \beta^{(m)}, \gamma$  ( $1 \le \gamma \le p-1$ ) such that

(1) 
$$\sum_{i=1}^{m} a_i p_{i}^{\alpha^{(i)}}(\beta^{(i)} + [\gamma]) = 0.$$

Set  $\alpha = \min(\alpha^{(1)}, \ldots, \alpha_{\underline{n}}^{(m)})$  and divide here by  $p^{\alpha}$ , then consider (1) mod p. We get

$$\sum_{i=1}^{m} a_i \varepsilon_i \gamma \equiv 0 \text{ (mod } p),$$

where  $\varepsilon_i = 1$  or 0 according as  $\alpha_i = \alpha [\text{or}[\alpha_i > \alpha]]$ . Hence

$$\sum_{i=1}^{m} a_i \varepsilon_i \equiv 0 \pmod{p}.$$

Since p is large (e.g. it suffices to take  $|p| > |a_1| + \ldots + |a_m|$ ), we get

$$\sum_{i=1}^m a_i \varepsilon_i = 0.$$

Since  $\alpha = \alpha_i$  for at least one  $i, \varepsilon_1, \ldots, \varepsilon_m$  is a non-trivial (0, 1) solution of (\*). This proves (i) [R. Rado, *Math. Zeitschr.* 36 (1933) 424–480].

22. We specify f(k) later. Let  $1, \ldots, k$  be k colors and color the integers with them at random, independently of each other and giving each integer each color with probability 1/k. Let  $A_j$  denote the event that S+j does not meet one of the colors. Then

$$P(A_j) \leq \sum_{i=1}^k P(S+j \text{ does not meet color } i) = k \left(1 - \frac{1}{k}\right)^{f(k)}$$
.

Now form a graph G on Z, the set of integers, by connecting  $j_1$  to  $j_2$ , if  $(S + j_1) \cap (S + j_2) \neq \emptyset$ . Then  $A_j$  is independent of

$$\{A_{\nu}:(j,\nu)\notin E(G)\}.$$

Finally, observe that the degree of each point in G is

$$\begin{aligned} d &= |\{v \colon (0, v) \in E(G)\}| = |\{v \colon (S + v) \cap S \neq 0\}| = \\ &= |\{s - s' \colon s, s' \in S\}| \le f(k)^2. \end{aligned}$$

Thus, if

(1) 
$$k \left(1 - \left| \frac{1}{k} \right|^{f(k)} \le \frac{1}{4f(k)^2},\right.$$

we have

$$P(A_j) \leq \frac{1}{4d}$$

and thus by 2.18,

$$P(\bar{A}_0 \dots \bar{A}_N) > 0$$

for any N. Of course, we cannot assert that  $P\left(\prod_{j=0}^{\infty} \bar{A}_{j}\right) > 0$ , but the existence of a coloration such that all translated copies of S meet all colors still follows. One possibility would be to apply compactness theorems (cf. 14.19) but here

we can construct this coloration directly.

Let r be the diameter of S ( $r = \max \check{S} - \min S$ ) and  $N > k^{r+1}$ . Consider a k-coloration for which  $A_0 \dots A_N$  holds, i.e.  $S+1, \dots, S+N$  meet all colors. Since  $N > k^{r+1}$ , there are two segments  $\{v, v+1, \dots, v+r\}$  and  $\{\mu, \mu+1, \dots, \mu+r\}$  ( $v < \mu$ ), which are colored similarly. Now, if we keep the coloration of  $v, v+1, \dots, \mu-1$  but extend this in both directions with period  $\mu-v$  then, as is easily verified, S+j will meet all colors for every integer j.

It only remains to find an f(k) for which (1) holds. Certainly  $f(k) = [ck \log k]$ 

will be appropriate for large enough c, because

$$\left(1-\frac{|1|}{k}\right)^{f(k)} \approx \left(1-\frac{1}{k}\right)^{ck\log k} < e^{-c\log k} = \frac{1}{k^c},$$

while

 $\frac{1}{4f(k)^2} = \frac{1}{4c^2k^2\log^2 k} > \frac{1}{k^{c-1}}.$ 

23. (a) Let us denote the elements of GF(q) by  $0, 1, \ldots, q-1$  (the operations between them will play no role.) Each element of the n-dimensional affine space over GF(q) can be represented as a vector  $(a_1, \ldots, a_n)$ ,  $0 \le a_i \le q-1$ , which in turn can be regarded as a (q-1)-collection of elements of  $\{1, \ldots, n\}$ . Thus if these are k-colored and n > N(k, r, q-1), where N(k, r, q-1) is defined as in 14.17, then there will be disjoint non-empty sets  $V_1, \ldots, V_r \subseteq \{1, \ldots, n\}$  and a (q-1)-collection  $\mathcal{B}$  of points of  $\{1, \ldots, n\} - V_1 - \ldots - V_r$  such that all collections  $\sum_{i=1}^r a_i V_i + \mathcal{B}$   $(0 \le a_i \le q-1)$  have the same color; let, say,

$$V_i = \{k_i + 1, \dots, k_{i+1}\}\ (i = 1, \dots, r; 0 = k_1 < k_2 < \dots < k_{r+1} = m)$$

and let j occur in  $\mathfrak{B}$   $b_j$  times (clearly  $b_j = 0$  for  $1 \leq j \leq m$ ), then all vectors the form

$$(a_1, \ldots, a_1, a_2, \ldots, a_2, a_3, \ldots, a_r, b_{m+1}, \ldots, b_n)$$

have the same color. But these vectors form an r-dimensional subspace.

(b) We prove the (formally) more general assertion formulated in the hint by simultaneous induction on  $r_1, \ldots, r_k$  and k. For k = 1 we clearly can take

(1) 
$$\overline{N}(1; r_1; q-1) = r_1.$$

On the other hand, if  $r_i = 0$  for some  $1 \le i \le k$ , say  $r_k = 0$ , then

(2) 
$$\overline{N}(k; r_1, \dots, r_{k-1}, 0; q-1) = \overline{N}(k-1; r_1, \dots, r_{k-1}, q-1)$$

will be an appropriate choice. For if any point has color k we are done; if color k does not occur we have the case with k-1 colors. So we may assume  $r_i > 0$  for  $1 \le i \le k$ . Then let

(3) 
$$\overline{N}(k; r_1, \ldots, r_k; q-1) = N(k; m, q-1),$$

where

$$m = 1 + \max_{1 \le i \le k} \overline{N}(k; r_1, \dots, r_{i-1}, r_i - 1, r_{i+1}, \dots, r_k; q - 1)$$

and N(k, m, q-1) is defined as in part (a). To prove that (3) is an appropriate choice let  $n \geq N(k; r_1, \ldots, r_k; q-1)$  and suppose that the points of the n-dimensional projective space over GF(q) are k-colored. Let  $\Sigma_0$  be a hyperplane (the "infinitely distant" hyperplane). Remove  $\Sigma_0$ , then the remaining affine space contains a monochromatic subspace  $\Phi$  of dimension m by (3) and (a). Let  $\Phi$  have, e.g. color 1. Let  $\Phi_1$  be the subspace of the projective space spanned by  $\Phi$  and put  $\Psi = \Phi_1 \cap \Sigma$ . Then  $\Psi$  is an (m-1)-dimensional projective space, which is k-colored. Thus by the choice of m,  $\Psi$  has a monochromatic subspace  $\Delta$  of dimension  $r_1 - 1$  in the first color or of dimension  $r_i$  in the i-th color  $1 \leq i \leq k$ . In the latter case we are done. If  $\Delta$  is  $1 \leq i \leq k$  is that i-dimensional and has color 1, we can take any i-dimensional subspace i-dimensional and i-dimensional subspace with color 1 [R.L. Graham—B.L. Rothschild, i-dimensional subspace i-dimensional subspace with color 1 [R.L. Graham—B.L. Rothschild, i-dimensional subspace i-dimensional subspac

**24.** First solution: Let  $n \ge R_{k+1}^{r-1}(r,\ldots,r)$ , where R is the Ramsey number as in 14.3. Let  $S = \{1,\ldots,n\}$ , and denote by V the set of all (r-1)-element subsets of S. Define a hypergraph on V by letting the r (r-1)-tuples of each r-subset of S form an edge of H. Then clearly H is r-uniform, any two edges of H have at most one point in common, and the fact that H is not k-colorable is just the assertion of Ramsey's theorem.

Second solution: Let  $n \ge N(k; 1, q - 1)$ , where q is a prime power. Let V(H) be the set of points of the n-dimensional affine space over GF(q) and let E(H) consist of the lines of this affine space. Then any two edges of H have at most one point in common and it is q-uniform. The assertion of 14.23a is just that H is not k-colorable. This settles the problem for the case when r is a prime power.

Now let r be arbitrary and select a prime power  $q \ge r$ . Carry out the above construction for q and then select an arbitrary r-subset of each edge of the resulting hypergraph. These r-element sets constitute a hypergraph with the desired properties.

25. For each  $a_i$ , let t(i) denote the maximum length of subsequences

$$a_i \leq a_{j_1} \leq \ldots \leq a_{j_{t-1}}$$

with

$$i < j_1 < \ldots < j_{t-1}$$

Suppose that max  $t(i) \leq k$  (otherwise we are done). Then there is a  $\tau$  such that  $t(i) = \tau$  holds for at least k + 1 indices i. Let

$$t(i_1) = \ldots = t(i_{k+1}) = \tau, \ i_1 < \ldots < i_{k+1}.$$

Then we claim that

$$a_{i_1} > a_{i_2} > \ldots > a_{i_{k+1}}.$$

Suppose indirectly that

$$a_{i_{\nu}} \leq a_{i_{\mu}} \ (\nu < \mu).$$

Then, since  $t(a_{i\mu}) = \tau$ , we have a sequence

$$i_{\mu} < j_1 < \ldots < j_{\tau-1}$$

with

$$a_{i\mu} \leq a_{j_1} \leq \ldots \leq a_{j_{\tau-1}}.$$

Then

$$a_{i_{\nu}} \leq a_{i_{\mu}} \leq a_{j_1} \leq \ldots \leq a_{j_{\tau-1}},$$

contradicting  $t(a_{i_v}) = \tau$ .

*Remark*: Define i < j if i < j and  $a_i \le a_j$ . Then we have a partial order on the set  $\{1, \ldots, n^2 + 1\}$ . Thus the result would follow from 9.32.

**26.** (a) Consider t(x) as in the hint. Observe that  $\max t(x) = t(1)$ . We use induction on n.

First we prove the assertion of the hint. Assume indirectly that  $f(x) > 2^k$  and  $t(x) \ge t(1) - k$ . Then  $x > 2^k$  and k < n - 1. Let

$$x = a_1 < \ldots < a_{t(x)},$$

$$f(a_1) \leq \ldots \leq f(a_{t(x)}).$$

By the induction hypothesis, there exist

$$1 \le b_0 < b_1 < \ldots < b_k \le 2^k$$

with

$$f(b_0) \leq f(b_1) \leq \ldots \leq f(b_k).$$

Then

$$f(b_0) \leq f(b_1) \leq \ldots \leq f(b_k) \leq b_k \leq 2^k < f(a_1) \leq \ldots \leq f(a_{t(x)}),$$

whence

$$t(1) \ge t(x) + k + 1,$$

a contradiction.

Now observe that if t(x) = t(y), then  $f(x) \neq f(y)$  (otherwise, the chain starting with y (x < y) could be extended to x) and thus,

$$|\{x: t(x) = t(1) - k\}| \le 2^k \quad (0 \le k \le t(1) - 1).$$

Hence,

$$2^{n-1} = \sum_{k=0}^{t(1)-1} |\{x : t(x) = t(1) - k\}| \le \sum_{k=0}^{t(1)-1} 2^k = 2^{t(1)} - 1.$$

Hence

$$t(1) \geq n$$
,

which was to be proved.

## (b) Let

$$\varphi_0(2^k+l)=2^k-l \ (k=0,\ldots,n-2;\ 0\leq l\leq 2^k-1).$$

Then  $\varphi_0(i) \leq i$   $(i = 1, \ldots, 2^{n-1} - 1)$  and  $\varphi_0$  consists of n-1 monotone decreasing intervals; any sequence

$$1 \leq a_1 < \ldots < a_t \leq 2^{n-1} - 1$$

with

$$f(a_1) \leq \ldots \leq f(a_t)$$

contains at most one number from each of these intervals. Hence  $t \leq n-1$  [E. Harzheim, *Publ. Math. Debrecen* 14 (1967) 45–51].

**27.** (a) Let  $\eta(x, y)$  be the maximum length of a path  $(x, y = y_1, y_2, \dots, y_{\eta})$  with

$$f(x, y_1) \le f(y_1, y_2) \le \ldots \le f(y_{\eta-1}, y_{\eta})$$

and also, let  $\varphi(x, y)$  be the maximum length of paths  $(x, y = y_1, y_2, \dots, y_{\varphi})$  with

$$f(x, y_1) > f(y_1, y_2) > \ldots > f(y_{\varphi-1}, y_{\varphi}).$$

Observe that

(\*) for any two (adjacent) edges (x, y) and (y, z), either  $\eta(x, y) > \eta(y, z)$  or  $\varphi(x, y) > \varphi(y, z)$ . In fact, let, e.g.  $f(x, y) \le f(y, z)$  and  $(y, z = z_1, z_2, \ldots, z_{\eta(y,z)})$  be a path with

$$f(y, z_1) \le f(z_1, z_2) \le \ldots \le f(z_{\eta(y,z)-1}, z_{\eta(y,z)}).$$

Then the path  $(x, y, z_1, \ldots, z_{\eta(y,z)})$  shows that  $\eta(x, y) \ge \eta(y, z_1) + 1$ . For  $w \in V(T_n)$ , let

$$P_{w} = \{ (\eta(w, x), \varphi(w, x)) : (w, x) \in E(T_{n}) \}$$

and let  $\overline{P}_{w}$  consist of all maximal elements of  $P_{w}$  (with respect to the partial ordering  $(\eta, \varphi) \leq (\eta', \varphi')$  iff  $\eta \leq \eta'$  and  $\varphi \leq \varphi'$ ). Then we claim that

$$(**) \qquad \overline{P_w} \neq \overline{P_v}, \text{ if } w \neq v.$$

For suppose, e.g.  $(v, w) \in E(T_n)$ . By the definition of  $\overline{P_v}$ , there is a pair  $(\eta_0, \varphi_0) \in \overline{P_v} = \overline{P_w}$  with  $\eta(v, w) \leq \eta_0$ ,  $\varphi(v, w) \leq \varphi_0$ . Let  $\eta_0 = \eta(w, x)$ ,  $\varphi_0 = \varphi(w, x)$ . Then  $\eta(v, w) \leq \eta(w, x)$ ,  $\varphi(v, w) \leq \varphi(w, x)$ , which contradicts (\*).

All that remains is to count how many sets come into consideration as sets  $\overline{P_w}$ . If  $\overline{P_w} = \{(\eta_1, \varphi_1), \dots, (\eta_k, \varphi_k)\}$ , then  $\eta_1, \dots, \eta_k$  are distinct (so are  $\varphi_1, \dots, \varphi_k$ ) and if, say,  $\eta_1 < \dots < \eta_k$ , then  $\varphi_1 > \dots > \underline{\varphi_k}$ . Hence, the two sets  $\{\eta_1, \dots, \eta_k\}$  and  $\{\varphi_1, \dots, \varphi_k\}$  uniquely determine  $\overline{P_w}$ . For a fixed k, there are at most  $\binom{p'}{k}$  ways to choose  $\eta_1, \dots, \eta_k$  and  $\binom{q'}{k}$  ways to choose  $\varphi_1, \dots, \varphi_k$ , where  $p' = \max \eta(x, y)$  and  $q' = \max \varphi(x, y)$ . Thus the number of distinct sets  $\overline{P_w}$  is

$$\leq \sum_{k=0}^{p'} \binom{p'}{k} \binom{q'}{k} = \sum_{k=0}^{p'} \binom{p'}{k} \binom{q'}{q'-k} = \binom{p'+q'}{p'}.$$

Thus

$$\binom{p'+q'}{p'} \ge n > \binom{p+q-2}{p-1},$$

whence either  $p' \geq p$  or  $q' \geq q_{\bullet}$ . This proves the assertion.

(b) With a slight change in notation, we are going to construct a transitive tournament  $T_n$  on  $n = \begin{pmatrix} p+q \\ p \end{pmatrix}$  points and a function

$$f: E(T_n) \to \mathbf{Z}$$

(which will be an injection) such that the maximum length of a monotone increasing path is p and the maximum length of a monotone decreasing path is q. The case p = 1 or q = 1 is trivial; so suppose  $p, q \ge 2$ .

We have

$$\binom{p+q}{p} = \left[\binom{p+q-1}{p} + \binom{p+q-1}{p-1}\right].$$

By induction on p, q, we construct a transitive tournament  $T_m$  on  $m=\begin{pmatrix} p+q-1\\ p \end{pmatrix}$  points and a tournament  $T_l$  on  $l=\begin{pmatrix} p+q-1\\ p-1 \end{pmatrix}$  points, together with two functions  $g:E(T_m)\to \mathbb{Z}$ ,  $h\colon E(T_l)\to \mathbb{Z}$  such that the maximum length of monotone increasing [decreasing] paths in  $T_m$  and  $T_l$  are p and p-1 [q-1 and q], respectively. Let  $V(T_n)=V(T_m)\cup V(T_l)$ ,

$$E(T_n) = E(T_m) \cup E(T_l) \cup (V(T_m) \times V(T_l))$$

and

$$f(x,y) = \begin{cases} g(x, y) & \text{if} \quad (x, y) \in E(T_m), \\ h(x, y) & \text{if} \quad (x, y) \in E(T_l), \\ -M & \text{otherwise,} \end{cases}$$

where  $M > \max_{(x,y)} (|g(x,y)|, |h(x,y)|)$ . It is easy to verify that  $T_n$  and f have the desired properties [V. Chvátal–J. Komlós, Canad. Math. Bull. 14 (1971) 151–157].

**28.** (a) If  $0 \le i \le m-1$ , let  $x_{i+1}$  be any element of

$$S - \{f(S - \{x_1, \ldots, x_{\nu}\}) : 0 \leq \nu \leq i\}$$

(which set is clearly non-empty), and let

$$T = S - \{x_1, \ldots, x_m\}.$$

Then by the choice of the  $x_i$ 's,

$$f(S - \{x_1, \ldots, x_{\nu}\}) \in T \quad (\nu = 0, \ldots, m-1)$$

and this holds trivially for v = m too.

(b) We select  $x_{2^n}, x_{2^{n-1}}, \ldots$  recursively. If  $x_{2^n}, \ldots, x_{\mu+1}$  are already selected, then let  $x_{\mu}$  be an element of  $S = \{x_{2^n}, \ldots, x_{\mu+1}\}$ , which occurs the least number of times in the form  $f(S = \{x_{2^n}, \ldots, x_{\nu+1}\})$ ,  $\mu \leq \nu \leq 2^n$ .

To handle this procedure, let  $\varphi(m)$  denote the number of indices  $v \ge m+1$ 

such that

$$f({x_1, \ldots, x_{\nu}}) \in {x_1, \ldots, x_m}.$$

Then we have

(1) 
$$\varphi(m) - \varphi(m-1) \le \frac{1}{m} \varphi(m) - 1$$

since the indices v counted in  $\varphi(m)$  but not in  $\varphi(m-1)$  are those with  $v \ge m+1$  and

$$f(\{x_1,\ldots,x_v\})=x_m$$

and by the choice of  $x_m$ , the number of such indices v is at most  $\varphi(m)/m$ ; and besides, m is counted in  $\varphi(m-1)$ , but not in  $\varphi(m)$ .

We want to show that  $\varphi(2^k) \geq (n-k)2^k$ ; more generally we show that

(2) 
$$\varphi(2^k+l) \ge (n-k)2^k + (n-k-2)l \quad (0 \le l \le 2^k).$$

Note that for  $l=2^k$  this inequality is the same as

$$\varphi(2^{k+1}) \ge (n-k-1)2^{k+1}.$$

Inequality (2) is easily proved by induction on  $n-2^k-l$ . If  $n-2^k-l=0$ , it is trivially true. Suppose  $2^k+l < n$ . We may assume  $l < 2^k$  since otherwise we could consider the decomposition  $2^k+0=2^{k-1}+2^{k-1}$ . By (1)

$$\varphi(m-1) \ge 1 + \frac{m-1}{m} \varphi(m)$$
,

whence

$$\varphi(2^k+l) \ge 1 + rac{2^k+l}{2^k+l+1} \varphi(m).$$

By induction hypothesis

$$\varphi(m) = \varphi(2^k + l + 1) \ge (n - k) 2^k + (n - k - 2) (l + 1) =$$

$$= (n - k) (2^k + l + 1) - 2(l + 1).$$

Thus

$$arphi(2^k+l) \geq 1 + (2^k+l) (n-k) - rac{2(l+1)(2^k+l)}{2^k+l+1} = \ = (2^k+l) (n-k) + 1 - 2l - 2 + rac{2l+2}{2^k+l+1}$$

and since the left-hand side is an integer,

$$\varphi(2^k + l) \ge (2^k + l) (n - k) + 1 - 2l - 2 + 1 =$$

$$= (n - k) 2^k + (n - k - 2) l,$$

as stated. Thus  $\varphi(0) = \varphi(1) + 1 \ge n + 1$ 

(c) Let  $\varphi_0$  be the mapping of  $\{1, \ldots, 2^n\}$  into itself defined in 14.26b. We define a mapping f associating with each k-element subset X of  $\{1, \ldots, 2^n - 1\}$  the  $\varphi_0(k)^{\text{th}}$  largest element of X. This is possible, because  $1 \leq \varphi_0(k) \leq k$ . Now suppose  $X_0 \subset X_1 \subset \ldots \subset X_n$  is a chain of subsets with  $f(X_0) = \ldots = f(X_n) = y$  (one may suppose  $X_0 = \{y\}$ , but this is irrelevant). Let  $k_i = |X_i|$ . Then, by the definition of f,

$$\varphi_0(k_i) = |X_i \cap \{y, \dots, 2^n - 1\}|,$$

thus  $\varphi_0(k_0) \leq \varphi_0(k_1) \leq \ldots \leq \varphi_0(k_n)$ . But this is impossible by 14.26.

Remark: Part (b) gives, of course, a new proof of part (a) of 14.26; but the proof given there is much simpler [E. Harzheim, Publ. Math. Debrecen 15 (1968) 19–22].

**29.** Consider the complete graph  $K_N$  on the given points; orient its edges upward and color an edge color i, if it forms with the positive half of the x-axis an angle between  $\frac{i}{n}\pi$  and  $\frac{i+1}{n}\pi$  ( $i=0,\ldots,n-1$ ) (we may assume that these angles themselves never occur or else we rotate the picture). Suppose there exists no sequence  $a_0,\ldots,a_n$  of points in which  $\not < a_{i-1}a_ia_{i+1} > \left(1-\frac{1}{n}\right)\pi$  for  $i=1,\ldots,k-1$ . Then the graph with edges of color i contains no directed path of length k; such a path would exactly mean an almost straight broken line. Therefore by 9.9 the graph formed by edges of color i is k-colorable. Since  $K_N$  is the union of n such graphs it follows by 9.3 that  $\chi(K_N) \leq k^n$ , a contradiction [P. Erdős–G. Szekeres,  $Compositio\ Math.\ 2\ (1965)\ 463-470$ ].

**30.** (a) Define a tournament on S with values associated with the edges as in the hint. By 14.27a there exists a monotone increasing path of length p+1

or a monotone decreasing path of length q+1. Let, e.g. the first case occur. Then there are points  $(x_0, y_0), \ldots, (x_{p+1}, y_{p+1})$  such that  $x_0 < x_1 < \ldots < x_{p+1}$  and

$$\frac{y_0 - y_1}{x_0 - x_1} < \dots < \frac{y_{p+1} - y_p}{x_{p+1} - x_p}.$$

Clearly these points form a p + 2-gon convex from below.

(b) We construct a set S of  $\binom{p+q}{p}$  points spanning no (p+2)-gon convex from below and no (q+2)-gon convex from above, by induction on p+q. The case p=1 or q=1 is trivial, so let  $p,q\geq 2$ . Let  $S_1$  be a set of  $\binom{p+q-1}{p-1}$  points spanning no (p+1)-gon convex from below and no (q+2)-gon convex from above. Also let  $S_2$  be a set of  $\binom{p+q-1}{p}$  points spanning no (p+2)-gon convex from below and no (q+1)-gon convex from above; both  $S_1, S_2$  should consist of points in general position.




Fig. 104

Let us place  $S_1$ ,  $S_2$  on the plane in such a way that the vector connecting some two points of them is  $(M, M^2)$ , where M is very large (Fig. 104). Let us consider any polygon P convex from below; let  $(x_1, y_1), \ldots, (x_t, y_t)$  be its points,  $x_1 < \ldots < x_t$ . If all these points belong to  $S_1$  or  $S_2$ , then  $t \le p+1$  by definition. So suppose  $(x_1, y_1), \ldots, (x_t, y_t) \in S_1$ ,  $(x_{t+1}, y_{t+1}), \ldots, (x_t, y_t) \in S_2$  (clearly any point of  $S_1$  is to the left from any point of  $S_2$  if M is large enough). Then

$$\frac{y_{i+1} - y_i}{x_{i+1} - x_i} \approx M$$


so if M is larger than

$$\max\left\{\frac{y'-y}{x'-x}:(x,y),(x',y')\in S_2\right\}$$

then our polygon cannot be convex unless i+1=t. Since  $i \leq p$  by the choice of  $S_1$  it follows that  $t \leq p+1$ . Similarly we conclude that any polygon

convex from above spanned by S has at most q+1 vertices [P, Erdős, G. Szekeres; ibid.].

31. (a) We show first that 5 points always determine a convex quadrilateral. If the convex hull of them is a quadrilateral we are done, so suppose this convex hull is a triangle abc. This contains two further points d, e. Suppose the line de intersects the edges ab and ac of the triangle. Then bcde is a convex quadrilateral (Fig. 105).



Now let us 2-color all quadruples of the given n points: let a quadruple be red, if it forms the vertices of a convex quadrilateral and blue otherwise. If

$$(1) n \geq R_2^4(m,5),$$

then we have either m points with all quadruples red or 5 points with all quadruples blue. The latter case is impossible by the above. Hence we have m points any 4 of which form a convex quadrilateral. Hence the m points form a convex m-gon.

(b) I. We show that (1) can be replaced by

$$n \geq {2m-4 \choose m-2} + 1.$$


In fact, given any  $\binom{2m-4}{m-2}+1$  points in general position, they always span an m-gon convex from below or from above by 14.30a; this is a convex m-gon.

II. We construct a set of  $2^{m-2}$  points spanning no convex m-gon. Let  $T_i$  be a set of points in general position such that  $|T_i| = \binom{n-2}{i}$  and  $T_i$  spans no (i+2)-gon convex from above and no (n-i)-gon convex from below. Let  $T_i$  have diameter less than 1  $(i=0,1,\ldots,n-2)$ . Also we may suppose that any line formed by two points of  $T_i$  forms an angle less than 45° with the x-axis (by applying an affinity perpendicular to the x-axis, if necessary).

Let P be a large regular (4n-4)-gon with centre at the origin and let  $v_0$ , ...,  $v_{n-2}$  be its vertices in the segment of  $\pm 45^{\circ}$  around the positive half of the x-axis (Fig. 106). Let us place  $T_i$  around  $v_i$  (i.e. let us translate one point of  $T_i$  into  $v_i$ ). The resulting set T has

$$|T| = \sum_{i=0}^{n-2} |T_i| = \sum_{i=0}^{n-2} {n-2 \choose i} = 2^{n-2}$$

elements. We claim that it contains no convex n-gon. Let P be any convex polygon spanned by T. If P is contained in a  $T_i$   $(0 \le i \le n-2)$ , let a and b



be the vertices of P with least and largest x-coordinate respectively. The diagonal ab splits P into two polygons, convex from above and below, respectively. Hence by the definition of  $T_i$ , the number of vertices of these polygons is  $\leq i + 1$  and  $\leq n - i - 1$ , respectively. Hence the number of vertices of P is  $\leq (i+1) + (n-i-1) - 2 = n-2$ .

Now suppose that P intersects  $T_i$  and  $T_j$ , i < j. Let i be the least and j be the largest index for which this holds. Now P can intersect  $T_{\mu}$ ,  $i < \mu < j$  in at most one point, and its pieces in  $T_i$  and  $T_j$  are convex from above and below, respectively (Fig. 107). Hence P has at most

$$(i+1)+(j-i-1)+(n-j-1)=n-1$$

points.

(c)  $K(4) \ge 5$  trivially;  $K(4) \le 5$  by the first paragraph of the solution. Thus K(4) = 5.

 $K(5) \ge 9$  by (b). So what we have to show is that any 9 points (in general position) contain a convex pentagon. If the convex hull of the 9 points is a pentagon we are done. So suppose it is a quadrilateral or a triangle.

1° Suppose the convex hull is a quadrilateral xyzu. Consider the 5 remaining points. If they form a convex pentagon we are done. Otherwise, there are 4 among them, which do not form a convex quadrilateral; let a point v be contained in a triangle abc (Fig. 108). Consider the angles avb, bvc, cva. One of these contains x and y. Then xyavb is a convex pentagon.

2° Suppose the convex hull is a triangle abc. Again we have two subcases: I. The convex hull of the remaining 6 points is a quadrilateral xyzu. Let p, q be two further points (inside xyzu). If the line pq intersects two adjacent edges of xyzu we have a pentagon trivially (Fig. 109). So suppose pq intersects the edges xy and uv. We may suppose the intersection with xy is nearer p than q (Fig. 109). Now consider the semilines px, py, qz, qu. They divide the




Fig. 110

outside of xyzu into 4 regions. If the region ypqz contains one of a, b, c we are done. Similarly if xpqu contains one of a, b, c.

If ypqz and xpqu contain none of a, b, c, then either xpy or zqu contains two of

those; then we find a convex pentagon again.

II. Finally, suppose the convex hull of the 6 points inside abc is another triangle xyz. Again let p, q be two points inside xyz. We may assume that the line pq intersects the edges xy and xz. Consider the angles xpy, xqz. Each of these may contain at most one of a, b, c otherwise we find a convex pentagon. Hence one of a, b, c lies in the region determined by the semilines py, qz and the segment pq; but this yields a convex pentagon again (Fig. 110, p. 505) [ibid.].

## § 15. Reconstruction

1. (a) Let  $e_1, \ldots, e_k$   $(k \ge 4)$  be the edges adjacent to a given point  $x \in V(G)$ . Then  $e_1, \ldots, e_k$  form a complete subgraph of L(G). Moreover, they form a maximal complete subgraph (clique) of L(G); for if  $e \in E(G)$  is any other edge, then it does not contain x and thus, it meets at most two of  $e_1, \ldots, e_k$ .

Conversely, let  $e_1, \ldots, e_k$  form a clique of L(G),  $k \ge 4$ , we claim that they are the edges adjacent to one of the points of G. For let x be the common point of  $e_1$  and  $e_2$ . If  $e_3, \ldots, e_k$  also contain x, then we are finished, since, by the maximality of the complete subgraph of G formed by  $e_1, \ldots, e_k$ , no other edge can contain x.

Assume indirectly that  $e_3$  does not contain x, then  $e_1$ ,  $e_2$ ,  $e_3$  form a triangle. Since  $k \geq 4$ , we also have an  $e_4$ ; obviously,  $e_4$  cannot meet all of  $e_1$ ,  $e_2$ ,  $e_3$ , a contradiction.

The above assertions show that, if G has degrees at least 4, then we obtain G from L(G) as follows: Let  $C_1, \ldots, C_n$  be those cliques of L(G) with at least 4 points; associate a point  $x_i$  with  $C_i$  and join  $x_i$  and  $x_j$  iff  $C_i$  and  $C_j$  have a point in common. The resulting graph G' will be isomorphic to G.

Thus, if  $G_1$ ,  $G_2$  are two graphs with all degrees at least 4 and  $L(G_1) \simeq L(G_2)$ 

then, "reconstructing"  $G_1$  and  $G_2$  as above, we get isomorphic graphs.

(b) Let  $S_1, \ldots, S_n$  be the stars of the points of  $G_1$  and let  $S'_1, \ldots, S'_m$  be the stars of the points of  $G_2$ . Consider an isomorphism  $\sigma: L(G_1) \to L(G_2)$ .


We may assume  $n \geq m$ .

 $\sigma(S_i)$  cannot be a star for each  $i=1,\ldots,n$ . For if  $\sigma(S_i)$  is a star, then  $\sigma(S_i)$ ,  $\sigma(S_j)$   $(i \neq j)$  are different stars (since  $S_i \neq S_j$ , except in the trivial case  $G_1 \cong K_2$ , which we may ignore) and so, it would follow that m=n,  $\sigma(S_i)=S_i'$   $(i=1,\ldots,n)$  with a suitable choice of indices. But then mapping the center of  $S_i$  onto the center of  $S_i'$   $(i=1,\ldots,n)$ , we get an isomorphism between  $G_1$  and  $G_2$ .

Thus  $\sigma(S_1)$  is not a star (say). We claim that there are three edges  $e_1$ ,  $e_2$ ,  $e_3$  in  $G_1$  and corresponding edges  $e'_1$ ,  $e'_2$ ,  $e'_3$  in  $G_2$ , which form a triangle in one graph

and a star in the other.

1° If  $|S_1| = 1$ . i.e.  $S_1 = \{e_1\}$ , then no endpoint of  $e'_1 = \varrho(e_1)$  has degree 1. So there are two edges  $e'_2$ ,  $e'_3 \neq e'_1$  starting from the two endpoints of  $e'_1$ .  $e_2 = \sigma^{-1}(e'_1)$  and  $e_3 = \sigma^{-1}(e'_3)$  meet  $e_1$ , which is only possible if they start from






Fig. 111

the same endpoint of  $e_1$ . But then  $e_2'$  and  $e_3'$  must meet too, i.e.  $e_1'$ ,  $e_2'$ ,  $e_3'$  form a triangle.

2° [If  $S_1 = \{e_1, e_2\}$ , then  $e'_1 = \sigma(e_1)$ ,  $e'_2 = \sigma(e_2)$  have a point x' in common. There must be a further edge  $e'_3$  starting from x'.  $e_3 = \sigma^{-1}(e'_3)$  meets both  $e_1$  and  $e_2$ , hence they form a triangle.

3° If  $S_1 = \{e_1, e_2, e_3\}$ , then  $e'_1$ ,  $e'_2$ ,  $e'_3$  must form a triangle; otherwise, they would start from the same point and since  $\sigma(S_i)$  is not a star, there would be a fourth edge meeting all of them, which is clearly impossible.

So we have three edges  $e_1$ ,  $e_2$ ,  $e_3$  forming a triangle, so that the corresponding edges  $e'_1$ ,  $e'_2$ ,  $e'_3$  start from the same point x.  $e_1 = (u_2, u_3)$ ,  $e_2 = (u_1, u_3)$ ,  $e_3 = (u_1, u_2, u_3)$ 

 $= (u_1, u_2), e'_i = (x, y_i) (i = 1, 2, 3).$ 

The above pair yields a solution:  $G_1 = (\{u_1, u_2, u_3\}, \{e_1, e_2, e_3\}) \not\cong G_2 = (\{x, y_1, y_2, y_3\}, \{e_1', e_2', e_3'\})$  but  $L(G_1) \cong L(G_2)$  We claim there is no further such pair. If there was, we would have  $|E(G_i)| \geq 4$ .


Let  $e_4 \in E(G_1)$ ,  $e_4 \neq e_1$ ,  $e_2$ ,  $e_3$ . We may assume  $e_4 = (u_1, v)$ . Then  $e'_4 = \sigma(e_4)$  meets  $e_1$  and  $e_3$ , hence  $e'_4 = (y_2, y_3)$ . Since  $G_1 \ncong G_2$ , we have a further edge  $e_5 \in E(G)$ . Again we may assume that it meets one of  $e_1$ ,  $e_2$ ,  $e_3$ ,  $e_4$ . Obviously,


 $u_1 \notin e_5$ .

If  $e_5$  does not meet  $e_4$ , then it meets  $e_1$  and  $e_2$  (say).  $e_5' = \sigma(e_5)$  must meet  $e_1'$  and  $e_2'$  but not  $e_3'$  and  $e_4'$ , which is impossible. If  $e_5$  meets  $e_4$  but not  $e_1$ , then  $e_5'$  meets  $e_4'$  but not  $e_1'$ ,  $e_2'$ ,  $e_3'$ , which is impossible again. Thus  $e_5 = (u_2, v)$  or  $(u_3, v)$ ; say  $e_5 = (u_2, v)$ , then  $\sigma(e_5) = e_5' = (y_1, y_3)$ . Since  $G_1 \not\cong G_2$ , we must have a sixth edge  $e_6$ . As above, we get  $e_6 = (u_3, v)$ ,  $\sigma(e_6) = (y_1, y_2)$ . Now  $G_1 \not\cong G_2$  implies that there must be a seventh edge, but the same argument as above shows this is impossible (Fig. 111).

- (c) Let  $\sigma: L(G_1) \to L(G_2)$  be a non-trivial isomorphism. Again let  $S_1, \ldots, S_n$  and  $S'_1, \ldots, S'_m$  be the stars of points in  $G_1$  and  $G_2$ , respectively. If  $\sigma(S_i)$  is a star of each  $1 \le i \le n$ , then we get, as before, that  $\sigma$  is a *trivial* isomorphism. Otherwise, we can follow the preceding solution but everywhere, where we argued "since  $G_1 \not\cong G_2$  we have a further edge" we must stop and list an example. So we get the pairs shown in Fig. 112, p. 508 [H. Whitney, *Amer. J. Math.* 54 (1932) 160–168].
- **2.** (a) Let E,  $F \in E(K_n^r)$ . Then the number of edges of  $K_n^r$  disjoint from E and F is

$$\binom{n-|E\cup F|}{r} = \binom{n-2r+|E\cap F|}{r}.$$





Similarly, the number of edges of  $K_n^r$  disjoint from  $\alpha(E)$  and  $\alpha(F)$  is

$$\binom{n-2r+|\alpha(E)\cap\alpha(F)|}{r}$$
.

Thus

$$\binom{n-2r+|E\cap F|}{r} = \binom{n-2r+|\alpha(E)\cap\alpha(F)|}{r}.$$

Since  $n-2r \ge r$ , this implies that  $|E \cap F| = |\alpha(E) \cap \alpha(F)|$ .

We want to show, as in 15.1, that the set of all r-tuples containing any given point x is mapped onto the set of r-tuples containing a point  $\beta(x)$ . This then defines a permutation  $\beta$  of  $V(K_n^r)$ , which induces  $\alpha$ .

So let  $x \in V(K_n^r)$ , and let E, F be two edges with  $E \cap F = \{x\}$ . Then, by the above,  $\alpha(E)$  and  $\alpha(F)$  have a unique point  $\beta(x)$  in common. We claim that, if any edge  $A \in E(K_n^r)$  contains x, then  $\alpha(A)$  contains  $\beta(x)$  (and conversely).

Let  $A \in E(K_n^r)$ ,  $x \in A$ , and let B be an r-tuple such that  $B \cap A = A - E - F$ . Then

(1) 
$$|B \cap A| = |A - E - F| = |A| - |A \cap E| - |A \cap F| + |A \cap E \cap F| = r + 1 - |A \cap E| - |A \cap F|.$$

Since

$$|\alpha(B) \cap \alpha(E)| = |B \cap E| = 0,$$
  
$$|\alpha(B) \cap \alpha(F)| = |B \cap F| = 0,$$

we have

$$\begin{aligned} |\alpha(B) \cap \alpha(A)| &\leq |\alpha(A) - \alpha(E) - \alpha(F)| = r - |\alpha(A) \cap \alpha(E)| - \\ &- |\alpha(A) \cap \alpha(F)| + |\alpha(A) \cap \alpha(E) \cap \alpha(F)|. \end{aligned}$$

Here  $|\alpha(B) \cap \alpha(A)| = |B \cap A|, |\alpha(A) \cap \alpha(E)| = |A \cap E|, |\alpha(A) \cap \alpha(F)| =$  $= A \cap F$ , thus (1) implies that

$$|\alpha(A) \cap \alpha(E) \cap \alpha(F)| \geq 1$$
,

i.e.  $\beta(x) \in \alpha(A)$ .

(b) We sharpen the preceding argument. The number of r-tuples disjoint from two given r-tuples E, F with  $|E \cap F| = r - 1$  is

$$\binom{n-r-1}{r} > 0,$$

and is less than this if  $|E \cap F| < r - 1$ . Therefore,  $|E \cap F| = r - 1$  iff

 $|\alpha(E) \cap \alpha(F)| = r - 1.$ 

Suppose now we know  $|E \cap F| = k \Leftrightarrow |\alpha(E) \cap \alpha(F)| = k$  for  $k = 0, 1, \ldots$ , ...,  $k_0$  (we certainly know this for k=0). Then  $|E \cap F| = k_0 + 1 \Leftrightarrow$  $\Leftrightarrow |E \cap F| > k_0$  and there is an r-tuple A with  $|E \cap A| = r - 1$ ,  $|F \cap A| = k_0 \Leftrightarrow |\alpha(E) \cap \alpha(F)| > k_0$  and there is an r-tuple A with  $|\alpha(E) \cap A| = k_0$  $\bigcap \alpha(A)| = r - 1, \ |\alpha(F) \cap \alpha(A)| = k_0 \Leftrightarrow |\alpha(E) \cap \alpha(F)| = k_0 + 1.$  This proves that  $|E \cap F| = |\alpha(E) \cap \alpha(F)|$  for any two r-tuples E, F.

Now let x be a point of  $K_n^r$  and E, F two r-tuples in  $K_n^r$  with  $E \cap F = \{x\}$ . Let  $\alpha(E) \cap \alpha(F) = \{\beta(x)\}\$ , we show that  $x \in A$  implies  $\beta(x) \in \alpha(A)$  or, equivalently,  $x \notin A$  implies  $\beta(x) \notin \alpha(A)$ .

Let  $x \notin A$ , and assume indirectly that  $\beta(x) \in \alpha(A)$ . Set  $|A \cap E| = i$ ,  $|A \cap A| = i$ 

 $\bigcap F = j, d = |V(K_n^r) - A - E - F|$ . Then

$$d = n - (3r - 1 - i - j) \le i + j + 1 \le r$$
.

Also,

$$|V(K_n^r) - A - \{x\}| = n - r - 1 \ge r.$$

So we can find an r-tuple B such that

$$V(K_n^r) - A - E - F \subseteq B \subseteq V(K_n^r) - A - \{x\}.$$

Consider  $\alpha(B)$ . Obviously,  $\alpha(B) \cap \alpha(A) = \emptyset$ , so  $\beta(x) \notin \alpha(B)$ . Moreover,

$$|\alpha(B) \cap \alpha(E)| = |B \cap E|$$
  
$$|\alpha(B) \cap \alpha(F)| = |B \cap F|,$$

thus

$$|\alpha(B) - \alpha(E) - \alpha(F)| = |B| - |B \cap E| - |B \cap F| = d.$$

Hence

$$|V(K_n^r) - \alpha(A) - \alpha(E) - \alpha(F)| \ge |\alpha(B) - \alpha(E) - \alpha(F)| = d.$$

But

$$|V(K_n^r) - \alpha(A) - \alpha(E) - \alpha(F)| = n - 3r + |\alpha(A) \cap \alpha(E)| + + |\alpha(A) \cap \alpha(F)| + |\alpha(E) \cap \alpha(F)| - |\alpha(A) \cap \alpha(E) \cap \alpha(F)| = = n - 3r + i + j + 1 - 1 = d - 1,$$

a contradiction [C. Berge, Hypergraph Seminar, Lecture Notes in Math. 411, Springer, 1974, 1-12].

3. Let E, F be two r-tuples,  $|E \cap F| = j$ . Let us count the number of r-tuples adjacent to both E and F. We get

(1) if j < 2i - r, there are no such r-tuples,

(2) if  $2i - r \leq j$ , the number of r-tuples adjacent to both E and F is

$$\sum_{r=2i-r}^{\min(i,j)} {j \choose r} {r-j \choose i-v}^2 {n-2r+j \choose r-2i+v} = f_j(n)$$

(we consider i and r fixed).

The polynomials  $f_j(n)$  are distinct for j = 2i - r,  $2i - r + 1, \ldots, r - 1$ , since their leading terms are

$$egin{aligned} inom{r-j}{i-j}^2 & rac{n^{r-2i+j}}{(r-2i+j)!} & ext{if } 2i-r \leq j \leq i, \\ inom{j}{i} & rac{n^{r-i}}{(r-i)!} & ext{if } i \leq j < r. \end{aligned}$$

Thus, if n is large enough the numbers  $f_j(n)$  are distinct for  $j=2i-r,\ldots,r-1$ . This implies that we have the characterization:

 $|E \cap F| = j$  iff the number of r-tuples adjacent to both E and F is equal to  $f_j(r)$ . Thus, if we define a graph on the points of  $L_i(K_n^r)$  by joining two of them when the number of neighbors they have in common is  $f_{i-1}(n)$  (note that  $2i - r \le i - 1$ ), we get  $L_{i-1}(K_n^r)$ .  $\alpha$  is therefore an automorphism of  $L_{i-1}(K_n^r)$ . Proceeding similarly, we get that  $\alpha$  is an automorphism of  $L_0(K_n^r)$ . But  $L_0(K_n^r) = \overline{L(K_n^r)}$ , so  $\alpha$  is an automorphism of  $L(K_n)$ . By 15.2, this implies that  $\alpha$  is induced by a permutation of  $V(K_n^r)$  [C. Berge; ibid.].

4. (a) Let  $S_1, \ldots, S_n$  be the stars of points of G, and consider the hypergraph

$$H = (E(G), \{S_1, \ldots, S_n\}).$$

It is obvious that  $L(H) \simeq G$ .

More generally, let  $C_1, \ldots, C_m$  be complete graphs such that  $C_1 \cup \ldots \cup C_m = G$ . For each  $x \in V(G)$ , let  $U_x$  be the set of those  $C_i$ 's containing x, and set

$$H = (\{C_1, ..., C_m\}; \{U_x : x \in V(G)\}).$$

Then  $L(H)_{\underline{a}} \simeq G$ . The natural correspondence is  $\sigma(U_x) = x$  and it is trivial to

verify that it is an isomorphism.

Conversely, suppose  $L(H) \cong G$  for a hypergraph H and let  $\sigma: L(H) \to G$  be an isomorphism. For each  $v \in V(H)$ , let  $C_v$  be the subgraph of G spanned by the points  $\sigma(E)$ , where  $v \in E \in E(H)$ . Obviously,  $C_v$  is a complete subgraph,  $\bigcup_{v \in V(H)} C_v = G$  and if  $U_x$  denotes the set of complete graphs  $C_v$  containing x, we have

$$H \simeq (\{C_v : v \in V(H)\}; \{U_x : x \in V(G)\}).$$

(b) We prove by induction on n that G is the union of  $\frac{n^2}{4}$  complete graphs.

For n=2 this is obvious. Let n>2, and let x, y be two adjacent points. G-x-y is the union of  $t=\left[\frac{(n-2)^2}{4}\right]=\left[\frac{n^2}{4}\right]-n+1$  complete graphs  $C_1$ , ...,  $C_t$  and certain isolated points.

Let  $z \in V(G) - x - y$ . It z is adjacent to both of x, y, then let  $C_z$  be the triangle xyz. If z is adjacent to x or y only, then define  $C_z$  as the edge joining z to x, y. Otherwise, let  $C_z = \{z\}$ . Also let  $C_0$  be the edge (x, y). Now adding the new complete graphs  $C_0$  and  $C_z$  ( $z \in V(G) - x - y$ ) we get

$$t+n-2+1=\left[\frac{n^2}{4}\right]$$

complete graphs whose union is G.

Thus we can cover the edges of G by  $\left[\frac{n^2}{4}\right]$  complete graphs and this yields a

hypergraph with line-graph G on  $\frac{n^2}{4}$  points.

The result is sharp, as is shown by a complete bipartite graph  $K_{\left[\frac{n}{2}\right]} \cdot \left[\frac{n+1}{2}\right]$  [P. Erdős–A.W. Goodman–L. Pósa, *Canad. J. Math.* 18 (1966) 106–112].

5. (a) Let x, y, z be three points of G such that, removing any combination of them, G remains connected (i.e. x, y, z are three endpoints of a spanning tree of G; if the only spanning tree of G is a path, then G is a path or circuit and therefore a line-graph).

Let  $G-x \cong L(H_x)$ ,  $G-y \cong L(H_y)$ , ...,  $G-x-y \cong L(H_{xy})$ , ...  $G-x-y-y \cong L(H_{xyz})$ . Then  $L(H_{xyz}) \cong L(H_{xy}-e_z)$  for the edge  $e_z$  corresponding to z. Since  $|E(H_{xyz})| > 6$ , this means that the isomorphism between  $L(H_{xyz})$  and  $L(H_{xy}-e_z)$  is induced by an embedding of  $H_{xyz}$  into  $H_{xy}$ . Similarly, we get a whole lot of embeddings (illustrated in Fig. 113), which commute with the isomorphisms between  $L(H_{xy})$ , etc. and the corresponding induced subgraphs of G (Fig. 113).

Since we also know that the embedding of  $H_{xyz}$  into  $H_x$ ,  $H_y$ ,  $H_z$  is unique, this diagram commutes. So we may identify all points mapped onto each other by embeddings, and form the union H. The edges of H correspond to the points




Fig. 113

of G in the natural way and it is easy to check that this correspondence is an isomorphism between L(H) and G.

(b) Let  $G_1, \ldots, G_k$  be those simple graphs which are not line-graphs but each proper induced subgraph of which is a line-graph. Since by the preceding

problem they have at most nine points, their number is finite.

Now if G is any simple graph which is not a line graph, then it contains a minimal induced subgraph, which is not a line graph, and this is one of  $G_1, \ldots, G_k$ . Conversely, if G contains one of  $G_1, \ldots, G_k$  then, obviously, it cannot be a line-graph.

**6.** (a) Suppose the points a, b, c, d, of G = L(H) form a 3-star with center a. Then the edge a is adjacent to b, c and d; some two of them b and c say, start

from the same endpoint of a. But then b and c are adjacent.

Now let abc, bcd be odd triangles, we claim a, b, c start from the same point of H. Suppose indirectly that a, b, c form a triangle of H. By definition, there is an edge x adjacent to one or three of a, b, c. But this is impossible since x must start from a point of the triangle formed by a, b, c in H and so, x is adjacent to two of a, b, c. Thus a starts from the common point of b, c. Similarly, d starts from the common point of b, c implying that a, d are adjacent.

(b) We may assume that G is connected since each component satisfies (i) and (ii) and if they are line-graphs, then so is G. Let  $(a,b) \sim (c,d)$  be as in the hint. Suppose  $(a,b) \sim (c,d) \sim (e,f)$  are distinct edges of G, we show that  $(a,b) \sim (e,f)$  (the other criteria for an equivalence relation are trivially fulfilled). Observe that  $\{a,b,c,d\}$  spans a complete graph. Consider a and e (say),  $a \neq e$ . If a = c or d, then a, e are trivially adjacent so suppose  $a \neq c,d$ ; similarly  $e \neq c,d$ . Then acd is an odd triangle (if b=c or d this follows immediately from the definition of equivalence, otherwise b is adjacent to three points of it), similarly cde is an odd triangle, whence a and e are adjacent by (ii). Now, if (a,b) and (e,f) have no common endpoint we immediately conclude that  $(a,b) \sim (e,f)$ . Suppose a=e. If c or d is different from a, b, f, then it is adjacent to three points of the triangle abf, so this triangle is odd; if (c,d) is also an edge of this triangle, then it must be odd again. So  $(a,b) \sim (e,f)$  follows again.


Let  $C_1, \ldots, C_N$  be the equivalence classes of the relation  $\sim$ . They obviously form complete graphs. We show that each point of G is contained in at most two of these complete graphs unless G is one of the exceptional graphs shown

in Fig. 114a, b, c.

Assume that  $x \in V(G)$  is contained in three complete graphs, i.e. there are three non-equivalent edges (x, y), (x, z) (x, u) adjacent to x. By (i) some two of y, z, u are adjacent; say  $(y, z) \in E(G)$ . Since xyz is not an odd triangle, u must be adjacent to one of y, say  $(u, y) \in E(G)$ , and then  $(u, z) \notin E(G)$ . If |V(G)| = 4,

then we get(a) in Fig. 114.

If G is larger, then there is a point v adjacent to one of x, y, z, u. It must be adjacent to an even number of points of the triangles xyz and xyu, therefore it must be adjacent to one of x, y. Let, say,  $(v, x) \in E(G)$  (the roles of x and y have been symmetric as, obviously, (y, x), (y, z), (y, u), are non-equivalent). By (i) v is adjacent to one of z and u, say  $(u, v) \in E(G)$ . Again, because xyz and xyu are not odd triangles  $(v, y) \in E(G)$  and  $(v, z) \in E(G)$ . If G has only five points, we get(b) in Fig. 114.



Obviously, (v, x) is non-equivalent to at least one of (x, z) and (x, u), say (v, x) is non-equivalent to (x, u), i.e. vxu cannot be an odd triangle; for if it were, then a point adjacent to an odd number of points of vxu would be adjacent to an odd number of points of one of the triangle vxz, xyz, xyu, as is easy to verify. Thus, the roles of y, z, u, v are symmetric.


Now suppose that G has a further point w, adjacent to one of the x, y, z, u, v. w cannot be adjacent to x; for if it were, then one of (y, v) as well as one of z, u ought to be a neighbor of w; let, say, (y, w),  $(z, w) \in E(G)$ , then xyz is odd, a contradiction. So w is adjacent to y (say) but not to x. Then it follows that it is adjacent to u, v, z also. So if G has 6 points, we have(c) in Fig. 114.

Observe that no two edges spanned by  $x, \ldots, w$  can be equivalent, hence the roles of  $x, \ldots, w$  are symmetric. Therefore, the argument of the last paragraph shows that no further edge can be adjacent to any of the points  $x, \ldots, w$ , i.e. indeed we only get the three graphs of Fig. 114.

Now it is easy to complete the proof. If G is one of the graphs of Fig. 114. then it is a line-graph of the corresponding graph in Fig. 114a', b', c'. Otherwise, G can be covered by edge-disjoint complete subgraphs so that each point is contained in exactly two of them (we add one-point graphs, if necessary). As in 15.40a, this implies that G is a line-graph.

(c) We claim that G is a line-graph of a simple graph iff it does not contain any of the graphs shown in Fig. 115 (p. 514) as an induced subgraph.

It is immediately clear that these graphs are not line-graphs since they violate (i) or (ii) of (a). Conversely, suppose G is not a line-graph. Then it either contains




the 3-star as an induced subgraph or it violates (ii), i.e. there are points a, b, c, d such that abc, bcd are odd triangles and a, d are non-adjacent. Let e and f be points adjacent to an odd number of points of abc and bcd, respectively. Then there are several cases to distinguish: e = f or  $e \neq f$ ; which points of the corresponding triangle are adjacent to e, f. The consideration of these cases is straightforward but lengthy; they yield the graphs shown in Fig. 115. We leave the details to the reader [L.W. Beineke; see B, H].

7. From 15.4 we know that a graph is the line-graph of a 3-uniform hypergraph without multiple edges iff its edges can be covered by complete subgraphs so that each point is contained in three of them and each edge is contained in at most two of them.

Now suppose that the graph shown in Fig. 20 (p. 138) is a line-graph, i.e. it has such a decomposition into complete graphs. Obviously,  $(x_0, z_1)$  and  $(y_0, z_1)$  must be complete graphs by themselves. Since  $z_1$  is contained in three complete graphs only,  $z_1$ ,  $x_1$ ,  $y_1$  must be one. Similarly,  $u_1$ ,  $x_1$ ,  $y_1$  must be one of the covering complete subgraphs. Thus  $(x_1, y_1)$  is already contained in two covering complete subgraphs, hence  $(x_1, z_2)$  and  $(y_1, z_2)$  must be covering complete subgraphs by themselves. From here on, the argument periodically repeats till we get that  $(x_n, z_{n+1})$  and  $(y_n, z_{n+1})$  are covering complete subgraphs by themselves. Since, obviously, so are  $(x_{n+1}, z_{n+1})$  and  $(y_{n+1}, z_{n+1})$ ,  $z_{n+1}$  is contained in four of them, a contradiction.

Now remove a point from G. There are essentially four kinds of points:  $x_i$  (or  $y_i$ ),  $z_i$ ,  $u_i$ ,  $v_i$  (or  $w_i$ ). Figure 116 shows how to decompose G-t into complete subgraphs, if  $t=x_i$ ,  $z_i$ ,  $u_i$ ,  $v_i$  [L. Nickel].



8. (a) See Fig. 117.

- (b) This is just a re-formulation of the corollary, which has been drawn from 6.69.
- **9.** (a) Let  $e_1=(x,y),\ e_2=(x,z)$  be adjacent edges of  $G_1$ ; we show that  $\psi(e_1),\ \psi(e_2)$  are adjacent as well. If  $e_1,\ e_2$  are parallel, clearly so are  $\psi(e_1),\ \psi(e_2)$ . Since  $G_1-x$  is 2-connected it contains a circuit  $C_x$  through y and z, if  $y\neq z$ . Then  $\psi(C)$

is a circuit in  $G_2$ . Since  $C+e_1$  contains no other circuit, neither does  $\psi(C)+\psi(e_1)$ . Therefore,  $\psi(e_1)$  is not a chord of  $\psi(C)$ , i.e. it has an endpoint u not on  $\psi(C)$ . But  $C+e_1+e_2$  contains a circuit C' through  $e_1$  and  $e_2$ . Then  $\psi(C')\subseteq \psi(C)+\psi(e_1)+\psi(e_2)$  is a circuit in  $G_2$  containing  $\psi(e_1)$  and  $\psi(e_2)$ . Thus there is an edge  $\psi(e_1)$  of  $\psi(C')$  adjacent to u. This edge must be  $\psi(e_2)$ . So  $\psi(e_1)$  and  $\psi(e_2)$  are adjacent.

- (b) Now let  $x \in V(G_1)$ , and let  $e_1, \ldots, e_k$  be all edges adjacent to x. Then  $\psi(e_1), \ldots, \psi(e_k)$  intersect by pairs and no three of them form a triangle. Hence there is a point  $\eta(x)$  contained in all of  $\psi(e_1), \ldots, \psi(e_k)$ . The point  $\eta(x)$  is unique; for if a  $u \neq \eta(x)$  also belonged to  $\psi(e_1), \ldots, \psi(e_k)$ , then  $\psi(e_1), \ldots, \psi(e_k)$  would be parallel; thus  $e_1, \ldots, e_k$  would be parallel, i.e. x would have a unique neighbor, which is impossible, as  $G_1$  is 3-connected. Hence  $\eta$  is well-defined. Let  $u \in V(G_2)$  and  $f_1, \ldots, f_k$  be all edges adjacent to u. As above, there is a unique point  $x \in V(G_1)$  contained in  $\psi^{-1}(f_1), \ldots, \psi^{-1}(f_k)$ . Then  $\eta(x)$  is contained in  $f_1, \ldots, f_k$  and hence  $u = \eta(x)$ . Therefore,  $\eta(x)$  is onto. Similarly, it follows that it is one-to-one. Also, if  $(x, y) = e \in E(G)$ , then  $\eta(x)$ ,  $\eta(y)$  are endpoints of  $\psi(e)$ . Therefore  $\eta$  is an isomorphism [H. Whitney, see loc. lit. p. 507].
- 10. If r=2 the assertion is trivial. Assume r>2. Let v(v') be the point next to  $x_r$  ( $x_r'$ ), which has degree at least 3. Remove  $x_r$  and all inner points of the path joining  $x_r$  to v from T, and remove  $x_r'$  and the inner points of the ( $x_r', v'$ )-path from T'. This way we get two trees  $T_1, T_1'$  with endpoints  $x_1, \ldots, x_{r-1}$  and  $x_1', \ldots, x_{r-1}'$ , respectively. Obviously, the distance between  $x_i$  and  $x_j$  is the same in T and  $T_1$  ( $1 \le i < j \le r-1$ ) and similarly for T' and  $T_1'$ . Thus, by the induction hypothesis,  $T_1 \simeq T_1'$ , and, in fact, there exists an isomorphism  $\varphi$  such that  $\varphi(x_i) = x_i'$ .

We show that  $\varphi(v) = v'$ . Consider the unique  $(\varphi(v), v')$ -path, and extend it to

a path joining  $x_i'$  to x'  $(1 \le i < j \le r - 1)$ . Then

$$d(x'_i, x'_j) + d(x'_i, x'_r) - d(x'_j, x'_r) = 2d(x'_i, v'),$$
  
 $d(x_i, x_j) + d(x_i, x_r) - d(x_j, x_r) = 2d(x_i, v).$ 

while

So by the assumption  $d(x_i, v) = d(x'_i, v')$ , which implies that  $v' = \varphi(v)$ . It also follows that  $d(x_r, v) = d(x'_r, v')$ ; for

$$d(x_r, x_i) + d(x_r, x_j) - d(x_i, x_j) = 2d(x_r, v), \ d(x_r', x_i') + d(x_r', x_j') - d(x_i', x_j') = 2d(x_r', v')$$

and the left-hand sides are equal by the assumption. Now  $\varphi$  extends to an isomorphism of T onto T' obviously [E. A. Smolenskii, *Zhurnal Vychisl. Mat. i Matem. Fiz.* (1962) 371–372].

11. (a) Let  $P_{ij}$  be the (unique)  $(x_i, x_j)$ -path in T. Let Q be the (unique)  $(P_{ij}, x_k)$ -path and denote by v its beginning point. Then, obviously

$$d(x_i, x_i) + d(x_i, x_k) - d(x_i, x_k) = 2d(x_i, v)$$

which proves (a).

Now consider the two paths  $P_{ij}$  and  $P_{kl}$ . They may have 0, 1 or at least 2 points in common. Suppose first that they do not meet, then let Q be the

 $(P_{ij}, P_{kl})$ -path with endpoints  $u \in V(P_{ij})$ ,  $v \in V(P_{kl})$ . Let  $R_i$ ,  $R_j$  be the  $(x_i, u)$ ,  $(x_i, u)$ -paths and  $R_k$ ,  $R_l$  be the  $(x_k, v)$ ,  $(x_l, v)$ -paths in T. Then

$$P_{ik} = R_i \cup Q \cup R_k, \qquad P_{jl} = R_j \cup Q \cup R_l$$
  
 $P_{jk} = R_j \cup Q \cup R_k, \qquad P_{il} = R_i \cup Q \cup R_l.$ 

So

$$d_{ik} + d_{jl} = d_{il} + d_{jk} = d_{ij} + d_{kl} + 2d(u, v),$$

which proves assertion (b) in this case.

Now suppose that  $P_{ij}$  and  $P_{kl}$  have exactly one point v in common. Then considering the form of paths joining v to  $x_i$ ,  $x_j$ ,  $x_k$   $x_l$  we get, in the same way as above,

$$d_{ij}+d_{kl}=d_{ik}+d_{jl}=d_{il}+d_{jk}.$$

Finally, if  $P_{ij}$  and  $P_{kl}$  have more than one point in common, then either  $P_{ik}$  and  $P_{jl}$  or  $P_{il}$  and  $P_{jk}$  do not meet and we have the first case by exchanging indices.

(c) Suppose a tree T' is constructed with endpoints  $x_1, \ldots, x_{r-1}$  such that  $d(x_i, x_j) = d_{ij}$  holds in T' for each  $1 \le i < j \le r-1$ . Let  $x_i, x_j$  be two points of T' such that

$$d_{ri} + d_{ri} - d_{ii}$$

is minimal. Let v be the point on the  $(x_i, x_i)$ -path in T' for which

$$d(v, x_i) = \frac{1}{2} (d_{ri} + d_{ij} - d_{rj})$$

and join  $x_r$  to v by a path of length  $\frac{1}{2}(d_{ri}+d_{rj}-d_{ij})$  (this is possible since these

numbers are natural numbers by (a)). We claim that the resulting tree T satisfies  $d(x_k, x_l) = d_{kl}$ , for any  $1 \le k < l \le r$ . This is trivial unless l = r. It is also easy to see, when k = i or k = j from the definition of u, so suppose  $k \ne i$ , j.

By (b), two of the numbers  $d_{ij} + d_{rk}$ ,  $d_{ir} + d_{jk}$ ,  $d_{jr} + d_{ik}$  are equal and not less than the third. We claim that  $d_{ij} + d_{rk}$  is not less than the other two. For suppose

$$d_{ij} + d_{rk} < d_{ir} + d_{ik}$$

then

$$d_{rk} + d_{rj} - d_{jk} < d_{ri} + d_{rj} - d_{ij}$$

which contradicts the choice of i and j. So, for a suitable choice of indices i, j,

(1) 
$$d_{ij} + d_{kr} = d_{ir} + d_{jk} \ge d_{jr} + d_{ik}$$
.

Now T itself satisfies (b), and, in fact, the  $(x_r, x_k)$ -path and  $(x_i, x_j)$ -path in T certainly intersect. Hence, by the solution of the preceding problem

(2) 
$$d(x_i, x_j) + d(x_k, x_r) = d(x_{i_0}, x_r) + d(x_{j_0}, x_k) \ge d(x_{j_0}, x_r) + d(x_{i_0}, x_k),$$

where  $(i_0, j_0)$  is a permutation of (i, j). Since we have  $d(x_i, x_j) = d_{ij}$ ,  $d(x_{i_0}, x_r) = d_{i_0r}$  etc., (1) and (2) imply that we may assume  $i_0 = i$ ,  $j_0 = j$  and then (1) and (2) imply

$$d(x_k, x_r) = d_{kr}$$

[K.A. Zaretskii, Usp. Matem. Nauk. 20 (1965) 6, 94-96].

12. (a) The number 2l occurs in  $\{a_i + a_j : 1 \le i < j \le n\}$  exactly

$$\sum_{v=0}^{l} {k+1 \choose 2v} {k+1 \choose 2l-2v}$$

times if l is odd and

$$\sum_{v=0}^{l} {k+1 \choose 2v} \left( {k+1 \choose 2l-2v} - \frac{1}{2} \left[ {k+1 \choose l}^2 + {k+1 \choose l} \right]$$

times if l is even. It occurs in  $\{b_i + b_j : 1 \le i < j \le n\}$  exactly

$$\sum_{v=0}^{l-1} \binom{k+1}{2v+1} \binom{k+1}{2l-2v-1} - \frac{1}{2} \left[ \binom{k+1}{l}^2 + \binom{k+1}{l} \right]$$

times if l is odd and

$$\sum_{\nu=0}^{l-1} {k+1 \choose 2\nu+1} {k+1 \choose 2l-2\nu-1}$$

times if l is even. Suppose, e.g. l is odd, then one has to verify that

$$\sum_{k=0}^{l} {k+1 \choose 2\nu} {k+1 \choose 2l-2\nu} = \sum_{\nu=0}^{l-1} {k+1 \choose 2\nu+1} {k+1 \choose 2l-2\nu-1} - \frac{1}{2} \left[ {k+1 \choose l}^2 + {k+1 \choose l} \right]$$

or, equivalently,

$$\sum_{\mu=0}^{2l} (-1)^{\mu} {k+1 \choose \mu} {k+1 \choose 2l-\mu} = -\frac{1}{2} \left[ {k+1 \choose l}^2 + {k+1 \choose l} \right],$$

which follows from identity (d) in 1.42.

(b) Suppose  $\{a_i + a_j : 1 \le i < j \le n\} = \{b_i + b_j : 1 \le i < j \le n\}$ , but  $\{a_i : 1 \le i \le n\} \neq \{b_i : 1 \le i \le n\}$ . We show that n is a power of 2. We may assume  $a_i, b_i \ge 0$ , since adding any constant to all  $a_i$  and  $b_i$  the condition remains valid. Set

$$f(x) = \sum_{i=1}^{n} x^{a_i}, \quad g(x) = \sum_{i=1}^{n} x^{b_i}.$$

Then

$$f^{2}(x) = \sum_{i=1}^{n} x^{2a_{i}} + 2 \sum_{1 \leq i < j \leq n} x^{a_{i} + a_{j}} = f(x^{2}) + 2 \sum_{1 \leq i < j \leq n} x^{a_{i} + a_{j}},$$

$$g^{2}(x) = g(x^{2}) + 2 \sum_{1 \leq i < j \leq n} x^{b_{i} + b_{j}}$$

and thus,

$$f^2(x) - g^2(x) = f(x^2) - g(x^2), f(x) + g(x) = \frac{f(x^2) - g(x^2)}{f(x) - g(x)}.$$

We want to substitute x = 1 in the last identity, since, then we will have 2n on the left-hand side. But x = 1 is a root of f(x) - g(x); so write

$$f(x) - g(x) = (1 - x)^{k+1} p(x) \quad (k \ge 0, \ p(1) \ne 0).$$

Then

$$\frac{f(x^2) - g(x^2)}{f(x) - g(x)} = (1 + x)^{k+1} \frac{p(x^2)}{p(x)}$$

and

$$2n = f(1) + g(1) = (1+1)^{k+1} \frac{p(1)}{p(1)} = 2^{k+1},$$

which proves the assertion [J. Selfridge-E.G. Straus, Pac. J. Math. 8 (1958) 847-856].

13. More generally, we prove that, if

$$h(s_1,\ldots,s_k) = \sum_{i=1}^M \alpha_i \begin{pmatrix} c_{i1} \\ s_1 \end{pmatrix} \ldots \begin{pmatrix} c_{ik} \\ s_k \end{pmatrix}$$

(where  $(c_{i1}, \ldots, c_{ik}) \neq (c_{j1}, \ldots, c_{ik})$  for  $i \neq j$ ) vanishes for any choice of  $s_1, \ldots, s_k \geq 0$ , then  $\alpha_1 = \ldots = \alpha_M = 0$ . The assertion of the problem will follow by considering h = f - g.

Let  $c_1, \ldots, c_n$  be the distinct values of  $c_{ik}$   $(1 \leq i \leq N)$ . Then

$$h(s_1,\ldots,s_k)=\sum_{i=1}^p h_i(s_1,\ldots,s_{k-1})\begin{pmatrix}c_i\\s_k\end{pmatrix},$$

where  $h_i$  is of the same form as h. Setting  $x_i = h_i(s_1, \ldots, s_{k-1})$ , we have

$$(1) \begin{array}{c} x_1 + \ldots + x_p = 0, \\ \binom{c_1}{1} x_1 + \ldots + \binom{c_p}{1} x_p = 0, \\ \vdots \end{array}$$

 $egin{pmatrix} c_1 \\ p-1 \end{pmatrix} x_1 + \ldots + egin{pmatrix} c_p \\ p-1 \end{pmatrix} x_p = 0.$ 

We claim that the determinant of this system of equations is non-zero. We use the formula

$$\sum_{k=0}^{n} S(n, k) k! \binom{x}{k} = x^{n},$$

where S(n, k) are Stirling numbers of second kind (see 1.7). Multiply the  $(i + 1)^{st}$  row of the determinant of (1) by i! and then add S(j, i) times the

 $(i+1)^{\rm st}$  row to the  $(j+1)^{\rm st}$  row  $(j=1,\ldots,p-1;\ i=j-1,\ldots,0)$ . Then we get

$$egin{array}{ccccc} egin{pmatrix} \begin{pmatrix} c_1 \ 0 \end{pmatrix} & \cdots & \begin{pmatrix} c_p \ 0 \end{pmatrix} \ dots & & dots \ egin{pmatrix} c_1 \ p-1 \end{pmatrix} & \cdots & \begin{pmatrix} c_p \ p-1 \end{pmatrix} \ \end{array} = rac{1}{0\,!\,1\,!\,\ldots\,(p-1)\,!} egin{pmatrix} 1 & \ldots & 1 \ c_1 & \ldots & c_p \ dots & & dots \ c_1^{p-1} & \ldots & c_p^{p-1} \ \end{array} = rac{1}{1\leq i < j \leq p} rac{c_i-c_j}{i-j} 
eq 0.$$

Thus (1) has only the trivial solution for  $x_1, \ldots, x_p$ , i.e.  $h_i(s_1, \ldots, s_{k-1}) = 0$  for any choice of non-negative integers  $s_1, \ldots, s_{k-1}$ . Using induction on k (k = 0 can be considered as true) we find that all coefficients  $h_i(s_1, \ldots, s_{k-1})$  are 0. Thus so are all coefficients in  $h(s_1, \ldots, s_k)$ .

## 14. Form

$$\sum_{x\neq y} |E(G_i - x - y)|.$$

Here each edge is counted at each pair x, y different from its endpoints, hence

$$\sum_{x \neq y} |E(G_i - x - y)| = {|V| - 2 \choose 2} |E(G_i)|.$$

This implies that  $|E(G_1)| = |E(G_2)|$ . Now form

$$\sum_{y \neq x_0} |E(G_i - x_0 - y)|,$$

where  $x_0$  is fixed. If e is an edge adjacent to  $x_0$ , then e is counted here |V|-2 times; otherwise, it is counted |V|-3 times. Thus,

$$\sum_{y \neq x_0} |E(G_i - x_0 - y)| - (|V| - 3) |E(G_i)|$$

is the degree of  $x_0$  in  $G_i$ . Since this value is the same for i=1 and 2,  $d_{G_i}(x_0) = d_{G_i}(x_0)$ .

Finally, we have

$$|E(G_i)| - |E(G_i - x - y)| = d_{G_i}(x) + d_{G_i}(y) - \varepsilon_i(x, y),$$

where  $\varepsilon_i(x, y)$  is the number of (x, y)-edges in  $G_i$ . Since the other terms do not depend no i, we have  $\varepsilon_1(x, y) = \varepsilon_2(x, y)$ , i.e.  $G_1$ ,  $G_2$  are identical.

15. (a) Let  $H_1$ ,  $H_2$  be two circuits with  $V(H_1) = V(H_2) = V$ . Then  $H_1 - x$  is a path of length |V| - 2 and so is  $H_2 - x$ . So  $H_1 - x \cong H_2 - x$  for all  $x \in V$ , but, of course,  $H_1$  and  $H_2$  need not be identical. Thus the assertion is false.

(b) Let  $G_i - x$  have  $f(H, G_i - x)$  subgraphs isomorphic to H. Form

$$\sum_{x \in V} f(H, G_i - x).$$

Then each subgraph of  $G_i$  isomorphic to H is counted here |V| - |V(H)| times, so

$$\sum_{x \in V} f(H, G_i - x) = (|V| - |V(H)|) f(H, G_i).$$

Since |V| > |V(H)| by the assumption and the left-hand side is the same for i = 1, 2, we get

 $f(H, G_1) = f(H, G_2)$ 

as stated. The same argument works for induced subgraphs.

(c) Suppose first that  $G_1$  consists of an isolated point and a connected component H on |V|-1 points. Then  $G_2$  has an induced subgraph H' isomorphic with H, by the preceding assertion. Since we have, by the preceding assertion again,  $|E(G_1)| = |E(G_2)|$ , it follows that  $G_2$  has no edge other than those of H', i.e.  $G_2$  consists of H' and an isolated point. Thus  $G_1 \cong G_2$ .

Conversely, if  $G_2$  has a component with |V|-1 points then, similarly,  $G_1 \cong G_2$ . Thus, in every case, the number of components of  $G_1$  and  $G_2$  isomorphic

with H, where |V(H)| = |V| - 1, is the same.

Suppose that this holds for each connected graph H with  $|V(H)| = |V| - 1, \ldots, k + 1$ . Let |V(H)| = k. To get the number of components of  $G_i$  isomorphic with H we count all induced subgraphs of  $G_i$  isomorphic with H and subtract the number of induced subgraphs, isomorphic with H, of larger components of  $G_i$ . By the preceding problem and the induction hypothesis, this number will be the same for i = 1 and 2.

So  $G_1$ ,  $G_2$  have the same number of components isomorphic with any given H, hence  $G_1 \cong G_2$ . [P.J. Kelly, *Pacific J. Math.* 7 (1957) 961–968; for a survey on the reconstruction problem see J.A. Bondy–R. Hemminger, J. Graph

Theory 1 (1977) 227–268.]

16. (a) First we remark that since

$$d_{T_1}(x) = |V| - 1 - |E(T_1 - x)| = |V| - 1 - |E(T_2 - x)| = d_{T_2}(x),$$

 $T_1$  and  $T_2$  have the same endpoints. Let W be the set of these endpoints. If |W|=2, then both  $T_1,T_2$  are paths and thus  $T_1 \cong T_2$ . We will assume in the sequel that  $|W| \geq 3$ .

Let  $\hat{P}$  be any maximum path in  $T_1$  and x an endpoint not on P. Then by  $T_2 \cong x = T_1 - x \supseteq P$  it follows that  $T_2$  also contains a path of the same length. Hence the diameter of  $T_2$  is at least as large as the diameter of  $T_1$ . This being true conversely as well, it follows they have the same diameter d.

(b) We discuss the case of even d. The odd case is simpler. Let  $c_i$  be the center of  $T_i$ . We may assume d > 2 since, if  $T_i$  is a star, then it contains a point of degree n-1 and hence, the other tree is also a star.

1° Suppose first that  $T_i - x$  has diameter d for each endpoint x.

Let us determine the number  $m_i(D)$  of branches of  $T_i$  attached to  $c_i$ , which are isomorphic to a specified tree D with a specified "root" (i.e. with a specified point to be attached to  $c_i$ ). We prove by "backwards" induction that  $m_1(D) = m_2(D)$ . If  $|V(D)| \ge |V|$  this is trivially true.

Let us count the number of all occurrences of branch D attached to the center of  $T_i-x$ ,  $x\in W$ . Then by  $T_1-x\simeq T_2-x$ , we obtain the same number. The branches of  $T_i-x$  isomorphic to D arise in two ways; they are either branches of  $T_i$  themselves or arise from a larger branch of  $T_i$  by removing x. The numbers of branches D of the graphs  $T_i-x$  arising from larger branches of  $T_i$  are the same for i=1, 2 since  $T_1$  and  $T_2$  have the same larger branches. Therefore, the numbers of branches D of the graphs  $T_i-x$ , which are branches D of  $T_i$  are the same for i=1, D. Since each branch D of D, this proves that  $T_1-x=1$  times, where  $T_1-x=1$  is the number of endpoints of  $T_1-x=1$ . This implies that  $T_1-x=1$ .

 $2^{\circ}$  Let us assume that there is exactly one point  $x_0$  such that the diameter of  $T_1 - x_0 = T_2 - x_0$  is less than d. Then there must be exactly two branches  $B_1^i$ ,  $B_2^i$  attached to  $c_i$  in  $T_i$ , which contain a  $\frac{d}{2}$ -path; there may be any number of other branches.  $B_1^i$  contains exactly one point at distance  $\frac{d}{2}$  from  $c_i$ ;  $B_2^i$  contains at least two such points.

 $2^{\circ}$  (a). Suppose first that neither of  $B_1^1$ ,  $B_1^2$  is a single path. Then  $B_2^i$  is the largest branch in the trees  $T_i - x$  ( $x \in W - \{x_0\}$ ) relative to the center, which has at least two points at distance  $\frac{d}{2}$  from the root and so,  $B_2^1$  and  $B_2^2$  are isomorphic

rooted trees. Hence it follows as in  $1^{\circ}$  that  $T_1$ ,  $T_2$  have the same branches attached to their centers.

 $2^{\circ}$  (b.) Assume  $B_1^1$  is a single path. Then the endpoint  $x_1$  of  $T_1 - x_0$  adjacent to  $x_0$  has the property that it is contained in all (d-1)-paths of  $T_1 - x_0$ . Hence  $T_2 - x_0$  also contains an endpoint  $x_2$ , which occurs in all (d-1)-paths. Now  $x_0$  must be adjacent to  $x_2$  in  $T_2$ ; since otherwise both  $x_2$  and  $x_0$  would be endpoints of  $T_2$  contained in all d-paths, a contradiction.  $x_0$  is connected to corresponding points of  $T_1 - x_0 \cong T_2 - x_0$ , i.e.  $T_1 \cong T_2$ .

3° The case when  $T_1 - x \cong T_2 - x$  has diameter d-1 for two points x (it clearly cannot have diameter d-1 for more than 2 endpoints) can be treated anologously [15.15; ibid.].

17. (a) Let H be any graph on n points and let  $|G \to H|$  denote the number of those mappings  $\varphi: V(G) \to V(H)$ , which are one-to-one and map adjacent points onto adjacent points. Then  $|G \to H|$  can be expressed by an inclusion-exclusion formula as follows. For each  $S \subseteq E(G)$  let  $G_s$  denote the graph (V(G), S). Then, by a method similar to the one in 5.18,

$$|G \to H| = \sum_{S \subseteq E(G)} (-1)^{|S|} |G_S \to \overline{H}|.$$

In particular,

$$|G_1 \to G_2| = \sum_{S \subseteq E(G_1)} (-1)^{|S|} |G_{1,S} \to \bar{G}_2|,$$

(2) 
$$|G_2 \to G_2| = \sum_{S \subseteq E(G_2)} (-1)^{|S|} |G_{2,S} \to \overline{G}_2|.$$

Here, the graphs  $G_{1,S}$   $(S \subset E(G_1))$  and  $G_{2,S}$   $(S \subset E(G_2))$  are pairwise isomorphic; this follows just like 15.14b. Moreover,

$$|G_{1, E(G_1)} \to \overline{G}_2| = |G_1 \to \overline{G}_2| = 0$$

since  $G_1$  has more edges than  $\overline{G}_2$ . Similarly,

$$|G_{2, E(G_2)} \to \bar{G}_2| = 0.$$

Thus, the right-hand sides of (1) and (2) have the same terms. This implies that

$$|G_1 \to G_2| = |G_2 \to G_2| > 0.$$

Thus  $G_1$  has a one-to-one mapping into  $G_2$ . Since they have the same number of edges,  $G_1 \simeq G_2$  [L. Lovász, J. Comb. Theory 13 (1972) 309–310].

(b) We may assume  $V(G_1) = V(G_2) = V$ . Consider a random permutation  $\pi$  of V. Denote by  $A_i(B_i)$  the event that  $e_i$  is mapped onto a pair of non-adjacent points in  $G_1$  (in  $G_2$ ). Define, as in 2.2a,

$$A_I = \prod_{i \in I} A_i,$$
  $B_I = \prod_{i \in I} B_i,$   $\sigma_j = \sum_{|I|=j} P(A_I),$   $\sigma'_j = \sum_{|I|=j} P(B_I).$ 

Note that as in 15.14b or part (a) it follows that

$$\sigma_{j} = \sigma'_{j}$$
 for  $j = 0, 1, ..., m - 1$ .

Denote by  $\eta$  and  $\xi$  the numbers of  $A_i$ 's and  $B_i$ 's that occur. Then by 2.7b,

$$E(x^{\eta}) = \sum_{j=0}^{m} (x-1)^{j} \, \sigma_{j}, \quad E(x^{\xi}) = \sum_{j=0}^{m} (x-1)^{j} \, \sigma'_{j}.$$

Hence

(1) 
$$E(x^{\eta}) - E(x^{\xi}) = (x-1)^m (\sigma_m - \sigma'_m).$$

Set here x = -1 and use the trivial relations

$$|E((-1)^{\eta})| \leq 1$$
 and  $|E((-1)^{\xi})| \leq 1$ .

Then we get by (1)

$$|\sigma_m - \sigma_m'| \leq \frac{1}{2^{m-1}} < \frac{1}{n!}$$

by hypothesis. But from the definition of the probability field,  $\sigma_m$  and  $\sigma'_m$  are rationals with denominator n!. Hence they must be equal.

Now, taking the constant terms in (1), we obtain

$$P(\eta=0)=P(\xi=0).$$

But, if  $\pi$  is the identity, then  $\xi = 0$ . So

$$P(\eta = 0) = P(\xi = 0) > 0,$$

i.e. there is a permutation, which maps each  $e_i$  onto an edge. This is then an isomorphism between  $G_1$  and  $G_2$  [W. Müller, J. Comb. Theory [B] 22 (1977) 281-283].

18. We are going to show that, if |Y| = k - 1, then

$$|E(H_1 - X)| = |E(H_2 - X)|.$$

Let  $X_1, \ldots, X_t$  be the k-tuples containing Y (t = |V| - k + 1). Then

(1) 
$$\sum_{i=1}^{t} |\mathbb{Z}(H_1 - X_i)| = (|V| - k - r + 1) |E(H_1 - X)|,$$

which follows by a simple counting. Similarly,

(2) 
$$\sum_{i=1}^{t} |E(H_2 - X_i)| = (|V| - k - r + 1) |E(H_2 - X)|.$$

Since the left-hand sides of (1) and (2) are equal and |V|-k-r+1>0, (1) and (2) imply that  $|E(H_1-Y)|=|E(H_2-Y)|$ . Similarly this follows for each Y with  $|Y|\leq k$ .

Let  $T \subseteq V$ , |T| = r. We want to calculate the multiplicity of T as an edge of  $H_i$ ; i.e. we want to determine the number of those edges of  $H_i$ , which do not belong to  $H_i - t$  for any  $t \in T$ . By inclusion-exclusion, this number is

$$\sum_{X \subseteq T} (-1)^{|X|} |E(H_i - X)|.$$

Since  $k \ge r$ , this number is the same for i = 1, 2. Hence  $H_1$ ,  $H_2$  are identical [V. Faber, Hypergraph Seminar, Lecture Notes in Math. 411, Springer, 1974, 85–94].

19. Let  $v_i(X)$  denote the multiplicity of  $X \subseteq V$  as an edge of  $H_i$  (i = 1, 2). The assumption implies that

$$(1) v_1(X - \{x\}) + v_1(X) = v_2(X - \{x\}) + v_2(X),$$

since  $v_i(X - \{x\}) + v_i(X)$  is the multiplicity of the edge  $X - \{x\}$  in  $H_i \setminus x$ . Assume now that  $H_1 \neq H_2$ . Then there are sets  $X \subseteq V$  with  $v_1(X) \neq v_2(X)$ . Let X be a minimal such set. Then by (1),  $v_1(X - \{x\}) \neq v_2(X - \{x\})$  for any  $x \in X$ . This is only possible, if  $X = \emptyset$ . So we may assume  $v_1(\emptyset) > v_2(\emptyset)$ .

We prove by induction on |X| that

$$u_1(X) > \nu_2(X), \quad \text{if} \quad |X| \text{ is even,}$$
 $\nu_1(X) < \nu_2(X), \quad \text{if} \quad |X| \text{ is odd.}$ 

This follows easily from (1). In particular,

$$\nu_1(X) > \nu_2(X) \ge 0$$
, if  $|X|$  is even,

i.e. every even subset of V is an edge of  $H_1$ . Similarly, every odd subset of V is an edge of  $H_2$  [see C. Berge, Hypergraph Seminar, Lecture Notes in Math. 411, Springer, 1974, 1–12].

**20.** (a) Let hom (A, b) and epi (A, B) denote the numbers of homomorphisms of A into B and epimorphisms of A onto B, respectively. Observe first:

(\*) If 
$$\operatorname{epi}(G_1, H) = \operatorname{epi}(G_2, H)$$
 for every  $H$ , then  $G_1 \simeq G_2$ .

In fact, we have epi  $(G_1, G_2) = \text{epi } (G_2, G_2) > 0$ , epi  $(G_2, G_1) = \text{epi } (G_1, G_1) > 0$ , i.e.  $G_1, G_2$  are epimorphic images of each other. Since they are finite, they must be isomorphic.

So we want to relate  $hom(G_i, H)$  and  $epi(G_i, H)$ . Obviously, each epimorphism is a homomorphism and a homomorphism is an epimorphism iff every edge of H is in the image. So we can use the inclusion-exclusion formula and get

$$\operatorname{epi}(G_i, H) = \operatorname{hom}(G_i, H) - \sum_{e \in E(H)} \operatorname{hom}(G_i, H - e) +$$

$$+\sum_{\{e_1,e_2\}\subseteq E(H)} \mathrm{hom}\,(G_i,H-e_1-e_2)-\ldots =\sum_{S\subseteq E(H)} (-1)^{|S|} \mathrm{hom}\,(G_i,H-S).$$

Since the right-hand sides are the same for i = 1, 2 and any fixed H, we conclude that

$$\operatorname{epi}(G_1, H) = \operatorname{epi}(G_2, H)$$

holds for every graph H. By (\*), this implies that  $G_1 \simeq G_2$ .

(b) Let mon (A, B) denote the number of monomorphisms (= one-to-one homomorphisms) of A into B. Again,

(\*) if mon 
$$(H, G_1) = \text{mon}(H, G_2)$$
 for every  $H$ , then  $G_1 = G_2$ .

In fact, mon  $(G_2, G_1) = \text{mon } (G_1, G_1) > 0$ , mon  $(G_1, G_2) = \text{mon } (G_2, G_2) > 0$ , i.e.  $G_1, G_2$  have monomorphisms into each other, whence  $G_1 \cong G_2$  follows because they are finite.

To relate mon  $(H, G_i)$  and hom  $(H, G_i)$ , observe that a homomorphism is a monomorphism if and only if it does not identify any pair of points. Let  $\Theta((x_1, y_1), \ldots, (x_k, y_k))$  be the least equivalence relation such that  $x_i, y_i$  belong to the same class  $(i = 1, \ldots, k)$  and denote, for any simple graph H and equivalence relation  $\Theta$  on V(H), by  $H/\Theta$  the simple graph obtained by identifying the points in every given class of  $\Theta$  and forgetting about multiplicities of edges in the resulting graph. Then we can express  $\text{mon}(H, G_i)$  by inclusion-exclusion.

$$\begin{split} & \operatorname{mon}\left(H,G_{i}\right) = \operatorname{hom}\left(H,G_{i}\right) - \sum_{\{x_{1},y_{1}\} \subseteq V(H)} \operatorname{hom}\left(H/_{\Theta(x_{1},y_{1})},G_{i}\right) + \\ & + \sum_{\{x_{1},y_{1}\} \neq \{x_{2},y_{2}\} \subseteq V(H)} \operatorname{hom}\left(H/_{\Theta((x_{1},y_{1}),(x_{2},y_{2}))},G_{i}\right) - \ldots = \\ & \binom{|V(H)|}{2} \\ & \sum_{k=0} \left(-1\right)^{k} \sum_{\{x_{1},y_{1}\} \neq \ldots \neq \{x_{k},y_{k}\} \subseteq V(H)} = \operatorname{hom}\left(H/_{\Theta((x_{1},y_{1}),\ldots,(x_{k},y_{k}))},G_{i}\right). \end{split}$$

Again, the right-hand side is the same for i = 1 and 2, so mon  $(H, G_1) =$ =mon  $(H, G_2)$  and (\*) proves the assertion [L. Lovász, *Periodica Math. Hung.* 1 (1971) 145–156].

**21.** Let  $\varphi, \psi: H \to G_1$  be homomorphisms, and set

$$\xi(v) = (\varphi(v), \psi(v)) (v \in V(H)).$$

15.21

Then, as is easily verified,  $\xi$  is a homomorphism of H into  $G_1 \times G_1$ ; conversely, each homomorphism of H into  $G_1 \times G_1$  arises uniquely in this way. Hence

(1) 
$$\operatorname{hom}(H, G_1 \times G_1) = (\operatorname{hom}(H, G_1))^2.$$

Now we have

 $(\text{hom } (H, G_1))^2 = \text{hom } (H, G_1 \times G_1) = \text{hom } (H, G_2 \times G_2) = (\text{hom } (H, G_2))^2$ and, since hom  $(H, G_i) \ge 0$ ,

$$hom (H, G_1) = hom (H, G_2).$$

This holds for any H, so by 15.19b, we get  $G_1 \simeq G_2$ . [L. Lovász; ibid.].

22. (a) We try  $F = K_2$ . Then observe that, if  $v \in V(G_1)$ , then  $(v, x) \in V(G_1 \times F)$  has the same degree as v. Therefore,  $G_1$  and  $G_2$  must have the same degrees. A simple non-isomorphic pair with the same degree sequence is shown in Fig. 118. Forming the direct products with  $K_2$  (we set  $V(K_2) = \{1, 2\}$ ), we get two disjoint 6-cycles in each case, as shown in Fig. 119.





Fig. 118






Fig. 119

(b) Let H be any graph. For reasons similar to those in the preceding solution,

(1) 
$$\begin{aligned} & \operatorname{hom}\left(H,G_{1}\right)\operatorname{hom}\left(H,F\right) = \operatorname{hom}\left(H,G_{1}\times F\right) = \\ & = \operatorname{hom}\left(H,G_{2}\times F\right) = \operatorname{hom}\left(H,G_{2}\right)\operatorname{hom}\left(H,F\right). \end{aligned}$$

We want to show that

(2) 
$$hom(H, G_1) hom(H, F') = hom(H, G_2) hom(H, F').$$

If hom (H, F') = 0, then (2) is trivially satisfied. If hom (H, F') > 0 then, obviously, hom (H, F) > 0, so (1) implies that hom  $(H, G_1) = \text{hom } (H, G_2)$ , whence (2) follows again.

(c) The relation  $(G \cdot H)^0 = G^0 \times H^0$  formulated in the hint is easily verified. Then

$$G_1^0 {\times} H^0 = (G_1 \cdot H)^0 {\, \cong \,} (G_2 \cdot H)^0 = G_2^0 {\times} H^0.$$

Let  $E^0$  denote the graph with one point and one loop. Then  $E^0$  is a subgraph of  $H^0$  and so by (b)

$$G_1^0 \simeq G_1^0 \times E^0 \simeq G_2^0 \times E^0 \simeq G_2^0$$
.

(d) The assertion formulated in the hint follows by an argument similar to the one in 15.20b except that  $\Theta((x_1, y_1), \ldots, (x_k, y_k))$  must be interpreted as the least congruence relation containing the pairs  $(x_1, y_1), \ldots, (x_k, y_k)$ .

Now suppose that  $G_1, G_2, F$  are finite groups and

$$G_1 \times F \simeq G_2 \times F$$
.

Denoting the number of homomorphisms of the group H into the group G by hom (H, G), we have

$$hom(H, G_1) hom(H, F) = hom(H, G_2) hom(H, F),$$

for each finite group H. Since hom  $(H, F) \neq 0$ , this yields

$$hom (H, G_1) = hom (H, G_2),$$

whence  $G_1 \simeq G_2$ .

For finite rings the proof is the same. In fact, it is the same for any three algebras  $G_1, G_2, F$  provided F has a 1-element subalgebra. [For a generalization to categories see L. Lovász,  $Acta\ Sci.\ Math.\ 33\ (1972)\ 319-322.$ ]

## **Dictionary**

of the combinatorial phrases and concepts used

Abel identities: see 1.44.

Adjacency matrix of a graph G with vertices  $v_1, \ldots, v_n$ : the matrix  $A_G = (a_{ij})_{i,j=1}^n$ , where  $a_{ij}$  is the number of  $(v_i, v_j)$ -edges.

Adjacent: see graph.

Arborescence: a digraph G with a specified vertex a called the *root* such that each point  $x \neq a$  has indegree 1 and there is a unique (a, x)-path for each point x. Arborescences can be obtained by specifying a vertex a of a tree and then orienting each edge e such that the unique path connecting a to e ends at the tail of e. An *inverse*  $\sim$  is the digraph obtained from an arborescence by inverting its edges.

Automorphism of a [di]graph: a permutation  $\alpha$  of V(G) such that the number of (x, y)-edges is the same as the number of  $(\alpha(x), \alpha(y))$ -edges  $(x, y \in V(G))$ . We also speak of the automorphisms of a graph G with colored edges. This means a permutation  $\alpha$  such that the number of (x, y)-edges is the same as the number of  $(\alpha(x), \alpha(y))$ -edges with any given color. The set of all automorphisms of a [di]graph forms a permutation group A(G).

Balanced circuit in a hypergraph: a circuit  $(x_1, E_1, \ldots, x_k, E_k)$  such that either k = 2 or there is an incidence  $x_i \in E_j$ , where  $j \neq i, i - 1$  and  $(i, j) \neq (1, k)$ .  $\sim$  hypergraph: a hypergraph in which every circuit of odd length is balanced.

totally ~ hypergraph: every circuit is balanced.

Bell number: see partition.

Binomial coefficient  $\binom{n}{k}$ : the number of ways to select k elements out of n.

One has

(1) 
$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n(n-1)\dots(n-k+1)}{k!}, \qquad (0 \le k \le n)$$

and by definition  $\binom{k}{2}$ 

$$\binom{k}{0} = 1, \ (k = 0, 1, \ldots).$$

Formula (1) defines  $\binom{n}{k}$  for any real (or complex) value of n.

DICTIONARY 529

**Bipartite** (2-chromatic) graph: A graph G admitting a bipartition or 2-coloration  $\{A, B\}$  of its points such that each edge connects a point of A to a point of B (cf. chromatic number).

Block of a graph G: a cut-edge or a maximal 2-connected subgraph. Each edge is contained in a unique block. Blocks could also be defined as classes of the equivalence relation on E(G) defined by "e and f are on a circuit or e=f". The blocks of a graph give a "cactus-like" structure; each point belonging to more than one block is a cutpoint, and the number of branches relative to any point is the same as the number of blocks containing this point (Fig. 120). Blocks containing one cutpoint are called endblocks.




Fig. 120

**Branch** of a graph G relative to a point x: a subgraph consisting of a component  $G_1$  of G-x, the point x and all edges connecting x to  $G_1$ .

**Bridge** of a subgraph  $G_1$ : A (connected) subgraph B such that either B consists of a single edge with both endpoints in  $G_1$  or B consists of a connected component of  $G-V(G_1)$  together with all edges connecting this component to  $G_1$  and their endpoints in  $G_1$ . The bridges of  $G_1$  partition  $E(G)-E(G_1)$ ; i.e. they could also be defined as classes determined by the equivalence relation " $e_1 \sim e_2$  iff  $e_1 = e_2$  or there is a path connecting  $e_1$  and  $e_2$  disjoint from  $G_1$ ".

Brooks' Theorem: see 9.13.

Brun's Sieve: see 2.13.

Burnside Lemma: see 3.23.

Cayley Formula: see 4.2.

Characteristic polynomial of a graph G: The polynomial  $p_G(\lambda) = \det(\lambda I - A_G)$ , where  $A_G$  is the adjacency matrix of G. This clearly does not depend on the labelling of points. The roots of the characteristic polynomial, i.e. the eigenvalues of  $A_G$ , are called the *eigenvalues* of the graph G.

Chord of a subgraph  $G_1 \subseteq G$ : an edge  $e \in E(G) - E(G_1)$  connecting two points of  $G_1$ .

Chromatic index of a hyper graph G: the least integer k for which the edges of G can be k-colored such that adjacent edges have different colors. We denote it by q(G). Clearly  $q(G) = \chi(L(G))$ .

Chromatic number of a graph  $\lceil digraph, hypergraph \rceil G$ : the least integer k for which G has a "good k-coloration" (see below), we denote this number by  $\chi(G)$ . Clearly,  $\chi(G) > 0$ , if G is non-empty;  $\chi(G) > 1$ , if E(G) is non-empty.  $\chi(G) = 0$  $=\infty$  if G has a loop [or, when G is a hypergraph, it has an edge with less than 1 endpoints].

Chromatic polynomial  $P_G(\lambda)$  of a graph G: the number of good  $\lambda$ -colorations of  $G(\lambda = 0, 1, \ldots)$ . This turns out to be a polynomial in  $\lambda$  (for fixed G) and so, its definition can be extended to all real (or complex) values of  $\lambda$ . Note that two  $\lambda$ -colorations differing in the labelling of colors count as different.

Circuit in a graph: a walk  $(x_1, e_1, \ldots, x_k, e_k, x_{k+1})$  such that  $x_1, \ldots, x_k$  are distinct points,  $e_1, \ldots, e_k$  are distinct edges and  $x_1 = x_{k+1}$ . If the graph is simple, we will denote it by  $(x_1, \ldots, x_k)$ .  $\sim in \ a \ digraph$ : a circuit in the graph obtained from G by replacing each

oriented edge by an unoriented one with the same endpoints. Cf. cycle.

 $\sim$  in a hypergraph: a sequence  $(x_1, E_1, \ldots, x_k, E_k)$ , where  $x_1, \ldots, x_k$  are distinct points,  $E_1, \ldots, E_k$  are distinct edges and  $x_i \in E_i$   $(i = 1, \ldots, k)$ ,  $x_{i+1} \in E_i$  (i = 1, ..., k-1) and  $x_1 \in E_k$ . k is the length of this circuit.

Clique: a maximal complete subgraph of a graph.

Clutter: a hypergraph in which no edge contains another.

Collection: a set S together with an assignment of positive integers called multiplicities to the elements of S; anything not in S can be said to have multiplicity 0. If  $\mathcal{A}$ ,  $\mathcal{B}$  are collections then  $\mathcal{A} + \mathcal{B}$  is defined as the collection in which

the multiplicity of x is the sum of its multiplicities in  $\mathcal A$  and  $\mathcal B$ .  $m\mathcal A=\mathcal A+\ldots$ 

 $\dots + \mathcal{H}$ . m-collection: a collection in which each element has multiplicity less than m.  $m^X$  is the set of all m-collections of elements of X.

Coloration: a (legitimate, good) k-coloration of a graph [digraph, hypergraph] is an assignment of "colors" (usually one of the integers  $1, \ldots, k$ ) to the points such that the endpoints of every edge have different "colors".

k-colorable graph [digraph, hypergraph]: it has a good k-coloration.

Complement of a simple graph G: the simple graph  $\bar{G}$  defined by

$$V(\bar{G})=V(G),\; E(\bar{G})=\{(x,y):x,y\in V(G),x
eq y,(x,y)\notin E(G)\}.$$
 Clearly  $(\overline{\bar{G}})=G.$ 

$$\sim$$
 of a simple digraph  $G$ : the simple digraph  $\overline{G}$  defined by  $V(\overline{G}) = V(G), \qquad E(\overline{G}) = V(G) \times V(G) - E(G).$ 

Complete graph: a simple graph in which any two distinct points are adjacent. A complete graph on n points will be denoted by  $K_n$ .

DICTIONARY 531

Complete bipartite graph: a simple graph whose points can be partitioned into two classes U, W such that two points are adjacent iff one of them belongs to U and the other to W. If |U| = n, and |W| = m, then the complete bipartite graph is denoted by  $K_{n,m}$ .

**Component** (connected  $\sim$ ) of a graph G: a maximal connected subgraph of G. Any two connected components of G are vertex-disjoint and each vertex (and edge) belongs to one of them. Their number is denoted by c(G);  $c_1(G)$  denotes the number of those connected components with an odd number of points.

strong ~ (of a digraph): maximal strongly connected subgraph.

**Connected** graph: a graph, which is not representable in the form  $G_1 \cup G_2$ , digraph where  $G_1$  and  $G_2$  are vertex-disjoint non-empty graphs. Equivalently: any two points are connected by a path in the graph.

weakly  $\sim digraph$ : a digraph not representable as  $G_1 \cup G_2$ , where  $G_1, G_2$ 

are vertex-disjoint non-empty digraphs.

strongly ~ digraph: a digraph in which any two points are connected by

a (directed) path.

~ hypergraph: it is not representable as  $H_1 \cup H_2$ , where  $H_1$ ,  $H_2$  are vertex-disjoint non-empty hypergraphs. Note that if  $\emptyset \in E(H)$ , H is not connected.

k-connected between a and b: if less than k points ( $\neq a, b$ ) and/or edges are removed, there still exists an (a, b)-path in the [di]graph (removal of edges

is only needed when a, b are adjacent).

 $\sim$  [di]graph G: on at least k+1 points, k-connected between any two points. Equivalent formulations:  $|V(G)| \geq k+1$  and G-X is [strongly] connected for any set  $X \subset V(G)$ ,  $|X| \leq k-1$ . Or, for a graph: G cannot be represented as  $G_1 \cup G_2$ , where  $V(G_1)$ ,  $V(G_2) \neq V(G)$  and  $|V(G_1) \cap V(G_2)| \leq k-1$ . The complete graph  $K_n$  is thus (n-1)-connected but non n-connected. Connected and 1-connected are equivalent for graphs with at least two points.

**Contraction** of an edge e of a [di]graph: removal of this edge and identification of its endpoints. Contracting a subgraph means contracting all edges of it (the order in which the contraction is made is irrelevant). Note that multiple edges may appear.

k-cover of a [hyper]graph: a collection of points such that each edge contains at least k of them. (Point-) cover: 1-cover: A k-cover can also be regarded as a mapping  $t: V(G) \to \{0, 1, \ldots\}$  such that  $\sum_{x \in E} t(x) \ge k$  for each edge E.

fractional cover: an assignment of a non-negative real weight t(x) to each point x such that  $\sum_{x \in E} t(x) \ge 1$  for every edge E.

**Critical:** a graph G is called (edge-)critical with respect to property P, or critically having property P if G has it but, on removing any edge, the resulting graph will not have property P. Point-critical is defined analogously.

 $\alpha$ -critical:  $\alpha(G - \{e\}) > \alpha(G)$ , for each edge e.

 $\chi$ -critical:  $\chi(G - \{e\}) < \chi(G)$ , for each edge e  $\tau$ -critical:  $\tau(G - \{e\}) < \tau(G)$ , for each edge e.

v-critical hypergraph: v(H-x) = v(H), for each  $x \in V(G)$ .

factor-critical graph: G has no 1-factor but G-x has a 1-factor for each point x.

(a, b)-cut: a set F of edges representing (covering) all (a, b)-paths. The cut determined by  $S \subseteq V(G)$  is the set of edges connecting S to V(G) - S. If C is a cut in the digraph G determined by S', then  $C^*$  is the cut determined by V(G) - S.

Cutset (separating set): a set of points [edges] in a connected graph whose removal results in a disconnected graph. A cutpoint [cut-edge or isthmus] is a point [edge] forming a cutset by itself.

Cycle in digraph: a walk  $(x_1, e_1, \ldots, e_k, x_{k+1})$ , where  $x_1, \ldots, x_k$  are distinct and  $x_{k+1} = x_1$ .

Cycle index of a permutation group  $\Gamma$ : the polynomial

$$p_{\Gamma}(x_1,\ldots,x_n)=\sum_{n\in\Gamma}x_1^{k_1(n)}\ldots x_n^{k_n(n)},$$

where n is the number of objects  $\Gamma$  acts on and  $k_i(\pi)$  is the number of i-cycles in the cycle decomposition of the permutation  $\pi$ .

**Degree** (valency) of a point x in a [hyper]graph G: the number of edges containing x (in the case of graphs a loop is counted twice). The degree of x is denoted by  $d_G(x)$ . d(G) denotes the maximum degree of G. A graph is k-regular if every point has degree k. The indegree [outdegree] of a point x in a digraph G is defined as the number of edges with their head [tail] in x; they are denoted by  $d_G(x)$  and  $d_G^+(x)$ , respectively.

Diameter of a graph G: the maximum distance between points of G.

**Digraph:** a set V(G) of points or vertices, a set E(G) of edges, and an assignment of an ordered pair of points with each edge; the first and second elements of this pair are called the tail and the head of the edge, respectively. Two edges are parallel, if they have the same heads and tails. The digraph is simple, if it contains no two parallel edges. In this case, E(G) can be considered as a subset of  $V(G) \times V(G)$ . If G is a graph and for each edge we declare one of its endpoints to be its head, the other one its tail, we obtain a digraph called an orientation of G. If  $e = (x, y) \in E(G)$  we use either of the phrases: e is directed from x to y; y can be reached from x on e; e leaves x and enters y. Cf. also graph.

**Distance** between two points x, y in a graph G: the minimum length of (x, y)-paths; if no path of G connects x to y, their distance is  $\infty$ . It is denoted by  $d_G(x, y)$  — or shortly d(x, y), if the graph G under consideration is well-determined.

**Dual** map of a connected planar map G: the map  $G^*$  constructed as follows. We select a point  $x_F$  in each of the faces F of G; these will be the vertices of  $G^*$ . Also we select a point  $p_e$  on each edge e of G. We connect each point  $p_e$  to the points  $x_F$ ,  $x_{F'}$ , by Jordan curves  $J_e$ ,  $J'_e$  interior to F and F', respectively, where F, F' are the two faces adjacent to e. If F = F' (i.e. the same face of G bounds e from both sides), then  $J_e$ ,  $J'_e$  should connect  $p_e$  to  $x_F$  such that they leave  $p_e$  on different sides of e (this happens, if e is a cutting edge). Moreover, let us choose  $J_e$ ,  $J'_e$  such that the arcs  $J_e$  connecting  $x_F$  to points  $p_e$  on the boundary of F should have no point in common other than  $x_F$ . Set  $e^* = J_e \cup J'_e$  and  $E(G^*) = e^* : e \in E(G)$ . Then  $G^*$  is a planar map and is, essentially, uniquely defined,

533 DICTIONARY

i.e. if  $\hat{G}^*$  is another dual map of G, then there is a homeomorphism  $\alpha$  of the plane onto itself such that  $\varphi(x) = x$  for each  $x \in V(G)$ ,  $\varphi(e) = e$  for each  $e \in E(G)$ ,  $\varphi(V(G^*)) = V(\hat{G}^*)$  and if  $\hat{e}^*$  is the edge corresponding to  $e^*$  in  $\hat{G}^*$ , then  $\varphi(e^*) =$  $=\hat{e}^*$ . The dual of  $G^*$  is G. The above construction and these last assertions involve much from plane topology that we accept here without proof (Fig. 121).




Fig. 121

Edge: see graph, digraph, hypergraph.

Edge-cover of a [hyper]graph: a set of edges containing all points.

k-edge-connected between a and b: the removal of no more than k-1 edges results in a [di]graph containing an (a, b)-path.

~ [di]graph: k-edge connected between any two points. Equivalently: the removal of no more than k-1 edges results in a [strongly] connected [di]graph.

Edmonds' Matching Algorithm: see 7.34.

Elementary bipartite graph: see 7.7.

Empty [hyper]graph: has no points and no edges.

**Endomorphism** of a [di]graph G: a homomorphism of G into itself. The set of all endomorphisms of G with the composition as multiplication, form a semigroup, denoted by  $\operatorname{End}(G)$ .

**Endpoint** of a graph G: a point with degree 1. ~ of an edge: see graph.

Erdős-de Bruijn Theorem: see 8.14.

Erdős-Ko-Rado Theorem: see 13.28.

Erdős-Stone Theorem: see 10.38.

Euler-trail: a trail containing all edges of a [di]graph.

Eulerian [di]graph: it has an Euler-trail.

534 DICTIONARY

Euler's Formula: see 5.24.

Face: see planar map.

**f-factor** of a graph: given a function f defined on V(G) an f-factor is a spanning subgraph G' such that  $d_{G'}(x) = f(x)$  for each point x. A 1-factor is, therefore, a system of independent edges covering all points.

Ferrer's diagram: see 1.16.

(a, b)-flow: a non-negative real valued function f defined on the edges of a digraph such that, for each point  $x_0 \neq a, b$ ,

$$\sum_{e=(x_0,y)\in E(G)} f(e) = \sum_{e=(y,x_0)\in E(G)} f(e)$$

(i.e. the amount of "water" entering  $x_0$  is the same as the amount leaving it;

"Kirchhoff's Law"). The value of an (a, b)-flow f is

$$w(f) = \sum_{e=(a,x)} f(e) = -\sum_{e=(x,b)} f(e)$$

Forest: a graph without circuits. The components of a forest are trees.

Frucht's Theorem: see 12.5.

Gallai-Edmonds Structure Theorem: see 7.32.

Generating function of a sequence  $\{a_n\}_{n=0}^{\infty}$ : the function

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

**exponential** 
$$\sim$$
:  $f(x) = \sum_{n=0}^{\infty} \frac{a_n}{n!} x^n$ .

Geometric lattice: a lattice L such that L is generated by its atoms and whenever x covers  $x \wedge y$ ,  $x \vee y$  covers y. The rank r(x) is defined as the maximum length of (0, x)-chains less one. This function satisfies:  $r(x) \geq 0$ ,  $x > x' \rightarrow r(x) \geq r(x')$ , if x covers y, then  $r(y) \leq r(x) \leq r(y) + 1$ , and  $r(x \vee y) + r(x \wedge y) \leq r(x) + r(y)$ . Examples of such lattices are the lattices of subspaces of any subset of a projective or affine plane.

Girth of a graph G: the length of its shortest circuit. The girth is 1 iff G has a loop and it is 2 iff G has multiple edges.

**Graph:** a graph G consists of a finite set V(G) of points (vertices) and a finite set E(G) of edges, and an assignment of an unordered pair of elements of V(G) to each edge  $e \in E(G)$ , called the endpoints of e. We write G = (V(G), E(G)). An edge is said to join or connect its endpoints. If e connects x to y then e is called an (x, y)-edge. An edge with two identical endpoints is a loop. Two edges with the same pair of endpoints are parallel or multiple. The graph is simple if it has no loops or parallel edges. In this case we may consider E(G) as a set of 2-subsets of V(G). An edge and a point are adjacent (incident) if the point is an endpoint of the edge. Two edges are adjacent if they share an endpoint. Two

points are adjacent (neighboring) if they are connected by an edge. The set of edges adjacent to a point is its star. The set of points adjacent to  $X \subseteq V(G)$  is denoted by  $\Gamma_G(X)$  or simply  $\Gamma(X)$  if the graph in question is clear from the context. Graphs without loops are special hypergraphs (cf. also digraph).

Hajós' Construction: see 8.16.

**Hamiltonian circuit** [cycle, path]: a circuit [cycle, path] containing all points of a [di]graph.

Hamiltonian [di]graph: it has a Hamiltonian circuit [cycle].

**Homomorphism** of  $[di]graph\ G_1$  into  $[di]graph\ G_2$ : a mapping  $\varphi:V(G_1)\to V(G_2)$  such that if  $(x,y)\in E(G_1)$ , then  $(\varphi(x),\ \varphi(y))\in E(G_2)$ .

**Hypergraph** (set-system): a hypergraph H consists of a finite set V(H) of points (vertices), a finite set E(H) of edges and an assignment of a subset of V(H) to each edge E, the set of endpoints (elements) of E. So we write H = (V(H), E(H)). Two edges with the same endpoints are parallel. The hypergraph is simple, if it contains ro parallel edges. In this case, E(H) can be considered as a set of subsets of V(H). An edge and a point are called incident, if the point is an endpoint of the edge. The hypergraph is r-uniform, if every edge has r endpoints. A 2-uniform hypergraph is a graph without loops. The complete r-uniform hypergraph on n vertices is a simple hypergraph having all r-subsets of its vertex set for its edges. It is denoted by  $K_r^r$ .

**Identification** of two points x, y of a [di]graph G results in a [di]graph G' with  $V(G') = V(G) - \{x, y\} \cup \{\overline{xy}\}$ , where  $z = \overline{xy}$  denotes a new point, E(G') = E(G) and each edge  $e \in E(G)$  should have the same endpoints in G' as in G except if it has had X or Y as endpoint it should have  $\overline{xy}$  instead. Thus each (x, y)-edge becomes a loop in  $\overline{xy}$ .

**Incidence matrix** of a graph G with vertices  $v_1, \ldots, v_n$  and edges  $e_1, \ldots, e_m$ : the matrix  $B_G = (b_{ij})_{i=1}^n \sum_{j=1}^m v_j$ , where  $b_{ij} = 1$ , if  $v_i$  and  $e_j$  are adjacent and 0 otherwise.

Incident: see graph, hypergraph.

Inclusion-exclusion Formula: see §2.

Independent points in a [di] graph: no two are adjacent. The maximum number of independent points in G is denoted by  $\alpha(G)$ .

 $\sim$  edges in a [hyper, di]graph: no two have an endpoint in common. The maximum number of independent edges in G is denoted by  $\nu(G)$ . Independent edges from a matching.

 $\sim$  paths in a [di]graph: paths having no points in common except possibly their endpoints.

Induced subgraph: see subgraph.

Interval graph: a simple graph whose vertices are intervals on a line, and two vertices are adjacent iff they intersect as intervals.

Isolated point of a graph: a point adjacent to no edge.

**Isomorphism** of  $G_1$  onto  $G_2$ : a one-to-one mapping  $\varphi$  of  $V(G_1)$  onto  $V(G_2)$  and a one-to-one mapping  $\tilde{\varphi}$  of  $E(G_1)$  onto  $E(G_2)$  such that if x is an endpoint [head, tail] of e, then  $\varphi(x)$  is an endpoint [head, tail] of  $\tilde{\varphi}(e)$ . If  $G_1, G_2$  are simple, then  $\varphi$  plays no important role and we define an isomorphism as a one-to-one mapping  $\varphi$  of  $V(G_1)$  onto  $V(G_2)$  such that  $(x, y) \in E(G_1) \leftrightarrow (\varphi(x), \varphi(y)) \in E(G_2)$ 

Isthmus: see cut-edge.

**Kernel:** an independent set S of points in a digraph such that, for each  $x \in V(G) - S$ , there exists a  $y \in S$  such that  $(x, y) \in E(G)$ .

König's Theorem: see 7.2.

Kuratowski's Theorem: see 5.38.

Line-graph of a [hyper]graph G: a simple graph L(G) defined by

$$V(L(G)) = E(G),$$

 $E(L(G)) = \{(e, f) : e, f \in E(G), e, f, \text{ have an endpoint in common}\}.$   $\sim of a digraph G: a simple digraph <math>L(G)$  defined by

$$V(L(G)) = E(G)$$

 $E(L(G)) = \{(e, f) : e, f \in E(G), \text{ the head of } e \text{ is the tail of } f\}.$ 

Loop: see graph.

**Matching:** a k-matching in a hypergraph G is a collection of edges of G such that each point belongs to at most k of them (note that repetition of edges is allowed). A 1-matching is also called a matching. A k-matching can be considered as a mapping  $w: E(G) \to \{0, 1, \ldots\}$  such that  $\sum w(E) \leq k$  for every point

x (w(E) is the multiplicity with which E occurs in the matching). A perfect k-matching is a k-matching such that each edge belongs to exactly k members of it (note the difference between this and a k-factor!). A fractional matching is an assignment of a non-negative real weight w(E) to each edge E such that  $\sum_{E\ni x} w(E) \le 1$  for every point x.

Max-Flow-Min-Cut Theorem: see 6.74.

Maximum [minimum] means maximal [minimal] in cardinality.

Maximal [minimal] means maximal [minimal] with respect to inclusion.

Menger's Theorem: see 6.39.

Moebius function: see 2.22.

Moebius Inversion Formula: see 2.26.

Orientation: see digraph.

Parallel edges: see graph, digraph, hypergraph.

Partial hypergraph of a hypergraph H: a hypergraph H' with  $V(H') \subseteq V(H)$ ,  $E(H') \subseteq E(H)$ .

**Partition** of a set S: a system  $\{A_1, \ldots, A_k\}$  of disjoint non-empty subsets of S (called *classes* of the partition) such that  $A_1 \cup \ldots \cup A_k = S$ . The number  $B_n$  of partitions of an n-element set is called a Bell-number.

 $\sim$  of a number n: a collection  $\{a_1,\ldots,a_k\}$   $(a_1\geq\ldots\geq a_k)$  of positive inte-

gers such that  $a_1 + \ldots + a_k = n$ .

**Path** in a [di]graph: a walk  $(x_1, e_1, \ldots, e_k, x_{k+1})$ , where  $x_1, \ldots, x_{k+1}$  are distinct points. It can be denoted by  $(x_1, \ldots, x_{k+1})$ , if the [di]graph is simple. (X, Y)-path: a path in a [di]graph, connecting a point of X to a point of Y

and having no other point in common with  $X \cup Y$ .

**Perfect graph:** a simple graph G such that each induced subgraph G' of Gsatisfies

 $\omega(G') = \gamma(G').$ 

Perfect Graph Theorem: see 13.57.

**Permanent** of a matrix  $(a_{ij})_{i=1}^n \stackrel{n}{\underset{i=1}{\longrightarrow}} :$ 

$$\operatorname{per} A = \sum_{\pi} a_{1\pi(1)} \dots a_{n,\pi(n)},$$

where  $\pi$  ranges over all permutations of  $\{1, \ldots, n\}$ 

**Permutation** of a set  $\Omega$ : a one-to-one mapping of  $\Omega$  onto itself. The number of permutations of an n-element set is  $n! = 1 \cdot 2 \cdot \dots \cdot n$ . The identity permutation is denoted by 1. If  $\gamma$  is a permutation of  $\Omega$ ,  $x \in \Omega$  and  $\gamma(x) = x$ , then x is called a fixed point of  $\gamma$ .

cyclic  $\sim$ : identify two orderings  $(x_1, \ldots, x_n), (y_1, \ldots, y_n)$  of the same set S, if  $y_1 = x_{k+1}, \ldots, y_{n-k} = x_n, y_{n-k+1} = x_1, \ldots, y_n = x_k$  for some  $1 \le k \le n$ , A class of orderings identified in this way is a cyclic permutation.

~ group: a group  $\Gamma$  such that to each  $\gamma \in \Gamma$  a permutation  $\tilde{\gamma} \in \Gamma$  of a finite set  $\Omega$  is assigned such that  $\gamma\delta(x) = \tilde{\delta}(\tilde{\gamma}(x))$   $(\gamma, \delta \in \Gamma)$ . If no confusion can arise we will assume the elements of  $\Gamma$  are permutations themselves, the same permutation occurring possibly several times. If  $\tilde{\gamma} = \delta$  implies  $\gamma = \delta$ , or, equivalently,  $\tilde{\gamma} \neq 1$  for  $\gamma \neq 1$ , the permutation group is called *effective*. If for any pair  $x, y \in \Omega$  there is a  $\gamma \in \Gamma$  such that  $\gamma(x) = y$  the group is transitive. If, for any pair  $x, y \in \Omega$ , there is at most one  $\gamma \in \Gamma$  with  $\gamma(x) = y$ , or, equivalently, no  $\gamma \in \Gamma$ ,  $\gamma \neq 1$  has fixed points, the permutation group is semiregular. If it is both semiregular and transitive it is called regular. In this case  $|\Gamma| = |\Omega|$ and the elements of  $\Omega$  can be identified with the elements of  $\Gamma$  such that  $\gamma(\delta) =$  $=\delta\gamma$  for all  $\gamma, \delta\in\Gamma$ .

Petersen graph: see Fig. 9, p. 71.

**Pfaffian** of a skew-symmetric matrix  $(a_{ij})_{i=1}^{2n} \stackrel{2n}{j=1} = A$  (i.e.  $a_{ii} = 0$ ,  $a_{ij} = -a_{ji}$ ):

$$\operatorname{Pf} A = \sum \varepsilon_{i_1 j_1, \dots, i_n j_n} a_{i_1 j_1}, \dots, a_{i_n j_n},$$

where  $\varepsilon_{i_1,j_1;\ldots;i_nj_n}$  is the sign of the permutation

$$\begin{pmatrix} 1 & 2 & \dots & 2n-1 & 2n \\ i_1 & j_1 & \dots & i_n & j_n \end{pmatrix}$$

and the summation is taken over all partitions of  $\{1, \ldots, 2n\}$  of the form  $\{\{i_1, j_1\}, \ldots, \{i_n, j_n\}\}$ . It is easy to see that the term corresponding to partition  $\{\{i_1, j_1\}, \ldots, \{i_n, j_n\}\}\$  does not depend on the order of classes and/or the order of the two elements within a class.

**Planar** map: a graph whose vertices are points in the plane and whose edges are Jordan curves in the plane (ending in their corresponding graphical endpoints) with no point in common except their endpoints. A connected component of the set obtained by removing the edges and vertices of a planar map from the plane is called a *face* (region, country). The boundary of a face is always the union of certain edges; if the map G is 2-connected as a graph, the boundary of each face is a closed Jordan curve, consisting of the edges of a circuit in G.

 $\sim$  graph: a graph G isomorphic to a planar map. Such a planar map is called an *embedding* of G in the plane.

Point: see graph, digraph, hypergraph.

Pólya's Enumeration Method: see 3.26-30.

**Product** of two simple [di]graphs: we consider three kinds of products: (weak) direct  $\sim G_1 \times G_2$  of  $G_1$  and  $G_2$ , defined by

$$V(G_1 \times G_2) = V(G_1) \times V(G_2),$$

$$E(G_1 \times G_2) = \{((x_1, x_2), (y_1, y_2)) : (x_1, y_1) \in E(G_1), (x_2, y_2) \in E(G_2)\}.$$

strong direct  $\sim G_1 \cdot G_2$  of  $G_1$  and  $G_2$ , defined by

$$V(G_1 \cdot G_2) = V(G_1) \times V(G_2),$$

 $E(G_1 \cdot G_2) =$ 

 $=\{((x_1,x_2),(y_1,y_2)):(x_1,y_1)\in E(G_1)\text{ and }(x_2,y_2)\in E(G_2),\text{ or }x_1=y_1\text{ and }(x_2,y_2)\in E(G_2),\text{ or }(x_1,y_1)\in E(G_1)\text{ and }x_2=y_2\}.$ 

Cartesian  $\sim G_1 \oplus G_2$ , defined by

$$V(G_1 \oplus G_2) = V(G_1) \times V(G_2),$$

$$E(G_1 \oplus G_2) = \{ ((x_1, x_2), (y_1, y_2)): x_1 = y_1 \text{ and } (x_2, y_2) \in E(G_2); \\ \text{or } (x_1, y_1) \in E(G_1) \text{ and } x_2 = y_2) \}.$$

Thus  $G_1 \cdot G_2 = (G_1 \times G_2) \cup (G_1 \oplus G_2)$  (see Fig. 122).  $\sim$  of two hypergraphs  $H_1, H_2$ : the hypergraph  $H_1 \times H_2$  defined by

$$V(H_1 \times H_2) = V(H_1) \times V(H_2),$$

 $E(H_1 \times H_2) = \{E_1 \times E_2 : E_1 \in E(H_1), E_2 \in E(H_2)\}.$ 




Fig. 122

Prüfer code: see 4.5.

Pseudosymmetric digraph: see symmetric.

Ramsey's theorem: see § 14.

Regular graph: see degree; ~ group: see permutation group.

**Removal** of a set  $X \subseteq V(G)$  from a [hyper-, di-]graph G: removal of points in X together with all edges adjacent to them. The resulting [hyper-, di-]graph is denoted by G - X; if  $X = \{x\}$  we write simply G - x.

**Restriction** of a hypergraph H onto  $X \subseteq V(H)$ : the hypergraph  $H_X$  on the set X, for which  $E(H_X)$  is the collection of sets  $E \cap X$  ( $E \in E(H)$ ). If X = V(H) - Y then we adopt the further notation  $H_X = H \setminus Y$  and  $H_X = H - Y$ , if  $Y = \{y\}$ .

Rigid graph: has no proper endomorphism.

Selberg Sieve see 2.14-17.

Semiregular group: see permutation group.

**Separate:** a set X of points and edges is said to separate A and  $B(A, B \subseteq V(G))$ , if it represents (covers) all (A, B)-paths (cf. also *cutset*).

Sieve Formula: see § 2.

Simple graph, digraph, hypergraph: see graph, digraph, hypergraph.

Spanning subgraph of G: subgraph G' such that V(G') = V(G).

**Spectrum** of a graph G: The spectrum (collection of eigenvalues) of the adjacency matrix  $A_G$  of G. Since  $A_G$  is symmetric, the eigenvalues of G (the elements of its spectrum) are real.

Sperner's Lemma: see 5.29.

Sperner's Theorem: see 13.21.

Splitting a point x of a graph G into points  $x_1, \ldots, x_k$ : we remove x and replace each (x, y)-edge  $(y \in V(G) - \{x\})$  by an  $(x_i, y)$ -edge for exactly one  $i, 1 \le i \le k$ . Star: a tree with one point connected to all other points.

∼ of a vertex of a graph: see graph.

Stirling number S(n, k) (first kind): the coefficient of  $x^k$  in the expansion of the product  $x(x-1) \dots (x-n+1)$ .

 $\sim S(n, k)$  (second kind): the number of partitions of n objects into exactly k classes.

Subdivision of a graph G: a graph G' arising from G by replacing each edge e by a path  $P_e$  (of lengths  $\geq 1$ ), connecting the endpoints of e and having no other point in G, such that the paths  $P_e(e \in E(G))$  are independent. We call the points of G principal points of G'.

Subgraph of G: a graph G' with  $V(G') \subseteq V(G)$  and  $E(G') \subseteq E(G)$ . We write  $G' \subseteq G$ .

 $\sim$  of G induced by a set  $X \subseteq V(G)$ : the graph G[X] with V(G[X]) = X,  $E(G[X]) = \{e \in E(G) : e \subseteq X\}$ .

Substitution of a graph G for a point x of a graph H (supposing G and H are vertex-disjoint): we remove x from H, and replace each (x, y)-edge  $(y \in V(H) - \{x\})$  by the |V(G)| edges connecting y to the points of G.

Symmetric digraph: a simple digraph such that  $(x, y) \in E(G)$ , whenever  $(y, x) \in E(G)$ . Pseudosymmetric:  $d^+(x) = d^-(x)$  at every point.

System of distinct representatives of a hypergraph H: a one-to-one mapping  $\varrho$ :  $E(H) \to V(H)$  such that  $\varrho(E) \in E$  for each  $E \in E(H)$ . If no confusion can arise, we also call the range  $\varrho(E(H))$  a system of distinct representatives.

**Tournament:** a (simple) digraph T without loops in which exactly one of (x, y) and (y, x) is an edge for every pair  $x \neq y$ ,  $x, y \in V(T)$ .

Trail: see walk.

**Transitive** tournament: a tournament T such that  $(x, y) \in E(T)$  and  $(y, z) \in E(T)$  imply  $(x, y) \in E(T)$ . The points of a transitive tournament have an ordering  $(x_1, \ldots, x_n)$  such that  $(x_i, x_j) \in E(G) \leftrightarrow i < j$ .

~ permutation group: see permutation group.

Tree: a connected graph without circuits. It may also be defined as a connected graph such that removing any edge disconnects it; or as a circuit-free graph in which the introduction of any new edge will produce a circuit. A tree on n points has exactly n-1 edges and it always has at least two points of degree 1, provided  $|V(G)| \ge 2$ .

**Triangulation** (planar): a planar map in which each face is a triangle.

 $\sim$  of a circuit C: a graph consisting of this circuit and n-3 non-crossing "interior" diagonals (n is the length of C).

Turán's Theorem: see 10.34.

Tutte's Theorem: see 7.27.

Uniform: see hypergraph.

Valency: see degree.

Vertex: see graph, digraph, hypergraph.

Walk in a [di]graph: a sequence  $(x_1, e_1, \ldots, x_k, e_k, x_{k+1})$  in which  $x_1, \ldots, x_k$  are points and  $e_i$  is an  $(x_i, x_{i+1})$ -edge  $(i = 1, \ldots, k)$ . If the [di]graph is simple we may describe the walk by the sequence  $(x_1, \ldots, x_{k+1})$ . The walk is open [closed] iff  $x_{k+1} \neq x_1[x_{k+1} = x_1]$ . The length of the walk is k above. A walk is a trail if no edge is used twice.

Wheel: a graph obtained from a circuit by connecting all points to a new point (the "center" of the wheel).

O-graph: a graph consisting of three independent paths connecting two points.

# Notation

$A_G$ :	adjacency matrix of graph $G$ .
A(G):	automorphism group of $G$ .
$B_G$ :	incidence matrix of $G$ .
c(G):	number of components of $G$ .
$c_1(G)$ :	number of odd components (i.e. components with an odd num-
$c_1(\sigma)$ .	ber of points) of $G$ .
$d_G(x)$ :	degree of point $x$ in $G$ .
d(G):	maximum degree in $G$ .
$d_G(x, y)$ :	distance between points $x, y$ in $G$ .
$d_G^+(x)$ :	outdegree of point $x$ in $G$ .
$d_{\overline{G}}(x)$ :	indegree of point $x$ in $G$ .
E(G):	set of edges of $G$ .
$\operatorname{End}(G)$ :	endomorphism semigroup of $G$ .
$\exp x$ :	$e^{x}$ .
$\mathbf{j}[J]$ :	a vector [matrix] all entries of which are 1's.
$k_i(\pi)$ :	number of i-cycles in permutation $\pi$ .
$K_n^r$ :	complete r-uniform hypergraph on n points. For $r = 2$ the super-
III.	script is omitted.
L(G):	line-graph of $G$ .
O(f(n)):	a function $g(n)$ such that $g(n)/f(n)$ is bounded.
o(f(n)):	a function $g(n)$ such that $g(n)/f(n) \to 0$ as $n \to \infty$ .
$p_G(\lambda)$ :	characteristic polynomial of $G$ .
$P_G(\lambda)$ :	chromatic polynomial of $G$ .
$p_{\Gamma}(x_1,,x_n)$ :	cycle index of permutation group $\Gamma$ .
per $A$ :	permanent of matrix A.
Pf A:	Pfaffian of matrix A.
q(G):	chromatic index of $G$ .
s(n, k),	
$\hat{S}(n, k)$ :	Stirling numbers of the first and second kind, respectively.
V(G):	set of points of $G$ .
<b>Z</b> :	the set of integers.
$\delta_G(X)$ :	number of $(X, V(G) - X)$ -edges in $G$ .
$\alpha(G)$ :	maximum number of independent points.
$\Gamma_G(X)$ :	set of points in graph G adjacent to at least one point of $X \subseteq V(G)$ .
$\nu(G)$ :	maximum number of independent edges in G (matching number).
$\varrho(G)$ :	minimum number of edges of G covering all points.

542

A + B:

moA:  $m^S$ :

#### NOTATION

minimum number of points of G representing all edges.  $\tau(G)$ :  $\gamma(G)$ : chromatic number of G. maximum number of points in a clique (or in a complete sub- $\omega(G)$ : graph) of G.  $d(x), d^{+}(x),$ abbreviations for  $d_G(x)$ ,  $d_G^+(x)$ ,  $\Gamma_G(x)$  in the case when the graph  $\Gamma(X)$ , etc.: G is understood. integer part of x: largest integer not greater than x. [x]least integer not less than x.  $[x]^*$ : (where G is a graph (digraph, hypergraph) and  $F \subseteq E(G)$ ) removal G-Fof all edges in  $\hat{F}$  (but removing no points). (where  $f \in E(G)$ ) shorthand for  $G - \{f\}$ , when no confusion can G-farise. G: the complement of [di]graph G.  $G_1 \cup G_2$ : the [di]graph defined by  $V(G_1 \cup G_2) = V(G_1) \cup V(G_2)$ ,  $E(G_1 \cup G_2) = V(G_1) \cup V(G_2)$  $\bigcup G_2 = E(G_1) \cup E(G_2)$  (here  $G_1, G_2$  may have points or edges in (where G is a graph [digraph, hypergraph] and  $X \subseteq V(G)$ ) G-X: removal of points in X and all edges adjacent to them. G-x: (where  $x \in V(G)$ ) shorthand for  $G - \{x\}$ . (where H is a hypergraph and  $X \subseteq V(H)$ ) the restriction of H  $H \setminus X$ : onto V(H) - X.  $H \setminus x$ : (where  $x \in V(H)$ ) shorthand for  $H \setminus \{x\}$ . the subgraph of G induced by  $X \subseteq V(G)$ . G[X]the restriction of hypergraph H onto  $X \subseteq V(H)$ .  $H_X$ : G/F. (where  $F \subseteq E(G)$ ) the graph arising from graph G by contracting all edges in F. G/f: (where  $f \in E(G)$ ) shorthand for  $G/\{f\}$ .  $G_1 \times G_2$ : (weak) direct product of [di]graphs  $G_1, G_2$ .  $G_1 \cdot G_2$ : strong direct product.  $G_1 \oplus G_2$ : Cartesian product.  $H_1 \times H_2$ : direct product of hypergraphs  $H_1$ ,  $H_2$ .

sum of the collections A, B.

(where  $\mathcal{A}$  is a collection)  $\mathcal{A} + \ldots + \mathcal{A}$ .

the set of all m-collections of elements of S.

## Index

## of the abbreviations of textbooks and monographs

- B: C. Berge, Graphs and Hypergraphs, North-Holland—American Elsevier, 1973.
- Biggs: N. Biggs, Algebraic Graph Theory, Cambridge Univ. Press, 1974.
   ES: P. Erdős—J. Spencer, Probabilistic Methods in Combinatorics, Akadémiai Kiadó, Budapest, 1974.
- Fe: W. Feller, An introduction to Probability Theory and its Applications, 2<sup>nd</sup> ed., Wiley, New York-Chapman Hall, London 1957.
- FF: L. R. Ford—D. R. Fulkerson, Flows in Networks, Princeton Univ. Press, 1962.
- H: F. Harary, Graph Theory, Addison-Wesley, 1969. Hall: M. Hall, Combinatorial Theory, Blaisdell, 1967.
- Hu: T. C. Hu, Integer Programming and Network Flows, Addison-Wesley, 1969.
- K: D. König, Theorie der endlichen und unendlichen Graphen, Leipzig, 1936.
- M: J. W. Moon, Topics on Tournaments, Holt, Rinehart and Wilson, 1968.
- Mi: L. Mirsky, Transversal Theory, Academic Press, 1971.
- O: O. Ore, Theory of Graphs, Amer. Math. Soc. Coll. Publ., 1962.
  OF: O. Ore, The Four Color Problem, Academic Press, 1967.
- OF: O. Ore, The Four Color Problem, Academic Press, 1967.
  R: J. Riordan, An introduction to Combinatorial Analysis, Wiley, 1958.
- Ré: A. Rényi, Foundations of Probability, Holden-Day, San Francisco-Cambridge-London-Amsterdam, 1970.
- S: H. Sachs, Einführung in die Theorie der endlichen Graphen, Teil I-II., Teubner, 1970–1972.
- W: K. Wagner, Graphentheorie, Bibliographisches Inst. AG. 1970.
- Wi: R. J. Wilson, Introduction to Graph Theory, Oliver & Boyd, 1972,
   WV: H. Walther-H.-J. Voß, Über Kreise in Graphen, VEB Deutscher Verlag der Wiss., 1974.



# Subject Index

Italics refer to main occurrences.

### Examples:

## Explanation:

5.18, 20, 22-26

5.0 87

\$5, Problems 18, 20 and 22 through 26, and/or the Hints or the Solutions to these problems Introduction to §5

Most of chapter 7 is relevant

Abel identities 1.44: 4.6 affine space 14.23 alternating path 7.3,8,26,34 - circuit 4.24-29; 7.28,31 analytic function 4.20 arborescence 2.23,33; 4.15-17 spanning - 4.15,16; 5.10,11 atom of a Boolean algebra 2.2,6 - of a lattice 2.28,37 automorphism group 4.17; 8.15; 11.8-10,

16,17; §12; 13.22; 15.2 averaging method 13.0,12,41

Bell-numbers  $(P_n)$  1.9,10,12 Bernstein's Theorem 13.6 Binet-Cauchy formula 4.9,10 bipartite graph 4.11,21,31; 5.1,3,22; 6.5; 7.1–21,40; 8.16; 9.30; 10.15; 11.11,19,22; 13.1,10,50 - elementary graph 7.7-9,15 block 5.35; 6.0; 9.40,55; 10.1,5 Bondy's Condition 10.21 Borsuk's Theorem 9.27 breaking a stick 1.37 bridge 6.69,70 Bruck's Theorem 9.13 Brun's Sieve 2.13

cancellation law 15.22 Cauchy Integral Formula 1.15; 4.7 Cauchy's Theorem 12.7 Cayley Formula 1.15; 4.2-9,14,18 chain 2.25,34-37; 13.20-24; 14.28 character 11.8,10,17

Burnside's Lemma 3.24-29

characteristic polynomial 1.29; §11 characterization, good 5.0 Chebyshev's Inequality 2.19 choice function 14.28 chord 5.37; 6.35,70; 8.17; 9.24,28,29; 10.2 chromatic index 7.10; 13.56 - number (see also coloration) §9; 10.35,38,39; 11.20,21; 13.33-47 - polynomial 9.0,36-49

root of the --9.44-48Chvátal's Condition 10.21 clique 9.2; 11.18; 12.1 coatom 2.37

coloration of a graph 3.23; 4.21; 5.26; 6.8; 7.3-7; §9; 10.35; 11.19-22; 12.11,20, 22: 13.12,33-47; 14.4-6

- of the natural numbers 14.11,12,15, 18 - 22

- of the plane 14.7-10 - of the power set 14.13-17

compactness 14.19 complete subgraphs, number of 10.40 conic section 10.15,36 connectivity of graphs §6; 12.15

edge - 9.21; 12.14,16

strongly connected digraph 5.3,5,7-9;

6.6,9-13,29; 8.8; 10.41-44 contraction 4.17,27; 9.38; 10.3,6-8; 12.12,20 convex polygon 1.39,41; 3.23; 14.30,31 - hull 6.65; 14.31

coset 3.24 cover 7.1-5,12; 8.2-4; 10.18,31; 13.25, 26,30,48-55

fractional - 13.30,48-55 k-cover 13.26,49,50

Fano plane 13.35

Ferrer diagram 1.16,19,23

Fisher's Inequality 13.15

Five Color Theorem 9.50

Fibonacci numbers 1.27; 4.22,28,32

covering number of circuits 10.16-18 flow 6.71-78 forest 4.14; 6.37; 7.3; 11.3,4; 13.5 maximal - 7.34,37 Cramer's Rule 1.27,28; 4.33; 7.20 critical α-critical 8.12-27; 9.0 Four Color Theorem 9.0 v-critical 13.29,53 Frobenius-Perron theory 11.0 τ-critical 13.32,52 Frucht's Theorem 12.5 y-critical 9.17-24 factor - 7.26,31,32,38 Gallai-Edmonds Structure Theorem 7.32 Galois field (GF(q)) 5.31,36; 10.15,36; critically k-connected 6.36-38,54,59-65 11.26; 13.37; 14.23.24 -k-edge-connected 6.49,63cut 5.33; 6.50,71-74; 8.18; 9.29; 10.5; game (on a graph) 8.8,9 game (of a graph) 8.3,9 generating function  $\S 1$ ; 3.14,19,29 exponential -  $-\S I$ ; 3.13–15,31; 4.7 girth 9.25–27; 10.10–17; 12.25 Golden Ratio Theorem 9.49 12.5.15 evele index (see also permutation group) 3.7-12,15,26,30: 4.19 cycles of a permutation 3.1-13,22,26-29 group (see also permutation group) alternating - 3.10 degree maximal - 2.18; 7.10,17,18,25; 8.1,23; 10.6,16,38; 11.14; 12.6; 13.17,30,56 minimal - 6.61-65; 7.10,12,24; 9.1; commutative - 3.9; 5.4; 11.8,9,17; 12.13,17 cyclic - 11.3,10,13,19 10.3,8,35; 11.14,20; 13.17,44 symmetric - 3.1,7,15; 4.19; 12.19 degree sequence 4.1; 5.1; 6.3; 7.47-53; group character 11.8,10,17 10.24; 12.9 - of automorphisms 11.8-10,16,17; §12; diameter of a graph 6.22; 10.13; 11.23; 13.22: 15.2 15.16 edge transitive - - 12.12,18 digraph 3.14; 4.10,15-17; 5.3-13,19; 6.6, 9-13,39,45,71; 7.44,49,51; 8.4-8; 9.9-Hajós' Construction 9.16.20 11,26; 12.4,5,24; 13.7 Hall Theorem 13.5 Hamiltonian circuit 5.7,21,22; 10.21-27; 12.17; 14.4; 15.15 acyclic - 4.15; 6.11; 8.5 strongly connected - see connectivity - cycle 6.12,13; 10.25 - path 5.19,20; 10.21-24,42,43 symmetric - 8.5 Dilworth's Theorem 9.32 Diophantine Equation 14.20,21 Helly property 13.56 Dirac's Condition 10.21,23 Hoffmann-Kruskal Theorem 4.64 dual of a graph 5.23,36; 15.8 homeomorphic graphs (see subdivision) duality theorem of linear programming homomorphism of graphs 9.15; §12: 15.20, 13.48 - of groups and rings 15.22 Edmonds' Matching Algorithm 7.34 Hungarian method 7.3 eigenvalue 1.29,30; 4.29; §11 hypergraphs *§13* balanced – 13.39,40,54 eigenvector 1.30; 11.1-5,8,13,14,18-24 endomorphism 12.22,24,25 normal - 13.55-57 enumeration §1-4 totally balanced - 13.2,3,4 Erdős-de Bruijn Theorem 9.14,27 uniform - 13.3,12,13,25-32,36,41-47; Erdős-Ko-Rado Theorem 13.28,31 14.24; 15.18 Erdős-Stone Theorem 10.38 Euclidean space 14.10 inclusion-exclusion formula (see also sieve) Euler trail 5.0,6,9-11,14; 7.39 2.0,2,4; 3.2; 4.12,23,35; 9.37; 15.17, Eulerian graph 5.7; 6.51,53,56 20 Euler's Formula 4.27; 5.24,34; 12.18 independent sets of vertices 2.12; 7.37; event 2.2,6,8,11,17,18,20; 3.18 \$8 expected number 2.7; 3.5,17 - events 2.0,18,20; 3.16 - paths 4.31; 5.19; 6.39-46; 8.2-4; 9.24; 10.5 factor 1-factor 4.0,21-28,32; 5.18,21; §7; 9.55; index (see cycle index) 11.11; 12.16 f-factor 7.16,39-45 interval graph 9.28,41 inversions of a permutation 3.20 factor critical graph 7.26,31,32,38

> Jensen's Inequality 10.36,37,40,41 K. Jordán's Formula 2.7

circuit - 15.9

isomorphism 3.9; 4.13,18; 10.0; 11.10; 12.13,25; 15.1-3,9,10; 15.15-22

Kempe chains 9.0,54 kernel of a digraph 8.5 Kirchhoff Equations 4.0 König's Theorem 7.2-4; 9.21 König-Hall Criterion 7.20 Kruskal algorithm 6.25 Kuratowski Criterion 5.0,37 Kuratowski Theorem 10.4,5

Labyrinth Problem 5.12 Lagrange Method 2.15 lattice 2.23,27-31,34-36 geometric - 2.37 Lefschetz Fixed Point Theorem 6.15 line-graph 9.26,52; 11.2,15,25; 12.1,18; 13.43,56,57; 15.1-7 linear space 5.0,31,38; 10.15

MacLane's Criterion 5.35,36 Markov's Inequality 2.19 matching 4.12,31; 5.18; 6.67; §7; 11.4-6; 13.48 - 55fractional - 13.48-55 k-matching 7.37; 13.49,50 f-matching 7.43 matching number of circuits (number of disjoint circuits) 10.17,18

matrix adjacency - 11.7,14 complex - 15.13diagonal - 4.16; 11.16 incidence - 4.9; 11.15; 13.37-39 orthogonal - 11.16,21 permanent of a -4.21,33,36skew symmetric - 4.24,26 symmetric - 11.16; 15.11,18

unimodular - 13.37-39 -, compatible with a partial order 2.21,

Max-Flow-Min-Cut Theorem 6.0,74,77 Menger's Theorem 6.0,39-44, 58,77; 7.2 Moebius function of a partially ordered set 2.22 - 24

number theoretical - - 2.3, 23 Moebius inversion formula 2.26,35 monoid 12.24,25 monotone mapping 1.32,33; 3.19 - subsequence 14.25-27

network 6.0,25 network-flow problem 7.16 Newton's Formula 1.40

orbit 3.24-28: 12.7.10.12 orientation of graphs 3.13; 4.10,17,24,26, 27,29; 5.13; 6.29,34,54,56; 7.45,46; 9.11,46; 12.10

partially ordered set 2.21-24,31,32; 9.32; 13.22partition of a set 1.2,6,10,12,14,16-19,24,35, 36; 2.29,30,36; 3.8,14,22,31; 4.8,12,

14,24; 9.23,36 - into even number of classes  $(Q_n)$  1.14 partition into classes of even cardinality  $(R_n)$  1.14 - of numbers  $(\pi_n)$  1.20,21,23,25; 3.1 of vertices 4.31; 5.17; 9.2,8,23 path of maximum length 5.37; 6.6,14,17, 19,21,75; 10.2,20,27; 14.5 Pentagonal Numbers Theorem 1.19 perfect graph 9.28-35; 13.56,57 Perfect Graph Theorem 13.57 permutation 1.3; §3; 4.17,32-36; 11.11; 12.2; 13.9,21,27,28,32; 15.17 conjugacy of -s 3.1 cycles of a - 3.2-6,13,22,27-29; 4.17,26; 12.7 - matrix 11.16 permutation group 3.0-2,7-10,15,24-29; 12.0,13,19,21 cycle index of -3.7-12,15,26,30;4.19regular - - 3.9; 11.8,9,17; 12.7,13,17 semiregular -12.10,12,20transitive - - 11.10,17; 12.13-17 Petersen graph 10.10; 11.2; 12.1,17,22 Petersen's Theorem 7.29 planar graph 5.23-30,34-38; 6.69; 9.49-57; 12.18,19; 15.8 - map 5.15,23-30,34-38; 6.57,69 - tree 4.13 planarity criterion Kuratowski's - - 5.37 MacLane's - -5.35Whitney's -5.36Platonic Bodies 8.15; 12.2,19 Pólya-Redfield Method 3.26,29,31; 4.19 Pósa's Condition 10.21,24 potential 5.4; 9.10,11,56 probability 2.2,6,11,14,16,18,20; 3.3-5,18 product of graphs Cartesian -6.4; 9.6; 11.7,9; 12.17,20; 14.10 direct - -6.4; 9.7; 15.21,22 lexicographic - - 12.11 strong direct - - 9.31-22; 11.7; 13.51; 15.21 - 22of hypergraphs 13.51 Kronecker – of matrices 11.7 topological – 9.14 wreath - (of groups) 3.8 projective plane 10.15; 11.25 space 10.36; 13.17; 14.23 Prüfer Code 4.5

radius of convergence 4.20 Ramsey numbers 14.1-4 Ramsey's Theorem 10.32; 14.1,2,3,24 random variable 2.7; 3.5,16 Reconstruction Conjecture 15.15-17 regular graph 3.13,14; 5.2,21,22,27; 6.52; 7.10,28,30,39; 7.40; 8.15,24; 10.10-15,23,32; 11.2,12,24; 12.8,15 resultant 4.29 rigid graph 12.23,25

ring 15.22

roots of the chromatic polynomial 9.44-48 rooted graph 4.13-19; 5.10; 7.34

Shoenflies Theorem 6.69 Second Moment Method 2.19 semigroup 12.24,25 sieve

Brun's - 2.13 Eratosthenes' - 2.0

- formula *§2*; 4.23,35; 9.37; 15.17 Selberg's -2.14,20

singularity of an analytic function 1.15; 4.20

spanning arborescence 4.15,16; 5.10,11 - subgraph 7.41,42, 46; 9.54; 10.5; 11.13

- tree 4.9-16; 5.23,24,33,36; 6.7,8,25 spectrum §11 Sperner's Lemma 5.0,29 Sperner's Theorem 13.21,22 splitting a set 1.2,35-37

stabilizer 3.25 step polygon 1.32 Stirling Formula 4.20

Stirling numbers 1.2,6-9; 2.4; 3.12,17; 4.31,36; 9.23

strategy 3.18 winning - 3.18; 8.8,9 subdivision 5.37; 6.52; 10.0,3-6,9; 12.18

Sylvester determinant 4.29

Tarry algorithm 5.12 Taylor series 1.0; 3.11,31 Tihonov's Lemma 9.14 tournament 5.20; 6.12,13; 8.6; 10.41-44;

12.7,12

number of cycles in a - 10.41,42 transitive - 4.31; 5.20; 9.27; 10.40-43; 14.27

transitive sub- 10.44 tree §4; 6.20-26; 7.46,47; 9.29,39; 11.6; 13.1; 15.10,11,16

baricenter of a -6.22bicenter of a -4.17; 6.21

center of a - 3.8; 4.17; 6.21; 15.16

optimal -6.25,26planar - 4.13

triangle-free graph 9.25; 10.30,31 triangles, number of 10.32,33 triangulation 1.41; 5.38; 9.49,52,54,55 - of a convex n-gon 1.39,40

Turán's Theorem 10.30,34,38; 15.4 Tutte's Theorem 7.27,28,32; 12.16

Van der Waerden's Theorem (see coloration of natural numbers) 14.18 Vandermonde determinant 1.28

wheel 9.39; 10.4,18 Whitney's Criterion 5.0,36

Zarankiewitz's Problem 10.37

## **Author Index**

\* Asterisks indicate references to books.

Aardenne, Ehrenfest, T. van 5.11 Aarts, J. M. 9.54 Ádám, A. 6.34 \*Aho, A. V. p. 10 Anderson, W. N. 8.9 Andrásfai, B. 8.25 Appel, K. 9.0

Babai, L. 12.6-9,11,12,19,20,25 Beineke, L. W. 7.24; 8.17; 15.6 \*Berge, C. [B] 4.1,2,4,12,14,16,24,25,27; 5.11, 12,19-22; 6.3,10,12,29,54,66; 7.4,5,10,11,27,47-49,51,52; 8.4,5,10,17,20, 21,23; 9.6,9,16,21,29,32,36; 10.21,22,24, 25,26,34; 13.1,5-7,32,37-40,45,54,56,57; 14.1-4; 15.6 BERGE, C. 7.27; 8.17; 9.29; 13.38,40,51, 54; 15.2,3,19 Bernstein, F. 13.6 \*Biggs, N. L. [Biggs] 4.9 BIGGS, N. L. 4.9; 11.2,20,23,26 \*BOGNÁR, J. 3.6,18 Bollobás, B. 10.18; 13.10,32 Bondy, A. 6.3,8; 10.21; 13.10; 14.6; 15.15.16 Borsuk, K. 9.27 Bosák, J. 5.22 BOUWER, I. Z. 12.21 Brooks, R. L. 9.13 Bruijn, N. G. de 3.7,27; 5.11; 9.14; 13.20,21

Calczynska-Karlowicz, M. 13.27 Camion, P. 6.12 Cayley, A. 4.2,14 Chao, C. Y. 11.16,17; 12.13 Chen, W. K. 5.17,32 Chvátal, V. p. 10; 8.7; 10.21,26; 14.27 Cook, S. A. p. 10; 8.0 Corrádi, K. 13.13 Crawley, G. P. 13.14 Csima, J. 7.9,14 Cvetković, D. M. 11.7 Czipszer, J. 10.2 Dambit, J. J. 6.11
Daniels, H. E. 1.34
Daykin, D. E. 13.8,18,31
Deza, M. 13.17
Dilworth, R. P. 9.32; 13.14,22
Dirac, G. A. 6.33,41,66; 9.17,21,22,29,54; 10.3,4,21,27,29
Djorović, D. Z. 11.10
Dziobek, O. 46.

EDMONDS, J. p. 10; 6.71,75; 7.20,32,34 EHRENFEST, P. 5.11 ELSPAS, B. 11.10 \*ERDŐS, P. [ES], p. 11; 9.27; 13.41,42; 14.1,2 ERDŐS, P. 2.18; 7.51,52; 8.20; 9.14,27; 10.0,10-13,18,26,28,35,38; 13.9,23,24, 28,41-44,46,47; 14.2,3,6,8-10,22,29-31; 15.4 EULER, L. 1.19,23; 5.6,24

Faber, V. 15.18 Fáry, I. 5.38 \*Ford, L. R. [FF] 6.72-78 Fisher, R. A. 13.15 Folkman, J. 14.15 Ford, L. R. 6.74 Frank, A. 6.11 Fréchet, M. 2.10 Frucht, R. 12.1,3-5,8 \*Fulkerson, D. R. [FF] Fulkerson, D. R. 6.71,74; 7.35,51

Gaddum, J. W. 9.5 Gale, D. 6.76; 7.16 Gallai, T. 5.17,28,32; 6.11,17; 7.1,26, 32,51,52; 8.4,19,20; 9.4,9; 10.28 \*Gantmacher, F. R. 11.0 Gerencsér, L. 14.5 Ghouila-Houri, A. 13.37 Goodman, A. W. 10.32,33; 15.4 Graham, R. L. 2.33; 11.22; 14.8–10,14, 16,18,23 Greenwell, D. 9.7, 19 Grinberg, E. J. 6.11 GROOT, J. DE 9.54 GUPTA, R. P. 7.12; 9.5 GYÁRFÁS, A. 14.5

HADWIGER, H. 14.7 HAJNAL, A. 8.10,21,23: 9,27,29: 13,29: Hajós, Gy. 6.43,44; 9.16,28 HAKEN, W. 9.0 \*HALBERSTAM, H. 2.13 HALES, A. 14.17 HALIN, R. 6.60 \*HALL, M. [HALL] 3.0; 7.15; 13.15 HALL, P. 7.4; 12.5; 13.5 \*HARARY, F. [H] 14.1,3; 15.6 HARARY, F. 8.17 Накинетм, Е. 14.26,28 HAVEL, V. 7.50 HEAWOOD, P. J. 9.52 HEDRLIN, Z. 12.21.23.24 HEMMINGER, R. 15.15,16 HERCZOG, J. 13.9 HETYEI, G. 7.7-9,24,28 Ноггман, А. J. 7.35,51; 11.15,21,24,25 \*Норскорт, J. E. р. 10 \*Hu, T. C. [Hu] 6.56,72-78

#### IMRICH, W. 12.13

JAEGER, F. 13.32 JEWETT, R. I. 14.17 JORDÁN, J. 2.7 JUNG, H. A. 6.57

Karp, R. M. p. 10; 6.75; 8.0 Kasteleyn, P. W. 4.24,25,27 Katona, G. O. H. 13.28,31,32 Kelly, J. B. 6.20; 13.16 Kelly, P. J. 15.15,16 Kirchhoff, G. 4.9 Klarner, D. A. 4.13 Kleitman, D. 6.25; 13.11,18 Ko, Chao 13.28 Komlós, J. 14.27 \*König, D. [K] 6.21,22; 7.29 König, D. 7.2 Kotzig, A. 6.47,52; 7.24 Kővári, T. 10.37 Kruljswijk, T. 13.20,21 Kruskal, J. B. 13.31 Kundu, S. 7.53

Lambek, J. 12.24 Las Vergnas, M. 13.45 Leeb, K. 14.14 Lehmer, D. H. 4.33 Levin, L. A. p. 10; 8.0 Lovász, L. 2.18; 6.8,58; 7.6,15,28,31,46, 53; 8.7,22,25; 9.7,19,35; 10.4;11.3-5.8; 12.9; 13.4,5.8; 13.18,30,36,43-47,52,53, 55-57; 14.22; 15.17,20-22 Lubell, D. 13.21 MacLane, S. 5.35
Mader, W. 6.49,60,63; 10.7,8; 12.15
Mani, P. 6.57
Mantel, W. 10.30
Marica, J. 13.8
McAndrew, M. H. 7.35,51
McEliece, R. J. 13.51
Melnikov, L. S. 8.26
Menger, K. 6.39
Minty, G. J. 6.10; 9.11
\*Mirsky, L. [Mi] 13.5,6,7
\*Mogyoródi, J. 3.6,18
Montgomery, P. 14.8-10
\*Montroll, E. W. 4.0,29
\*Moon, J. W. [M] 10.41-44
Moon, J. W. [M] 10.41-44
Moon, J. W. 4.4; 10.40; 12.7
Moser, L. 1.9; 10.40
Mowshowitz, A. 11.3-5,16,17
Müller, W. 15.17
\*Mullin, R. 1.45
Mycielsky, J. 9.18

NASH-WILLIAMS, C. St. J. A. 6.54,56; 10.23,25 NEŠETŘIL, J. 12.10,25 NICKEL, L. 15.7 NORDHAUS, E. A. 9.5 NOWITZ, L. A. 12.13

\*Ore, O. [O] 7.2,4-5,16; 9.36 \*Ore, O. [OF] 9.51,52,54,56 Ore, O. 7.5,16 O'Neil, P. V. 4.12

Payan, C. 13.32
Pelikán, J. 11.3-5; 12.17
Perfect, H. 6.42
Petersdorf, M. 11.16,17
Petersen, J. 7.29
Plesnik, J. 7.30
Plummer, M. D. 6.35-37,68; 7.15,24; 8.17,19
Pollak, O. 2.33; 11.22
Pólya, G. 3.7-9,29; 4.7,13
\*Pontryagin, L. S. 5.29
Pósa, L. 5.17; 8.3; 9.14; 10.2,5,18,20,21; 15.4
Posner, E. C. 13.51
\*Prékopa, A. 3.6,18
Prüfer, A. 4.5
Pultr, A. 12.21,23,24

\*RIORDAN, J. [R] 1.24 RADO, R. 13.7,28; 14.3,21 \*RADEMACHER, H. 1.22 RAYNAUD, H. 14.4 \*READ, R. C. 4.0 READ, R. C. 9.36 RÉDEI, L. 5.20 REIMAN, I. 10.36 \*RÉNYI, A. 3.6,18 \*RÉNYI, A. [Ré] 2.6-8,10,11 RÉNYI, A. 1.14; 2.12; 3.16,17; 4.3,8 RÉNYI, C. 3.16,17 ROBBINS, H. E. 6.29 \*ROTA, G. C. 1.45; 2.0 ROTA, G. C. 2.21-29 \*ROTH, K. F. 2.13 ROTHSCHILD, B. L. 13.29; 14.8-10,14,16, 18,23 ROY, B. 9.9

Ryser, H. J. 7.16 Sabidussi, G. 12.8,12,13 \*Sachs, H. [S] 5.37,38; 6.25; 7.4,29; 9.16, 18,25,51,52,54; 10.11-13,34,41 SACHS, H. 7.21; 9.57; 10.11-13; 11.2,12, 16,17 Sanders, J. 14.15 SAUER, N. 13.10 SCHÄUBLE, M. 9.55 SCHÖNHEIM, J. 13.8,9,18 SCHUR, I. 14.11 SCHÜTZENBERGER, M. P. 2.30 SCHWENK, A. J. 11.6 Selfridge, J. 15.12 \*SESHU 4.0 SHANNON, C. E. 7.10 SHRIKHANDE, S. S. 11.26 SIMMONS, G. J. 6.52 Simonovits, M. 10.16-18,38,39; 13.51

\*Spencer, J. [ES] p. 11; 9.27; 13.41,42; 14.1,2

Spencer, J. 14.8-10

Spencer, E. 5.29; 13.21

Spitzer, F. 3.22

Stanley, R. 9.46

Stone, A. H. 10.38

Strauss, E. G. 14.8-10; 15.1

SMITH, C. A. B. 5.21 SMOLENSKI, E. A. 15.10

Sós, V. T. 10.37

\*SPANIER, E. 6.15

SURÁNYI, J. 9.29 SURÁNYI, L. 8.22 \*Szász, D. 3.6,18 Szekeres, G. 14.29-31 Szele, T. 5.19 Szemerédi, E. 8.27


Tait, P. G. 9.51 Tarry, G. 5.12 Temperley, H. N. V. 4.12,30 Tengbergen, C. A. van E. 13.20,21 Toft, B. 9.17 Turán, P. 10.34,37 Turner, J. 11.10 \*Tutte, W. T. 6.64 Tutte, W. T. 6.70; 7.27,37,43; 9.25,36; 10.11

\*Ullman, J. D. p. 10

VESZTERGOMBI, K. 4.36 VIZING, V. G. 7.10; 8.26; 9.6 \*Voss, H.-J. [WV] 6.17; 10.7,8,12,13,16, 17,29 Voss, H.-J. 10.16–18,29

Waerden, V. L. Van der
\*Wagner, K. [W] 5.35
Wagner, K. 5.38
Walther, H. J. 6.17
Watkins, M. E. 6.60; 12.13,15
Weinberg, L. 4.12
Werra, V. de 7.11
Wessel, W. 8.13,19
Whitney, H. 5.36; 6.33,69; 9.56; 15.1,9
\*Wielandt, H. 3.0
Wilf, H. S. 2.31, 11.20
\*Wilson, R. J. [Wi] 5.36; 6.29; 9.36,51,56
Woopall, D. R. 13.45
\*Walther, H. J. [WV] 6.17; 10.7,8,12, 13,16,17,29
Wyman, M. 1.9

ZAKS, J. 7.36 ZARANKIEWICZ, K. 10.37 ZARETSKI, K. A. 15.11







## We recommend

# B. ANDRÁSFAI INTRODUCTORY GRAPH THEORY

In English — 267 pages 264 figures  $16 \times 23$  cm — Cloth

Graph theory is a newly developed topic of mathematics and has been spreading over an increasingly wider field in the last one or two decades. It offers the solutions to a multiplicity of naturally arising problems of interest while requiring no great preliminary mathematical knowledge; it does, however, frequently require intensive thinking. It is therefore likely to become especially suitable for young people interested in mathematics enabling them to spread their wings. The author has compiled the material of his book with this object in mind. He offers his material not in a rigid prearranged systematization, but follows a certain way of discovering the solutions, i.e. he sets the reader simple exercises with tasks based on these.

Distributors

KULTURA

H-1389 Budapest, P.O.B. 149