

Tk 32.934



KFKI 1/1970

1970 MAR 19

I. Nagy

L. Pál

ELECTRICAL RESISTIVITY AND THERMOELECTRIC
POWER OF NI NEAR THE CURIE POINT

Hungarian Academy of Sciences

CENTRAL
RESEARCH
INSTITUTE FOR
PHYSICS

BUDAPEST

The following information is provided for the purpose of
 providing a general overview of the project and its
 objectives. The information is not intended to be
 used as a substitute for the full project report.
 The project is a study of the effects of the
 implementation of the new curriculum on the
 performance of students in the subject of
 mathematics. The study is being conducted in
 a number of schools in the district and the
 results will be reported in a separate report.
 The project is being funded by the
 Department of Education and the results will
 be used to inform the development of the
 curriculum for the future.

Electrical resistivity and thermoelectric power of Ni
near the Curie point

I. Nagy and L. Pál

Central Research Institute for Physics

Budapest

In the last few years there has been a growing interest in the critical behaviour of the transport properties of matter /1/. The primary aim of the research has been to study the appearance and character of the singularities in transport coefficients. In this paper some new results of high precision electrical resistivity and thermoelectric power measurements on high purity nickel in the vicinity of its Curie point will be reported.

It was noticed a long time ago /2/ that the temperature coefficient of the resistivity $\alpha(T) = \frac{d\rho(T)}{dT}$, where $\rho(T)$ is the resistivity/ of various d-metals has a sharp maximum at T_c , although this fact has only recently attracted greater attention /3/. Three years ago Craig et al. /4/ measured directly $\alpha(T)$ in five-9's purity Ni and found a logarithmic divergence for $T > T_c$ and for $\epsilon = \frac{|T-T_c|}{T_c} \geq 5 \times 10^{-3}$. Closer to the Curie point they observed a stronger than logarithmic divergence. It was suggested by Craig et al. that the change in the strength of the divergence is related to the fact that the range of the critical fluctuations equals the mean free path of the electrons at a temperature near the Curie point.

These observations stimulated Fisher and Langer /5/ to improve the existing theoretical treatments of the resistivity anomaly such as that of de Gennes and Friedel /6/. It was clearly shown that the main contribution to the resistivity anomaly in magnetic metals near the Curie point is due to the short-range spin fluctuations and hence $\alpha(T)$ should have the same temperature dependence as that of the magnetic specific heat above but near the Curie temperature. The measured values of $\alpha(T)$ on Ni are consistent with this conclusion in the temperature region of logarithmic divergence, but the change in the order of the divergence observed by Craig et al. cannot be explained by the Fisher-Langer theory.

In order to obtain more precise data and to study the most sensitive electronic transport property of metals it was decided to perform careful electrical resistivity and thermoelectric power measurements in the vicinity of the Curie point on nickel samples of 99,999 per cent purity produced by Johnson and Matthey.

For the measurements three samples were prepared from the same Ni. Each of the samples was a 50 cm long Ni wire of 0,1 cm diameter wound as a small sphere which was inserted into the middle of a precisely shielded spherical vacuum furnace. The temperature inhomogeneity in the sample was less than 0,01 °K.

The resistivity measurements were carried out using d.c. and a manual potentiometer with a photo-galvanometer amplifier of high stability. During the measurement the current in the sample was kept constant within 10^{-6} . The voltage on the specimen was almost exactly compensated by the potentiometer and only the noncompensated part of the voltage was fed after a suitable amplification into the y-input of an x-y recorder. A Pt-PtRh thermocouple welded on the sample measured the temperature. The amplified non-compensated voltage of the thermocouple was fed into the x-input of the recorder which directly recorded the $\rho(T)$ curve in arbitrary units.

The thermoelectric power was measured on the same samples as the resistivity. In order to produce a small temperature difference between the ends of the Ni wire direct current of 0,2 A was applied to the sample through the current leads for a short period /5-10 sec/. The generated Peltier heat resulted in a temperature difference of a few tenths centigrade on the sample. After the switching off of the current the temperature difference begins to decrease immediately and disappears in about 2 minutes. By measuring the decreasing voltages of the Pt-Ni-Pt and PtRh-Ni-PtRh differential thermocouples the thermoelectric power of Ni can be determined.

The voltage of one of the differential thermocouples versus that of the other was recorded after a suitable amplification by an x-y recorder. If the parasitic voltages do not change during the measurement a straight line can be expected from the slope of which it is easy to calculate the thermoelectric power of Ni. Since the temperature difference on the sample was not larger than $0,2^{\circ}$ even at temperatures near the Curie point, the just described method allowed to determine the absolute thermoelectric power of Ni to a better accuracy than 5 per cent.

In order to determine the divergence of the temperature coefficient of the resistivity in the vicinity of the Curie point $\alpha(T)$ was represented in a manner proposed by Fisher for other divergent quantities:

$$\alpha(T) = A \frac{\epsilon^{-\lambda} - 1}{\lambda} + B$$

where $\epsilon = \frac{|T - T_c|}{T_c}$ and A, λ and B are temperature independent constants. From the measured $\rho(T)$ curve the $\alpha(T) = \frac{d\rho(T)}{dT}$ curve was calculated by graphical differentiation using intervals $\Delta T = 0,1^{\circ}$. In Fig. 1 the temperature dependence of $\alpha(T)$ for the sample No. 1 is shown.

Calculating the second derivative of $\rho(T)$ for $T > T_c$ we used the log-log plot of the curve

$$\frac{d\alpha(T)}{dT} = -A \epsilon^{-(\lambda+1)}$$

for the evaluation of the parameter λ . The log-log plot of $d\alpha/dT$ can be seen in Fig. 2. The agreement with Craig's curve is quite satisfactory. The only difference is that the break point at which the strength of the divergence is changing lies nearer the Curie point than in Craig's curve which gives $\epsilon \sim 5.10^{-3}$ as compared with our $\epsilon \sim 1,6.10^{-3}$. Below the Curie point the strength of the divergence could not be clearly determined.

The thermoelectric power measurements on the sample No. 1 resulted in a very unexpected anomaly. It can be seen in Fig. 3 that the thermoelectric power versus temperature curve has a sharp minimum followed by a sharp maximum in the same very narrow temperature interval in which the deviation from the logarithmic divergence of the $\alpha(T)$ curve was observed. Far from the Curie point the temperature dependence of the thermoelectric power does not differ from the results of other measurements /7/.

In our experiment however, the most important result is the disappearance of the break point on the $\alpha(T)$ curves measured on the other two samples. Fig. 4 shows the $\alpha(T)$ curve for the sample No. 2. Precisely the same curve was found in the case of the sample No. 3. In Fig. 5 the log-log plot of $\frac{d\alpha(T)}{dT}$ clearly demonstrates that the logarithmic divergence is conserved on both sides of the Curie temperature at temperatures very near the critical point, too. More exactly, it was found that the temperature coefficient of the resistivity


in Ni has a logarithmic divergence in the immediate neighbourhood of the Curie temperature ($|T-T_c| \geq 0,2 \text{ } ^\circ\text{K}$).

The temperature dependence of the thermoelectric power on these samples /No. 2 and No. 3/ is completely different from that measured on the sample /No. 1/ with a break point on the $\alpha(T)$ curve. In the samples No. 2 and No.3 the thermoelectric power versus temperature curves do not show any surprising anomaly in the vicinity of the Curie point. As can be seen in Fig. 6, the curve for the sample No. 2 is quite similar to that reported earlier /7/.

In the light of these results a few comments can be made concerning the Fisher-Langer theory and the Craig experiment. First, Fisher's conclusion that $\alpha(T)$ should vary as the magnetic specific heat above the Curie point is consistent with the experiment covering the immediate neighbourhood of the Curie point ($|T-T_c| \geq 0,2 \text{ } ^\circ\text{K}$). Second, the prediction that $\alpha(T)$ should have a critical exponent of $2\beta-1$ below the Curie point / β is the critical exponent for the magnetization/ seems to be not true because the experiment suggests a logarithmic divergence. Third, Craig's suggestion that the crossover between the two types of singular behaviour occurs when the correlation length equals the electronic mean free path is rather questionable since the onset of the crossover may depend on several factors the nature of which is not yet understood. The only thing which can be regarded as a real physical consequence of the short electronic mean free

path is that which was pointed out by Fisher, and this is the importance of the short-range part of the spin-correlation function in the critical behaviour of the resistivity of magnetic metals.

Finally, one has to note here, that the critical behaviour of the thermoelectric power unfortunately cannot be analysed theoretically because of the lack of any theory in this field.




References

1. P. P. Craig, J. Appl. Phys., 40 964 /1969/
2. W. Gerlach, Physik Z., 24 953 /1932/
H. Bittel and W. Gerlach, Ann. Physik, 33 661 /1938/
3. Ya. A. Kraftmakher, Fiz. Tverd. Tela, 7 8 /1965/
4. P. P. Craig, W. I. Goldberg, T. A. Kitchens and
I. I. Budnick, Phys. Rev. Letters, 19 1334 /1967/
5. M. E. Fisher and J. S. Langer, Phys. Rev. Letters,
20 665 /1968/
6. P. G. de Gennes and J. Friedel, J. Phys. Chem. Solids,
4 71 /1958/
7. M. V. Vedernikov and N. V. Kolomoets, Sov. Phys. Solid
State, 2 2420 /1961/

Figure captions

- Fig. 1 $\alpha(T) = d\rho/dT$ the temperature coefficient of resistivity of the Ni sample No. 1 versus temperature in the vicinity of the Curie point.
- Fig. 2 The log-log plot of $d\alpha/dT$ versus $\epsilon = |T-T_C|/T_C$. λ is the critical exponent describing the divergence of $\alpha(T)$.
- Fig. 3 Variation of the absolute thermoelectric power with temperature of the Ni sample No. 1 in the immediate neighbourhood of the Curie point.
- Fig. 4 The temperature coefficient of resistivity of the Ni sample No. 2 versus temperature.
- Fig. 5 The log-log plot of $d\alpha/dT$ versus $\epsilon = |T-T_C|/T_C$ for the sample No. 2.
- Fig. 6 The temperature dependence of the absolute thermoelectric power of the sample No. 2.






Fig. 1






Fig. 2





Fig. 3






Fig. 4






Fig. 5





Fig. 6

G1. 832





Kiadja a KFKI Könyvtár- és Kiadói Osztály
O.v.: Dr. FARKAS Istvánné
Példányszám: 185 Munkaszám: 4903
Készült a KFKI házi sokszorosítójában
F.v.: Gyenes Imre
Budapest, 1970. január 28.