

KFKI-1982-83

A. SÜTŐ  
T. YALCIN  
C. GRUBER

A PROBABILISTIC APPROACH TO THE MODELS  
OF SPIN GLASSES

*Hungarian Academy of Sciences*

CENTRAL  
RESEARCH  
INSTITUTE FOR  
PHYSICS

BUDAPEST



A PROBABILISTIC APPROACH TO THE MODELS  
OF SPIN GLASSES

A. Sütő, T. Yalcin\*, C. Gruber\*

Central Research Institute for Physics  
H-1525 Budapest 114, P.O.B. 49, Hungary

\*Institut de Physique Théorique  
Ecole Polytechnique Fédérale de Lausanne  
PHB - Ecublens - CH-1015 Lausanne, Switzerland

*Submitted to J. Stat. Phys.*



## ABSTRACT

Introducing the notions of quenched and annealed probability measures, a systematic study of some problems in the description of spin glasses is attempted. Inequalities and variational principles for the free energies are derived. The absence of spontaneous breakdown of the gauge symmetry is discussed and some high temperature properties are studied. Examples of annealed models with more than one phase transition are shown.

## АННОТАЦИЯ

Для изучения моделей спиновых стекол нами введены понятия мер вероятности, относящихся к случаям быстрого и медленного охлаждения. Выведены неравенства и вариационные принципы для соответствующих свободных энергий. Обсуждено отсутствие спонтанного нарушения калибровочной симметрии, и изучены высокотемпературные свойства. Приведены примеры для моделей с медленным охлаждением, имеющих несколько фазовых переходов.

## KIVONAT

A spinűveg-modellek leírására bevezetjük a "gyorshűtött" és "hőtemperált" esetekhez rendelt valószínűségi mértékeket. Egyenlőtlenségeket és variációs elvet vezetünk le a megfelelő szabadenergiákra. Targyaljuk a gauge-szimmetria megőrzését, magas hőmérsékleti tulajdonságokat. Példát adunk olyan hőtemperált modellekre, melyekben egynél több fázisátmenet történik.



## 1. Introduction

A commonly accepted way to describe disordered systems is to represent them by statistical ensembles in which the degrees of freedom are coupled by random interactions. This additional randomness may be treated in different ways: Random interactions may be considered as new degrees of freedom and in extreme cases they may be in thermal equilibrium with the rest of the system (annealed state) or completely frozen in some random position (quenched state). Many years after the pioneering work of Brout<sup>1</sup> the quenched state of certain spin models became the center of interest of the theoretical research on spin glasses. Edwards and Anderson<sup>2</sup> pointed out that these systems can properly be described by the quenched state of randomly interacting spin models on regular lattices. Meanwhile, one encounters two main difficulties: the first is to calculate, in a respectable approximation, the quenched free energy and the second is to give a reliable proof that there exists a phase transition - in the static sense - between the high temperature paramagnetic and the low temperature spin glass state. Neither of these problems has got so far a reassuring solution, excepted probably in the case of the so called Sherrington-Kirkpatrick model<sup>3</sup>.

While not pretending to contribute to the solution of these great questions, in the present paper we attempt a systematic study of what were called the "annealed" and "quenched" states. To this end we define in sec 2 the annealed and quenched probability measures which play the same role as the Gibbs measure in equilibrium systems. In Section 3 we derive some inequalities for the quenched free energy and establish in sec 4 a variational principle characterising both the annealed and quenched free energies in the space of the joint probability distributions for spins and bonds. In Section 5 we raise the question of the uniqueness of the quenched state and show that the gauge invariance cannot be broken by different choices

for the boundary condition. This suggests that the "order parameter" of the spin glass state must be the expectation value of a gauge invariant and non-local observable. The order parameter, proposed by Edwards and Anderson<sup>2</sup> has indeed these properties, as we point out in Section 6. A discussion of high temperature properties is also given there and the functional relationship between order parameter and free energy is established. Finally, in Section 7 we return to the study of annealed models and give examples for one, two and three consecutive phase transitions in such models.



## 2. Definition of the system

Let  $L$  be a lattice; at each site  $i$  of  $L$  is associated a single spin space  $S$  where  $S$  is a subset of  $\mathbb{R}^v$ . The spin configurations are defined by  $\sigma: L \rightarrow S$  and the formal hamiltonian of the system is given as

$$H(J, \sigma) = - \sum_{b \subset L} J_b \phi_b(\sigma) \quad (2.1)$$

Here the  $\phi_b$ 's are bounded, real valued functions depending on  $\sigma_b = \{\sigma_i; i \in b\}$  and the  $J_b$ 's are real random variables with probability distribution  $d\mathcal{G}_b$  the mean value of which is finite. The finite partial sums of (2.1) are well defined for any  $\sigma$  with  $\mathcal{G}$ -probability 1. In particular

$$H_V(J, \sigma) = - \sum_{b \subset V} J_b \phi_b(\sigma) \quad (2.2)$$

exists for all finite  $V \subset L$ .

On  $S$  is given an a priori finite measure  $d\mu_0$  not necessarily normalised and  $d\mu_V(\sigma)$  denotes  $\prod_{i \in V} d\mu_0(\sigma_i)$  while  $d\mathcal{G}_V(J) = \prod_{b \subset V} d\mathcal{G}_b(J_b)$ . In the following discussion we consider only finite volumes and we omit the label  $V$ .

For given interactions  $J$ , the free energy  $F(J)$  and the equilibrium state at inverse temperature  $\beta$  are defined as usual by:

$$e^{-\beta F(J)} = \int d\mu(\sigma) e^{-\beta H(J, \sigma)} \quad (2.3)$$

and by the Gibbs probability measure

$$dG_J(\sigma) = g(J, \sigma) d\mu(\sigma)$$

where

$$g(J, \sigma) = e^{\beta [F(J) - H(J, \sigma)]} \quad (2.4)$$

(2.4) can also be written as



$$F(J) = H(J, \sigma) + \frac{1}{\beta} \ln g(J, \sigma) \quad (2.5)$$

which yields (since the Gibbs measure is normalised)

$$\begin{aligned} F(J) &= \int d\mu(\sigma) g(J, \sigma) \left[ H(J, \sigma) + \frac{1}{\beta} \ln g(J, \sigma) \right] \quad (2.6) \\ &= E(J) - \frac{1}{\beta} S(J) \end{aligned}$$

The quenched free energy  $\bar{F}$  and the quenched state are defined by

$$\bar{F} = \int d\mathcal{J}(J) F(J) = -\frac{1}{\beta} \int d\mathcal{J}(J) \ln \left[ \int d\mu(\sigma) e^{-\beta H(J, \sigma)} \right] \quad (2.7)$$

and by the quenched probability measure

$$dQ(J, \sigma) = g(J, \sigma) d\mu(\sigma) d\mathcal{J}(J) \quad (2.8)$$

which yields

$$\bar{F} = \int dQ(J, \sigma) \left[ H(J, \sigma) + \frac{1}{\beta} \ln g(J, \sigma) \right] = \bar{E} - \frac{1}{\beta} \bar{S} \quad (2.9)$$

We shall impose that the distribution  $d\mathcal{J}_b(x)$  falls off sufficiently rapidly so that  $\bar{F}$  is well defined.

The annealed free energy  $F_{\text{an}}$  and the annealed state are defined by the prescription that the average over the interactions has to be performed in the partition function, i.e.

$$F_{\text{an}} = -\frac{1}{\beta} \ln \left[ \int d\mathcal{J}(J) e^{-\beta F(J)} \right] = -\frac{1}{\beta} \ln \left[ \int d\mathcal{J}(J) d\mu(\sigma) e^{-\beta H(J, \sigma)} \right] \quad (2.10)$$

and by the annealed probability measure

$$dA(J, \sigma) = h(J, \sigma) d\mu(\sigma) d\mathcal{J}(J)$$

where 
$$h(J, \sigma) = e^{\beta [F_{\text{an}} - H(J, \sigma)]} \quad (2.11)$$

i.e. 
$$F_{\text{an}} = H(J, \sigma) + \frac{1}{\beta} \ln h(J, \sigma) \quad (2.12)$$

which yields

$$F_{an} = \int dA(J, \sigma) [H(J, \sigma) + \frac{1}{\beta} \ln h(J, \sigma)] = E_{an} - \frac{1}{\beta} S_{an} \quad (2.13)$$

Introducing the space  $\mathcal{B}$  of joint probability distributions

$$\mathcal{B} = \left\{ f = f(J, \sigma); f(J, \sigma) \geq 0 \int dJ d\mu f = 1 \right\} \quad (2.14)$$

and the free energy functional  $F[f]$  defined on  $\mathcal{B}$  by

$$F[f] = \int dJ d\mu(\sigma) f(J, \sigma) [H(J, \sigma) + \frac{1}{\beta} \ln f(J, \sigma)] \quad (2.15)$$

we can express  $\bar{F}$  and  $F_{an}$  respectively as

$$\bar{F} = F[g] \qquad F_{an} = F[h] \quad (2.16)$$

Finally for any subset  $B = (b_1, \dots, b_k)$  of  $b$ 's in  $V$  we introduce  $J_B = \{J_b\}_{b \in B}$  and

$$\bar{F}(J_B) = \int \prod_{\substack{b \in V \\ b \notin B}} dJ_b(J_b) F(J) \quad (2.17)$$

which implies in particular

$$\bar{F}(J_B) = F(J) \quad \text{if } B = \{b; b \in V\}$$

$$\bar{F}(J_B) = \bar{F} \quad \text{if } B = \emptyset$$

Let us note that for any  $J'_B = \{J'_b\}_{b \in B}$ ,  $\bar{F}(J'_B)$  represents the quenched free energy with respect to the new measure  $dJ'(J)$  where

$$dJ'(J) = \prod_{\substack{b \in V \\ b \notin B}} dJ_b(J_b) \prod_{b \in B} \delta(J_b - J'_b) dJ_b$$



### 3. Inequalities for the quenched free energy

In this section we first collect inequalities for  $\bar{F}(J_B)$ ; we then discuss the dependence of  $\bar{F}$  on the distribution  $d\mathcal{P}$ .

#### Proposition 3.1

$$i) \quad \bar{F}(J_B) \leq \bar{F}(J_{B'}) \quad \text{for any } B \subset B' \quad (3.1)$$

$$\text{where } J_b = \int d\mathcal{P}_b(x) x$$

$$ii) \quad F_{\text{an}} \leq \bar{F} \leq F(\bar{J}) \quad (3.2)$$

The proof of i) follows from Jensen's inequality using the known fact that  $F(J)$  is a concave function of each  $J_b$ 's and thus  $F(J_B)$  is also a concave function of each  $J_b$ 's,  $b \in B$ . The proof of ii) follows from Jensen's inequality using the fact that  $F = -\beta \ln Q$  is a convex functional of the partition function  $Q$ .

This proposition was earlier published by Rosa<sup>4</sup>; as mentioned in Sec. 2 the inequalities (3.1) can be regarded as the comparison of two different averages of the same function  $F(J)$ ; we thus have

$$\int d\mathcal{P}_1 F \leq \int d\mathcal{P}_2 F \quad (3.3)$$

where the probability measure  $d\mathcal{P}_2$  is sharper than  $d\mathcal{P}_1$ .

The question naturally arises whether (3.3) is generally true, i.e. whether a "sharpening" of the distribution causes the quenched free energy to increase. The inverse problem is also of interest: does a broadening decrease  $F$ ? At first we show that the broadening problem can always be solved.



Lemma 3.1

Let  $f = \mathbb{R} \rightarrow \mathbb{R}$  be concave and let  $\rho$  and  $\nu$  be two probability measures on  $\mathbb{R}$  such that

i)  $\int d\rho(x) f(x)$  exists

ii)  $\int d\nu(x) x = 0$  and

$\int d\nu(x) f(x+y)$  exists for all  $y$  in  $\mathbb{R}$

then  $\int d(\rho * \nu)(x) f(x) \leq \int d\rho(y) f(y)$  (3.4)

where  $d(\rho * \nu)$  denotes the convolution of the measures.

Proof:

Using Jensen's inequality together with  $\int d\nu(x) x = 0$ , implies

$$\int d\rho(x) f(x) \geq \int d\rho(x) \int d\nu(y) f(x+y) = \int d(\rho * \nu)(y) f(y)$$

This lemma can be immediately extended to concave functions of several variables and implies the following result.

Proposition 3.2

Let  $\{d\nu_b\}$ ,  $b \in \mathbb{C}$ , be a set of probability measures with zero means and such that  $\int \prod_{b \in \mathbb{C}} d\nu_b(J_b) F(J_b + \tilde{J}_b)$  exists for all  $\tilde{J}$

then  $\int d(\rho * \nu)(J) F(J) \leq \int d\rho(J) F(J) = \bar{F}$  (3.5)

Remarks

1. If  $E$  denotes the mean value and  $\Delta^2$  the mean square deviation then  $E_{\rho * \nu} = E_{\rho} + E_{\nu}$  and  $\Delta_{\rho * \nu}^2 = \Delta_{\rho}^2 + \Delta_{\nu}^2$ . Hence in proposition



3.2 we have  $E_{\rho^{*\nu}} = E_{\rho}$  and  $\Delta_{\rho^{*\nu}}^2 \geq \Delta_{\rho}^2$ , so that  $\rho^{*\nu}$  is indeed broadened in comparison to  $\rho$  and does decrease  $\bar{F}$ .

2. Replacing  $d\rho_b$  by  $d(\rho_b * \nu_b)$  corresponds to replacing  $J_b$  by  $J_b + \xi_b$  where  $\xi_b$  is a random variable independent of  $J_b$  and distributed according to  $d\nu_b$ .

3. The inequalities (3.1) follow from this theorem if in the latter we replace  $\rho_b(x)$  by  $\delta(x - \tilde{J}_b)$  for  $b \in B'$  and take  $\nu_b(x) = \delta(x)$  for  $b \notin B'$  and  $b \in B$

$$\nu_b(x) = \rho_b(x + \tilde{J}_b) \text{ for } b \in B'/B$$

In view of the second remark, the sharpening problem can be formulated in this way: given a random variable  $J_b$  we have to find a non trivial decomposition of  $J_b$  into the sum of two independent random variables, one of them having zero mean. This problem can not be generally solved. However if the  $J_b$ 's are Gaussian Random variables then such a decomposition is possible<sup>5</sup> and we have the following result.

Proposition 3.3

Consider the quenched free energy associated with two different set of Gaussian distributions  $\{\rho_b^{(1)}\}$  and  $\{\rho_b^{(2)}\}$  such that  $E_{\rho_b^{(1)}} = E_{\rho_b^{(2)}}$  and  $\Delta_{\rho_b^{(1)}}^2 > \Delta_{\rho_b^{(2)}}^2$  for all  $b$ .

then 
$$\int d\rho^{(1)}(J) F(J) \leq \int d\rho^{(2)}(J) F(J) \tag{3.6}$$

Proof:

Let  $\nu_b$  be the gaussian measure with zero mean and mean square deviation  $\Delta_{\rho_b^{(1)}}^2 - \Delta_{\rho_b^{(2)}}^2$ . Then  $\rho_b^{(1)} = \rho_b^{(2)} * \nu_b$  and the statement follows from proposition 3.2.

There is another way to generalize the inequality (3.1) which shows that the approximation of a given distribution with a



suitably chosen discrete distribution results in the increase of the quenched free energy. The better the discrete distribution approaches the original one the smaller is the upper bound.

Proposition 3.4

Let  $d\varphi_b(x) = d\varphi_b(-x)$  for all  $b$  and let

$$d\nu_b(x) = \frac{1}{2} [\delta(x - |\overline{J}_b|) + \delta(x + |\overline{J}_b|)] \quad \text{where} \quad |\overline{J}_b| = \int d\varphi_b(x) |x|$$

then:  $\overline{F} \leq \int d\nu(J) F(J) \leq F(\overline{J})$ .

Proof:

We may fix the interactions with the exception of the single  $J_b$  and it is sufficient to show that for the concave function  $F(J_b)$  the inequalities

$$\int_{-\infty}^{\infty} F(x) d\varphi_b(x) \leq \frac{1}{2} (F(|\overline{J}_b|) + F(-|\overline{J}_b|)) \leq F(0)$$

hold. Here the second inequality comes from the definition of a concave function. Now  $2d\varphi_b$  is a probability measure on both  $\mathbb{R}^+$  and  $\mathbb{R}^-$  and hence

$$2 \int_{-\infty}^0 F(x) d\varphi_b(x) \leq F\left(2 \int_{-\infty}^0 x d\varphi_b(x)\right)$$

$$2 \int_0^{\infty} F(x) d\varphi_b(x) \leq F\left(2 \int_0^{\infty} x d\varphi_b(x)\right)$$

But

$$2 \int_0^{\infty} x d\varphi_b(x) = -2 \int_{-\infty}^0 x d\varphi_b(x) = \int_{-\infty}^{\infty} |x| d\varphi_b(x)$$

which concludes the proof.

The generalisation of this proposition for non-even distributions



and for more detailed partitions of the support of  $g$  is easy and we leave it to the reader.

#### 4. Variational Principles

The existence of a variational principle for the free energy is a manifestation of equilibrium. Since the quenched system is not in equilibrium we cannot expect that a variational principle completely analogous to that of equilibrium systems will also hold for  $\bar{F}$ . In mean field calculation (Edwards and Anderson<sup>2</sup>, Sherrington and Kirkpatrick<sup>3</sup>) the free energy of the quenched state at low temperature is above the continuation from high temperature, this continuation being essentially the annealed free energy (see Proposition 3.1). In this section we give some insight on this mean field result by showing that the quenched free energy satisfies a variational principle on a subspace  $\mathcal{B}_0$  of the space  $\mathcal{B}$  (2.14); on the other hand the annealed free energy satisfies a variational principle on the whole space  $\mathcal{B}$ .

Let us recall that for given interaction  $J$ , the variational principle for the finite volume free energy is expressed by the inequality

$$F(J) = F[J; g_J] \leq F[J; f] \quad (4.1)$$

where the free energy functional is defined by

$$F[J; f] = \int d\mu(\sigma) f(\sigma) \left[ H(J, \sigma) + \frac{1}{\beta} \ln f(\sigma) \right] \quad (4.2)$$

on the space of distribution functions  $f = f(\sigma)$ , and  $g_J = g(J, \sigma)$  is the Gibbs function (2.4).

Indeed the inequality (4.1) is a consequence of the inequality

$$F[J; f] - F(J) = -\frac{1}{\beta} \int d\mu(\sigma) f(\sigma) \ln \frac{g(J, \sigma)}{f(\sigma)} \geq 0.$$



For random systems we consider the free energy functional (2.15) defined on  $\mathcal{B}$  and we also introduce  $\mathcal{B}_0$  the subspace of  $\mathcal{B}$  defined by:

$$\mathcal{B}_0 = \left\{ f = f(J, \sigma); f(J, \sigma) \geq 0, \int d\mu(\sigma) f(J, \sigma) = 1, \forall J \right\} \quad (4.7)$$

Proposition 4.1

$$1) F_{an} = F[h] = \min_{f \in \mathcal{B}} F[f] \quad (4.4)$$

2) For any  $f \in \mathcal{B}$

$$F[f] - \int d\rho(J) d\mu(\sigma) f(J, \sigma) F(J) \geq 0 \quad (4.5)$$

and the equality holds for  $f = g(J, \sigma)$

$$3) \bar{F} = F[g] = \min_{f \in \mathcal{B}_0} F[f] \quad (4.6)$$

Proof:

1) Using the definitions (2.12), (2.13), (2.15), we have

$$F[f] - F_{an} = -\frac{1}{\beta} \int d\rho(J) d\mu(\sigma) f(J, \sigma) \ln \frac{h(J, \sigma)}{f(J, \sigma)} \geq 0$$

$$2) F[f] - \int d\rho(J) d\mu(\sigma) f(J, \sigma) F(J) =$$

$$= -\frac{1}{\beta} \int d\rho(J) d\mu(\sigma) f(J, \sigma) \ln \frac{g(J, \sigma)}{f(J, \sigma)} \geq 0$$

3) Using (4.5) we have for any  $f \in \mathcal{B}_0$

$$F[f] \geq \bar{F}$$

which concludes the proof.

Another way to formulate the above result is the following: in the space  $\mathcal{B}$  of the joint probability distributions the minimi-



zation of  $F[f]$  yields the annealed free energy, while the quenched free energy is obtained by minimizing the difference (4.5).

To conclude this section we remark that the quenched entropy  $\bar{S}$  and the annealed entropy  $S_{an}$  are not necessarily positive for the scales of the entropies depend on the norm chosen for the a priori measure  $\mu_0$ . Indeed changing  $\mu_0$  to  $C\mu_0$  will change  $S(J)$  (2.6) to  $S(J) + |V| \ln C$ . If the normalisation of  $\mu_0$  is chosen so that  $S(J) \gg 0$  (e.g. if  $g(J, \sigma) \leq 1$  for all  $(J, \sigma)$ ) then the quenched entropy  $\bar{S}$  will be non-negative; however the annealed entropy may still be negative, in particular for large values of  $\beta$ . As an example we mention that for the Ising spin  $\frac{1}{2}$  models, the choice

$$d\mu_0(\sigma) = [\delta(\sigma-1) + \delta(\sigma+1)] d\sigma \quad (4.6)$$

$$d\varphi_b(x) = d\varphi_b(-x)$$

will give

$$F_{an} = -\frac{1}{\beta} \left[ |V| \ln 2 + \sum_{bcv} \ln \int e^{\beta x} d\varphi_b(x) \right] \quad (4.7)$$

and

$$S_{an} = |V| \ln 2 + \sum_b \left[ \ln \int d\varphi_b(x) e^{\beta x} - \frac{\int d\varphi_b(x) e^{\beta x} x}{\int d\varphi_b(x) e^{\beta x}} \right]$$

This shows that  $S_{an}$  goes to  $|V| \ln 2$  if  $\beta$  goes to 0,  $S_{an}$  changes sign with increasing  $\beta$  and goes to  $-\infty$  with  $\beta$  going to infinity\*. Therefore one can imagine that underestimating  $\bar{F}$  with an improper choice of  $f$  in  $\mathcal{B}$  may result that  $F[f]$  approaches  $F_{an}$  sufficiently closely that a negative entropy will be obtained for large values of  $\beta$ . A similar mistake really occurred in the original paper written on the Sherrington-Kirkpatrick model<sup>3</sup>.

\* At least for distributions  $d\varphi_b(x) = \varphi_b(x) dx$ , with  $\varphi_b \in C^1$ .



## 5. Stability of the Gauge Symmetry under Boundary Perturbation

The boundary conditions play a predominant role in the definition of infinite volume equilibrium states. In particular if the hamiltonian has some internal symmetries the interaction with the fixed boundary spins breaks the symmetry and may lead to a "spontaneous breakdown" of this symmetry in the thermodynamic limit.

The random Hamiltonian (2.1) has also internal symmetries. However we shall show in this section that in "pure" models of spin glasses the gauge symmetry cannot be broken by the boundary conditions. This implies that a spin glass cannot generally be characterized by a local order parameter; in fact the order parameter proposed by Edwards and Anderson<sup>2</sup> is the expectation value of a non-local quantity taken with the measure (2.8) (see 6.2). This order parameter can be considered as a local observable only if one introduces the so-called replicas; in this case however one loses the clear description of the quenched states as probability measures. This explains somewhat the difficulty to prove the existence of phase transitions in quenched models.

In this section we consider hamiltonian of the form

$$H(J, \sigma) = - \sum_b \sum_{\alpha=1}^2 J_{b,\alpha} \prod_{i \in b} \sigma_{i,\alpha} = - \sum_b J_b \sigma_b \quad (5.1)$$

where  $\sigma_{i,\alpha}$  is the  $\alpha^{\text{th}}$  component of the spin at site  $i$ , and the  $J_{b,\alpha}$ 's are independent random variables.

We assume that the interactions have finite range, i.e. there exists some  $R > 0$  such that  $J_{b,\alpha}$  is strictly zero for any  $b$  with  $\text{diam}(b) > R$ . Furthermore we consider only "pure" models of Spin glasses, i.e. models with even distribution for each interactions

$$dJ_{b,\alpha}(x) = dJ_{b,\alpha}(-x) \quad (5.2a)$$



Finally we assume that the a priori measure  $\mu_0$  on  $S \subset \mathbb{R}^V$  is even in all components

$$d\mu_0(s_1\sigma_1, \dots, s_V\sigma_V) = d\mu_0(\sigma_1, \dots, \sigma_V) \quad (5.2b)$$

$$\forall (s_1, \dots, s_V) \quad s_i = \pm 1$$

Let  $\mathcal{G} = \left\{ s = \{s_{i,\alpha}\}_{\substack{i \in L \\ 1 \leq \alpha \leq \nu}} ; s_{i,\alpha} = \pm 1 \right\}$  ; for any  $s$  in  $\mathcal{G}$  we introduce the automorphism  $\mathcal{T}_s$  defined on the algebra of local observables by

$$(\mathcal{T}_s f)(J, \sigma) = f(sJ, s\sigma) \quad (5.3)$$

where:  $(sJ)_{b,\alpha} = J_{b,\alpha} \prod_{i \in b} s_{i,\alpha}$

$$(s\sigma)_{i,\alpha} = s_{i,\alpha} \sigma_{i,\alpha}$$

These transformations are internal symmetries of the system, since they leave the Hamiltonian and the measure invariant, i.e.

$$\mathcal{T}_s H = H$$

and  $d\mu(s\sigma) = d\mu(\sigma) \quad d\varrho(sJ) = d\varrho(J)$

Let us note that  $\mathcal{G}$  is a group of gauge transformations since it is generated by transformations involving only a finite number of lattice sites. We shall not consider other symmetries the system might also have.

We are interested in local observables of the form

$$f(J, \sigma) = \prod_{b,\alpha} J_{b,\alpha}^{n_{b,\alpha}} \prod_{i,\beta} \sigma_{i,\beta}^{m_{i,\beta}} \quad (5.4)$$

where  $n_{b,\alpha}$  and  $m_{i,\beta}$  are non-negative integers which are different from zero only for a finite number of  $b$ 's and  $i$ 's. We introduce the notation



$$\text{supp } f = \bigcup_{b: n_{b,\alpha} \neq 0} b \cup \bigcup_{i: m_{i,\beta} \neq 0} \{i\} \subset L$$

For any  $f$  of the form 5.4 we have

$$\begin{aligned} \mathcal{T}_s f &= (-1)^{N_f(s)} f \\ N_f(s) &= \sum n_{b,\alpha} |V_\alpha(s) \cap b| + \sum m_{i,\beta} |V_\beta(s) \cap \{i\}| \\ V_\alpha(s) &= \{j \in L ; s_{j,\alpha} = -1\} \end{aligned}$$

and therefore  $f$  is gauge breaking, i.e.  $\mathcal{T}_s f \neq f$ , if  $N_f(s)$  is odd for some  $s_0$  in  $\mathcal{G}$  in which case

$$\mathcal{T}_s f = -f$$

From the invariance property of  $H_V$  and  $dQ_V$  we have immediately the following result.

Proposition 5.1

Let  $f$  be any gauge-breaking observable of the form (5.4).

Then

$$\int f dQ_V = 0$$

if  $V \supset \text{supp } f$ .

(A similar result was obtained by Avron et al<sup>6</sup> in the case where  $f$  is a pure product of Ising spins).

To show the absence of spontaneous breakdown of this gauge symmetry we consider quenched states with boundary conditions  $dQ_V^{(b.c)}$ . Such states are defined through (2.8) by the hamiltonian

$$H_V = -\sum_{b \cap V \neq \emptyset} J_b \sigma_b$$

together with the boundary conditions

$$\begin{cases} \sigma_i = \hat{\sigma}_i & \text{if } i \notin V \\ \hat{\beta}_{b,\alpha}(x) & \text{if } b \not\subset V \end{cases} \quad (5.5)$$



For example for "free" boundary conditions  $\hat{J}_{b,\alpha}(x) = \delta(x)$  ;  
for "fixed" boundary conditions  $\hat{J}_{b,\alpha}(x) = \delta(x - \hat{J}_{b,\alpha})$  .

Proposition 5.2

Let  $f$  be any gauge-breaking local observable of the form (5.4).  
Then for any boundary conditions (5.5)

$$\langle f \rangle = \int f dQ_V^{(b.c.)} = 0$$

if  $\text{dist}(\text{supp} f, V^c) > R$

Proof:

Let us take  $s_0$  in  $\mathcal{G}$  such that  $\tau_{s_0} f = -f$  and  $s_i = 1$  for  $i$  outside  $\text{supp} f$ . The measure  $dQ_V^{(b.c.)}$  is then invariant under this transformation and yields  $\langle f \rangle = -\langle f \rangle = 0$ .

We notice that this theorem does not exclude the existence of a phase transition in the sense that different weak limits of the quenched measures  $dQ_V^{(b.c.)}$  can be obtained, i.e. the possibility still stands for the non-uniqueness of the expectation value of some local gauge-invariant quantity. However, as far as we know, such a hypothesis has never appeared in the the literature and can be qualified as "unphysical". An eventual spin-glass transition is expected to be characterised by a singularity in the thermodynamic functions and in a non-local order parameter. This we discuss in the following section .

6. The Edwards-Anderson Order Parameter

The order parameter  $q$  proposed by Edwards and Anderson<sup>2</sup> to describe the spin glass transition is defined as  $\langle |\langle \sigma_0 \rangle|^2 \rangle$  where the first average is the thermal one and the second is taken over the interactions. In order to obtain a non zero value



one has to choose some boundary conditions. We take for the distribution of interactions  $J_{b,\alpha}$  across the boundary the same distribution  $d\mathcal{J}_{b,\alpha}$  as the one defining the system and for the external spins some configuration  $\hat{\sigma}$ ; we denote  $dQ_{\hat{\sigma}}^V$  the quenched state associated with this boundary condition. With our notation the order parameter  $q$  is the thermodynamic limit of

$$q_{\hat{\sigma}}^V = \int d\mathcal{J}(\mathcal{J}) \sum_{\alpha=1}^{\nu} \left[ \int d\mu_{\nu}(\sigma) g_{\hat{\sigma}}^V(\mathcal{J}, \sigma) \sigma_{0,\alpha} \right]^2 = \langle |\langle \sigma_0 \rangle_{\hat{\sigma}}^V(\mathcal{J})|^2 \rangle \quad (6.1)$$

provided this limit exists. In (6.1)  $g_{\hat{\sigma}}^V(\mathcal{J}, \sigma)$  is the probability density (2.4) associated with the boundary condition  $\hat{\sigma}$ .

We note that  $q_{\hat{\sigma}}^V$  has two particular features: Firstly  $q_{\hat{\sigma}}^V$  is a non-local observable with respect to the quenched measure. Indeed

$$q_{\hat{\sigma}}^V = \int dQ_{\hat{\sigma}}^V \sigma_0 \langle \sigma_0 \rangle_{\hat{\sigma}}^V(\mathcal{J}) \quad (6.2)$$

and  $\sigma_0 \langle \sigma_0 \rangle_{\hat{\sigma}}^V(\mathcal{J})$  is non-local since its support is the whole volume occupied by the system.

Secondly  $q_{\hat{\sigma}}^V$  is the expectation value of an observable which is invariant with respect to  $\mathcal{G}_V = \{s; s \in \mathcal{G} \quad s_i = 1 \quad \forall i \notin V\}$ . Indeed for any  $s$  in  $\mathcal{G}$  we have

$$\mathcal{E}_s [\sigma_{0,\alpha} \langle \sigma_{0,\alpha} \rangle_{\hat{\sigma}}^V(\mathcal{J})] = \sigma_{0,\alpha} \langle \sigma_{0,\alpha} \rangle_{\hat{\sigma}}^{s\hat{\sigma}}(\mathcal{J})$$

therefore  $\sigma_{0,\alpha} \langle \sigma_{0,\alpha} \rangle_{\hat{\sigma}}^V(\mathcal{J})$  is invariant under any gauge transformation  $s$  in  $\mathcal{G}_V$  (see also Avron et al<sup>6</sup>).

Because of this gauge-invariance character  $q$  will not distinguish between different gauge breaking phases. It is however an order



parameter in the sense that it is zero at high temperatures (Proposition 6.1) and is perhaps non zero at low temperatures if the dimension of the lattice exceeds some finite value. As we shall see below (Proposition 6.2)  $q$  is independent of the boundary conditions  $\hat{\sigma}$ . This behaviour of  $q$  implies a new type of low temperature phase - the spin glass - for the local moment  $\langle \langle \sigma_0 \rangle \rangle$  vanishes at all temperatures in any model satisfying (5.2) (Proposition 6.3).

Proposition 6.1

Let us consider Ising models ( $\sigma_i = \pm 1$ ) with finite range even interactions\* and even distribution of spins and bonds (5.2). Suppose the interactions are independent random variables and the number of different distributions is finite. Then the order parameter  $q$  vanishes for sufficiently high temperature.

Proof:

Using Griffith's inequality<sup>7</sup> and  $|\sigma_0| \leq 1$  we find that

$$q_{\hat{\sigma}} = \langle |\langle \sigma_0 \rangle_{\hat{\sigma}}(J)|^2 \rangle \leq \langle \langle \sigma_0 \rangle_V^+ (|J|) \rangle$$

where  $|J| = \{|J_b|\}$  and + means that the boundary spins are positive. Now let  $R$  be the range of the interaction and  $\lambda = \text{dist}(0, V^c)$ . A generalisation of Fisher's estimate for pair correlations using self-avoiding walks<sup>8</sup> gives

$$\langle \sigma_0 \rangle_V^+ (|J|) \leq \sum_{n \geq \lambda/R} \sum'_{\{b_1 \dots b_n\}} \tanh \beta |J_{b_1}| \dots \tanh \beta |J_{b_n}| \quad (6.3)$$

Here the prime indicates that  $(b_1 \circ \dots \circ b_n) \cap V = \{0\}$  and there is no non-empty subset  $\{b_{i_1}, \dots, b_{i_k}\} \subset \{b_1, \dots, b_n\}$  such that  $b_{i_1} \circ \dots \circ b_{i_k} = \emptyset$  ( $A \circ B = A \setminus B \cup A \cap B$ ). Under reasonable assumption on the lattice and the set of  $\{b\}$  there exists a constant  $c$ , depending on  $L$  and on the interacting sets, such

\* i.e.  $J_b = 0$  if  $|b|$  is odd.



that the number of sets  $\{b_1, \dots, b_n\}$  in (6.3) is smaller than  $C^n$ . Moreover,

$$0 \leq \int_{-\infty}^{\infty} \tanh \beta |x| d\varrho_b(x) \leq \mathcal{E}(\beta) \leq 1$$

where  $\mathcal{E}(\beta)$  goes to zero with  $\beta$  going to zero.

Therefore, for small enough  $\beta$ ,  $C \cdot \mathcal{E}(\beta) < 1$  and

$$\langle \langle \sigma_i \rangle_V^+ (|J|) \rangle \leq \sum_{n \geq \lambda/R} [C \cdot \mathcal{E}(\beta)]^n \leq \frac{[C \cdot \mathcal{E}(\beta)]^{\lambda/R}}{1 - C \cdot \mathcal{E}(\beta)} \quad (6.4)$$

If  $V$  increases then  $\lambda$  tends to  $\infty$  and the r.h.s. goes to zero.

In the following part of this section we discuss some formulas obtained for Ising models by gauge fixing.

Proposition 6.2

Consider an Ising models ( $\sigma_i = \pm 1$ ) with even distribution (5.2) and let  $f(J, \sigma)$  be a function of  $\sigma_i$ 's for  $i \in V$  and of  $J_b$ 's for  $b \cap V \neq \emptyset$ . Then

i) If  $f$  is invariant under  $\mathcal{G}_V$

$$\int dQ_V^{\hat{\sigma}} f = 2^{|V|} \int d\varrho(J) g_V^+(J, +) f(\hat{\sigma}J, +) \quad (6.5)$$

where in  $\hat{\sigma}J$ ,  $\hat{\sigma}$  is extended to  $V$  with the value  $\hat{\sigma}_i = 1$  for  $i \in V$  and  $+$  denotes the configuration  $\sigma_i = +1$ .

ii) If  $f$  is gauge invariant

$$\int dQ_V^{\hat{\sigma}} f = 2^{|V|} \int d\varrho(J) g_V^+(J, +) f(J, +) \quad (6.6)$$

Proof:

Using the invariance property (5.2) and  $\mathcal{T}_s f = f$  for all  $s \in \mathcal{G}_V$  we have:

$$\begin{aligned}
 \int dQ_V^{\hat{\sigma}} f &= \int d\rho(J) d\mu(\sigma) g_V^{\hat{\sigma}}(J, \sigma) f(J, \sigma) \\
 &= \int d\rho(J) d\mu(\sigma) g_V^{\hat{\sigma}}(J, +) f(J, +) \\
 &= \int d\rho(J) d\mu(\sigma) g_V^+(J, +) f(\hat{\sigma}J, +) \\
 &= 2^{|V|} \int d\rho(J) g_V^+(J, +) f(\hat{\sigma}J, +)
 \end{aligned}$$

which concludes the proof of (i) and (ii).

Let us note that according to (6.6) the expectation value of any gauge-invariant quantity is independent of the boundary condition and can be obtained by "fixing the gauge" at  $\sigma_i = +1$  and then averaging with the probability distribution

$$dg_V^+(J) = 2^{|V|} g_V^+(J, +) d\rho(J) \quad (6.7)$$

Proposition 6.3

For Ising models ( $\sigma_i = \pm 1$ ) with even distributions (5.2)

$$\begin{aligned}
 \langle \langle \sigma_A \rangle_V^{\hat{\sigma}} \langle \sigma_B \rangle_V^{\hat{\sigma}} \rangle &= \delta_{A,B} 2^{|V|} \int d\rho(J) g_V^+(J, +) \langle \sigma_A \rangle_V^+ \quad (6.8) \\
 &= \delta_{A,B} \langle \langle \sigma_A \rangle_V^+ \rangle_{g_V^+}
 \end{aligned}$$

where A and B are subsets of V and  $\sigma_A$  denotes  $\prod_{i \in A} \sigma_i$ .

Proof:

For A = B (6.8) follows immediately from (6.6) since

$$\begin{aligned}
 \langle \langle \sigma_A \rangle_V^{\hat{\sigma}} \langle \sigma_B \rangle_V^{\hat{\sigma}} \rangle &= \int dQ_V^{\hat{\sigma}} \sigma_A \langle \sigma_B \rangle_V^{\hat{\sigma}}(J) \quad (6.9) \\
 \mathcal{C}_S(\sigma_A \langle \sigma_A \rangle_V^{\hat{\sigma}}(J)) &= \sigma_A \langle \sigma_A \rangle_V^{\hat{\sigma}}(J)
 \end{aligned}$$

and

$$\langle \sigma_A \rangle_V^{\hat{\sigma}}(\hat{\sigma}J) = \langle \sigma_A \rangle_V^+(J)$$



For  $A \neq B$  we can write

$$\langle \langle \sigma_A \rangle_{\hat{\sigma}_V} \langle \sigma_B \rangle_{\hat{\sigma}_V} \rangle = \int d\mu_V(\sigma) \sigma_A \sigma_B \left\{ \int d\rho(J) g_{\hat{\sigma}_V}^{\hat{\sigma}}(J, \sigma) \sigma_A \langle \sigma_A \rangle_{\hat{\sigma}_V}^{\hat{\sigma}}(J) \right\}$$

using the fact that

$$\sigma_A \langle \sigma_A \rangle_{\hat{\sigma}_V}^{\hat{\sigma}}(J) = \langle \sigma_A \rangle_{\hat{\sigma}_V}^{\hat{\sigma}}(\sigma J)$$

and

$$g_{\hat{\sigma}_V}^{\hat{\sigma}}(J, \sigma) = g_{\hat{\sigma}_V}^{\hat{\sigma}}(\sigma J, +)$$

we find

$$\begin{aligned} \langle \langle \sigma_A \rangle_{\hat{\sigma}_V} \langle \sigma_B \rangle_{\hat{\sigma}_V} \rangle &= \int d\mu_V(\sigma) \sigma_A \sigma_B \left\{ \int d\rho(J) g_{\hat{\sigma}_V}^{\hat{\sigma}}(J, +) \langle \sigma_A \rangle_{\hat{\sigma}_V}^{\hat{\sigma}}(J) \right\} \\ &= 2^{|V|} \delta_{A,B} \left\{ \int d\rho(J) g_{\hat{\sigma}_V}^{\hat{\sigma}}(J, +) \langle \sigma_A \rangle_{\hat{\sigma}_V}^{\hat{\sigma}}(J) \right\} \end{aligned}$$

which concludes the proof.

### Consequences of Proposition 6.3

1. For  $A = B = \{0\}$  we find

$$q_{\hat{\sigma}_V}^{\hat{\sigma}} = \langle \langle \sigma_0 \rangle_{\hat{\sigma}_V}^+ \rangle_{g_V^+} \quad (6.10)$$

which shows that  $q$  is independent of the boundary conditions  $\hat{\sigma}$ .

2. For  $A = \{i\}$  and  $B = \{j\}$  with  $i \neq j$  we have

$$\langle \langle \sigma_i \rangle_{\hat{\sigma}_V} \langle \sigma_j \rangle_{\hat{\sigma}_V} \rangle = 0 \quad (6.11)$$

This heuristically obvious result was used earlier (see e.g. Fischer<sup>9</sup>) to conclude that the quenched susceptibility  $\bar{\chi}$  is proportional to  $1-q$ ; indeed

$$\bar{\chi}_V \sim \frac{1}{|V|} \sum_{i,j \in V} (\langle \langle \sigma_i \sigma_j \rangle \rangle - \langle \langle \sigma_i \rangle \rangle \langle \langle \sigma_j \rangle \rangle)$$

$$= \frac{1}{|V|} \sum_{i \in V} (1 - \langle\langle \sigma_i \rangle^2 \rangle) + \frac{1}{|V|} \sum_{i \neq j} (\langle\langle \sigma_i \sigma_j \rangle \rangle - \langle\langle \sigma_i \rangle \langle\langle \sigma_j \rangle \rangle)$$

Using Proposition 6.3 it follows that both terms in the second summation vanish while  $\langle\langle \sigma_i \rangle^2 \rangle$  tends to  $q$  in the thermodynamic limit, at least for a translationally invariant state.

3. For any A  $\langle\langle \sigma_A \rangle \rangle = 0$  independent of the boundary conditions<sup>6</sup> (see also Prop. 5.2).

The inequality

$$\langle (\langle \sigma_A \rangle_V^{\hat{\sigma}})^2 \rangle = \int dg_V^+(J) \langle \sigma_A \rangle_V^+(J) \geq 0$$

suggests that  $dg_V^+(J)$  favors the ferromagnetic interactions. This is correct in the following sense.

Proposition 6.4

For Ising models with even distributions (5.2),

$$\int J_b dg_V^+(J) \geq 0 \tag{6.12}$$

holds for all interactions.

Proof:

It is sufficient to show that  $g_V^+(J, +)$  is an increasing function of  $J_b$ . Indeed,

$$\begin{aligned} \frac{\partial g_V^+(J, +)}{\partial J_b} &= \frac{\partial}{\partial J_b} \frac{\exp\{\beta \sum_{b' \in V, b' \neq b} J_{b'}\}}{\sum_{\sigma_i: i \in V} \exp\{\beta \sum_{b' \in V, b' \neq b} J_{b'} \sigma_{b'}\}} \\ &= \beta g_V^+(J, +) - \beta g_V^+(J, +) \langle \sigma_b \rangle_V^+ \geq 0. \end{aligned}$$



Corollary

The averaged energy of any bond is non-positive. Indeed, the averaged energy of the bond  $b$  is

$$-\int J_b \sigma_b dQ_V^{\hat{\sigma}} = -\int J_b dg_V^+(\mathcal{J}) \leq 0, \quad (6.13)$$

according to (6.6) and (6.12).

To conclude this section we establish the connection between the order parameter  $q$  and the derivative of the quenched free energy, with respect to an external field  $h$ . Let  $F_V^{\hat{\sigma}}(\mathcal{J})$  be the free energy in volume  $V$  with boundary condition  $\hat{\sigma}$ . Since  $F_V^{\hat{\sigma}}(\mathcal{J})$  is gauge invariant, (6.6) implies

$$\int F_V^{\hat{\sigma}} dQ_V^{\hat{\sigma}} = 2^{|\mathcal{V}|} \int F_V^+(\mathcal{J}) g_V^+(\mathcal{J}, +) d\rho(\mathcal{J}) \quad (6.14)$$

However

$$\begin{aligned} \int F_V^{\hat{\sigma}} dQ_V^{\hat{\sigma}} &= \int F_V^{\hat{\sigma}}(\mathcal{J}) \int g_V^{\hat{\sigma}}(\mathcal{J}, \sigma) d\mu_V(\sigma) d\rho(\mathcal{J}) \\ &= \int F_V^{\hat{\sigma}}(\mathcal{J}) d\rho(\mathcal{J}) = \int F_V^+(\mathcal{J}) d\rho(\mathcal{J}) = \bar{F}_V \end{aligned} \quad (6.15)$$

Therefore the quenched free energy is independent of the boundary condition and

$$\bar{F}_V = 2^{|\mathcal{V}|} \int F_V^+(\mathcal{J}) g_V^+(\mathcal{J}, +) d\rho(\mathcal{J}) \quad (6.16)$$

Let now  $F_V^+(\mathcal{J}, h)$  be the free energy defined by the equation

$$\exp(-\beta F_V^+(\mathcal{J}, h)) = \sum_{\sigma_i: i \in \mathcal{V}} \exp\left(\beta \sum_{b \in \mathcal{V} \neq \phi} J_b \sigma_b + \beta h \sum_{i \in \mathcal{V}} \sigma_i\right)$$

where  $\sigma_i = +1$  if  $i \in \mathcal{V}^c$ .

Let moreover

$$\bar{F}_V(h) = 2^{|\mathcal{V}|} \int F_V^+(\mathcal{J}, h) g_V^+(\mathcal{J}, +) d\rho(\mathcal{J}) \quad (6.17)$$



We should emphasize that in the above definition  $g_v^+$  does not depend on  $h$ . Therefore,  $\bar{F}_v(h)$  is not equal to the quenched free energy in the presence of a non-random external field, though  $\bar{F}_v(0) = \bar{F}_v$ . The reason for the introduction of  $\bar{F}_v(h)$  through Eq. (6.17) is that it is coupled to the averaged order parameter

$$Q_v = \frac{1}{|V|} \sum_{i \in V} q_v(i) \quad (6.18)$$

where  $q_v(i) = 2^{|V|} \int d\beta(J) g_v^+(J,+) \langle \sigma_i \rangle_v^+(J)$

just in the same way as in non-random models the free energy is coupled to the average magnetization: the comparison of Eqs. (6.18) and (6.17) yields

$$-\frac{\partial}{\partial h} \left( \frac{1}{|V|} \bar{F}_v(h) \right) = Q_v \quad (6.19)$$

Therefore, in a translationally invariant phase the order parameter  $q$  can be obtained as the thermodynamic limit of the l.h.s. of Eq. (6.19) - provided that this limit exists.

### 7. Annealed models with one, two and three phase transitions

The annealed models (see Section 2) are usually considered to be trivial and hence of no further interest. This opinion comes from the fact that an annealed model with even distributions for the bonds is in fact equivalent to a model without any interaction (see (4.7) and (7.2)).

Non trivial results can be obtained either by introducing interactions among the bonds or by destroying the symmetry of their a priori distributions. As an example to the former possibility we mention the Ashkin-Teller model<sup>10</sup> in which two consecutive phase transitions were conjectured by Wegner<sup>11</sup> and proved rigorously by Pfister<sup>12</sup>.

Here we exhibit simple examples of annealed models with asymmetric



bond distribution, in which one, two or three phase transitions take place as the temperature changes, depending on the choice of the lattice and of certain parameters.

We shall consider only Ising models, i.e.  $\sigma_i = \pm 1$ ; our discussion is based on the following simple observation:

Let  $f = f(\sigma)$  be a local observable; then

$$\langle f \rangle_{an} = \int dA(J, \sigma) f(\sigma) = \int d\mu(\sigma) f(\sigma) e^{\beta F_{an}} \prod_b \int d\varphi_b(J_b) e^{\beta J_b \sigma_b}$$

But  $\sigma_b = \prod_{i \in b} \sigma_i = \pm 1$  implies

$$\langle f \rangle_{an} = \frac{\int d\mu(\sigma) f(\sigma) e^{\sum_b K_b \sigma_b}}{\int d\mu(\sigma) e^{\sum_b K_b \sigma_b}} \quad (7.1)$$

where 
$$K_b = \frac{1}{2} \ln \left( \frac{\int d\varphi_b(x) e^{\beta x}}{\int d\varphi_b(x) e^{-\beta x}} \right) = K_b(\beta) \quad (7.2)$$

Equation (7.1) shows that the annealed system is equivalent to a spin  $\frac{1}{2}$ -system with the same lattice and bond structure with fixed interactions  $J_b = \frac{1}{\beta} K_b(\beta)$ . Therefore the possible phase transitions of the annealed system can be investigated using the known phase transition of the spin  $\frac{1}{2}$  system. It should be stressed that the interactions  $J_b$  in the corresponding model are  $\beta$  dependent and this will lead to the existence of several phase transitions. One should also note that

$$K_b(0) = 0$$

and  $\text{sign } K_b = \text{sign} \left( \int d\varphi(x) \text{sh}(\beta x) \right)$

In particular for small  $\beta$

$$\text{sign } K_b = \text{sign} \left( \int d\varphi(x) \cdot x \right)$$

To illustrate the possible existence of several phase transitions we restrict ourselves to the simplest annealed Ising models

$$H = - \sum_{\langle ij \rangle} J_{ij} \sigma_i \sigma_j \quad (7.3a)$$

with nearest neighbour interactions distributed according to

$$g(J_{ij}) = p \delta(J_{ij} - a) + (1-p) \delta(J_{ij} + b); a, b > 0 \quad (7.3b)$$

and the a priori even measure  $\mu_0$


$$\mu_0(\sigma) = \delta(\sigma - 1) + \delta(\sigma + 1) \quad (7.3c)$$

In this case (7.2) yields

$$K_{ij} = K = \frac{1}{2} \ln \left( \frac{p e^{\beta a} + (1-p) e^{-\beta b}}{p e^{-\beta a} + (1-p) e^{\beta b}} \right) \quad (7.4)$$

Therefore for small  $\beta$   $\text{sign } K = \text{sign} \left( a - \frac{1-p}{p} b \right)$

$$\begin{aligned} \text{for large } \beta \quad K &\sim \frac{1}{2} \beta \cdot \ln(a-b) && \text{if } a > b \\ &-\frac{1}{2} \beta \cdot \ln(b-a) && \text{if } a < b \\ &\frac{1}{2} \ln \left( \frac{p}{1-p} \right) && \text{if } a = b \end{aligned}$$





I)  $a > b$

II)  $a = b$

III)  $a < b$

i) ———  $a > \frac{1-p}{p} b$

i) ———  $p > \frac{1}{2}$

i) ———  $a > \frac{1-p}{p} b$

ii) - - - -  $a < \frac{1-p}{p} b$

ii) - - - -  $p < \frac{1}{2}$

ii) - - - -  $a < \frac{1-p}{p} b$

In conclusion if the spin  $\frac{1}{2}$ -system has an ordered phase for  $|K| > K_c$ , then the annealed system will have at least one phase transition in the case I and III; it will have at least one phase transition in the case II if  $\frac{1}{2} |\ln \frac{p}{1-p}| > K_c$ ; it will have at least three phase transition in the case I-ii if  $K_{\min} < -K_c$  and in the case III-i if  $K_{\max} > K_c$ .

Proposition 7.1

Let us consider the annealed Ising model (7.3) with  $a = b = \bar{J}$  on a  $d$ -dimensional simple cubic lattice with  $d \geq 2$ . Let  $K_c$  be the critical value of the spin  $\frac{1}{2}$  model with  $J_{ij} = J > 0$ .

Then:

i) For  $p > p_c = \frac{e^{2K_c}}{1 + e^{2K_c}}$  (7.5)

there exists two ferromagnetically ordered phases for

$$\beta > \beta_p = \frac{1}{2\bar{J}} \ln \left( \frac{p(1 + e^{2K_c}) - 1}{p(1 + e^{2K_c}) - e^{2K_c}} \right)$$

ii) for  $p < 1-p_c$  there exists two antiferromagnetically ordered phases for  $\beta > \beta_{(1-p)}$ .

Let us recall that for  $d = 2$  we have  $\text{sh } 2 K_c = 1$  which yields  $p_c = \frac{1}{\sqrt{2}}$ .

Proof:

For a given  $p$ ,  $K(\beta)$  given by (7.4) is positive monotonically increasing if  $p > \frac{1}{2}$  (resp. negative monotonically decreasing if



$p < \frac{1}{2}$ ) and (7.5) implies that  $K(\beta) > K_c$  for  $\beta > \beta_p$  (resp.  $K(\beta) < -K_c$  for  $\beta > \beta_{1-p}$ ) which concludes the proof.

Proposition 7.2

Let us consider the annealed Ising model (7.3) with  $b > a > 0$ ; then there exists some  $p_c > \frac{1}{2}$  such that for  $p > p_c$

i) on the  $d$ -dimensional simple cubic lattice with  $d \geq 2$ , there exists  $0 < \beta_0 < \beta_1 < \beta_2 < \infty$  such that for  $\beta < \beta_0$  and for  $\beta_1 < \beta < \beta_2$  there exists a unique (paramagnetic) equilibrium state, for  $\beta_0 < \beta < \beta_1$ , there exists a ferromagnetic ordering and for  $\beta > \beta_2$  there exists an anti-ferromagnetic ordering.

ii) on the 2-dimensional triangular lattice there exists  $0 < \beta_0 < \beta_1 < \infty$  such that for  $\beta < \beta_0$  and for  $\beta > \beta_1$  there exists a unique (paramagnetic) equilibrium state while for  $\beta_0 < \beta < \beta_1$ , there exists a ferromagnetic ordering.

Proof:

Let  $\xi = \frac{p}{1-p}$ ; for fixed  $\beta$ ,  $K(\xi, \beta)$  Eq. (7.4) is an increasing function of  $\xi$  which tends to  $\beta a$  as  $\xi$  tends to infinity; therefore for any  $\alpha$   $\max_p K(\beta, \xi) > K_c$  if  $\xi > \xi_c$ . Furthermore  $\frac{\partial K}{\partial \beta} = 0$  if  $\beta$  is the solution of

$$\text{ch}(a + b)\beta = \frac{a\xi^2 - b}{(b - a)\xi}$$

which is uniquely specified by  $(a, b, p)$ ; therefore for given  $p$ ,  $K(\beta)$  is a concave function which shows that there exists exactly two values  $\beta_0$  and  $\beta_1$  such that  $K(\beta) = K_c$ .

Now on simple cubic lattices the critical temperatures are determined by  $K(\beta_0) = K(\beta_1) = K_c$  and  $K(\beta_2) = -K_c$ . For the triangular lattice  $K(\beta_0) = K(\beta_1) = K_c$  and there is some  $\tilde{\beta}$  so that  $K(\beta) < 0$  for  $\beta > \tilde{\beta}$ . However, the antiferromagnetic



model does not undergo any phase transition at  $\beta < \infty$  (Wannier<sup>13</sup>).


#### Acknowledgement

One of the authors (A.S.) wishes to thank the Institut de Physique Théorique, Université de Lausanne and particularly Profs P. Erdős, J.-J. Loeffel, G. Wanders and F. Rothen for their kind hospitality during his stay in Lausanne. He is also indebted to the members of the Institut de Physique Théorique, EPF Lausanne for many useful discussions.

References

1. R. Brout, Phys. Rev. 115, 824 (1959)
2. S.F. Edwards and P.W. Anderson, J. Phys. F 5, 965 (1975)
3. D. Sherrington and S. Kirkpatrick, Phys. Rev. Letters 35,  
1792 (1975)  
G. Parisi, Phys. Reports 67, 25 (1980)  
G. Parisi and G. Toulouse, J. Physique Lettres 41, L361 (1980)
4. S.G. Rosa, Physics Letters 78A, 468 (1980)
5. W. Feller, An Introduction to probability theory and its  
application, New York-London: Wiley (1971)
6. J.E. Avron, G. Roepstorff and L.S. Schulman, J. Stat. Phys. 26,  
25 (1981)
7. R. Griffiths, in Statistical Mechanics and Quantum Field Theory,  
(Les Houches) New York, London, Paris: Gordon and  
Breach (1970)
8. M.E. Fisher, Phys. Rev. 162, 475 (1967)
9. K.H. Fischer, Phys. Rev. Letters 34, 1438 (1975); Solid St. Commun.  
18, 1515 (1976)
10. J. Ashkin, E. Teller, Phys. Rev. 64, 178 (1943)
11. F.J. Wegner, J. Phys. C 5, L131 (1972)
12. C.-E. Pfister, to appear (1982)
13. G.H. Wannier, Phys. Rev. 79, 357 (1950)





63.297



Kiadja a Központi Fizikai Kutató Intézet  
Felelős kiadó: Kroó Norbert  
Szakmai lektor: Dr. Forgács Gábor  
Nyelvi lektor: Dr. Túttó István  
Példányszám: 570 Törzsszám: 82-573  
Készült a KFKI sokszorosító üzemében  
Felelős vezető: Nagy Károly  
Budapest, 1982. november hó