



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



Math 5058.32 . 3



SCIENCE CENTER LIBRARY

BOUGHT WITH THE INCOME

FROM THE BEQUEST OF

PROF. JOHN FARRAR, LL.D.


AND HIS WIDOW


ELIZA FARRAR

FOR

"BOOKS IN THE DEPARTMENT OF MATHEMATICS,
ASTRONOMY, AND NATURAL PHILOSOPHY"







IOANNIS BOLYAI DE BOLYA

APPENDIX

SCIENTIAM SPATII ABSOLUTE VERAM EXHIBENS.

0

IOANNIS BOLYAI DE BOLYA
APPENDIX

SCIENTIAM SPATII ABSOLUTE VERAM EXHIBENS: A VERITATE AUT FALSITATE
AXIOMATIS XI. EUCLIDEI, A PRIORI HAUD UNQUAM DECIDENDA, INDEPENDENTEM:
ADIECTA AD CASUM FALSITATIS QUADRATURA CIRCULI GEOMETRICA.

EDITIO NOVA

OBLATA AB ACADEMIA SCIENTIARUM HUNGARICA
AD DIEM NATALEM CENTESIMUM AUCTORIS CONCELEBRANDUM.

EDIDERUNT

IOSEPHUS KÜRSCHÁK MAURITIUS RÉTHY
BÉLA TÓTÖSSY DE ZEPETHNEK

ACADEMIÆ SCIENTIARUM HUNGARICÆ SODALES.



LIPSIAE,
IN ÆDIBUS B. G. TEUBNERI.

*

[BUDAPESTINI, SUMPTIBUS ACADEMIÆ SCIENTIARUM HUNGARICÆ.]

MCMIII.

Math ~~5038.32.8~~
v 5858.32.3




Favor fund

510

INDEX TABULARUM

ET PAGINARUM, QUIBUS FIGURÆ TABULARUM TRACTANTUR.

Tab.	Fig.	Pag.	Tab.	Fig.	Pag.	Tab.	Fig.	Pag.
I.	1	3	III.	8	8	V.	14	15
	1	4		9	11		14	24
	2	3		9	22		14	29
	2	4		10	13		14	33
	3	4		10	23		15	16
	4	5		10	28		15	32
II.	5	5	IV.	11	13		15	33
	5	9		12	14	VI.	16	18
	5	10		12	21		17	20
	6	6		12	24		18	27
	6	11		12	27	VII.	19	29
	7	6		13	15		20	30
	7	9					21	31
	7	10					22	31
	7	28					23	34



A P P E N D I X.

SCIENTIAM SPATII *absolute veram* exhibens:
a veritate aut falsitate Axiomatis XI Euclidei
(a priori haud unquam decidenda) in-
dependentem; adjecta ad casum fal-
sitatis, quadratura circuli
geometrica.

Auctore JOHANNE BOLYAI de eadem, Geometrarum
in Exercitu Caesareo Regio Austriaco Ca-
strensium Capitaneo.



EXPLICATIO SIGNORUM.

- \overline{ab} denotet complexum *omnium* punctorum cum punctis a, b in recta sitorum.
- \overline{ab} • rectæ \overline{ab} in a bifariam sectæ dimidium illud, quod punctum b complectitur.
- \overline{abc} • complexum *omnium* punctorum, quæ cum punctis a, b, c (non in eadem recta sitis) in eodem plano sunt.
- $ab\overline{c}$ • plani \overline{abc} per \overline{ab} bifariam secti dimidium, punctum c complectens.
- abc • portionum, in quas \overline{abc} per complexum rectarum $b\overline{a}, b\overline{c}$ dividitur, *minorem*; sive *angulum*, cuius $b\overline{a}, b\overline{c}$ crura sunt.
- $abcd$ • (si d in abc sit et $b\overline{a}, c\overline{d}$ se invicem non secent) portionem ipsius abc inter $b\overline{a}, bc, c\overline{d}$ comprehensam; $bac\overline{d}$ vero portionem plani \overline{abc} inter $\overline{ab}, c\overline{d}$ sitam.
- R • angulum rectum.
- $ab \triangleq cd$ • $cab = acd$.
- \equiv • congruens.*
- $x \rightsquigarrow a$ • x tendere ad limitem a .
- $\bigcirc r$ • peripheriam circuli radii r .
- $\odot r$ • aream circuli radii r .

* Sit fas, signo hocce, quo summus Geometra GAUSS numeros congruos insignivit, congruentiam geometricam quoque denotare: nulla ambiguitate exinde metuenda.



§. 1.

Si rectam am non secet plani eiusdem recta $b\bar{n}$, at secet quævis $b\bar{p}$ Fig. 1.
(in abn): designetur hoc per

$$bn \parallel am.$$

Dari talem $b\bar{n}$, et quidem *unicam*, e quovis puncto b (extra \bar{am}),
atque

$$bam + abn \text{ non } > 2R$$

esse patet; nam bc circa b mota, donec

$$bam + abc = 2R$$

fiat, $b\bar{c}$ aliquando *primo* non secat \bar{am} , estque tunc $bc \parallel am$.

Nec non patet esse $bn \parallel em$, ubivis sit e in \bar{am} (supponendo in omnibus talibus casibus esse $am > ae$).

Et si, puncto c in \bar{am} abeunte in infinitum, semper sit $cd = cb$: erit semper

$$cdb = (cbd < nbc);$$

ast $nbc \rightarrow 0$; adeoque et $adb \rightarrow 0$.

§. 2.

Si $bn \parallel am$; *est quoque* $cn \parallel am$.

Fig. 2.

Nam sit d ubicunque in $macn$. Si c in $b\bar{n}$ sit; $b\bar{d}$ secat \bar{am} (propter $bn \parallel am$), adeoque et $\bar{c}\bar{d}$ secat \bar{am} ; si vero c in $b\bar{p}$ fuerit; sit $bq \parallel c\bar{d}$: cadit $b\bar{q}$ in abn (§. 1.) secatque \bar{am} , adeoque et $\bar{c}\bar{d}$ secat \bar{am} . Quævis $\bar{c}\bar{d}$ igitur (in acn) secat in utroque casu \bar{am} absque eo, ut $c\bar{n}$ ipsam \bar{am} secet. Est ergo semper $cn \parallel am$.

§. 3.

Fig. 2. *Si tam br quam cs sit $\parallel am$, et c non sit in \overline{br} ; tum $b\overline{r}$, $c\overline{s}$ se invicem haud secant.*

Si enim $b\overline{r}$, $c\overline{s}$ punctum d commune haberent; (per §. 2.) essent dr et ds simul $\parallel am$, caderetque (§. 1.) $d\overline{s}$ in $d\overline{r}$ et c in \overline{br} (contra hyp.).

§. 4.

Fig. 3. *Si $man > mab$; pro quovis puncto b ipsius \overline{ab} datur tale c in \overline{am} , ut sit $bcm = nam$.*

Nam datur (per §. 1.) $b\overline{dm} > nam$, adeoque $m\overline{dp} = man$, caditque b in $n\overline{adp}$. Si igitur nam iuxta am feratur, usquequo $a\overline{fi}$ in $d\overline{p}$ veniat; aliquando $a\overline{fi}$ per b transiisse, et aliquod $bcm = nam$ esse oportet.

§. 5.

Fig. 1. *Si $bn \parallel am$, datur tale punctum f in \overline{am} , ut sit $f\overline{m} \triangleq bn$.*

Nam (per §. 1.) datur $bcm > cbn$; et si $ce = cb$, adeoque $ec \triangleq bc$; patet esse $bem < ebn$. Feratur p per ec , angulo bpm semper u , et angulo pbn semper v dicto; patet u esse prius ei simultaneo v minus, posterius vero esse maius. Crescit vero u a bem usque bcm continuo; cum (per §. 4.) nullus angulus $> bem$ et $< bcm$ detur, cui u aliquando = non fiat; pariter decrescit v ab ebn usque cbn continuo: datur itaque in ec tale f , ut $b\overline{fm} = f\overline{bn}$ sit.

§. 6.

Si $bn \parallel am$, atque ubivis sit e in \overline{am} et g in \overline{bn} : tum $gn \parallel em$ et $em \parallel gn$.

Nam (per §. 1.) est $bn \parallel em$, et hinc (per §. 2.) $gn \parallel em$.

Si porro $f\overline{m} \triangleq bn$ (§. 5.); tum $m\overline{f}bn = n\overline{b}fm$, adeoque (cum $bn \parallel f\overline{m}$ sit) etiam $f\overline{m} \parallel bn$, et (per præc.) $em \parallel gn$.

§. 7.

Si tam bn quam cp sit || am, et c non sit in bn̄: est etiam bn || cp. Fig. 4.

Nam bn̄, cp̄ se invicem non secant (§. 3.); sunt vero am, bn, cp aut in plano, aut non; atque in casu primo am aut in bn̄cp est, aut non.

Si am, bn, cp in plano sint, et am in bn̄cp cadat; tum quævis bq̄ (in nbc̄) secat am̄ in aliquo puncto d̄ (quia bn || am); porro cum dm̄ || cp sit (§. 6.), patet d̄q̄ secare cp̄, adeoque esse bn || cp.

Si vero bn, cp in eadem plaga ipsius am sint; tum aliqua earum ex. gr. cp *intra* duas reliquas bn̄, am̄ cadit; quævis bq̄ (in nba) autem secat am̄, adeoque et ipsam cp̄. Est itaque bn || cp.

Si mab, mac *angulum* efficiant: tum cbn cum abn non nisi bn̄, am̄ vero (in abn) cum bn̄, adeoque nbc̄ quoque cum am̄, nihil commune habent. Per quamvis bd̄ (in nba) autem positum bc̄d̄ secat am̄, quia (propter bn || am) bd̄ secat am̄. Moto itaque bc̄d̄ circa bc, donec ipsam am̄ *prima vice* deserat, postremo cadet bc̄d̄ in bc̄n̄. Eadem ratione cadet idem in bc̄p̄; cadit igitur bn in bc̄p̄. Porro si br || cp; tum (quia etiam am || cp) pari ratione cadit br in bam; nec non (propter br || cp) in bcp. Itaque b̄r̄ ipsis mab, pcb commune, nempe ipsum bn̄ est, atque hinc bn || cp.

Si igitur cp || am, et b extra am̄ sit: tum sectio ipsorum bam, bcp, nempe bn̄ est || tam ad am, quam ad cp.*

§. 8.

Si bn || et = cp (vel brevius bn || = cp), atque am (in nbc̄p) rectam bc̄ perpendiculariter bisecet; tum bn || am. Fig. 5.

Si enim bn̄ secaret am̄, etiam cp̄ secaret am̄ in eodem puncto (cum mabn = macp), quod et ipsis bn̄, cp̄ commune esset, quamvis bn || cp sit. Quævis bq̄ (in cbn) vero secat cp̄; adeoque secat bq̄ etiam am̄. Consequenter bn || am.

* Casu tertio *praemisso* duo priores, adinstar casus secundi §. 10. brevius ac elegantius simul absolvi possunt. (Ed. I. Tom. I. Errata Appendicis).

§. 9.

Fig. 6. Si $bn \parallel am$, $map \perp mab$, atque *angulus, quem* nbd cum nba (in ea plaga ipsius $mabn$, ubi map est) *facit, sit* $< R$: tum map et nbd *se invicem secant.*

Nam sit

$$bam = R, \text{ ac } \perp bn$$

(sive in b cadat c , sive non), et

$$ce \perp bn \text{ (in } nbd);$$

erit (per hyp.) $ace < R$, et $af (\perp ce)$ in ace cadet. Sit $a\tilde{p}$ sectio (punctum a commune habentium) $ab\tilde{f}$ et $am\tilde{p}$; erit

$$bap = bam = R$$

(cum sit $bam \perp map$). Si denique $ab\tilde{f}$ in abm ponatur (a et b manentibus); cadet $a\tilde{p}$ in am ; atque cum

$$ac \perp bn \text{ et } af < ac$$

sit, patet af *intra* bfi terminari, adeoque bf in abn cadere. Secat autem $b\tilde{f}$ ipsam $a\tilde{p}$ in *hoc* situ (quia $bn \parallel am$), adeoque etiam in situ *primo* $a\tilde{p}$ et $b\tilde{f}$ se invicem secant; estque punctum sectionis ipsis map et nbd commune: secant itaque map et nbd se invicem.

Facile exhinc sequitur map et nbd se mutuo secare, si summa inter-norum, quos cum $mabn$ efficiunt, $< 2R$ sit.

§. 10.

Fig. 7. Si tam bn quam cp sit $\parallel \simeq am$; est etiam $bn \parallel \simeq cp$.

Nam mab et mac aut *angulum* efficiunt, aut in plano sunt.

Si prius; bisecet qdf rectam ab perpendiculariter; erit $dq \perp ab$, adeoque $dq \parallel am$ (§. 8.); pariter si ers bisecet rectam ac perpendiculariter, est $er \parallel am$; unde $dq \parallel er$ (§. 7.). Facile hinc (per §. 9.) consequitur, qdf

et $\overline{a\Gamma s}$ se mutuo secare, et sectionem \overline{fs} esse $\parallel \delta q$ (§. 7.), atque (propter $bn \parallel \delta q$) esse etiam

$$fs \parallel bn.$$

Est porro (pro quovis puncto ipsius \overline{fs})

$$fb = fa = fc,$$

caditque \overline{fs} in planum \overline{tgf} , rectam bc perpendiculariter bisecans. Est vero (per §. 7.) (cum sit $fs \parallel bn$) etiam

$$gt \parallel bn.$$

Pari modo demonstratur $gt \parallel cp$ esse. Interim gt bisecat rectam bc perpendiculariter; adeoque $tgbn = tgcp$ (§. 1.) et

$$bn \parallel \sphericalangle cp.$$

Si bn , am , cp in plano sint; sit (*extra hoc planum cadens*) $fs \parallel \sphericalangle am$; tum (per præc.) $fs \parallel \sphericalangle$ tam ad bn quam ad cp , adeoque et $bn \parallel \sphericalangle cp$.

§. 11.

Complexus puncti a , atque *omnium* punctorum, quorum quodvis b tale est, ut si $bn \parallel am$ sit, sit etiam $bn \sphericalangle am$; dicatur F : sectio vero ipsius F cum quovis plano rectam am complectente nominetur L .

In quavis recta, quæ $\parallel am$ est, F gaudet puncto, et nonnisi uno; atque patet L per am dividi in duas partes congruentes; dicatur am *axis* ipsius L ; patet etiam, in quovis plano rectam am complectente, pro *axe* am unicum L dari. Quodvis eiusmodi L , dicatur L *ipsius* am (in plano, de quo agitur, intelligendo). Patet per L circa am revolutum, F describi, cuius am *axis* vocetur, et vicissim F *axi* am *attribuatur*.

§. 12.

Si b ubivis in L ipsius am fuerit, et $bn \parallel \sphericalangle am$ (§. 11.); tum L ipsius am et L ipsius bn coincidunt.

Nam dicatur L ipsius $h\bar{n}$ distinctionis ergo l : sitque c ubivis in L , et $cp = bn$ §. 11.: erit cum et $bn = am$ sit $cp = am$ §. 12., adeoque c etiam in L cadet. Et si c ubivis in L sit, et $cp = am$: tum $cp = bn$ §. 12.; caditque c etiam in l §. 11.. Itaque L et l sunt eadem: ac quævis $h\bar{n}$ est etiam axis ipsius L , et inter omnes axes ipsius L , \equiv est. Idem de F eodem modo patet.

§. 13.

Fig. 2. Si $bn = am$, $cp = dq$, et $bam + abn = 2R$ sit; tum etiam $dcp + cdq = 2R$. Sit enim $ea = eb$ et $efm = dcp$ §. 4.: erit cum

$$\begin{aligned} \text{sit} \quad & bam + abn = 2R = abn + abg \\ & ebg = eaf; \\ \text{adeoque si etiam } & bg = af \text{ sit,} \end{aligned}$$

$$\triangle ebg \cong \triangle eaf, \quad beq = aef,$$

cadetque g in $f\bar{e}$. Est porro $gfm + fgn = 2R$ quia $eqb = efa$. Est etiam $gn = fm$ §. 6.; itaque si $mfrs = pcdq$, tum $rs = gn$ §. 7., et r in vel extra fg cadit (si cd non $= fg$, ubi res iam patet).

I. In casu primo est frs non $> 1/2R - rfm = fgn$, quia $rs = fm$; ast cum $rs = gn$ sit, est etiam frs non $< fgn$; adeoque $frs = fgn$, et

$$rfm + frs = gfm + fgn = 2R.$$

Itaque et $dcp + cdq = 2R$.

II. Si r extra fg cadat; tunc $ngr = mfr$, sitque $mfgn = nghl = lhfo$ et ita porro, usquequo $fl =$ vel prima vice $> fr$ fiat. Est heic $fo = hl = fm$ (§. 7.). Si f in r cadat; tum fo in rs cadit (§. 1.); adeoque

$$rfm + frs = ffm + ffo = ffm + fgn = 2R;$$

si vero r in $h\bar{f}$ cadat, tum (per I.) est

$$rhl + hrs = 2R = rfm + frs = dcp + cdq.$$

§. 14.

Si $bn \parallel am$, $cp \parallel dq$, et $bam + abn < 2R$ sit; tum etiam $dcp + cdq < 2R$.

Si enim $dcp + cdq$ non esset $<$, adeoque (per §. 1.) esset $= 2R$; tum (per §. 13.) etiam $bam + abn = 2R$ esset (contra hyp.).

§. 15.

Perpensis §§. 13. et 14. *Systema Geometriae hypotesi veritatis Axiomatis Euclidei XI. insistens dicatur Σ ; et hypotesi contrariae superstructum sit S . Omnia, quae expresse non dicentur, in Σ vel in S esse; absolute enuntiari, i. e. illa, sive Σ sive S reipsa sit, vera asseri intelligatur.*

§. 16.

Si am sit axis alicuius L ; tum L in Σ recta $\perp am$ est.

Fig. 5.

Nam sit e quovis puncto b ipsius L axis bn ; erit in Σ

$$bam + abn = 2bam = 2R,$$

adeoque $bam = R$. Et si c quodvis punctum in \overline{ab} sit, atque $cp \parallel am$; est (per §. 13.) $cp \cong am$, adeoque c in L (§. 11.).

In S vero nulla 3 puncta a, b, c ipsius L vel F in recta sunt.

Nam aliquis axium am, bn, cp (ex. gr. am) intra duos reliquos cadit; et tunc (per §. 14.) tam bam quam $cam < R$.

§. 17.

L est etiam in S linea, et F superficies.

Fig. 7.

Nam (per §. 11.) quodvis planum ad axem am (per punctum aliquod ipsius F) perpendiculare secat ipsum F in peripheria circuli, cuius planum (per §. 14.) ad nullum alium axem $bñ$ perpendiculare est. Revolvatur F circa bn ; manebit (per §. 12.) quodvis punctum ipsius F in F , et sectio ipsius F cum plano ad $bñ$ non perpendiculari describet super-

Let a, b, c be the sides of a triangle $\triangle ABC$ and A, B, C the angles opposite to them respectively. Then we have

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

§ 12

If a line is drawn from the vertex of a triangle perpendicular to the base, then the sum of the squares of the two sides is equal to the square of the base plus four times the square of the distance from the vertex to the foot of the perpendicular.

Let ABC be a triangle with the perpendicular AD from A to BC . Then $AB^2 = AD^2 + BD^2$ and $AC^2 = AD^2 + DC^2$. Adding these two equations, we get $AB^2 + AC^2 = 2AD^2 + BD^2 + DC^2$. But $BD + DC = BC$, so $BD^2 + DC^2 = BC^2 - 2BD \cdot DC$. Therefore $AB^2 + AC^2 = 2AD^2 + BC^2 - 2BD \cdot DC$. Since $AD \perp BC$, $AD \cdot DC = BD \cdot DC$. Hence $AB^2 + AC^2 = 2AD^2 + BC^2 - 2AD \cdot DC = BC^2 + 4AD^2$.

This is the well-known theorem of Apollonius. It can be used to find the length of the altitude of a triangle if the lengths of the two sides and the base are known.

§ 13

The perpendicular bisectors of the three sides of a triangle are concurrent at a point called the circumcenter, which is the center of the circumcircle of the triangle.

Let ABC be a triangle and M, N, P the midpoints of the sides BC, CA, AB respectively. The perpendicular bisectors AM, BN, CP are concurrent at a point O . To see this, note that $AM \perp BC$ and $BN \perp AC$. The intersection O of AM and BN is equidistant from A, B, C . Hence O lies on CP as well. Therefore O is the circumcenter.

§ 14

The perpendicular bisectors of the three sides of a triangle are concurrent at a point called the circumcenter, which is the center of the circumcircle of the triangle.

* This is a special case of a more general theorem which states that the perpendicular bisectors of the sides of any polygon are concurrent if and only if the polygon is cyclic.

quivis angulus Llineus in F angulo planorum ad F per crura perpendiculariarum aequalis est.

§. 21.

Duae lineae Lformes $a\bar{p}$, $b\bar{d}$ in eodem F, cum tertia Lformi ab summam internorum $< 2R$ efficientes, se mutuo secant (per $a\bar{p}$ in F intelligendo L per a, p ductum, per $a\bar{p}$ vero dimidium illud eius ex a incipiens, in quod p cadit). Fig. 6.

Nam si am , bn axes ipsius F sint; tum $am\bar{p}$, $bn\bar{d}$ secant se invicem (§. 9.); atque F secat eorundem sectionem (per §§. 7. et 11.); adeoque et $a\bar{p}$, $b\bar{d}$ se mutuo secant.

Patet exhinc Axioma XI. et omnia, quæ in Geometria Trigonometrica-que (plana) asseruntur, absolute constare in F, rectarum vices lineis L subeuntibus: idcirco functiones trigonometricæ abhinc eodem sensu accipientur, quo in Σ' veniunt; et periphæria circuli, cuius radius Lformis $= r$ in F, est $= 2\pi r$, et pariter $\odot r$ (in F) $= \pi r^2$ (per π intelligendo $\frac{1}{2} \odot 1$ in F, sive notum 3.1415926 . . .).

§. 22.

Si $a\bar{b}$ fuerit L ipsius $a\bar{m}$, et c in $a\bar{m}$; atque angulus cab (e recta $a\bar{m}$ et Lformi linea $a\bar{b}$ compositus) feratur prius iuxta $a\bar{b}$, tum iuxta $b\bar{a}$ semper porro in infinitum: erit via $c\bar{d}$ ipsius c linea L ipsius $c\bar{m}$. Fig. 9.

Nam (posteriore l dicta) sit punctum quodvis d in $c\bar{d}$, $dn \parallel cm$, et b punctum ipsius L in \bar{dn} cadens; erit $bn \triangleq am$, et $ac = bd$, adeoque $dn \triangleq cm$, consequ. d in l. Si vero d in l et $dn \parallel cm$, atque b punctum ipsius L ipsi \bar{dn} commune sit; erit $am \triangleq bn$ et $cm \triangleq dn$, unde manifesto $bd = ac$, cadetque d in viam puncti c, et sunt l et $c\bar{d}$ eadem. Designetur tale l per l \parallel L.

§. 23.

Si linea Lformis $c\bar{d}f \parallel abe$ (§. 22.), et $ab = be$, atque $a\bar{m}$, $b\bar{n}$, $e\bar{p}$ sint axes; erit manifesto $c\bar{d} = d\bar{f}$; et si quælibet 3 puncta a, b, e fuerint ipsius

1. The first part of the document discusses the importance of maintaining accurate records of all transactions and activities. It emphasizes that this is crucial for ensuring transparency and accountability in the organization's operations.

2. The second part of the document outlines the various methods and tools used to collect and analyze data. It highlights the need for consistent data collection procedures and the use of advanced analytical techniques to derive meaningful insights from the data.

3. The third part of the document focuses on the role of technology in data management and analysis. It discusses how modern software solutions can streamline data collection, storage, and processing, thereby improving efficiency and reducing the risk of human error.

4. The fourth part of the document addresses the challenges associated with data security and privacy. It stresses the importance of implementing robust security measures to protect sensitive information from unauthorized access and data breaches.

5. The fifth part of the document discusses the ethical considerations surrounding data collection and analysis. It emphasizes the need for transparency in data handling practices and the importance of obtaining informed consent from individuals whose data is being collected.

6. The sixth part of the document provides a summary of the key findings and recommendations. It reiterates the importance of a data-driven approach to decision-making and offers practical advice on how to implement the discussed strategies effectively.

7. The final part of the document concludes with a call to action, encouraging all stakeholders to embrace a culture of data literacy and to work together to maximize the value of the organization's data assets.

unde $ambn \equiv amep$ erit, etsi hoc illius qualevis multipulum sit; quod singulare quidem est, sed absurditatem ipsius S evidenter non probat.

§. 25.

In quovis rectilineo triangulo sunt peripheriae radiorum lateribus aequalium, uti sinus angulorum oppositorum. Fig. 10.

Sit enim $abc = R$, et $am \perp bac$, atque sint $bn, cp \parallel am$; erit $cab \perp ambn$, adeoque (cum $cb \perp ba$ sit) $cb \perp ambn$, consequ. $cpbn \perp ambn$. Secet F ipsius cp rectas $\overline{bn}, \overline{am}$ (respective) in d, e , et fascias $cpbn, cpam, bnam$ in lineis L formibus cd, ce, de ; erit (§. 20.) $cde =$ angulo ipsorum ndc, nde , adeoque $= R$; atque pari ratione est $ced = cab$.

Est autem (per §. 21.) in L lineo triangulo ced (heic radio semper $= 1$ posito)

$$ec : dc = 1 : \sin. dec = 1 : \sin. cab.$$

Est quoque (per §. 21.)

$$\begin{aligned} ec : dc &= \odot ec : \odot dc \text{ (in } F) \\ &= \odot ac : \odot bc \text{ (§. 18.);} \end{aligned}$$

adeoque est etiam

$$\odot ac : \odot bc = 1 : \sin. cab;$$

unde assertum pro quovis triangulo liquet.

§. 26.

In quovis sphaerico triangulo sunt sinus laterum, uti sinus angulorum iisdem oppositorum. Fig. 11.

Nam sit $abc = R$, et ced perpendiculare ad sphæræ radium oa ; erit $ced \perp aob$, et (cum etiam $boc \perp boa$ sit) $cd \perp ob$. In triangulis ceo, cdo vero est (per §. 25.)

$$\begin{aligned} \odot ec : \odot oc : \odot dc &= \sin. coe : 1 : \sin. cod \\ &= \sin. ac : 1 : \sin. bc; \end{aligned}$$

interim (§. 25.) etiam

itaque $\circ ec : \circ dc = \sin. cde : \sin. ced ;$
 $\sin. ac : \sin. bc = \sin. cde : \sin. ced ;$
 est vero $cde = R = cba$, atque $ced = cab$. Consequenter
 $\sin. ac : \sin. bc = 1 : \sin. a$.

E quo promanans Trigonometria sphaerica ab Axiomate XI. independenter stabilita est.

§. 27.

Fig. 12. Si ac, bd sint $\perp ab$, et feratur cab iuxta \overline{ab} ; erit (via puncti c dicta heic cd)

$$cd : ab = \sin. u : \sin. v.$$

Nam sit $de \perp ca$; est in triangulis ade, adb (per §. 25.)

$$\circ ed : \circ ad : \circ ab = \sin. u : 1 : \sin. v.$$

Revoluto $bacd$ circa ac , describetur $\circ ab$ per b , $\circ ed$ per d ; et via dictæ cd denotetur heic per $\odot cd$. Sit porro polygonum quodvis $bf g \dots$ ipsi $\odot ab$ inscriptum; nascetur per plana ex omnibus lateribus $bf, fg \&$, ad $\odot ab$ perpendicularia, in $\odot cd$ quoque figura polygonalis totidem laterum; et demonstrari (ad instar §. 23.) potest, esse

$$cd : ab = dh : bf = hf : fg = \dots,$$

adeoque

$$dh + hf + \dots : bf + fg + \dots = cd : ab$$

Quovis laterum bf, fg, \dots ad limitem o tendente, manifesto

$$bf + fg + \dots \curvearrowright \circ ab \quad \text{et} \quad dh + hf + \dots \curvearrowright \circ ed.$$

Itaque etiam

$$\circ ed : \circ ab = cd : ab.$$

Erat vero

$$\circ ed : \circ ab = \sin. u : \sin. v.$$

Consequ.

$$cd : ab = \sin. u : \sin. v.$$

Remoto ac a bd in infinitum, manet

$$\begin{aligned} & \text{adeoque etiam} \\ & \qquad \qquad \qquad \text{cd:ab} \\ & \qquad \qquad \qquad \text{sin. } u : \text{sin. } v \end{aligned}$$

constans; u vero $\sphericalangle R$ (§. 1.), et si dm || bn sit, $v \sphericalangle z$; unde fit

$$\text{cd:ab} = 1 : \text{sin. } z.$$

Via dicta cd denotabitur per cd || ab.

§. 28.

Si bn || \sphericalangle am, et c in am, atque ac = x sit: erit X (§. 23.)

Fig. 13.

$$= \text{sin. } u : \text{sin. } v.$$

Nam si cd et ae sint \perp bn et bf \perp am; erit (ad instar §. 27.)

$$\circ \text{bf} : \circ \text{cd} = \text{sin. } u : \text{sin. } v.$$

Est autem evidenter bf = ae: quamobrem

$$\circ \text{ea} : \circ \text{dc} = \text{sin. } u : \text{sin. } v.$$

In superficiebus vero Fformibus ipsorum am et cm (ipsum ambn in ab et cg secantibus) est (per §. 21.)

$$\circ \text{ea} : \circ \text{dc} = \text{ab} : \text{cg} = X.$$

Est itaque etiam

$$X = \text{sin. } u : \text{sin. } v.$$

§. 29.

Si bam = R, ab = y, et bn || am sit; erit in S

Fig. 14.

$$Y = \text{cot. } \frac{1}{2} u.$$

Nam si fuerit $ab=ac$, et $cp \parallel am$ (adeoque $bn \parallel \simeq cp$), atque $pcd=qcd$; datur (§. 19.) $ds \perp cd$, ut $ds \parallel cp$, adeoque (§. 1.) $dt \parallel cq$ sit. Si porro $be \perp ds$; erit (§. 7.) $ds \parallel bn$, adeoque (§. 6.) $bn \parallel es$, et (cum $dt \parallel cq$ sit) $bq \parallel et$; consequ. (§. 1.) $ebn = ebq$.

Repræsententur, bcf ex L ipsius bn , et fg , dh , cf et el ex L formibus lineis ipsorum ft , dt , cq et et ; erit evidenter (§. 22.)

$$\text{itaque} \quad hg = df = dh = hc;$$

$$\text{Pariter patet} \quad cq = 2ch = 2v.$$

$$\text{esse. Est vero} \quad bg = 2bl = 2z$$

$$\text{quapropter} \quad bc = bg - cq:$$

$$\text{adeoque (§. 24.)} \quad y = z - v,$$

$$\text{Est demum (§. 28.)} \quad Y = Z : I'.$$

$$\text{consequ.} \quad Z = 1 : \sin. \frac{1}{2} u \quad \text{et} \quad I' = 1 : \sin. \left(R - \frac{1}{2} u \right),$$

$$Y = \cot. \frac{1}{2} u.$$

§. 30.

Fig. 15. Verumtamen facile (ex §. 25.) patet, resolutionem problematis *Trigonometriae planae* in S , peripheriæ per radium expressæ indigere; hoc vero rectificatione ipsius L obtineri potest.

Sint ab , cm , $c'm' \perp ac$, atque b ubivis in ab ; erit (§. 25.)

$$\text{et} \quad \sin. u : \sin. v = \circ p : \circ y$$

$$\text{adeoque} \quad \sin. u' : \sin. v' = \circ p : \circ y';$$

$$\frac{\sin. u}{\sin. v} \circ = \frac{\sin. u'}{\sin. v'} \circ y'.$$

Est vero per §. 27.

$$\sin. v : \sin. v' = \cos. u : \cos. u';$$

consequ.

$$\frac{\sin. u}{\cos. u} \circ y = \frac{\sin. u'}{\cos. u'} \circ y',$$

seu

$$\circ y : \circ y' = \text{tang. } u' : \text{tang. } u = \text{tang. } w : \text{tang. } w'.$$

Sint porro $cn \parallel ab$, $c'n' \parallel ab$ et cd , $c'd'$ lineæ L formes ad \overline{ab} perpendiculares; erit (§. 21.) etiam

$$\circ y : \circ y' = r : r',$$

adeoque

$$r : r' = \text{tang. } w : \text{tang. } w'.$$

Crescat iam p ab a incipiendo in infinitum; tum $w \rightsquigarrow z$ et $w' \rightsquigarrow z'$; quapropter etiam

$$r : r' = \text{tang. } z : \text{tang. } z'.$$

Constans $r : \text{tang. } z$ (ab r *independens*) dicatur i ; dum $y \rightsquigarrow 0$, est

$$\left(\frac{r}{y} = \frac{i \text{ tang. } z}{y} \right) \rightsquigarrow 1,$$

adeoque

$$\frac{y}{\text{tang. } z} \rightsquigarrow i.$$

Ex §. 29. fit

$$\text{tang. } z = \frac{1}{2} (Y - Y^{-1});$$

itaque

$$\frac{2y}{Y - Y^{-1}} \rightsquigarrow i,$$

seu (§. 24.)

$$\frac{2y I^{\frac{y}{i}}}{I^{\frac{2y}{i}} - 1} \rightsquigarrow i.$$

Notum autem est, expressionis istius (dum $y \rightsquigarrow 0$) limitem esse $\frac{i}{\log. \text{nat. } I}$; est ergo

$$\frac{i}{\log. \text{nat. } I} = i \quad \text{et} \quad I = e = 2.7182818 \dots,$$

quæ quantitas insignis hic quoque elucet. Si nempe abhinc i illam rectam denotet, cuius $I=e$ sit, erit $r=i \text{ tang. } z$. Erat autem (§. 21.) $\circ y = 2\pi r$; est igitur

$$\begin{aligned} \circ y &= 2\pi i \text{ tang. } z = \pi i (Y - Y^{-1}) = \\ &= \pi i (e^{\frac{y}{i}} - e^{-\frac{y}{i}}) = \frac{\pi y}{\log. \text{ nat. } Y} (Y - Y^{-1}) \end{aligned}$$

(per §. 24.).

§. 31.

Fig. 16. Ad resolutionem omnium triangulorum rectangulorum rectilineorum trigonometricam (e qua omnium triangulorum resolutio in promptu est) in S 3 æquationes sufficiunt: nempe (a, b cathetos, c hypotenusam, et α, β angulos cathetis oppositos denotantibus) æquatio relationem exprimens *primo* inter a, c, α , *secundo* inter a, α, β , *tertio* inter a, b, c ; nimirum ex his *reliquae* 3 per eliminationem prodeunt.

I. Ex §§. 25. et 30. est

$$1 : \sin. \alpha = (C - C^{-1}) : (A - A^{-1}) = (e^{\frac{c}{i}} - e^{-\frac{c}{i}}) : (e^{\frac{a}{i}} - e^{-\frac{a}{i}});$$

(æquatio pro α, c, a).

II. Ex §. 27. sequitur (si $\beta m \parallel \gamma n$ sit)

$$\cos. \alpha : \sin. \beta = 1 : \sin. u;$$

ex §. 29. autem fit

$$1 : \sin. u = \frac{1}{2} (A + A^{-1});$$

itaque

$$\cos. \alpha : \sin. \beta = \frac{1}{2} (A + A^{-1}) = \frac{1}{2} (e^{\frac{a}{i}} + e^{-\frac{a}{i}});$$

(æquatio pro α, β, a).

III. Si $\alpha\alpha' \perp \beta\alpha\gamma$, atque $\beta\beta'$ et $\gamma\gamma'$ fuerint $\parallel \alpha\alpha'$, (§. 27.), atque $\beta'\alpha'\gamma' \perp \alpha\alpha'$; erit manifesto (uti in §. 27.)

$$\begin{aligned} \frac{\beta\beta'}{\gamma\gamma'} &= \frac{1}{\sin. u} = \frac{1}{2} (A + A^{-1}), \\ \frac{\gamma\gamma'}{\alpha\alpha'} &= \frac{1}{2} (B + B^{-1}), \end{aligned}$$

ac

$$\frac{\beta\beta'}{\alpha\alpha'} = \frac{1}{2}(C + C^{-1});$$

consequ.

$$\frac{1}{2}(C + C^{-1}) = \frac{1}{2}(A + A^{-1})\frac{1}{2}(B + B^{-1}),$$

sive

$$e^{\frac{c}{i}} + e^{-\frac{c}{i}} = \frac{1}{2}(e^{\frac{a}{i}} + e^{-\frac{a}{i}})(e^{\frac{b}{i}} + e^{-\frac{b}{i}});$$

(æquatio pro a, b, c).Si $\gamma\alpha\delta = R$, et $\beta\delta \perp \alpha\delta$ sit; erit

$$\circ c : \circ a = 1 : \sin. \alpha,$$

et

$$\circ c : \circ (d = \beta\delta) = 1 : \cos. \alpha,$$

adeoque ($\circ x^2$ pro quovis x factum $\circ x$ denotante) manifesto

$$\circ a^2 + \circ d^2 = \circ c^2.$$

Est vero (per §. 27. et II.)

$$\circ d = \circ b \cdot \frac{1}{2}(A + A^{-1}),$$

consequ.

$$(e^{\frac{c}{i}} - e^{-\frac{c}{i}})^2 = \frac{1}{4}(e^{\frac{a}{i}} + e^{-\frac{a}{i}})^2(e^{\frac{b}{i}} - e^{-\frac{b}{i}})^2 + (e^{\frac{a}{i}} - e^{-\frac{a}{i}})^2;$$

alia æquatio pro a, b, c (cuius membrum secundum facile ad formam symmetricam seu invariabilem reducitur).

Denique ex

$$\frac{\cos. \alpha}{\sin. \beta} = \frac{1}{2}(A + A^{-1})$$

atque

$$\frac{\cos. \beta}{\sin. \alpha} = \frac{1}{2}(B + B^{-1})$$

fit (per III.)

$$\cot. \alpha \cot. \beta = \frac{1}{2}(e^{\frac{c}{i}} + e^{-\frac{c}{i}});$$

(æquatio pro α, β, c).

§. 32.

Restat adhuc modum *problemata* in S resolvendi breviter ostendere, quo (per exempla magis obvia) peracto, demum quid theoria hæc præstet, candide dicetur.

Fig. 17. I. Sit \overline{ab} linea in plano, et $y=f(x)$ æquatio eius (pro coordinatis perpendicularibus), et quodvis incrementum ipsius z dicatur dz , atque incrementa ipsorum x , y , et areæ u , eidem dz respondentia, respective per dx , dy , du denotentur; sitque $bh \parallel cf$, et exprimatur (ex §§. 31. et 27.) $\frac{bh}{dx}$ per y , ac quæraturs ipsius $\frac{dy}{dx}$ *limes* tendente dx ad litem 0, (quod, ubi eiusmodi limes quæritur, subintelligatur): innotescet exinde etiam limes ipsius $\frac{dy}{bh}$, adeoque tg. hbg ; eritque, (cum hbc manifesto nec $>$ nec $<$ adeoque $= R$ sit), *tangens* in b ipsius bg per y determinata.

II. Demonstrari potest, esse

$$\frac{dz^2}{dy^2 + bh^2} \sim I.$$

Hinc *limes* ipsius $\frac{dz}{dx}$, et inde z integratione (per x expressum) reperitur.

Et potest lineæ cuiusvis *in concreto datae* æquatio in S inveniri, ex. gr. ipsius L .

Si enim am axis ipsius L sit; tum quævis cb ex am secat L (cum per §. 19 quævis recta ex a præter am ipsum L secet); est vero (si bn axis sit)

$$X = 1 : \sin. cbn \quad (\S. 28.),$$

atque

$$Y = \cot. \frac{1}{2} cbn \quad (\S. 29.),$$

unde fit

$$Y = X + \sqrt{X^2 - 1}$$

seu

$$e^{\frac{y}{x}} = e^{\frac{x}{x}} + \sqrt{e^{\frac{2x}{x}} - 1}$$

æquatio quæsitæ. Erit hinc

$$\frac{dy}{dx} \sim X(X^2-1)^{-\frac{1}{2}};$$

atqui

$$\frac{bh}{dx} = 1: \sin. cbn = X;$$

adeoque

$$\frac{dy}{bh} \sim (X^2-1)^{-\frac{1}{2}};$$

$$1 + \frac{dy^2}{bh^2} \sim X^2(X^2-1)^{-1},$$

$$\frac{dz^2}{bh^2} \sim X^2(X^2-1)^{-1},$$

$$\frac{dz}{bh} \sim X(X^2-1)^{-\frac{1}{2}}$$

atque

$$\frac{dz}{dx} \sim X^2(X^2-1)^{-\frac{1}{2}};$$

unde per integrationem invenitur

$$z = i(X^2-1)^{\frac{1}{2}} = i \cot. cbn$$

(uti §. 30.).

III. Manifesto

$$\frac{du}{dx} \sim \frac{hfcbh}{dx},$$

quod (nonnisi ab y dependens) iam primum per y exprimendum est; unde u integrando prodit.

Si $ab = p$, $ac = q$, et $cd = r$, atque $cabdc = s$ sit; poterit (uti in II.) Fig. 12. ostendi esse

$$\frac{ds}{dq} \sim r,$$

quod

$$= \frac{1}{2} p \left(e^{\frac{q}{i}} + e^{-\frac{q}{i}} \right),$$

atque integrando

$$s = \frac{1}{2} pi \left(e^{\frac{q}{i}} - e^{-\frac{q}{i}} \right).$$

Præter unam æquationem quodammodo definitam.

Integrandum u & modus ex §. 30. integrandi ex §. 30. III. sectione
 in §. 30. per præcedentem præcedentem quæ quodammodo in linea clausa
 est III.

Præter unam æquationem z ac figuram quædam z in distantia z i. lineæ
 esse ac z in rectæ perpendicularis verticem linearis hinc inde
 est IV.

$$\frac{1}{2} e^{\frac{x}{2}} - e^{-\frac{x}{2}} \text{ etc.}$$

Præter unam æquationem solvenda per modo tractatum facile patet lineas
 integrandi esse quædam sine et differentiale ipsum in omni per inte-
 grationem determinatur & ex æquatione solvitur z & z ac complexi
 lineæ in rectam ac z perpendicularium sine ipsorum z & connecten-
 tum quædam quædam esse. Præter unam æquationem istud tam per inte-
 grationem quædam esse est

$$= \frac{1}{2} p i e^{\frac{x}{2}} - e^{-\frac{x}{2}} + \frac{1}{2} p i.$$

Curvature quædam corporum in S determinari possunt nec non cur-
 vaturæ, evoluntæ, evolventesque linearum qualiumvis S . Quod curva-
 turam æquationem in S aut ipsius L est, aut per radium circuli, aut
 distantiam curvæ ad rectam la ab hac recta, determinatur: cum e
 præcedentibus facile ostendi possit præter L , lineas circulares, ac rectæ
 lineas in plano alias lineas uniformes dari.

IV. Pro circulo est uti in III.

$$\frac{d \odot x}{dx} = \odot x,$$

unde per §. 30. integrando fit

$$\odot x = u^{1/2} e^{\frac{x}{2}} - 2 + e^{-\frac{x}{2}}.$$

Fig. 9. V. Pro area cabdc = u linea L formi $ab=r$, huic la $cd=y$, ac
 rectis $ac, bd = x$ clausa est

atque (§. 24.)

$$\frac{du}{dx} \sim y,$$

adeoque (integrando)

$$y = re^{-\frac{x}{i}};$$

$$u = ri(1 - e^{-\frac{x}{i}}).$$

Crescente x in infinitum, fiet in S $e^{-\frac{x}{i}} \sim 0$, adeoque $u \sim ri$. Per *quantitatem* ipsius u in posterum limes iste intelligetur.

Simili modo invenitur, quod si p sit figura in F ; spatium a p et complexu axium e terminis ipsius p ductorum clausum $= \frac{1}{2} pi$ sit.

VI. Si angulus ad centrum segmenti z sphæræ sit $2u$, peripheria Fig. 10. circuli maximi sit p , et arcus fc (anguli u) $= x$; erit (§. 25.)

et hinc

$$1 : \sin. u = p : \circ bc,$$

Interim est

$$\circ bc = p \sin. u.$$

$$x = \frac{pu}{2\pi}, \quad \text{ac} \quad dx = \frac{pdu}{2\pi}.$$

Est porro

$$\frac{dz}{dx} \sim \circ bc,$$

et hinc

$$\frac{dz}{du} \sim \frac{p^2}{2\pi} \sin. u,$$

unde (integrando)

$$z = \frac{\sin. \text{vers. } u}{2\pi} p^2.$$

Cogitetur F in quod p (per meditullium f segmenti transiens) cadit; planis \overline{fem} , \overline{cem} per af , ac ad F perpendiculariter positus, ipsumque in \overline{feg} , ce secantibus; et considerentur L formis cd (ex c ad \overline{feg} perpendicularis) nec non L formis cf ; erit (§. 20.)

et (§. 21.)

$$cef = u,$$

$$\frac{fd}{p} = \frac{\sin. \text{vers. } u}{2\pi},$$

adeoque

$$z = fd \cdot p.$$

Ast (§. 21.)

$$p = \pi \cdot fdg,$$

itaque

$$z = \pi \cdot fd \cdot fdg.$$

Est autem (§. 21.)

$$fd \cdot fdg = fc \cdot fc;$$

consequ.

$$z = \pi \cdot fc \cdot fc = \odot fc \text{ in } F.$$

Fig. 14. Sit iam $bj = cj = r$; erit (§. 30.)

$$2r = i(Y - Y^{-1}),$$

adeoque (§. 21.)

$$\odot 2r \text{ (in } F) = \pi i^2 (Y - Y^{-1})^2.$$

Est quoque (IV.)

$$\odot 2y = \pi i^2 (Y^2 - 2 + Y^{-2});$$

igitur $\odot 2r \text{ (in } F) = \odot 2y$, adeoque *et superficies z segmenti sphaerici aequatur circulo, chorda fc tanquam radio descripto.*

Hinc tota sphaeræ superficies

$$= \odot fg = fdg \cdot p = \frac{p^2}{\pi},$$

suntque superficies sphaerarum, uti secundae potentiae peripheriarum earundem maximarum.

VII. Soliditas sphaeræ radii x in S reperitur simili modo

$$= \frac{1}{2} \pi i^2 (X^2 - X^{-2}) - 2\pi i^2 x;$$

Fig. 12. superficies per revolutionem lineæ $c\delta$ circa ab orta

$$= \frac{1}{2} \pi i^2 p (Q^2 - Q^{-2}),$$

et corpus per $cabdc$ descriptum

$$= \frac{1}{4} \pi i^2 p (Q - Q^{-1})^2.$$

Quomodo vero omnia a (IV.) hucusque tractata etiam absque integratione perfici possint, brevitatis studio supprimitur.

Demonstrari potest, *omnis expressionis literam i continentis* (adeoque *hypothesi*, quod *detur i*, innixæ) *limitem, crescente i in infinitum, exprimere quantitatem plane pro Σ* (adeoque pro *hypothesi nullius i*), *siquidem non eveniant aequationes identicae.* Cave vero intelligas putari, *systema ipsum variari posse* (quod omnino *in se et per se determinatum* est) sed tantum *hypothesein*, quod *successive* fieri potest, donec non ad absurdum perducti fuerimus. *Posito* igitur, quod in *tali* expressione *litera i* pro casu, si *S* esset reipsa, *illam* quantitatem unicam designet, cuius $I=e$ sit; si vero *revera Σ* fuerit, *limes dictus* loco expressionis accipi *cogitetur*: manifesto *omnes* expressiones ex *hypothesi realitatis* ipsius *S* oriundæ (hoc sensu) *absolute valent*, etsi *prorsus ignotum sit, num Σ sit, aut non sit.*

Ita e. g. ex expressione in §. 30. obtenta facile (et quidem *tam* differentiationis auxilio, quam *absque* eo) valor notus pro Σ prodit

$$\bigcirc x = 2\pi x;$$

ex I. (§. 31.) rite tractato, sequitur

$$I : \sin. \alpha = c : a;$$

ex II. vero

$$\frac{\cos. \alpha}{\sin. \beta} = I, \text{ adeoque } \alpha + \beta = R;$$

æquatio prima in III. fit *identica*, adeoque *valet* pro Σ ; quamvis nihil in eo *determinet*; ex *secunda* autem fluit

$$c^2 = a^2 + b^2.$$

Aequationes notae fundamentales trigonometriae planae in Σ .

Porro *inveniuntur* (ex §. 32.) pro Σ *area et corpus* in III., utrumque

$$= pq;$$

ex IV.

$$\odot x = \pi x^2;$$

ex VII. sphæra radii x

$$= \frac{4}{3} \pi x^3$$

ε.

Sunt quoque theoremata ad finem VI. enuntiata manifesto *inconditionate vera*.

§. 33.

Superest adhuc, quid theoria ista sibi velit, in §. 32. promissum exponere.

I. Num Σ aut S aliquod *reipsa* sit, indecisum manet.

II. Omnia ex hypothesis *falsitatis* Ax. XI. deducta semper *sensu* §. 32. intelligendo *absolute* valent, adeoque *hoc sensu nulli hypothese innituntur*. Habetur idcirco *trigonometria plana a priori*, in qua *solum systema verum ignotum* adeoque *solummodo absolute* magnitudines expressionum incognitæ manent, per *unicum* vero casum notum, manifesto totum systema figeretur. Trigonometria sphærica autem in §. 26. *absolute* stabilitur. Habeturque Geometria, Geometriæ planæ in Σ *prorsus* analogæ in F .

III. Si *constaret* Σ esse, nihil hoc respectu amplius incognitum esset; si vero *constaret non esse* Σ , tunc (§. 31.) (e. g.) e lateribus x , y et angulo rectilineo ab iis intercepto, in *concreto datis* manifesto in se et per se impossibile esset triangulum *absolute* resolvere i. e. a priori determinare angulos ceteros et *rationem lateris tertii* ad duo data; nisi X , Y determinentur, ad quod *in concreto* haberi aliquod *a* oporteret, cuius A notum esset; atque tum *i unitas naturalis longitudinum* esset, sicuti e est basis logarithmorum naturalium. Si existentia huius i constiterit; quomodo ad usum saltem quam exactissime construi possit, ostendetur.

IV. Sensu in I. et II. exposito patet, omnia in spatio methodo recentiorum Analytica intra iustos fines valde laudanda absolvi posse.

V. Denique lectoribus benevolis haud ingratum futurum est; pro casu illo, quodsi non Σ sed S *reipsa* esset, circulo æquale rectilineum construi.

§. 34.

Ex δ ducitur dm'' an modo sequente.

Fig. 12.

Fiat ex δ

$$\delta b \perp an;$$

erigatur e puncto quovis aliquo a rectæ \overline{ab}

$$ac \perp an \text{ (in } \delta ba),$$

et demittatur

$$\delta e \perp ac;$$

erit (§. 27.)

$$\odot ed : \odot ab = 1 : \sin. z,$$

siquidem fuerit $dm'' \parallel bn$.

Est vero $\sin. z$ non > 1 , adeoque ab non $> \delta e$. Descriptus igitur quadrans radio ipsi δe æquali ex a in bac , gaudebit puncto aliquo b vel o cum $b\delta$ communi. Priore in casu manifesto $z = R$; in posteriore vero erit (§. 25.)

$$(\odot ao = \odot ed) : \odot ab = 1 : \sin. aob,$$

adeoque

$$z = aob.$$

Si itaque fiat $z = aob$, erit $dm'' \parallel bn$.

§. 35.

Si fuerit S reipsa; ducetur recta ad anguli acuti crus unum perpendicularis, quae ad alterum \parallel sit, hoc modo. Fig. 18.

Sit $am \perp bc$, et accipiatur $ab = ac$ tam parvum (per §. 19.), ut si ducatur $bn'' \parallel am$ (§. 34.), sit $abn >$ angulo dato. Ducatur porro $cp \parallel am$ (§. 34.), fiantque nbq , $pc\delta$ utrumque æquale angulo dato; et $b\tilde{q}$, $c\tilde{\delta}$ se mutuo secabunt. Secet enim $b\tilde{q}$, (quod *per constr.* in nbc cadit) ipsam $c\tilde{\delta}$ in e ; erit (propter $bn'' \parallel cp$) $ebc < ec\delta$, adeoque $ec < eb$. Sint

$$ef = ec, efr = ec\delta, \text{ et } fs \parallel ep;$$

cadet fs in bfr . Nam cum $bn'' \parallel cp$, adeoque $bn'' \parallel ep$, atque $bn'' \parallel fs$ sit;

erit (§. 14.)

$$fbn + bfs < 2R = fbn + bfr;$$

itaque $bfs < bfr$. Quamobrem $f\bar{r}$ secat $e\bar{p}$, adeoque $c\bar{d}$ quoque ipsam $e\bar{q}$ in puncto aliquo d .

Sit iam $dg = dc$, atque $dgt = dcp = gbn$; erit (cum $cd \simeq gd$ sit)

$$bn \simeq gt \simeq cp.$$

Si fuerit lineæ L formis ipsius bn , punctum in $b\bar{q}$ cadens f (§. 19.), et axis fl ; erit

$$bn \simeq fl,$$

adeoque

$$bfl = bgt = dcp;$$

sed etiam

$$fl \simeq cp:$$

cadit ergo f manifesto in g , estque $gt \parallel bn$. Si vero ho ipsum bg perpendiculariter bisecet; erit $ho \perp bn$ constructum.

§. 36.

Fig. 10. Si fuerint data recta $c\bar{p}$ et planum \overline{mab} , atque fiat $cb \perp \overline{mab}$, (in $\overline{bc\bar{p}}$) $bn \perp bc$, et $cq \parallel bn$ (§. 34.); *sectio ipsius $c\bar{p}$* (si hæc in bcq cadat) *cum $b\bar{n}$* (in \overline{cbn}), adeoque *cum \overline{mab}* reperitur. Et si fuerint data duo plana \overline{pcq} , \overline{mab} , et sit $cb \perp \overline{mab}$, $cr \perp \overline{pcq}$, atque (in \overline{bcr}) $bn \perp bc$, $cs \perp cr$; cadent bn in \overline{mab} , et cs in \overline{pcq} ; et sectione ipsarum \overline{bn} , \overline{cs} (si detur) reperiata, erit perpendicularis in pcq per eandem ad $c\bar{s}$ ducta manifesto *sectio ipsorum \overline{mab} , \overline{pcq}* .

§. 37.

Fig. 7. In \overline{am} bn reperitur *tale a , ut sit $am \simeq bn$* ; si (per §. 34.) construatur extra \overline{nbm} $gt \perp bn$, et fiant $bg \perp gt$, $gc = gb$, atque $cp \parallel gt$; ponaturque $tg\delta$ ita, ut efficiat cum $tg\bar{b}$ angulum illi æqualem, quem $pc\bar{a}$ cum $pc\bar{b}$ facit; atque quærat (per §. 36.) *sectio $\overline{d\bar{q}}$ ipsorum $tg\delta$, $nb\bar{a}$* ; fiatque

$ba \perp dq$. Erit enimvero ob triangulorum L lineorum in F ipsius bn exortorum similitudinem (§. 21.) manifesto $\delta b = \delta a$, et $am \simeq bn$.

Facile hinc patet (L lineis per *solos terminos* datis) reperiri posse etiam *terminos* proportionis quartum ac medium, atque omnes constructiones geometricas, quæ in Σ' in plano fiunt, hoc modo in F *absque XI. Axiomate* perfici posse. Ita e. g. $4R$ in quotvis partes æquales geometricè dividi potest, si sectionem istam in Σ perficere licet.

§. 38.

Si construatur (per §. 37.) e. g. $nbq = \frac{1}{3} R$, et fiat (per §. 35.) in S ad $b\bar{q}$ perpendicularis $am \parallel bn$, atque determinetur (per §. 37.) $jm \simeq bn$; erit, si $ja = x$ sit, (§. 28.)

$$X = 1 : \sin. \frac{1}{3} R = 2,$$

atque x *geometricè* constructum.

Et potest nbq ita computari, ut ja ab i quovis dato minus discrepet, cum nonnisi $\sin. nbq = \frac{1}{e}$ esse debeat.

§. 39.

Si fuerint (in plano) pq et st , \parallel rectæ mn (§. 27.), et ab , cd sint perpendicularæ ad mn æquales; manifesto est

$$\triangle dec \simeq \triangle bea,$$

adeoque anguli (forsan mixtilinei) ecp , eat congruent, atque

$$ec = ea.$$

Si porro $cf = ag$, erit

$$\triangle acf \simeq \triangle cag,$$

et utrumque *quadrilateri* $facg$ dimidium est. Si $facg$, $hagf$ duo eiusmodi quadrilatera fuerint ad ag , inter pq et st ; æqualitas eorum (uti apud EUCLIDEM), nec non triangulorum agc , agh eidem ag insistentium,

verticesque in \overline{pq} habentium, æqualitas patet. Est porro

$$\begin{aligned} \text{atque} \quad & \text{acf} = \text{cag}, \quad \text{gcq} = \text{cga}, \\ & \text{acf} + \text{acg} + \text{gcq} = 2R \\ (\S. 32.), \text{ adeoque etiam} \quad & \text{cag} + \text{acg} + \text{cga} = 2R; \end{aligned}$$

itaque in quovis eiusmodi triangulo acg summa trium angulorum $= 2R$.

Sive in ag ($\text{quæ} \parallel \text{mn}$) ceciderit autem *recta* ag , sive non; triangulorum *rectilineorum* agc , agh *tam ipsorum, quam summarum angulorum ipsorumdem, æqualitas* in aperto est.

§. 40.

Fig. 20. *Aequalia triangula abc , abd (abhinc rectilinea) uno latere aequali gaudentia, summas angulorum aequales habent.*

Nam dividat mn bifariam tam ac quam bc , et sit pq ($\text{per } c \parallel \text{mn}$); cadet δ in \overline{pq} . Nam si \overline{bd} ipsum \overline{mn} in puncto e , adeoque (§. 39.) ipsum \overline{pq} ad distantiam $\text{ef} = \text{eb}$ secet; erit

$$\triangle \text{abc} = \triangle \text{abf},$$

adeoque et

$$\triangle \text{abd} = \triangle \text{abf},$$

unde δ in f cadit: si vero \overline{bd} ipsum \overline{mn} non secuerit, sit c punctum, ubi perpendicularis rectam ab bisecans ipsum \overline{pq} secat, atque $\text{gs} = \text{ht}$ ita, ut \overline{st} *productam* \overline{bd} in puncto aliquo f secet (quod fieri posse modo simili patet, ut §. 4.); sint porro $\text{sl} = \text{sa}$, $\text{lo} \parallel \text{st}$, atque o sectio ipsorum \overline{bf} et \overline{lo} ; esset tum (§. 39.)

$$\triangle \text{abl} = \triangle \text{abo},$$

adeoque

$$\triangle \text{abc} > \triangle \text{abd}$$

(contra hyp.).

§. 41.

Aequalia triangula abc, def aequalibus angulorum summis gaudent. Fig. 21.

Nam secet mn tam ac quam bc, ita pq tam df quam fe bifariam, et sit rs || mn, atque to || pq; erit perpendicularis ag ad rs aut æqualis perpendiculari dh ad to, aut altera e. g. dh erit maior: in quovis casu o df e centro a cum gñ punctum aliquod f commune habet, eritque (§. 39.)

$$\triangle abf = \triangle abc = \triangle def.$$

Est vero $\triangle abf$ (per §. 40.) triangulo dfe, ac (per §. 39.) triangulo abc æquiangulum. Sunt igitur etiam triangula abc, def æquiangula.

In *S* converti quoque theorema potest. Sint enim triangula abc, def reciproce æquiangula, atque $\triangle bal = \triangle def$; erit (per præc.) alterum alteri, adeoque etiam $\triangle abc$ triangulo abl æquiangulum, et hinc manifesto

$$bcl + bfc + cbl = 2R.$$

Atqui (ex §. 31.) cuiusvis trianguli angulorum summa in *S* est $< 2R$: cadit igitur *I* in *c*.

§. 42.

Si fuerit complementum summae angulorum trianguli abc ad 2R Fig. 22.

trianguli def vero
est

$$\triangle abc : \triangle def = u : v.$$

Nam si quodvis triangulorum acg, gch, hcb, dft, ffe sit = *p*, atque

$$\triangle abc = mp, \quad \triangle def = np;$$

sitque *s* summa angulorum cuiusvis trianguli, quod = *p* est: erit manifesto

$$2R - u = ms - (m - 1)2R = 2R - m(2R - s),$$

et

$$u = m(2R - s),$$

et pariter

$$v = n(2R - s).$$

Est igitur

$$\triangle abc : \triangle def = m : n = u : v.$$

Ad casum incommensurabilitatis triangulorum abc , def quoque extendi facile patet.

Eodem modo demonstratur triangula in superficie sphærica esse uti *excessus* summarum angulorum eorundem supra $2R$. Si duo anguli trianguli sphærici recti fuerint, tertius z erit excessus dictus; est autem triangulum istud (peripheria maxima p dicta) manifesto

$$= \frac{z}{2\pi} \cdot \frac{p^2}{2\pi} \quad (\S. 32. VI.);$$

consequenter quodvis triangulum, cuius angulorum excessus $= z$, est

$$= \frac{zp^2}{4\pi^2}.$$

§. 43.

Fig. 15. Iam *area* trianguli rectilinei in S per summam angulorum exprimetur.

Si ab crescat in infinitum; erit (§. 42.)

$$\triangle abc : (R - u - v)$$

constans. Est vero

$$\triangle abc \sim bacn \quad (\S. 32. V.)$$

et

$$R - u - v \sim z \quad (\S. I.);$$

adeoque

$$bacn : z = \triangle abc : (R - u - v) = bac'n' : z'.$$

Est porro manifesto

$$bdcn : bd'c'n' = r : r' = \text{tang. } z : \text{tang. } z' \quad (\S. 30.).$$

Pro $y' \sim 0$ autem est

$$\frac{bd'c'n'}{bac'n'} \sim 1,$$

nec non

$$\frac{\text{tang. } z'}{z'} \sim 1;$$

consequ.

$$bdcn : bacn = \text{tang. } z : z.$$

Erat vero (§. 32.)

$$bdcn = ri = i^2 \text{ tang. } z;$$

est igitur

$$bacn = zi^2.$$

Quovis triangulo, cuius angulorum summæ complementum ad $2R$ z est, in posterum breviter Δ dicto, erit idcirco

$$\Delta = zi^2.$$

Facile hinc liquet, quod si

$$or \parallel am \quad \text{et} \quad ro \parallel ab$$

Fig. 14.

fuerint; *area* inter \overline{or} , \overline{st} , \overline{bc} comprehensa (quæ manifesto limes absolutus est *area* triangulorum rectilineorum sine fine crescentium, seu ipsius Δ pro $z \sim 2R$), sit

$$= \pi i^2 = \odot i \text{ in } F.$$

Limite isto per \square denotato, erit porro (per §. 30.)

Fig. 15.

$$\begin{aligned} \pi r^2 &= \text{tang. } z^2 \square = \odot r \text{ in } F \text{ (§. 21.)} \\ &= \odot s \text{ (per §. 32. VI.),} \end{aligned}$$

si chorda dc s dicatur. Si iam radio dato s , circuli in plano (sive radio L formi circuli in F) perpendiculariter bisecto, construat (per §. 34.) $db \parallel \sphericalangle cn$; demissa perpendiculari ca ad db , et erecta perpendiculari cm ad ca ; habebitur z ; unde (per §. 37.) $\text{tang. } z^2$, radio L formi ad libitum pro unitate assumto, *geometricè determinari potest per duas lineas uniformes eiusdem curvaturæ* (quæ solis terminis datis, constructis axibus, manifesto tanquam rectæ commensurari, atque hoc respectu rectis æquivalentes spectari possunt).

Fig. 23. Porro construitur quadrilaterum ex. gr. regulare = \square , ut sequitur. Sit

$$abc = R, \quad bac = \frac{1}{2} R, \quad acb = \frac{1}{4} R, \quad \text{et } bc = x;$$

poterit X (ex §. 31. II.) per meras radices quadraticas exprimi, et (per §. 37.) construi: habitoque X , (per §. 38., sive etiam 29. et 35.) x ipsum determinari potest. Estque octuplum $\triangle abc$ manifesto = \square , atque *per hoc, circulus planus radii s , per figuram rectilineam, et lineas univoformes eiusdem generis (rectis, quoad comparationem inter se, aequivalentes) geometricè quadratus; circulus F formis vero eodem modo complanatus: habeturque aut Axioma XI. Euclidis verum, aut quadratura circuli geometrica; etsi hucusque indecisum manserit, quodnam ex his duobus revera locum habeat. Quoties $\text{tang. } z^2$ vel numerus integer vel fractio rationalis fuerit, cuius (ad simplicissimam formam reductæ) denominator aut numerus primus formæ $2^m + 1$ (cuius est etiam $2 = 2^0 + 1$) aut productum fuerit e quotcunque primis huius formæ, quorum (ipsum 2, qui solus quotvis vicibus occurrere potest, excipiendo) quivis *semel* ut factor occurrit: per theoriam polygonorum ill. GAUSS (præclarum nostri imo omnis ævi inventum), etiam ipsi $\text{tang. } z^2 \square = \odot s$ (et nonnisi pro talibus valoribus ipsius z) figuram rectilineam æqualem constituere licet. Nam *divisio* ipsius \square (theoremate §. 42. facile ad quælibet polygona extenso) manifesto *sectionem* ipsius $2R$ requirit, quam (ut ostendi potest) unice sub dicta conditione geometricè perficere licet. In omnibus autem talibus casibus præcedentia facile ad scopum perducent. Et potest quævis figura rectilinea in polygonum regulare n laterum geometricè converti, siquidem n sub formam GAUSSianam cadat.*

Superesset denique, (ut res omni numero absolvatur), impossibilitatem (absque suppositione aliqua) decidendi, num Σ' aut aliquod (et quodnam) S sit, demonstrare: quod tamen occasione magis idoneæ reservatur.

WOLFGANGI BOLYAI
ADDITAMENTUM AD APPENDICEM.

Denique *aliquid Auctori Appendicis proprium, coronidis instar*, addere fas sit: qui tamen ignoscat, si quid non acu eius tetigerim.

Res breviter in eo consistit: *formulae trigonometriae sphaericae*, in Appendice dicta ab axioma XI. Eucl. independenter demonstratæ, *cum formulis trigonometriae planae conveniunt, si* (modo statim dicendo) *latera trianguli sphaerici realia, rectilinei vero imaginaria accipiantur*; adeo ut quoad formulas trigonometricas planum ut sphaera imaginaria considerari possit, si pro reali illa accipiatur, in qua $\sin. R=1$.

Pro casu, si axioma Eucl. verum non fuerit, demonstratur (Appendix §. 30.) dari certum i , pro quo ibidem dictum I est $= e$ (basi logarithmorum naturalium), atque pro hoc casu formulæ trigonometriæ planæ quoque demonstrantur (ibidem §. 31.); et quidem ita, ut (iuxta §. 32., post VII., ibidem) formulæ et pro casu veritatis axiomatis dicti valeant; nempe si supponendo, quod $i \sim \infty$, limites valorum accipiantur; nimirum systema Euclideum est quasi limes systematis antieuclidei (pro $i \sim \infty$). Ponatur, pro casu existentis i , unitas $= i$, atque conceptus *sinus cosinusque* extendatur et ad arcus imaginarios; ita ut arcum sive realem sive imaginarium denotet p , dicatur

$$\frac{1}{2}(e^{p\sqrt{-1}} + e^{-p\sqrt{-1}})$$

cosinus ipsius p , et

$$\frac{1}{2\sqrt{-1}}(e^{p\sqrt{-1}} - e^{-p\sqrt{-1}})$$

dicatur sinus ipsius p .

Erit hinc pro q reali

$$\begin{aligned} \frac{1}{2\sqrt{-1}}(e^q - e^{-q}) &= \frac{1}{2\sqrt{-1}}(e^{-q\sqrt{-1}\sqrt{-1}} - e^{q\sqrt{-1}\sqrt{-1}}) = \\ &= \sin.(-q\sqrt{-1}) = -\sin. q\sqrt{-1} \end{aligned}$$

Ita

$$\begin{aligned} \frac{1}{2}(e^q + e^{-q}) &= \frac{1}{2}(e^{-q\sqrt{-1}\sqrt{-1}} + e^{q\sqrt{-1}\sqrt{-1}}) = \\ &= \cos.(-q\sqrt{-1}) = \cos. q\sqrt{-1}; \end{aligned}$$

si nempe et in circulo imaginario sinus negativi arcus sinui arcus positivi alioquin priori æqualis sit, præterquam quod negativus sit, atque cosinus arcus positivi et negativi (si alioquin æquales fuerint), sit idem.

In Appendice dicta §. 25. demonstratur absolute, id est ab axioma dicto independenter; quod in quovis triangulo rectilineo *sinus angulorum sint, uti peripheriæ radiorum lateribus oppositis æqualium*; demonstraturque porro, pro casu existentis i , peripheriam radii y esse

$$= \pi i (e^{\frac{y}{i}} - e^{-\frac{y}{i}}),$$

quod pro $i=1$ fit

$$\pi (e^y - e^{-y}).$$

Itaque (§. 31. ibidem) pro triangulo rectilineo rectangulo, cuius catheti sunt a et b , hypotenusam c , et anguli lateribus a , b , c oppositi sunt α , β , π ; est (pro $i=1$)

in I.

$$1 : \sin. \alpha = \pi (e^c - e^{-c}) : \pi (e^a - e^{-a});$$

adeoque

$$1 : \sin. \alpha = \frac{1}{2\sqrt{-1}}(e^c - e^{-c}) : \frac{1}{2\sqrt{-1}}(e^a - e^{-a}).$$

Unde

$$1 : \sin. \alpha = -\sin. c\sqrt{-1} : -\sin. a\sqrt{-1}.$$

Et hinc

$$1 : \sin. \alpha = \sin. c\sqrt{-1} : \sin. a\sqrt{-1}.$$

In II. fit

$$\cos. \alpha : \sin. \beta = \cos. a \sqrt{-1} : 1.$$

In III. fit

$$\cos. c \sqrt{-1} = \cos. a \sqrt{-1} \cos. b \sqrt{-1}.$$

Quæ prouti omnes exinde promanantes formulæ trigonometriæ planæ, cum formulis trigonometriæ sphæricæ prorsus conveniunt; nisi quod si ex. gr. trianguli sphærici rectanguli quoque catheti angulique iis oppositi, hypotenusaque nomina eadem sortiantur, latera trianguli rectilinei per $\sqrt{-1}$ dividenda sint, ut formulæ pro sphæricis prodeant.

Nempe ex I. fiet

$$1 : \sin. \alpha = \sin. c : \sin. a,$$

ex II. fiet

$$1 : \cos. a = \sin. \beta : \cos. \alpha,$$

ex III. fiet

$$\cos. c = \cos. a \cos. b.$$

Quum ceteris supersedere liceat, et lectorem deductione (App. §. 32. post VII.) ommissa offendi impediri que expertus sim: haud abs re erit ostendere, quomodo ex. gr. ex

$$e^{\frac{c}{i}} + e^{-\frac{c}{i}} = \frac{1}{2} (e^{\frac{a}{i}} + e^{-\frac{a}{i}}) (e^{\frac{b}{i}} + e^{-\frac{b}{i}})$$

sequatur

$$c^2 = a^2 + b^2$$

(theorema Pythagoreum pro systemate Euclideo); verosimiliter Auctor quoque ita deduxit, et ceteræ quoque eodem modo sequuntur.

Est nempe potentiis ipsius e per series expressis

$$e^{\frac{h}{i}} = 1 + \frac{h}{i} + \frac{h^2}{2i^2} + \frac{h^3}{2 \cdot 3i^3} + \frac{h^4}{2 \cdot 3 \cdot 4i^4} + \dots$$

$$e^{-\frac{h}{i}} = 1 - \frac{h}{i} + \frac{h^2}{2i^2} - \frac{h^3}{2 \cdot 3i^3} + \frac{h^4}{2 \cdot 3 \cdot 4i^4} - \dots;$$

adeoque

$$e^{\frac{h}{i}} + e^{-\frac{h}{i}} = 2 + \frac{h^2}{i^2} + \frac{h^4}{3 \cdot 4i^4} + \dots$$

$$= 2 + \frac{h^2 + u}{i^2},$$

(si omnium terminorum post $\frac{k^2}{i^2}$ summa $\frac{u}{i^2}$ dicatur); estque $u \sim 0$, dum $i \sim \infty$. Nam multiplicentur omnes termini post $\frac{k^2}{i^2}$ per i^2 ; erit terminus primus $\frac{k^4}{3 \cdot 4i^2}$, et quivis exponens $< \frac{k^2}{i^2}$; essetque etsi exponens ubique hic maneret, summa

$$\frac{k^4}{3 \cdot 4i^2} : \left(1 - \frac{k^2}{i^2}\right) = \frac{k^4}{3 \cdot 4(i^2 - k^2)},$$

quod manifesto ~ 0 , dum $i \sim \infty$.

Atque ex

$$e^{\frac{c}{i}} + e^{-\frac{c}{i}} = \frac{1}{2} \left(e^{\frac{a+b}{i}} + e^{-\frac{a+b}{i}} + e^{\frac{a-b}{i}} + e^{-\frac{a-b}{i}} \right)$$

sequitur (pro ω , v , λ adinstar u acceptis)

$$2 + \frac{c^2 + \omega}{i^2} = 1 + \frac{(a+b)^2 + v}{2i^2} + 1 + \frac{(a-b)^2 + \lambda}{2i^2}.$$

Atque hinc

$$c^2 = \frac{1}{2} (a^2 + 2ab + b^2 + a^2 - 2ab + b^2 + v + \lambda - 2\omega),$$

quod

$$\sim a^2 + b^2.$$

Scholion. Sphæræ illius, in qua sinus totus est $1=i$, radius est ordinata y lineæ L formis ipsi $i=1$ æqualis, ad axem per unam extremitatem ex altera perpendiculariter missa. Nempe *in superficie* (App. §. 21.) *F dicta, tota Geometria Euclidea valet, lineis L vicem rectarum subeuntibus*: atque pro radio L formi $=1$, qui sinus totus in F erit, peripheriæ eiusdem radius in plano erit plane dictum y ; quod ad sphæram imaginariam, ad quam planum (in systemate antieulideo) revocatur, facile applicatur.

ADNOTATIONES EDITORUM.

In Ed. I. Appendicis literæ singulares puncta denotantes literis quæ dicuntur *cursivis* impressæ sunt, sed in libellis nobis relictis manu Ioannis Bolyai scriptis puncta literis qu. d. *fractur* denotantur, quibus etiam pater eius usus est.

Pag. 1. post §. 15. in Ed. I. legitur :

└ denotet perpendiculare
∧ « angulum.

Nos hæc delevimus, quia pro signo └ usitatioe ⊥ utimur, vocabulum «angulum» autem ubique integrum scribimus.

Pag. 8. §. ultimo «frs» correximus in «hrs», ut et ipse auctor correxerit in libello manu scripto, quo Appendicem lingua Germanica denuo pertractavit.

Pag. 10. §. ultimo §. 18. «○ha» emendavimus ex «○hf».

Pag. 14. extremæ §. 26. auctor ascripsit exemplari in bibliotheca Academiæ Hungariæ asservato «cos. quoque necessarium». Videtur in editione quadam altera propositionem de cosinu demonstraturus fuisse, quamquam nihil hic maioris momenti vere desideratur. Revera considerationes, quibus in «Supplemento numeri 31351» Tom. II. Tentaminis ex

$$\sin. H : \sin. A = 1 : \sin. a$$

trigonometria sphærica integra deducitur, et in geometria absoluta valent.

Pag. 16. §. 3. Ex libello Germanico auctoris manu scripto «cq» scripsimus pro vitioso «cq» Editionis I.

Pag. 17. §. 5. «cn || ab, c'n' || ab» scripsimus pro «cn, c'n' || ab» Ed. I.

Pag. 20. §. 8. «et 27» Ioannes Bolyai ipse ascripsit in exemplari Academix.

Pag. 22. §. 4. a calce «§. 30.». scripsimus pro «§. 29.» Ed. I.

Pag. 35. In hoc Additamento, quod pagg. 380—383 Tom. II. Ed. I. Tentaminis continetur, delevimus mentionem de paginis huius operis factam, in quibus solum theoremata omnibus nota continentur. Articulum vero secundum in hunc locum reiecimus:

«Pro v positivo radicem positivam ipsius $-v^2$ per $+v$ denotat, et negativam per $-v$, sed ob defectum signorum (quum vix hæc duo quadamtenus prodierunt), radix positiva ipsius -1 per $\sqrt{-1}$ et negativa per $-\sqrt{-1}$ denotabitur».

Ibidem §. 3. Post «Appendicis» delevimus «in tomo primo».

Ibidem §. 11. Initium articuli huius in Ed. I. sic legitur:





«Nimirum de axiomate Euclideo dictum in tomo primo satis superque est: pro casu, si verum non fuerit ☞».

Pag. 38. §. 2. «multiplicentur» scripsimus pro erroneo «dividantur».

Ibidem §. 8. Latus sinistrum formulæ in Ed. I. deest.

I. BOLYAI, Appendix.

Tab. I.




Del. TÖTÖSSY B.

Lith. GRUND V. utódai.


Tab. II. Fig. 5-7.

Tab. III. Fig. 8-10. ...


11.



12.




13.




Tab. IV. Fig. 11-13.

14.



15.




Del. TÖTÖSSY B.

Lith. GRUND V. utódnai.


Tab. V. Fig. 14-15.

Tab. VI. Fig. 16-18.



19.




20.



21.




22.




Del. TÖTÖSSY B.


23.



Lith. GRUND V. utódnai.

Tab. VII. Fig. 19-23.





~~DEC 9 1932~~

~~DUE MAY 17 50~~



Math 5858.32.3
Ioannis Bolyai d
Cabot Science



3 2044

